

Epistatic interactions between *Tgfb1* and genetic loci, *Tgfbm2* and *Tgfbm3*, determine susceptibility to an asthmatic stimulus

Julia Freimuth^{a,1}, Frederic F. Clermont^{a,b,1}, Xiaozhu Huang^c, Angela DeSapio^a, Taku A. Tokuyasu^{a,d}, Dean Sheppard^c, and Rosemary J. Akhurst^{a,e,2}

^aHelen Diller Family Comprehensive Cancer Center, ^cLung Biology Center, Department of Medicine, ^dComputational Biology Core, and ^eDepartment of Anatomy and Institute of Human Genetics, University of California, San Francisco, CA 94158; and ^bVIB (Vlaams Instituut voor Biotechnologie) Laboratory for Molecular Cancer Biology, Katholieke Universiteit Leuven, 3000 Leuven, Belgium

Edited by David A. Schwartz, University of Colorado, Aurora, CO, and accepted by the Editorial Board September 10, 2012 (received for review April 6, 2012)

TGF β activation and signaling have been extensively studied in experimental models of allergen-induced asthma as potential therapeutic targets during chronic or acute phases of the disease. Outcomes of experimental manipulation of TGF β activity have been variable, in part due to use of different model systems. Using an ovalbumin (OVA)-induced mouse model of asthma, we here show that innate variation within TGF β 1 genetic modifier loci, *Tgfbm2* and *Tgfbm3*, alters disease susceptibility. Specifically, *Tgfbm2*¹²⁹ and *Tgfbm3*^{C57} synergize to reverse accentuated airway hyperresponsiveness (AHR) caused by low TGF β 1 levels in *Tgfb1*^{+/-} mice of the NIH/OlaHsd strain. Moreover, epistatic interaction between *Tgfbm2*¹²⁹ and *Tgfbm3*^{C57} uncouples the inflammatory response to ovalbumin from those of airway remodeling and airway hyperresponsiveness, illustrating independent genetic control of these responses. We conclude that differential inheritance of genetic variants of *Tgfbm* genes alters biological responses to reduced TGF β 1 signaling in an experimental asthma model. TGF β antagonists for treatment of lung diseases might therefore give diverse outcomes, dependent on genetic variation.

Asthma is an allergic disease of the airways affecting more than 5% of the US population. It is characterized by airway hyperresponsiveness (AHR), inflammatory infiltration, increased mucus production, elevated serum IgE levels, and airway remodeling (1). Asthma can present within a wide range of disease severity, from mild and intermittent to severe, persistent, and drug refractory. It is considered a multifactorial disorder in which complex interplay between environmental and genetic factors determines disease risk and severity. Here we investigated the contribution of genetic factors that have previously been shown to interact with transforming growth factor β (TGF β) in vivo, to disease severity in a mouse model of asthmatic response.

Genetic variants of the human *TGFBI* gene are associated with asthma severity (2–5) and TGF β is synthesized by, and has effects on, several cell types of the lung in response to an asthmatic stimulus. Thus, the TGF β signaling pathway has been considered a potential therapeutic target in lung disease (6). It is a potent suppressor of inflammation, illustrated by lethal T-cell-mediated multifocal inflammation in *Tgfb1*^{-/-} mice (7, 8). It also regulates epithelial cell growth and differentiation and stimulates smooth muscle and myofibroblast differentiation and extracellular matrix deposition (6). TGF β appears protective in acute models of asthmatic pathology, seen both genetically and pharmacologically (9–12). Conversely, excess active TGF β can exacerbate chronic asthma pathology by induction of fibrosis (13, 14). It can also stimulate pulmonary inflammation and accumulation and contraction of smooth muscle through induction of T_H17 cells (15) and effects on intraepithelial mast cells (16), leading to airway obstruction and decreased lung function.

We have reported genetic loci, *Tgfbm2* and *Tgfbm3*, which dramatically alter phenotypic outcome of low TGF β 1 levels in mice, with effects on both vascular development and skin tumor susceptibility (17–20). We also showed that, as for human *TGFBI*, the mouse *Tgfb1* gene is polymorphic, with allelic variants that drive different expression levels in diverse mouse species, consequently

conferring strain-specific variation in tumor susceptibility (20). Interestingly, the biological outcome of *Tgfb1* genetic variation in terms of tumor risk is dependent on interaction with an unlinked genetic locus, *Skts15* (20), illustrating the power of epistasis in masking single gene effects and determining disease risk. Significantly, *Skts15* colocalizes with *Tgfbm3* on the genome (19). This locus is thus synonymous with *Skts15* and a potent modifier of two distinct TGF β -dependent phenotypes.

In the current study, we demonstrate that different components of the asthmatic response to the allergen, ovalbumin (OVA), are dependent on mouse strain background. More specifically, we show that potentiation of AHR by loss of a single *Tgfb1* allele is dependent on synergistic interaction between variant alleles of the two TGF β 1 modifier loci, *Tgfbm2* and *Tgfbm3*. Moreover, we demonstrate uncoupling of the inflammatory vs. the AHR response to an asthmatic stimulus, mediated by these two genetic variants.

Results

***Tgfb1* Haploinsufficiency Potentiates AHR in a Mouse-Strain-Specific Manner.** Several reports have suggested that TGF β 1 is protective against allergen-induced lung pathology (9–12). We compared wild-type with *Tgfb1*^{+/-} mice on two different strains, NIH/OlaHSD (NIH) and C57BL/6NTac, and investigated their physiological and cellular responses to acute exposure to the allergen OVA after a 3-wk period of OVA sensitization. As assessed by acute AHR, C57 wild-type mice were relatively resistant to the asthmatic challenge compared with NIH wild-type mice (Fig. 1*A* and *B*). Interestingly, we showed no effect of *Tgfb1* gene dosage on the C57 genetic background, such that there were identical cellular and physiological responses to the asthmatic stimulus regardless of *Tgfb1* genotype (Fig. 1*A*). However, we found strain differences with respect to *Tgfb1* gene dosage on asthmatic response; loss of one *Tgfb1* allele exacerbated AHR in NIH (Fig. 1*B*) but not C57 mice (Fig. 1*A*). We conclude that the effect of reduced TGF β 1 levels on asthmatic pathology was dependent on mouse strain background.

***Tgfbm2*¹²⁹ and *Tgfbm3*^{C57} Synergize to Reduce AHR in NIH-*Tgfb1*^{+/-} Mice.** In two independent mouse models of pathology, namely tumor susceptibility (20) and vascular dysplasia (21), we previously demonstrated that the phenotypic outcome of *Tgfb1* gene dosage is mouse strain dependent and strongly influenced by endogenous variation within *Tgfbm2* and *Tgfbm3* (20, 21). To examine the influence of these genetic loci on pulmonary responses

Author contributions: J.F., F.F.C., D.S., and R.J.A. designed research; J.F., F.F.C., X.H., and A.D. performed research; J.F., F.F.C., T.A.T., D.S., and R.J.A. analyzed data; and J.F., F.F.C., and R.J.A. wrote the paper.

The authors declare no conflict of interest.

This article is a PNAS Direct Submission. D.A.S. is a guest editor invited by the Editorial Board.

¹J.F. and F.F.C. contributed equally to this work.

²To whom correspondence should be addressed. E-mail: rakhurst@cc.ucsf.edu.

This article contains supporting information online at www.pnas.org/lookup/suppl/doi:10.1073/pnas.1205374109/-DCSupplemental.




Fig. 3. *Tgfbm2* and *Tgfbm3* together compensate for *Tgfb1* haploinsufficiency in *Tgfbm2/3* mice. (A) Respiratory resistance measured in response to escalating doses of ACh after challenge with OVA, in *Tgfb1*^{+/-} and wild-type mice on a *Tgfbm2/3* strain background. Data shown as mean ± SEM from each group (*n* = 10). (B and C) Western blot analysis using indicated antibodies was done on protein lysates from lung tissue after saline (S) or OVA (O) challenge. Of 18 independent pairwise comparisons made between lung tissues, 11 showed a higher elevation in P-Smad2 in *Tgfbm2/3* compared with NIH mice. (D) Immunohistochemistry for P-Smad2 with positive nuclear staining of bronchiolar epithelial cells of parental NIH and *Tgfbm2/3* lungs after OVA or saline exposure. Enlargement of stained section (red and black box) demonstrates positive staining of bronchiolar epithelial cells (yellow arrow), macrophages (green arrow), and some smooth muscle cells (black arrow) but not eosinophils (red arrow). (E) NIH ImageJ was used to quantify positively stained nuclei for P-Smad2. Quantification of P-Smad2 staining used five fields of view (*n* = 5–6 mice per group, 5 sections per lung). (F) *Inhba* mRNA expression analysis of parental NIH and *Tgfbm2/3* lungs after either saline or OVA challenge. Results are represented compared with GAPDH Ct value. Data shown as mean ± SEM from each group (*n* = 5). *P* values were determined using Student *t* test (**P* ≤ 0.05; ***P* ≤ 0.01; ****P* ≤ 0.001). (G) Immunohistochemistry for activinA with positive cytoplasmic staining of bronchiolar epithelial cells, macrophages (black arrow), and some smooth muscle cells but not eosinophils (red arrow) of parental NIH and *Tgfbm2/3* lungs after OVA or saline exposure.

superfamily. Interrogation of microarray data (see *Molecular Analysis of Differential Asthmatic Responses in NIH vs. Tgfbm2/3 Mice*) for differentially expressed TGFβ signaling genes (Dataset S14), suggested higher OVA-induction of *Inhba* in *Tgfbm2/3* than NIH mice. *Inhba* encodes activinA, another Smad2-activating ligand. This finding was validated by qRT-PCR, confirming significantly higher induction of *Inhba* gene expression in lungs of *Tgfbm2/3* (15.02-fold) vs. NIH mice (10.22-fold) (Fig. 3E). Immunohistochemical staining for activinA showed a very similar distribution to that of P-Smad2, with most activinA localized to bronchial epithelium and macrophages (Fig. 3F).

Enhanced Muscularization of Bronchiolar Submucosa in NIH vs. *Tgfbm2/3* Mice. H&E staining of the lungs suggested an apparent increase in subbronchiolar smooth muscle between OVA-treated NIH and *Tgfbm2/3* mice (Fig. 4A). To further investigate these changes, anti-α-smooth muscle actin (α-SMA) immunofluorescence staining was undertaken on paraffin sections to identify myofibroblasts and smooth muscle cells (Fig. 4B). In both saline control groups, the submucosal smooth muscle layer of the bronchioles was discontinuous, with no difference between NIH and *Tgfbm2/3* mice in the percent investment of bronchiolar epithelium with smooth muscle (Fig. S3). However, quantification of bronchiolar smooth muscle thickness demonstrated that both NIH and *Tgfbm2/3* mice showed a thickening of the smooth muscle layer in response to OVA challenge, and this effect was significantly greater in the NIH than in the *Tgfbm2/3* mice (Fig. 4B).

***Tgfbm2/3* Mice Exhibit Enhanced Airway Inflammation and Eosinophilia Compared with Parental NIH Mice.** Because asthma is considered an inflammatory disease, and the *Tgfbm2/3* mice were relatively protected from OVA-induced AHR, we hypothesized that this strain would show a reduced pulmonary inflammatory infiltrate compared with NIH mice after OVA challenge. On the contrary, although both mouse strains showed a robust increase in total bronchoalveolar lavage (BAL) cell counts after OVA challenge (Fig. 5A) there was a larger influx of inflammatory cells into the lungs of *Tgfbm2/3* compared with NIH mice (Fig. 5A). These findings were most evident when comparing eosinophil cell numbers (Fig. 5B; NIH: 145 ± 27.6; vs. *Tgfbm2/3*: 390 ± 105.8; *P* = 0.03). Comparison of infiltrating numbers of macrophages, lymphocytes, or non-eosinophil polymorphonuclear cells did not differ significantly between the mouse strains (Fig. 5C–E). Thus, the *Tgfbm2/3* mice paradoxically showed greater pulmonary eosinophilia than parental NIH, despite the relative resistance of the NIH strain in terms of AHR.

Robust T_H2-Mediated Inflammatory Response in both NIH and *Tgfbm2/3* Mice. Because the *Tgfbm2/3* mice had more pronounced eosinophilia than NIH mice, we expected the cytokine profiles of these two strains to differ considerably. IgE-dependent mast cell activation induces the T_H2 allergic response, via release of T_H2-associated cytokines IL-4, IL-5, and IL-13. To investigate the cellular mechanisms of action of the TGF-β modifier loci in enhancing eosinophilia but protection from AHR, we examined

Tgfbm3 control fundamental differences in the biosynthetic machinery of the cell even in the absence of OVA exposure.

Because the extensive OVA-induced inflammatory response in *Tgfbm2/3* mice might mask gene expression differences responsible for the higher OVA-induced AHR in NIH, we mined the dataset for transcripts exhibiting a significantly larger differential in gene expression between saline and OVA treatment (Δ_{S-O}) in NIH mice vs. *Tgfbm2/3* mice (Dataset S1B). As a control, we undertook the converse analysis, comparing genes that had a significantly higher saline-to-OVA differential (Δ_{S-O}) in *Tgfbm2/3* than NIH mice (Dataset S1A). As expected, transcripts showing significantly larger Δ_{S-O} differentials in *Tgfbm2/3* than NIH mice were enriched for inflammatory signatures including chemokines and chemokine receptors, as well as markers of epithelial keratinization (Dataset S4A). In contrast, genes exhibiting greater Δ_{S-O} differentials in NIH than *Tgfbm2/3* mice were enriched for protein biosynthesis and posttranslational modification, including phosphoproteins, acetylation, and methylation, as well as nucleotide metabolism and chromosomal proteins (Dataset S4B). Once again this indicates a fundamental difference in the overall cellular biosynthetic machinery, not only between baseline characteristics of NIH vs. *Tgfbm2/3* lungs (Dataset S2B), but also in the pulmonary responses of NIH to OVA (Dataset S4B).

Discussion

Because TGF β 1 levels are elevated and implicated in the pathology of both acute and chronic asthma models, and this signaling pathway is a potential therapeutic target for reducing excessive airway remodeling and pulmonary fibrosis, many studies have addressed the role of this growth factor in experimental asthma. Manipulation of TGF β activity either genetically or pharmacologically has had variable outcomes. In a model of chronic OVA-induced asthma, therapeutic anti-TGF β antibody dosing reduced peribronchiolar extracellular matrix, airway smooth muscle cells, and mucus production without stimulating airway inflammation or T_H2 cytokine production (11). In contrast, in a house dust-mite-induced model, anti-TGF β antibodies had no effect on airway remodeling but exacerbated eosinophilia resulting in potentiated AHR (23). More recently, it was shown that TGF β produced by bone marrow stromal cells that home to the lung during an asthmatic episode, is implicated in reducing the pulmonary allergic inflammatory response (24), which is consistent with other reports that TGF β plays a protective role in the asthmatic response (9, 10, 25, 26).

Here we have shown strain-specific differences in the magnitude of responses of C57 and NIH mice to an asthmatic stimulus. Both mouse strains showed elevated AHR to an OVA challenge, yet the C57 strain was relatively resistant compared with NIH. More importantly, whereas in NIH mice innate TGF β 1 was protective from OVA-induced AHR, on the C57 genetic background, loss of a single *Tgfb1* allele had no effect on AHR. Thus, the functional response to reduced TGF β 1 levels in a murine asthma model is determined by genetic background. Contrasting with our findings, others found that *Tgfb1* haploinsufficiency potentiates AHR in C57 mice (10). However, it is possible that the strain used in the earlier study was not a pure inbred C57 background, but carried genetic contaminants from other sources that might influence AHR (22). In the current study, *Tgfb1*^{+/-} mice had been backcrossed through >20 generations to C57 and found to be pure by whole genome scan 500K SNP array (Jax mouse diversity genotyping array).

In an attempt to identify genetic variants that might regulate the differential responses to reduced TGF β 1 signaling on AHR, we hypothesized that the *Tgfb1* genetic modifier loci, *Tgfbm2* and *Tgfbm3* might play a role. When we used NIH mice into which we had bred variant *Tgfbm2* and/or *Tgfbm3*, we found that neither locus alone affected AHR response but the two synergized to significantly reduce AHR. These two variant loci functionally compensated for reduced TGF β 1 levels in NIH-*Tgfb1*^{+/-} mice, in that mice carrying both variant loci no longer showed increased AHR in response to loss of a *Tgfb1* allele. Thus, dependent on genetic variation within *Tgfbm2* and *Tgfbm3*, the asthma-protective effect of TGF β 1 may be completely suppressed.

Another finding of the current study is the uncoupling between regulation of eosinophilia, airway remodeling, and AHR during an asthmatic episode, which also depends on *Tgfbm2* and *Tgfbm3*. Recent studies on human asthma have demonstrated clinical stratification of asthmatics into T_H2 low and T_H2 high populations (27, 28), implicating a noneosinophilic mechanism of disease pathology in T_H2 low asthmatics. However, the molecular etiology of asthma in the T_H2 low population has yet to be elucidated. In mice, variation in genetic background has previously been found to affect different components of the asthmatic response in different manners. For example, in an acute model of OVA-induced asthma, C57 and BALB/c mice both developed similar extents of eosinophilic inflammation. However, the BalbC strain shows a significantly larger elevation in AHR and peribronchiolar smooth muscle expansion compared with C57 (29). Moreover, in C57 mice, the asthmatic response is totally dependent on eosinophils, as C57 Δ dbiGATA1 knockout mice that lack eosinophils are completely resistant to OVA-induced AHR, whereas BALB/c Δ dbiGATA1 knockouts still show elevated AHR in response to OVA (30). In the current study, we found that synergy between *Tgfbm2* and *Tgfbm3* regulates the eosinophilic and AHR responses to OVA challenge but these two physiological responses are regulated in oppositional directions by this genetic interaction, resulting in reduced AHR in the presence of increased eosinophilia in NIH mice homozygous for *Tgfbm2* and *Tgfbm3*.

Tgfbm2 and *Tgfbm3* might functionally compensate for reduced TGF β 1 of *Tgfb1*^{+/-} mice by stimulating TGF β signaling downstream of the ligand, therefore reducing AHR. Indeed, mice carrying both variant modifier loci exhibited higher levels of OVA-induced pulmonary P-Smad2 than mice harboring the NIH alleles. ActivinA may contribute to this effect, because it was the only Smad2-activating ligand showing a greater induction in *Tgfbm2/3* than NIH mice. Both P-Smad2 and activinA were predominantly localized to bronchiolar epithelium and alveolar macrophages, with some staining in smooth muscle cells but none in eosinophils. Intriguingly, activinA has been reported, like myostatin, to suppress muscle mass (31) and may therefore contribute to the lack of bronchiolar smooth muscle expansion and consequent reduced AHR in *Tgfbm2/3* compared with NIH mice.

Two outstanding issues are the identity of the *Tgfbm2* and *Tgfbm3* causative genetic variants that compensate for TGF β 1 and elicit differential asthmatic responses. It is now widely accepted that genetic modifier loci that influence multifactorial diseases frequently constitute clusters of functionally related polymorphic genes. This has been observed in mouse models of lupus (32) and cancer (33), making it difficult to dissect the underlying molecular mechanisms. With respect to *Tgfbm2*, *Cenpf* harbors an amino acid polymorphism between mouse strains at an evolutionarily conserved residue (18) (Fig. S1A). *Cenpf* has been proposed to regulate muscle determination (34, 35), although its major function is regulation of the centrosome and cytokinesis. Conceptually, distinct *Cenpf* isoforms might drive variable proliferation and/or differentiation in distinct tissue subsets.

To address the issue further, we undertook microarray gene expression analysis of NIH vs. *Tgfbm2/3* mice. Consistent with the near genetic identity of NIH and *Tgfbm2/3* mice, microarray analysis revealed no large differentials in gene expression between the two strains. Both mouse strains showed induction of an inflammatory signature after OVA treatment, albeit to differing extents. Because *Tgfbm2/3* mice exhibited significantly more inflammation than NIH, this confounded interrogation of microarray data for genes that might contribute to the higher AHR in NIH vs. *Tgfbm2/3*. Around 500 genes showed greater differential gene expression (Δ_{S-O}) in NIH than in *Tgfbm2/3* mice. Intriguingly, this set of genes was enriched in proteins involved in protein biosynthesis and posttranslational modifications, including acetylation, phosphorylation, and methylation. In this respect, it is provocative that several genes encoding regulators of DNA or chromatin modification are localized in the *Tgfbm3* region, including *Asxl2*, a polycomb protein; *Dnmt3a*, DNA methyl transferase 3a; *DNAJC27* and *Ddx1*; as well as a regulator of translation and message stability, *Pum2* (Fig. S1B). In particular, it was recently demonstrated that DNA methyltransferase 3a acts directly on the *Il-13* gene within T_H2 cells to

down-regulate IL-13 expression, and hence allergic airway inflammation in mice (36). It is certainly possible that genetic variation at *Dnm3a* may contribute to the phenotypic differences conferred by *Tgfbm3*, acting in concert with *Tgfbm2*. Microarray gene expression analysis of whole lung may be insufficiently sensitive to detect differentials in gene expression within minor cell type subsets of the lung, such as bone marrow stromal cells (24) or T_H2 subsets (36), that nevertheless have a large impact on the outcome of the allergic response.

To summarize, the major conclusion from this study is that the biological responses to reduced TGFβ1 signaling in an experimental asthma model are dependent not only on the model (acute vs. chronic), or the cell types affected (smooth muscle vs. inflammatory vs. epithelial), but on the differential inheritance of endogenous genetic variants of wild-type modifier genes. This effect is pronounced, because differential inheritance of *Tgfbm2* and *Tgfbm3* can reverse the biological outcome of reduced *Tgfb1* levels. These findings present a cautionary note for the use of TGFβ agonists or antagonists for the treatment of acute or chronic lung diseases, indicating a more personalized approach to the use of such agents. However, further dissection of the genes and molecules involved in modifying the outcome of reduced TGFβ activity may shed light on alternative pathways that could be targeted for treatment of lung diseases.

Materials and Methods

Animals. The *Tgfb1* null allele used in this study was *Tgfb1^{Tm1n}* (7), referred to as *Tgfb1^{+/-}* throughout the text and figures. These mice were bred as previously described (19). NIH.C57-*Tgfbm3* congenic mice were generated by backcrossing B6NIHF1 mice through more than seven generations to recipient NIH mice. At each generation, selection was made for C57 genetic markers spanning from the centromere of chromosome 12 to exon 3 of the *Lipn1* gene. NIH.129-*Tgfbm2* congenic mice were the result of biological coselection with the *Tgfb1* null allele of 129 genomic DNA at *Tgfbm2* through

multiple backcross generations to NIH mice (22). The *Tgfbm2¹²⁹* region spans from exons 57–60 of *Ush2a* to exons 2–6 of *Ptpn14*.

All experiments involved 10-wk-old male animals in groups of 10 mice per arm except for experiment 4, where 8 animals were used in the OVA arm and 5 animals were used in the saline control group. All animal husbandry practices were in full compliance with the recommendations published in *Guide for the Care and Use of Laboratory Animals* (37). All proposed protocols have been approved by the University of California San Francisco Institutional Animal Care and Use Committee.

Analysis of MicroArray Dataset. Affymetrix Mouse Gene 1.0 ST Array CEL files were preprocessed (background correction, quantile normalization, probelet summarization) using the *aroma.affymetrix* R package (38). Differential expression analysis using moderated *t* statistic with Benjamini–Hochberg false discovery rate (adjusted *P* value) correction was performed using the *Limma* R package (39).

Genes responding to treatment with adjusted *P* value of <0.05 were initially selected. To focus on genes with substantial differences in log fold change (FC) between NIH and *Tgfbm2/3*, the following two quantities, *M* = difference in log FC between NIH and *Tgfbm2/3*, and *A* = average log FC, were evaluated. Genes with absolute *M/A* > 0.33 were retained. These were segregated into a group with stronger NIH response, where [abs (logFC NIH/logFC *Tgfbm2/3*) > 1] and one with stronger *Tgfbm2/3* response [abs (logFC NIH/logFC *Tgfbm2/3*) < 1].

Functional enrichments for the four gene lists were derived by uploading MicroArray ProbeID into DAVID using Affymetrix_Exon_Gene_ID as identifier and MoGene-1-0 as background setting. The output of SP_PIR keywords (SP_PIR = Swiss Prot Protein Information Resource) within the functional categories was determined with a *P* value threshold set to 0.05.

ACKNOWLEDGMENTS. Microarray analysis was performed by the Gladstone Institute's Microarray Core. Work was funded by National Institutes of Health Grants R01-HL078564 and R01-GM60514 (to R.J.A.) and the Bouque Foundation (R.J.A.). F.F.C. was the recipient of fellowships from the Belgian American Educational Foundation and the American Heart Association.

- McFadden ER, Jr., Gilbert IA (1992) Asthma. *N Engl J Med* 327:1928–1937.
- Pulleyn LJ, Newton R, Adcock IM, Barnes PJ (2001) TGFβ1 allele association with asthma severity. *Hum Genet* 109:623–627.
- Silverman ES, et al. (2004) Transforming growth factor-β1 promoter polymorphism C-509T is associated with asthma. *Am J Respir Crit Care Med* 169:214–219.
- Gentile DA, et al. (2004) Association between TNF-α and TGF-β genotypes in infants and parental history of allergic rhinitis and asthma. *Hum Immunol* 65:347–351.
- Mak JC, et al. (2006) Analysis of TGF-β(1) gene polymorphisms in Hong Kong Chinese patients with asthma. *J Allergy Clin Immunol* 117:92–96.
- Howell JE, McAnulty RJ (2006) TGF-β: Its role in asthma and therapeutic potential. *Curr Drug Targets* 7:547–565.
- Kulkarni AB, et al. (1993) Transforming growth factor β 1 null mutation in mice causes excessive inflammatory response and early death. *Proc Natl Acad Sci USA* 90:770–774.
- Diebold RJ, et al. (1995) Early-onset multifocal inflammation in the transforming growth factor β 1-null mouse is lymphocyte mediated. *Proc Natl Acad Sci USA* 92:12215–12219.
- Schramm C, et al. (2003) TGF-β regulates airway responses via T cells. *J Immunol* 170:1313–1319.
- Scherf W, Burdach S, Hansen G (2005) Reduced expression of transforming growth factor β 1 exacerbates pathology in an experimental asthma model. *Eur J Immunol* 35:198–206.
- McMillan SJ, Xanthou G, Lloyd CM (2005) Manipulation of allergen-induced airway remodeling by treatment with anti-TGF-β antibody: Effect on the Smad signaling pathway. *J Immunol* 174:5774–5780.
- Alcorn JF, et al. (2007) Transforming growth factor-β1 suppresses airway hyper-responsiveness in allergic airway disease. *Am J Respir Crit Care Med* 176:974–982.
- Sime PJ, Xing Z, Graham FL, Csaky KG, Gauldie J (1997) Adenovector-mediated gene transfer of active transforming growth factor-β1 induces prolonged severe fibrosis in rat lung. *J Clin Invest* 100:768–776.
- Kenyon NJ, Ward RW, McGrew G, Last JA (2003) TGF-β1 causes airway fibrosis and increased collagen I and III mRNA in mice. *Thorax* 58:772–777.
- Kudo M, et al. (2012) IL-17A produced by αβ T cells drives airway hyper-responsiveness in mice and enhances mouse and human airway smooth muscle contraction. *Nat Med* 18:547–554.
- Sugimoto K, et al. (2012) The αvβ6 integrin modulates airway hyper-responsiveness in mice by regulating intraepithelial mast cells. *J Clin Invest* 122:748–758.
- Bonyadi M, et al. (1997) Mapping of a major genetic modifier of embryonic lethality in TGF β 1 knockout mice. *Nat Genet* 15:207–211.
- Tang Y, et al. (2003) Genetic modifiers interact with maternal determinants in vascular development of *Tgfb1(-/-)* mice. *Hum Mol Genet* 12:1579–1589.
- Tang Y, et al. (2005) Epistatic interactions between modifier genes confer strain-specific redundancy for *Tgfb1* in developmental angiogenesis. *Genomics* 85:60–70.
- Mao JH, et al. (2006) Genetic variants of *Tgfb1* act as context-dependent modifiers of mouse skin tumor susceptibility. *Proc Natl Acad Sci USA* 103:8125–8130.
- Tian F, et al. (2003) Reduction in Smad2/3 signaling enhances tumorigenesis but suppresses metastasis of breast cancer cell lines. *Cancer Res* 63:8284–8292.
- Benzinou M, et al. (2012) Mouse and human strategies identify PTPN14 as a modifier of angiogenesis and hereditary haemorrhagic telangiectasia. *Nat Commun* 3:616.
- Fattouh R, et al. (2008) Transforming growth factor-β regulates house dust mite-induced allergic airway inflammation but not airway remodeling. *Am J Respir Crit Care Med* 177:593–603.
- Nemeth K, et al. (2010) Bone marrow stromal cells use TGF-β to suppress allergic responses in a mouse model of ragweed-induced asthma. *Proc Natl Acad Sci USA* 107:5652–5657.
- Gordon ED, et al. (2012) A protective role for periostin and TGF-β in IgE-mediated allergy and airway hyperresponsiveness. *Clin Exp Allergy* 42:144–155.
- Presser K, et al. (2008) Coexpression of TGF-β1 and IL-10 enables regulatory T cells to completely suppress airway hyperreactivity. *J Immunol* 181:7751–7758.
- Woodruff PG, et al. (2009) T-helper type 2-driven inflammation defines major sub-phenotypes of asthma. *Am J Respir Crit Care Med* 180:388–395.
- Choy DF, et al. (2011) Gene expression patterns of Th2 inflammation and intercellular communication in asthmatic airways. *J Immunol* 186:1861–1869.
- Van Hove CL, et al. (2009) Comparison of acute inflammatory and chronic structural asthma-like responses between C57BL/6 and BALB/c mice. *Int Arch Allergy Immunol* 149:195–207.
- Walsh ER, et al. (2008) Strain-specific requirement for eosinophils in the recruitment of T cells to the lung during the development of allergic asthma. *J Exp Med* 205:1285–1292.
- Lee SJ, et al. (2010) Regulation of muscle mass by follistatin and activins. *Mol Endocrinol* 24:1998–2008.
- Morel L, Blenman KR, Croker BP, Wakeland EK (2001) The major murine systemic lupus erythematosus susceptibility locus, *Slc1*, is a cluster of functionally related genes. *Proc Natl Acad Sci USA* 98:1787–1792.
- Samuelson DJ, et al. (2007) Rat *Mcs5a* is a compound quantitative trait locus with orthologous human loci that associate with breast cancer risk. *Proc Natl Acad Sci USA* 104:6299–6304.
- Papadimitou E, Ménard C, Grey C, Puceat M (2005) Interplay between the retinoblastoma protein and LEK1 specifies stem cells toward the cardiac lineage. *EMBO J* 24:1750–1761.
- Pabón-Peña LM, Goodwin RL, Cise LJ, Bader D (2000) Analysis of CMF1 reveals a bone morphogenetic protein-independent component of the cardiomyogenic pathway. *J Biol Chem* 275:21453–21459.
- Yu Q, et al. (2012) DNA methyltransferase 3a limits the expression of interleukin-13 in T helper 2 cells and allergic airway inflammation. *Proc Natl Acad Sci USA* 109:541–546.
- National Research Council, Committee for the Update of the Guide for the Care and Use of Laboratory Animals, and Institute for Laboratory Animal Research (2011) *Guide for the Care and Use of Laboratory Animals* (National Academies Press, Washington, DC), 8th Ed.
- Bengtsson H, Simpson K, Bullard J, Hansen K (2008) *Aroma.affymetrix*: A generic framework in R for analyzing small to very large Affymetrix data sets in bounded memory (Department of Statistics, Univ of California, Berkeley), Technical Report 745.
- Smyth GK (2005) *Limma*: Linear models for microarray data. *Bioinformatics and Computational Biology Solutions Using R and Bioconductor*, eds Gentleman R, Carey V, Dudoit S, Irizarry R, Huber W, Springer, New York, pp 397–420.