



Published in final edited form as:

Sex Transm Infect. 2012 November ; 88(7): 484–489. doi:10.1136/sextrans-2012-050618.

Condom Effectiveness against Non-Viral Sexually Transmitted Infections: A Prospective Study Using Electronic Daily Diaries

Richard Crosby, PhD¹, Richard A. Charnigo, PhD¹, Chandra Weathers, MPH¹, Angela M. Caliendo, MD, PhD², and Lydia A. Shrier, MD, MPH³

¹College of Public Health at the University of Kentucky

²Emory University School of Medicine, Department of Pathology and Laboratory Medicine (Infectious Diseases)

³Division of Adolescent/Young Adult Medicine, Children's Hospital Boston and Department of Pediatrics, Harvard Medical School

Abstract

Objectives—To prospectively evaluate the protective value of consistent and correct use of latex condoms against the acquisition of *Chlamydia trachomatis*, *Neisseria gonorrhoeae*, and *Trichomonas vaginalis*.

Methods—Patients (N=929) attending clinics that treat sexually transmitted infections (STIs) were prospectively followed for up to six months. Urine STI nucleic acid amplification testing was performed at baseline, three months, and six months. Participants were instructed to respond to daily prompts from a hand-held device by completing a report for each penile-vaginal sexual intercourse event. Generalized estimating equation models examined associations of consistent as well as consistent and correct condom use with STI incidence over 3-month intervals.

Results—Consistent condom use was not significantly associated with STI incidence (Estimated Odds Ratio [EOR]=.75; 95% confidence interval [CI]=.43-1.30; *P*=.31). However, individuals who used condoms both correctly and consistently were estimated to have 59% lower odds of acquiring an STI (EOR = .41; 95% CI = .19-.90; *P* = .026), compared to those who did not.

Conclusions—The correct as well as the consistent use of condoms greatly reduces the odds of non-viral STI acquisition.

Introduction

More than a decade ago, the United States Department of Health and Human Services issued a report about the effectiveness of latex condoms for the prevention of sexually transmitted infections (STIs).¹ Beyond HIV transmission and female-to-male transmission of gonorrhea, the report concluded evidence was insufficient to judge the protective value of condoms

Correspondence: Richard Crosby, PhD, College of Public Health, 151 Washington Ave., Lexington, KY 40506-0003, Telephone: 859-218-2039/Fax: 859-323-2933/crosby@uky.edu.

Author Contributions

Dr. Crosby acquired funding for the study, served as PI, provided oversight of all data collection procedures, drafted the initial manuscript, and revised the final manuscript. Dr. Charnigo served as the study statistician, provided oversight of data collection and cleaning, conducted all analyses, and participating in manuscript preparation.

Mrs. Chandra Weather served as the Project Director and thus provided oversight for data collection and assured adherence to study protocols in all location.

Dr. Caliendo supervised the laboratory responsible for NAAT testing of all urine specimens.

Dr. Shrier assisted Dr. Crosby in obtaining funding, provided scientific guidance, designed protocols for use of PDAs and CERTAS, and participated in manuscript preparation.

against other STIs. Since the report was issued, findings from a few well-designed studies suggest a protective value of condoms against male-to-female transmission of genital herpes,² Chlamydia/gonorrhea,³ syphilis,⁴ and human papillomavirus.⁵ However, most studies conducted in the past ten years have failed to show a significant association between condom use and STI acquisition.⁶ The majority of these studies have been designed with several inherent forms of bias toward the null hypothesis (i.e., condoms are not protective). For example, a misclassification bias is created when a study does not measure and adjust for the incorrect use of condoms (failure to use the device from start to finish of penetrative sex) or for the events of breakage or slippage.⁶⁻⁸ In essence, failure to control for condom breakage and slippage may produce the analytical equivalent of condom nonuse, which confounds study findings. For example, a prospective study of clinic attendees found 13% incidence of Chlamydia and gonorrhea among people reporting consistent condom use but also reporting at least one problem with correct use.⁹ In contrast, among those reporting consistent use and a lack of problems (slippage, breakage, leaking, early removal, late application), no incident infections were found.

Past studies have also relied on the validity of retrospective recall, sometimes over periods of time measured in months rather than days or weeks.^{6,7} Fortunately, the science of collecting valid self-reported data on condom use behaviors has greatly improved in recent years.¹⁰ In particular, mobile electronic devices enable daily reporting, which may dramatically reduce recall bias and minimize social desirability bias.¹¹ In parallel, improved technology has enhanced validity of nucleic acid amplification testing for STI assessment. Also, single-dose therapy to treat prevalent cases provides assurance that subsequent infections are truly incident cases.

Evidence suggests that people may use condoms as a consequence of a correct position that they are likely to have an STI.¹² This post-infection condom use confounds data analyses by deflating the expected negative association between use and STI acquisition.¹³ The issue is overcome by a prospective design that establishes an infection-free cohort initially and by measuring behaviors over relatively short time intervals. In condom effectiveness studies, the confound of post-infection condom use is eliminated from data analyses based on subjects' consistency (100% use) or lack thereof, instead of their relative frequencies of condom use. For example, someone who uses condoms 40% of the time, becomes infected, and then starts using condoms 80% of the time because of the infection is (appropriately) not deemed a superior condom user in the analytic approach of comparing 100% users to the remainder.

The purpose of this study was to prospectively determine the protective value of consistent and correct condom use against urethral/vaginal acquisition of three common STIs (*Chlamydia trachomatis*, *Neisseria gonorrhoeae*, and *Trichomonas vaginalis*) using daily electronic assessments and NAAT technology. First, we tested the hypothesis that consistent condom use would have a significant protective effect against the three STIs. Subsequently, we tested whether consistent *and correct* condom use would have a protective effect.

Methods

Study Design

Outpatients were recruited from five clinics caring for individuals at high-risk for STI in three U.S. cities: a publicly-funded STI clinic in the Southern US; a publicly-funded STI clinic in the Midwestern US; and an STI clinic of a large teaching hospital and two adolescent medicine clinics affiliated with a children's hospital, all in the Northeastern US. The STI clinics enrolled individuals aged 18 years and older; the adolescent clinics enrolled individuals as young as 15. Eligibility criteria included reporting penile-vaginal intercourse

in the preceding 3 months; willing to be tested for Chlamydia, gonorrhea, and trichomoniasis by providing a urine specimen; speaking English; willing to provide contact information; and providing written informed consent. Institutional review boards at the participating universities approved the study protocol with a waiver of parental consent (only assent was required) for adolescents less than 18 years of age.

Recruitment procedures varied slightly across the five clinics. At the adolescent clinics, the study was listed on a research recruitment flag attached to the appointment paperwork of age-eligible patients. The research assistant used the flag to identify eligible patients. This chart-flagging system at adolescent clinics precluded us from calculating a participation rate for those sites. Across the three remaining clinics, 1,424 patients agreed to be screened for eligibility. Of these, 1,297 were eligible and invited to participate; 794 enrolled, yielding a participation rate of 61.2%. With the remaining patients from the Boston clinics (n=135), the participant sample size was 929. Data were collected from December 2007 through April 2011.

At baseline, participants completed a gender-specific audio computer-assisted self-interview (A-CASI) assessing their sociodemographic characteristics, sexual history, and condom use behaviors, then provided a first-catch urine specimen for STI testing. To optimize their condom use, all participants were offered a brief (30 to 45 minutes) gender-specific educational session about using condoms, adapted from a safer sex intervention¹⁴ designated “best evidence” by the Centers for Disease Control and Prevention.¹⁵ A key feature of this interactive counseling program was learning about the “fit and feel” of condoms and the pleasure-based aspects of finding the right size and type of condoms, as well as the use of water-based lubricants. Participants were then offered their choice from an extensive selection of condoms and water-based lubricants.

Participants were trained in the use of an electronic diary report using a password-protected personal digital assistant (PDA) programmed with the Configurable Electronic Real-Time Assessment System (CERTAS; Personal Improvement Computer Systems, Inc. Reston, VA, USA). Each day, participants were prompted to answer the question of whether they had sex in the past 24 hours. Because the research question pertained to urethral/vaginal acquisition of STIs, our assessments concerned penile-vaginal sex. Thus, oral sex was not assessed in this study. Based on formative research,¹⁶ we defined sex as “putting the penis in the vagina” and the end of sex as the male orgasm. If participants responded affirmatively, they were asked questions about each instance of sexual intercourse. They were also asked to enter a report about their sexual behavior directly into the PDA after each time that they had sex to maximize capture of data on every sex event. To avoid duplicative reporting, participants were prompted to indicate on the daily diary if they had already completed a report about a sex event. To minimize loss of data, participants were asked to return the PDA memory card every 30 days, at which time a new one was provided.

At the 3-month and 6-month follow-up interviews, participants completed an ACASI regarding their sexual behaviors in the past 3 months and provided urine specimens for STI testing. Incident STIs detected at the 3-month visit were treated prior to the start of the second three months of daily reporting. Participants were offered remuneration in gift cards based on the study activities completed (maximum amount was approximately USD 2.50 per day of observation).

Assessments

STI testing—At baseline, three months, and six months, first-catch urine specimens were tested using the Becton Dickinson (Sparks, MD) ProbeTec ET *C. trachomatis* and *N. gonorrhoeae* Amplified DNA Assay¹⁷ and the Taq-Man PCR-ELISA for *T. vaginalis*

(developed and validated by the research laboratory¹⁸). All urine specimens were processed within 48 hours and shipped to the study laboratory within seven days. Participants testing positive for any of the three STIs were contacted immediately and asked to schedule an appointment for treatment. Single-dose therapy with CDC-recommended medications was used to maximize treatment effectiveness.¹⁹ Finally, to capture data on STIs that occurred between follow-up study visits, we conducted a chart review at the end of the observation period for each participant.

Daily measures—On the daily electronic report, participants who indicated sex in the past 24 hours were asked about each sex event, “Did you and your partner use a condom for the penis-in-vagina part of this sex event?” If they responded yes, they were asked questions about the condom use, including the occurrence of the following five errors and problems: breakage, slippage during sex, slippage after sex was over, putting the condom on after sex began, and taking the condom off before sex ended.

Data Analysis

Data were summarized across each three-month interval (0 to 3 months and 3 to 6 months), with participants contributing either one or two intervals to the analyses. Analyses were restricted to three-month intervals with at least one sex event report. *Consistent condom use* was defined as condom use with every sex event. *Correct condom use* was defined as a “no” answer to each of the five questions on errors and problems when a condom was used. The association between condom use and incident STI was examined using generalized estimating equation (GEE) models to account for within-individual correlation of observation intervals.^{20,21} Model 1 tested the association between consistent condom use (yes, no) and 3-month incident STI. Model 2 examined the association between condom use and incident STI when correct as well as consistent condom use (yes, no) was required. Both models adjusted for gender, age group (15-19, 20-24, 25+ years), and history of STI (STI reported on baseline ACASI or positive STI test result during the preceding 3-month interval), which were examined individually for significant interactions with the condom use variable. Significance was defined by an alpha level of .05. Power calculations suggested that 80% power was available to detect a protective odds ratio as small as .31 for the effect of consistent and correct use and .43 for the effect of only consistent use.

Results

The mean+SD age of the participants was 29.2+10.8 years. Two-thirds (n=617, 66.5%) identified as African American/Black and more than one-half (n=512, 55.1%) were women. The mean+SD number of lifetime sex partners was 29.7+38.2. Table 1 provides this demographic information stratified by city. Most (65.7%) of those 18 and older reported earning less than \$1,000 per month in income or social assistance and 54.4% of those under 18 reported they qualified for a free lunch at school. Just under one-third of the sample (30.7%) reported ever being diagnosed with a STI. Seven hundred and eight participants (76.2%) returned for the initial follow-up assessment visit and 523 returned for a second follow-up assessment.

Three hundred-eighty participants contributed data for two 3-month intervals and 200 contributed data for one 3-month interval, yielding a total of 960 3-month intervals. Of the 380 participants providing data for both intervals, only 99 were classified discrepantly on consistent use between the two intervals. Of these same 380 participants, only 52 were classified discrepantly on consistent and correct use between the two intervals. Of the 960 3-month intervals, 14 (1.5%) involved indeterminate STI assay results, resulting in 946 3-

month intervals for analysis. Table 2 displays further descriptive information for each 3-month interval, stratified by city.

Participants reported a total of 14,970 penile-vaginal sex events, of which 9,545 (63.8%) involved use of a condom. Of the sex events involving a condom, 2,285 (23.9%) included one or more condom use errors or problems, leaving 7,260 events when condoms were used correctly. Table 3 displays further descriptive information for each 3-month interval, stratified by gender.

Three-month and 6-month assays for incident STIs yielded 116 positive test results (44 for Chlamydia, 15 for gonorrhea, and 57 for trichomoniasis). In addition, chart reviews of clinic diagnoses made during study enrollment yielded 2 cases that were not detected by NAAT at follow-up. Of these 118 cases, 81 (69%; 31 for Chlamydia, 10 for gonorrhea, and 40 for trichomoniasis) had electronic diary data over the corresponding 3-month interval. Accounting for co-infections, there were 74 analysis intervals during which a subject acquired at least one of the three STIs.

To determine whether systematic differences existed between the 74 observation intervals remaining in the analysis and the 29 intervals that were excluded, a comparison was conducted based on the ACASI baseline data. These data allowed us to test for differences in consistent condom use (but not correct use as this had to be assessed in real time) and the three covariates (gender, age, and history of STIs). Significant differences for gender ($P=.27$), age ($P=.99$), and STI history ($P=.30$) were not found. However, the 29 intervals excluded were more likely ($P=.023$) to include consistent condom use than the 74 intervals in the analysis.

Bivariate Associations

Of the 946 3-month intervals, 603 (63.7%) involved less-than-consistent use of condoms. STIs occurred in 51 of these 603 intervals, yielding 8.46% incident infections among those not reporting consistent use during a 3-month interval. Conversely, 343 intervals (36.3%) involved consistent use of condoms. STIs occurred in 23 of these intervals, yielding 6.71% incident infections among those who reported consistent condom use. The difference between these two percentages (absolute difference = 1.75%; percent relative difference = 20.7%) was not significant (estimated odds ratio = .73; 95% confidence interval [CI] = .42 - 1.26; $P=.26$).

Of the 946 observation intervals, 777 (82.1%) involved less-than-consistent and/or incorrect use of condoms. Within these 777 intervals, 68 cases of at least one STI occurred, yielding 8.75% incident infections among those not reporting consistently *and correct* use during an observation interval. Conversely, 169 (17.9%) involved consistent and correct use of condoms. Within these intervals, 6 cases of at least one STI occurred, yielding 3.35% incident infections among those reporting consistent and correct use. The difference between these two percentages (absolute difference = 5.20%; percent relative difference = 59.0%) was significant (estimated odds ratio = .41; 95% CI = .19 - .88; $P=.023$).

We also investigated the possibility of a linear relationship between the number of unprotected events and the log odds of incident STIs. Bivariate analyses showed that incidence jumped from 3.6% (6/169) with no unprotected or imperfectly protected events, to 9.0% (11/122) with one, and to 8.9% (20/224) with between two and four such events, suggesting that, for example, one such event is not 25% as bad as four such events.

Multivariate Analyses

Table 4 displays the findings from GEE analyses modeling the effect of condom use on incident STI over the observation intervals. Adjusted for gender, age group, and history of STI, consistent condom use was not significantly protective against acquiring an incident STI (Model 1). Significant interactions were not found between consistent condom use and age group ($P=0.27$), gender ($P=0.43$), or STI history ($P=0.22$).

In contrast to Model 1, participants who used condoms both correctly and consistently were estimated to have 59% smaller odds of acquiring an STI over three months (see Model 2 in Table 4) compared to participants who did not use condoms both correctly and consistently, adjusting for gender, age group, and history of STI. Significant interactions were not found between consistent and correct condom use and age group ($P=0.60$), gender ($P=0.59$), or STI history ($P=0.81$).

Discussion

Findings provide a striking contrast between testing condom effectiveness based on consistent use versus a more refined measure accounting for errors and problems. This contrast parallels the concepts of “typical use effectiveness” and “perfect use effectiveness” from contraceptive studies.²² The discrepancy between typical and perfect use in this study was dramatic, with point estimates being .75 and .41, respectively. The implication is that global efforts to promote condom use should be augmented with efforts to promote their correct use. A recent review suggests that condom use errors and problems are a global issue.²³ Incomplete use of condoms is a problem requiring targeted education. Rectifying issues such as poor fit and feel of condoms and using oil-based lubricants may substantially reduce slippage and breakage.²³⁻²⁹

A methodological point of emphasis is that all potential forms of bias to studies of condom effectiveness favor the null hypothesis.^{6,7} Thus, the perfect use point estimate of .41 might have been even lower (but not higher) without bias. Indeed, the six incident cases observed for people using condoms consistently and correctly may be a result of an unprotected sex event, breakage event, etc. that was not reported. Collecting self-reported measures is a science that can only be improved, but never perfected, as tendencies to forget, fabricate, exaggerate, and under-report are inevitable.¹⁰

Several analytic points warrant discussion. In cases where partners are uninfected, inconsistent and incorrect condom use are irrelevant because participants cannot acquire infection. In cases with infected partners, pinpointing the transmission date is impractical, thereby creating the question of whether condoms were used consistently prior to, or after, infection. This question is mute when condoms are used consistently throughout the observation period or not used at all. Thus, using the frequency of unprotected sex as a predictor variable, rather than consistent use or correct and consistent use is inherently flawed. Finally, the absence of sexual event data for 29 intervals in which STIs were acquired was an unfortunate reality. That these intervals included disproportionate numbers of consistent condom users suggests further bias toward the null.

The protective odds ratio of .41 is remarkably similar to the .42 reported by Warner and colleagues against Chlamydia and gonorrhea.⁹ Only our estimate was obtained with accounting for the correct use of condoms. Further, Warner and colleagues did not assess *T. vaginalis* infection. Thus, our findings support and extend their findings.

An important implication of our findings pertains to the lack of significance for consistent use (unadjusted for correct use). This suggests that efficacy trials of safer sex programs

should be evaluated using the metric of consistent use and correct condom use. Failure to account condom use problems/errors could result in an effective intervention being incorrectly deemed ineffective. Also, based on imperfect protection offered by typical condom use, findings support the case of using biomarker outcomes (i.e., STI acquisition) in efficacy trials of HIV prevention programs. Failure to do so may result in an ineffective intervention being incorrectly deemed effective.

Limitations

The study was not adequately powered to analyze condom effectiveness separately for Chlamydia, gonorrhea, and trichomoniasis. Also, any prospective study of condom effectiveness against non-viral STIs precludes the possibility of knowing the infection status of sex partners. Thus, the estimate of .41 must be considered conservative. Further, misclassification bias from inaccurate reporting may have also led to conservative point estimates.^{7,9} Finally, it is well worth noting that sample bias is not a primary issue in this type of study given that the basic research question is biological/physical rather than behavioral.

Conclusion

In this largest study to-date designed specifically to test the effectiveness of condoms against non-viral STIs, we observed that consistent and correct use of condoms reduces the estimated odds of an infection by almost 60%, with no significant differences in effectiveness by age group, gender, or STI history. Magnified over an entire population, this level of risk reduction for sexually active people is substantial. In this study, almost 18% of analysis intervals entailed using condoms consistently and correctly - intervention efforts to raise that value to 59% (corresponding to conversions on half of the analysis intervals not entailing consistent and correct use) would be anticipated to avert approximately 21 infections per 1000 people over a three-month interval.

Acknowledgments

We wish to thank our advisory board members for the guidance: Drs. Ward Cates, Jonathan Zenilman, and Ralph DiClemente. Also, we are grateful to our research assistants - Margaret Reed, Rachel Vickers, Lisa Sunner, Christopher Lops, Amanda Wallace, Ashley Kendall, and Devika Bhushan.

Funding: Supported by a grant from NIAID to Dr. Crosby (RA1068119A) and this work was supported in part by the Center for AIDS Research (P30 AI050409)

The Corresponding Author has the right to grant on behalf of all authors and does grant on behalf of all authors, an exclusive licence (or non exclusive for government employees) on a worldwide basis to the BMJ Publishing Group Ltd to permit this article (if accepted) to be published in STI and any other BMJ PGL products and sub-licences such use and exploit all subsidiary rights, as set out in our licence <http://group.bmj.com/products/journals/instructions-for-authors/licence-forms>

References

1. United States Department of Health and Human Services. [Accessed July 5, 2011] Workshop Summary: Scientific evidence on condom effectiveness for sexually transmitted disease (STI) prevention. Available on-line at <http://www.niaid.nih.gov/dmid/STIs/condomreport.pdf>
2. Wald and A, Langenberg AGM, Link K, et al. Effect of condoms on reducing the transmission of herpes simplex virus type 2 from men to women. *JAMA*. 2001; 285:3100–3106. [PubMed: 11427138]
3. Warner L, Macaluso M, Austin HD, et al. Application of the Case-Crossover Design to reduce unmeasured confounding in studies of condom effectiveness. *Am J Epidemiol*. 2005; 161:765–773. [PubMed: 15800269]

4. Koss CA, Dunne EF, Warner L. A systematic review of epidemiologic studies assessing condom use and risk of syphilis. *Sex Transm Dis*. 2009; 7:401–405. [PubMed: 19455075]
5. Winer RL, Hughes JP, Feng Q, et al. Condom use and the risk of genital human papillomavirus infection in young women. *NEJM*. 2006; 354(25):2645–2654. [PubMed: 16790697]
6. Crosby RA, Bounse S. Condom effectiveness: Where are we now? *Sex Health*. 2011; 9:10–17. [PubMed: 22348628]
7. Crosby RA, DiClemente RJ, Holtgrave DR, Wingood GM. Design, measurement, and analytic considerations for testing hypotheses relative to condom effectiveness against non-viral STIs. *Sex Transm Infect*. 2002; 78:228–231. [PubMed: 12181456]
8. Crosby RA, Salazar LF, DiClemente RJ, et al. Accounting for failures may improve precision: evidence supporting improved validity of self-reported condom use. *Sex Transm Dis*. 2005; 32:513–515. [PubMed: 16041255]
9. Warner L, Newman DR, Fishbein M, et al. Problems with condom use among patients attending sexually transmitted disease clinics: prevalence, predictors, and relation to incident gonorrhea and chlamydia. *Am J Epidemiol*. 2008; 167:341–349. [PubMed: 17989058]
10. Zimmerman, RS.; Atwood, KA.; Cupp, PK. Improving validity of self-reports for sensitive behaviors. In: DiClemente, RJ.; Crosby, RA.; Salazar, LF., editors. *Research methods for health promotion*. Jossey-Bass Wiley; San Francisco, CA: 2006. p. 229-259.
11. Fairley CK, Sze JK, Vodstrcil LA, Chen MY. Computer-assisted self interviewing in sexual health clinics. *Sex Transm Dis*. 2010; 37:665–668. [PubMed: 20975481]
12. Peterman TA, Lin LS, Newman DR, et al. Does measured behavior reflect STD risk? An analysis of data from a randomized controlled behavioral intervention study. *Sex Transm Dis*. 2000; 27:446–451. [PubMed: 10987449]
13. Warner L, Newman DR, Austin HD, et al. Condom effectiveness for reducing transmission of gonorrhea and Chlamydia: The importance of assessing partner infection status. *Am J Epi*. 2004; 159:242–251.
14. Crosby RA, DiClemente RJ, Charnigo R, et al. A brief, clinic-based, safer sex intervention for African American men at-risk of HIV acquisition: A randomized controlled trial. *Am J Public Health*. 2009; 99:S96–S103. [PubMed: 19218185]
15. Centers for Disease Control and Prevention. [Accessed October 21, 2009] Compendium of Evidence-Based HIV Prevention Interventions. 2009. http://www.cdc.gov/hiv/topics/research/prs/promising_evidence-interventions.htm#completelist
16. Mehta CM, Sunner L, Head S, Crosby RA, Shrier LA. “Sex isn’t something you do with someone you don’t care about”: Adolescent females definitions of sex. *J Pediatr Adolesc Gynecol*. 2011; 24:266–271. [PubMed: 21715191]
17. Van der Pol BD, Ferrero L, Buck-Barrington E, Hook E, et al. Multicenter evaluation of the BDProbeTec ET system for detection of *Chlamydia trachomatis* and *Neisseria gonorrhoeae* in urine specimens, female endocervical swabs, and male urethral swabs. *J Clin Micro*. 2001; 39:1008–1016.
18. Caliendo AM, Jordan JA, Green AM, et al. Real-time PCR improves detection of *Trichomonas vaginalis* infection compared with culture using self-collected vaginal swabs. *Infect Dis Obstet Gyn*. 2005; 13:145–150.
19. Workowski KA, Berman S. Sexually transmitted diseases treatment guidelines, 2010. *MMWR*. 2010; 59(RR-12):1–110. [PubMed: 21160459]
20. Kung-Yee, Liang; Scott, Zeger. Longitudinal data analysis using generalized linear models. *Biometrika*. 1986; 73:13–22.
21. Hardin, J.; Hilbe, J. *Generalized estimating equations*. Chapman and Hall/CRC; London: 2003.
22. Hatcher, RA.; Trussel, J.; Nelson, AL.; Cates, W.; Kowal, D., editors. *Contraceptive technology*. 19th revised edition. Ardent Media; New York, NY: 2007.
23. Sanders SA, Yarber WL, Kaufman EL, Crosby RA, Graham CA, Milhausen RM. Condom use errors and problems: A global view. *Sexual Health*. 2011; 9:81–85. [PubMed: 22348636]
24. Crosby RA, Yarber WL, Sanders SA, et al. Men with Broken Condoms: Who and Why? *Sex Transm Infect*. 2007; 83:71–75. [PubMed: 16870644]

25. Crosby RA, Graham CA, Yarber WL, Sanders SA. If the condom fits, wear it: A qualitative study of young African American men. *Sex Transm Infect.* 2004; 80:306–309. [PubMed: 15295131]
26. Crosby RA, Yarber WL, Sanders SA, Graham CA. Condom discomfort and Associated problems with their use among university students. *Am J College Health.* 2005; 54:143–148.
27. Reece M, Herbenick D, Sanders SA, et al. Breakage, slippage and acceptability outcomes of a condom fitted to penile dimensions. *Sex Transm Infect.* 2008; 84:143–149. [PubMed: 17971373]
28. Crosby RA, DiClemente RJ, Yarber WL, et al. An event-specific analysis of condom breakage among African American men at risk of HIV acquisition. *Sexually Transmitted Disease.* 2008; 35:174–178.
29. Yarber WL, Graham CA, Sanders SA, Crosby RA. Correlates of condom Breakage and slippage among university undergraduates. *International Journal of STDs and AIDS.* 2004; 15:467–472.

Key Messages

1. Whether condoms confer significant protection against non-viral STIs is a function of their correct use. Incomplete use is especially common and therefore problematic.
2. The consistent and correct use of condoms provides excellent protection against non-viral STI acquisition.
3. The point estimate of .41 is conservative in that multiple forms of bias toward the null are common and unavoidable in studies of condom effectiveness.
4. Trials of safer sex programs should always be evaluated using the metric of consistent and correct condom use.

Table 1

Demographic Information of the Sample Stratified by City of Enrollment (N = 928)

	Northeastern City (n = 269)	Midwestern City (n = 248)	Southern City (n = 411)
Mean Age/SD and IQR ¹	22.8 (7.1) (18-25)	36.3 (11.6) (25.5-47)	29.0 (9.0) (22-35)
Female gender	172 (63.9%)	126 (50.6%)	214 (52.1%)
Black race	95 (35.3)	198 (79.8)	214 (52.1)
Mean # sex partners, 3M ²	2.0 (1.7) (1-2)	2.9 (5.0) (1-3)	3.5 (8.6) (1-3)
Mean # sex partners, LT ³	14.9 (25.9) (4-15)	39.2 (44.3) (1--50)	33.7 (38.2) (10-40)

¹Mean age in years, standard deviation, and interquartile range

²Mean number of penile-vaginal sex partners in the three months preceding enrollment, standard deviation, and interquartile range

³Mean number of penile-vaginal sex partners over the lifetime (LT), standard deviation, and interquartile range

Table 2

Descriptive Information of the Sample, Shown by 3-Month Interval of Observation, Stratified by City of Enrollment

	Northeastern	Midwestern	Southern	Total
<u>Sexual Behaviors</u>				
Interval 1 ¹	(n = 203)	(n = 148)	(n = 217)	(n = 568)
Mean # sex partners, 3M ²	2.4 (2.4) (1-3)	3.9 (3.4) (2-5)	3.8 (3.5) (1-5)	3.3 (3.2) (1-4)
PVI events per person dy ³	3008/21491 (.14)	2293/13620 (.17)	3762/21735 (.17)	9063/56847 (.16)
Interval 2 ⁴	(n = 130)	(n = 115)	(n = 147)	(n = 392)
Mean # sex partners, 3M ²	2.5 (2.6) (1-3)	3.2 (3.5) (1-4)	3.0 (2.7) (1-4)	2.9 (3.0) (1-3)
PVI events per person dy ³	1893/10885 (.13)	1884/10441 (.18)	2260/13492 (.17)	6037/34788 (.17)
<u>Sexually Transmitted Infections</u>				
Baseline Measures	cases/n (%)	cases/n (%)	cases/n (%)	cases/n (%)
Combined prevalent cases	19/268 (7.1)	52/244 (21.3)	38/265 (14.3)	169/920 (18.4)
Prevalent gonorrhea cases	0/268 (0.0)	3/244 (1.2)	32/408 (7.8)	35/920 (3.8)
Prevalent chlamydia cases	14/268 (5.2)	13/244 (5.3)	49/408 (12.0)	76/920 (8.3)
Prevalent trichomonas cases	5/268 (1.9)	36/244 (14.8)	38/408 (9.3)	79/920 (8.6)
Interval 1 ¹				
Combined incident cases	9/236 (3.8)	17/198 (8.5)	38/265 (14.3)	64/699 (9.2)
Incident gonorrhea cases	1/236 (.4)	3/198 (1.5)	6/264 (2.2)	10/698 (1.4)
Incident chlamydia cases	5/236 (2.1)	37/198 (3.5)	21/265 (7.9)	33/699 (4.7)
Incident trichomonas case	s 3/236 (1.3)	10/198 (5.1)	19/264 (7.2)	32/698 (4.6)
Cases w/in CCC users ⁴	1/39 (2.6)	2/40 (5.0)	1/30 (3.3)	4/109 (3.7)
Cases w/in <CCC users ⁵	7/164 (4.3)	10/106 (9.4)	25/182 (13.7)	42/452 (9.3)
Interval 2 ⁴				
Combined incident cases	3/164 (1.8%)	14/162 (8.6%)	22/186 (11.8%)	39/512 (7.6)
Incident gonorrhea cases	0/163 (0.0)	1/162 (1.6)	4/186 (2.2)	5/512(1.0)
Incident chlamydia cases	3/164 (1.8)	3/162 (1.9)	5/186 (2.7)	11/512 (2.1)
Incident trichomonas case	s 0/164 (0.0)	11/162 (6.8)	14/186 (7.5)	25/512 (4.9)
Cases w/in CCC users ⁴	0/28 (0.0)	2/34 (5.9)	0/21 (0.0)	2/83 (2.4)
Cases w/in <CCC users ⁵	2/123 (1.6)	10/96 (10.4)	19/145 (13.1)	31/364 (8.5)

¹Data from first three months of the study

²Data from daily diaries collected over a three-month period

³Total number of penile-vaginal sex events divided by total number of observation days

⁴Data from months four through six of the study

⁵CCC = Consistent and Correct Condom

⁶<CCC = not reporting CCC

Table 3

Descriptive Information, by Gender, Regarding Sexual Behaviors Reported by Interval of Data Collection Using Daily Electronic Dairies

Behavior	Males	Females
<u>Interval 1</u> ¹		
Valid n, Mean PVI (SD) and (IQR) ²	231 15.6 (16.1) (4-22)	337 16.2 (16.1) (5-22)
Valid n, Mean PVI partners (SD) and (IQR) ³	231 4.0 (3.9) (1-5)	337 2.9 (2.5) (1-4)
Number (%) of inconsistent condom use ⁴	141/231 (61.0)	228/337 (67.7)
Number (%) of < CCC ⁵	180/231 (77.1)	279/337 (82.8)
Condom Errors/Problems (231 males, 337 females)		
Breakage: Mean (SD)	.4 (1.2)	.5 (1.0)
Slippage during withdrawal: Mean (SD)	.4 (1.3)	.6 (1.5)
Slippage during PVI: Mean (SD)	.3 (1.1)	.6 (1.4)
Incomplete ⁶ condom use: Mean (SD)	1.8 (3.3)	1.5 (3.0)
Not using a new condom: Mean (SD)	.5 (1.8)	.5 (3.1)
<u>Interval 2</u> ⁷		
Valid n, Mean PVI (SD) and (IQR) ²	154 13.6 (14.7) (3-19)	238 16.6 (17.9) (4-22)
Valid n, Mean PVI partners (SD) and (IQR) ³	154 3.1 (3.2) (1-4)	238 2.7 (2.8) (1-3)
Number (%) of inconsistent condom use ⁴	86/154 (55.8)	154/238 (64.7)
Number (%) of < CCC ⁵	124/154 (80.5)	208/238 (87.4)
Condom Errors/Problems (154 males, 238 females)		
Breakage: Mean (SD)	.4 (1.2)	.4 (1.3)
Slippage during withdrawal: Mean (SD)	.3 (1.1)	.5 (1.6)
Slippage during PVI: Mean (SD)	.3 (.8)	.5 (1.6)
Incomplete ⁶ condom use: Mean (SD)	1.3 (3.6)	1.4 (3.3)
Not using a new condom: Mean (SD)	.5 (2.8)	.7 (6.2)

¹ Assessed during the first three months of the study

² Mean frequency of penile-vaginal sex during three-month observation interval, with standard deviation and interquartile range

³ Mean number of penile-vaginal sex partners during three-month observation interval, with standard deviation and interquartile range

⁴ Defined as using condoms for less than 100% of all penile-vaginal sex acts during the interval

⁵ Defined as not using condoms consistently and correctly for all penile-vaginal sex acts during the interval

⁶ Defined as putting condoms on after sex had begun or taking condoms off before sex ended

⁷ Assessed during the second three months of the study

Table 4

Generalized Estimation Equation (GEE) Estimates for the Effect of Condom Use on Incident Infections with Chlamydia, Gonorrhea, and Trichomoniasis

Model 1: Effects of Consistent Condom Use Only

Predictor Variable	Point Estimate	95% CI ^a	P-value
Consistent condom use	.75	.42-1.32	.32
Male gender	.81	.44-1.50	.50
Less than 20 years of age ^b	2.00	.99-4.03	.05
20 to 24 years of age ^b	1.11	.57-2.17	.76
Past sexually transmitted infections	2.96	1.58-5.56	.0001

Model 2: Effects of Consistent and Correct Condom Use

Consistent and correct condom use	.41	.19-.90	.026
Male gender	.83	.45-1.53	.55
Less than 20 years of age ^b	2.05	1.01-4.14	.047
20 to 24 years of age ^b	1.12	.57-2.17	.75
Past sexually transmitted infections	2.98	1.61-5.52	.0005

^aConfidence interval

^bReference group is persons 25 years of age or older