

# The Drosophilidae (Diptera) of the Scattered Islands, with the description of a novel association with *Leptadenia madagascariensis* Decne. (Apocynaceae)

Amir Yassin,<sup>1,\*</sup> Nelly Gidaszewski,<sup>2</sup> Béatrice Albert,<sup>3</sup> Jean Hivert,<sup>4</sup> Jean R. David,<sup>2,5</sup> Virginie Orgogozo<sup>1</sup> and Vincent Debat<sup>2</sup>

<sup>1</sup>CNRS UMR7592; Institut Jacques Monod; Université Paris VII Diderot; Paris, France; <sup>2</sup>CNRS UMR7205; Origine; Structure et Evolution de la Biodiversité (OSEB); Muséum National d'Histoire Naturelle; Paris, France; <sup>3</sup>UMR8079; Laboratoire Ecologie, Systématique et Evolution; Univ. Paris-Sud; Orsay, France; <sup>4</sup>Laboratoire Conservatoire Botanique National de Mascarin & Centre Permanent d'Initiatives pour l'Environnement; La Réunion, France; <sup>5</sup>CNRS UPR9034; Laboratoire Evolution, Génomes et Spéciation (LEGS); Gif-sur-Yvette Cedex, France

**Keywords:** taxonomy, DNA barcoding, island biogeography, myophily, conservation

Thirteen drosophilid species belonging to seven genera and two subfamilies are reported from three coral islands (namely Europa, Juan de Nova and Glorioso) that belong to the Scattered Islands in the Indian Ocean. Five species are cosmopolitan and five are African. Three are endemic to the insular Western Indian Ocean, including a presumably new *Scaptodrosophila* species. On the island of Juan de Nova, most captured flies had pollinia attached to the bases of their proboscis. DNA analysis using the *rbcl* gene revealed that these pollinia belong to the genus *Leptadenia* (Apocynaceae), of which a single species *L. madagascariensis*, endemic in Madagascar and Comoros, is present in this island. This is the first reported association between this plant and drosophilids.

## Introduction

The islands of the Western Indian Ocean harbor a distinctive drosophilid fauna that is mainly dominated by African species.<sup>1-3</sup> Among 158 species,<sup>3</sup> 25 are endemic to Madagascar, 12 to Seychelles, 31 to Mauritius and 2 to La Réunion. However, the fauna of the islands encircling Madagascar that are called Scattered Islands (Îles éparses) by the French administration have not been investigated. The Scattered Islands consist of four small coral islands (Europa, 30 km<sup>2</sup>; Juan de Nova, 5 km<sup>2</sup>; Glorioso islands, 7 km<sup>2</sup>; Tromelin, 1 km<sup>2</sup>) and an atoll (Bassas da India). All are located in the Mozambique Channel, except Tromelin which is located 470 km east of Madagascar and 560 km west of Mauritius and La Réunion (Fig. 1). The maximum altitude of these islands ranges from 7 to 14 m. Bassas da India is an atoll that becomes totally submerged by water everyday at high tides. Each of the remaining four islands is presently inhabited by a small military camp only.<sup>4</sup>

Previous entomological surveys were conducted on all islands except Juan de Nova and Bassas da India. The presence of 154 insect species, of which 24 were endemic, has been reported.<sup>5-8</sup> However, no drosophilid species has ever been recorded from these islands. In April 2011, a CNRS-funded scientific expedition was joined by four of us (JRD, NG, JH and AY). One of us (JH), with Benoît Dumeau, spent 52 d on Europa from October


to December 2011. We report here the presence of 13 drosophilid species including a probably new *Scaptodrosophila* species and we describe a new association between drosophilids and the flowering plant *Leptadenia madagascariensis* Decne. on the island of Juan de Nova.

## Results and Discussion

**Taxonomy and biodiversity.** Table 1 shows the list of drosophilid species collected and their geographical distribution. In total, 13 species belonging to seven genera and two subfamilies were collected. Of which, 10 species are also found on the African mainland and five outside Africa. Three species are endemic to the insular Western Indian Ocean.

Species diversity differed greatly among islands. No drosophilid was collected on the island of Tromelin as would be expected from the dry and stormy climate of this small island. In Europa and in spite of its floral richness and semi-arid climate, we could not find any drosophilid during our first visit in April 2011, which lasted three days. However, a later 52-d expedition in October–December 2011 yielded two species using fermented fruit traps: many specimens of the steganine *Cacoxenus polydorus* whose larvae prey on mealy bugs,<sup>9</sup> and a single female of *Drosophila melanogaster*. *Cacoxenus* adults are usually only accidentally attracted to fruit baits, and the trap was located in a sisal (Agavaceae)

\*Correspondence to: Amir Yassin; Email: yassin@ijm.univ-paris-diderot.fr  
Submitted: 06/27/12; Revised: 07/20/12; Accepted: 07/23/12  
<http://dx.doi.org/10.4161/fly.21853>



**Figure 1.** Maps of the Western Indian Ocean showing the geographical position of the Scattered Islands (underlined) and the trajectory of the Marion Duffresne ship (dashed line). Gray areas indicate locations where the traps were placed. The trap where drosophilids were collected in Europa is indicated with a dot.

field infested by a large colony of undetermined mealy bugs. *D. melanogaster* is a cosmopolitan, human commensal species that is usually found near human settlements and easily captured using fruit traps. However, the presence of only a single specimen in December near the military camp and its absence in April suggest that *D. melanogaster* may not be well established on Europa.

The greatest species diversity is encountered on the islands of Juan de Nova and Glorioso. The 11 species collected on these islands can be classified into three categories: four are cosmopolitan (i.e., found on at least three continents including the Americas) and generalists: *Drosophila malerkotliana*, *Drosophila simulans*, *Scaptodrosophila latifasciaeformis* and *Zaprionus indianus*; four are pan-African: the fig-breeder *Lissocephala sanu* (bred from *Ficus grevei* Baill. on Glorioso), the flower-breeder *Scaptodrosophila caliginosa*, the palm-breeder *Scaptodrosophila*

*bangi*, and the generalist *Zaprionus tuberculatus*; and three are endemic to the insular Indian Ocean: *Chymomyza bicolor*, *Scaptodrosophila* sp and *Leucophenga malgachensis* (which may also be present in South Africa<sup>10</sup>).

**Association with *Leptadenia*.** An unexpected phenomenon was observed on flies collected using fermented fruit traps on the island of Juan de Nova. Ninety percent of these flies, belonging to six species (Table 2) harbored large yellow bodies at the lower margin of the head capsule and the proboscis (Fig. 2A). Each yellow body consisted of a pair of cellular aggregates connected and attached to the fly body by a transparent peduncle (Fig. 2B). Microscopic observation suggested that these bodies might be pollinia. In order to determine the origin of these bodies, their DNA was extracted and the chloroplastic gene, *ribulose-1,5-bisphosphate carboxylase/oxygenase large subunit (rbcL)*, which is used in plant DNA barcoding,<sup>11</sup> was amplified. PCR amplicons of the correct size were obtained, and a GenBank similarity search using BLAST retrieved one sequence with 100% sequence identity, from the species *Leptadenia reticulata* (Retz.) Wight and Arn. (Apocynaceae) (accession number EU196282). However, *L. reticulata* is endemic to India and likely absent in the insular Indian Ocean.<sup>12</sup> In Juan de Nova, the only representative of the genus *Leptadenia* is *L. madagascariensis* (Fig. 2C; V. Boulet, J. Hivert and M. Lacoste, in lit.; not mentioned by Perrier de la Bâthie in 1921<sup>13</sup> but cited by Bosser in 1952<sup>14</sup>). Pollinia of *L. madagascariensis* specimens collected at the same time as the flies (Fig. 2D) were exactly similar to the yellow bodies present on the fly probosces. We also sequenced *rbcL* from conserved flowers of a *L. madagascariensis* specimen, and found 100% sequence identity between *rbcL* from fly-attached pollinia and from *L. reticulata*. A high similarity in *rbcL* sequences is frequent in plant molecular taxonomy studies.<sup>11</sup> This example shows the utility of DNA barcoding in identifying plants at the genus level by non-specialist, but also rise the concerns of its limitations at the species level.

Many drosophilid species are known to breed in flowers,<sup>15-17</sup> but plant pollination by flies, so-called myophily, was previously observed for only a few drosophilid species.<sup>18</sup> Apocynaceae plants are usually myophilic and known to emit odors that mimic fruit substrates.<sup>19</sup> Flies are usually attracted by the flower odor, but once on the flower they do not oviposit. According to the ASCLEPOL database ([http://www.bio.uni-bayreuth.de/planta2/research/pollina/as\\_pol\\_d.htm](http://www.bio.uni-bayreuth.de/planta2/research/pollina/as_pol_d.htm)), some drosophilid species were observed visiting different Apocynaceae hosts and carrying their pollinia on their probosces, but never from a species belonging to the genus *Leptadenia*. In South Africa, Agnew<sup>20</sup> described in detail the association between several drosophilid species (namely *D. melanogaster*, *D. simulans*, *D. immigrans* and *Zaprionus vittiger*) and *Orbea schweinfurthii* (A. Berger) Bruyns (Apocynaceae). Bhatnagar<sup>21</sup> recorded *D. seguyi* on *Cosmostigma cordatum* (Poir.) Almeida in India, and Yamashiro et al.<sup>22</sup> found *D. melanogaster* on *Cynanchum*

**Table 1.** Geographical distribution and number of drosophilid species collected in the Sparse Islands

Geographical region	EUR	JDN	GLO	TRO	COM	MAD	SEY	MAU	REU	AFR	COS	
Date of collection	Apr. 2011	Nov. 2011	Apr. 2011	Apr. 2011	Apr. 2011							
<b>Subfamily Drosophilinae</b>												
Genus <i>Chymomyza</i>	0	0	0	1	0		•					
Group <i>procnemis</i>												
<i>Chymomyza bicolor</i>												
Genus <i>Drosophila</i>												
Subgenus <i>Sophophora</i>	0	0	10	379	0		•	•		•	•	
Group <i>ananassae</i>												
<i>Drosophila malerkotliana</i>												
Group <i>melanogaster</i>	0	1	0	0	0	•	•	•	•	•	•	
<i>Drosophila melanogaster</i>												
<i>Drosophila simulans</i>	0	0	4	0	0	•	•	•		•	•	
Genus <i>Lissocephala</i>												
Group <i>sanu</i>	0	0	0	12	0				•	•		
<i>Lissocephala sanu</i>												
Genus <i>Scaptodrosophila</i>												
Group <i>aterrima</i>	0	0	1	0	0		•	•	•		•	
<i>Scaptodrosophila caliginosa</i>												
Group <i>brunnea</i>	0	0	10	20	0							
<i>Scaptodrosophila sp</i>												
Group <i>latifasciaeformis</i>	0	0	18	55	0		•	•	•	•	•	
<i>Scaptodrosophila latifasciaeformis</i>												
Group <i>saba</i>	0	0	0	3	0	•		•	•	•		
<i>Scaptodrosophila bangi</i>												
Genus <i>Zaprionus</i>												
Group <i>inermis</i>	0	0	0	21	0	•	•	•	•	•		
<i>Zaprionus tuberculatus</i>												
Group <i>vittiger</i>	0	0	50	109	0	•	•	•	•	•	•	
<i>Zaprionus indianus</i>												
<b>Subfamily Steganinae</b>												
Genus <i>Cacoxenus</i>	0	5	0	0	0			•	•	•		
Group <i>perspicax</i>												
<i>Cacoxenus polyodous</i>												
Genus <i>Leucophenga</i>												
Group <i>mutabilis</i>	0	0	0	2	0		•			?		
<i>Leucophenga malgachensis</i>												
Number of species	0	2	6	9	0	5	7	9	8	8	10	6


Dots indicate species that have been reported in other Indian Ocean islands or in mainland Africa, as well as cosmopolitan species.<sup>3</sup> Question mark indicates probable presence. EUR, Europa; JDN, Juan de Nova; GLO, Glorioso; TRO, Tromelin; COM, Comoros; MAD, Madagascar; SEY, Seychelles; MAU, Mauritius; REU, La Réunion; AFR, mainland Africa; COS, cosmopolitan.

*japonicum* (Morren and Decaisne) Hemsl. and *D. immigrans* and *Scaptodrosophila coracina* on *V. katoi* (Ohwi) Kitag. in Japan.

It is important to note that *L. madagascariensis* is also found in different localities in Madagascar and Comoros and that most of the drosophilid species observed to carry its pollinia in Juan de Nova also exist in these places. However, no pollinia have ever been reported on drosophilid probosces in these locations and we did not find any pollinia on drosophilid flies during our previous field trips in Madagascar. Two hypotheses may explain the strong association with *L. madagascariensis* on the island of Juan de Nova: either the plants on this island belong to a certain variety or subspecies that have evolved attractive smells specifically

targeting drosophilids, or that at the time of collection (the beginning of the dry season in Juan de Nova<sup>4</sup>), only *L. madagascariensis* flowers presented an appropriate sugar resource for drosophilid flies. Future seasonal surveys of drosophilids on Juan de Nova and genetic and biochemical comparisons of *L. madagascariensis* in different islands of the Western Indian Ocean will certainly help to elucidate the nature of its association with drosophilids.

A last interesting problem is the role of myophily for the reproduction of *Leptadenia*. Although this problem should deserve further studies, we would like to point out that *Leptadenia* fructification success is very low: a single plant usually produces very many flowers but very few fruits (our observations). Allogamy



**Figure 2.** Association between drosophilids and *Leptadenia madagascariensis* on the island of Juan de Nova. (A) A *Scaptodrosophila latifasciaeformis* female carrying several yellow pollinia on its proboscis. (B) Microscopic structure of a pollinium (pl) and three peduncles (pd) attached to a fly proboscis. (C) *L. madagascariensis* flowers on a shrub. (D) pollinia in a *L. madagascariensis* flower.

**Table 2.** Species list and proportion of flies collected on Juan de Nova bearing yellow pollinia on their probosces

Species	No. of flies	Proportion of flies with pollinia	No. of pollinia per fly
<i>Zaprionus indianus</i>	50	> 90%	1–6
<i>Drosophila malerkotiana</i>	10	30%	1–2
<i>Scaptodrosophila</i> sp	10	10%	1
<i>Scaptodrosophila latifasciaeformis</i>	18	6%	1
<i>Drosophila simulans</i>	4	0%	0
<i>Scaptodrosophila caliginosa</i>	1	0%	0

and inefficient pollinia transportation by flies may explain the low fructification rate in *Leptadenia*.

### Materials and Methods

Flies were collected either using fermenting banana traps hanged in shadowed shrubs or by net sweeping on moist litter. To make sure that there is no accidental importation of drosophilids, unripe bananas were bought in La Réunion and conserved in a refrigerator on the boat. Banana traps were prepared several days in advance and left fermenting in a closed box. We then made sure that there were no flies in the traps when we set them up in the field. We checked for flies in the traps for the next three days. Sweeping with a net over rotten fruits and litter was also used. Identification was made by AY and JRD according to Lamb,<sup>23</sup>

Tsacas et al.,<sup>24</sup> Tsacas and Chassagnard,<sup>9,25</sup> Bächli et al.,<sup>10</sup> Cariou et al.<sup>2</sup> and Yassin and David.<sup>26</sup> Flies were kept alive on standard laboratory medium or killed in absolute alcohol for subsequent analyses.

Yellow bodies carried by flies collected on the island of Juan de Nova were investigated and photographed using a Leica binocular stereoscope. They were carefully separated from fly heads and preserved in absolute ethanol. DNA was extracted using DNeasy Blood and Tissue Kit (QIAGEN) and universal primers (*rbcLa\_f* and *rbcLa\_rev*) of the chloroplast gene *rbcL*<sup>1</sup> were used. PCR amplification conditions were: 5 min at 95°C, followed by 5 cycles of 30 sec at 94°C, 1 min at 55°C and 1 min at 72°C, then 30 cycles of 30 sec at 94°C, 1 min at 54°C and 1 min at 72°C, followed by 10 min at 72°C. Amplicons were sequenced by Beckman-Coulter Genomics and sequence similarity was searched in GenBank using BLAST ([blast.ncbi.nlm.nih.gov/Blast.cgi](http://blast.ncbi.nlm.nih.gov/Blast.cgi)).

### Disclosure of Potential Conflicts of Interest

No potential conflicts of interest were disclosed.

### Acknowledgments

The Marion Dufresne 2011 expedition was funded by the CNRS-INEE 'Iles Eparses' project to VD. Morphological and molecular taxonomy was funded by an ATIP-AVENIR to VO and IDEV grant to BA. We are also grateful to the Terres Australes et Antarctiques Françaises (TAAF) for the logistic organization of the expedition. Amélie Rast provided valuable assistance in the lab.



## References

- Lachaise D, Harry M, Solignac M. Affinités biogéographiques des Drosophilidae de Madagascar et des îles de l'Océan Indien. In: Lourenco WR (ed), *Biogéographie de Madagascar*, 1996, pp. 467-78, Orstom, Paris.
- Cariou ML, Lachaise D, Matyot P, Gerlach J, Montchamp C, Legrand D, et al. Family Drosophilidae. *Faunistica* 2009; 85:355-85.
- Yassin A, David JR. Phylogenetic biogeography of Afrotropical Drosophilidae. In: Gailis M, Kalniš (eds), *Biogeography*, 2010, pp. 105-136, Nova Science Publishers, New York.
- Caceres S. Etude préalable pour le classement en réserve naturelle des Iles Eparses. Mémoire de DESS Sciences et Gestion de l'Environnement Tropical de l'Université de la Réunion 2003; [http://etec.univ-reunion.fr/get/documents/Iles%20Eparses/IlesEparses\\_SarahCACERES\\_2002.pdf](http://etec.univ-reunion.fr/get/documents/Iles%20Eparses/IlesEparses_SarahCACERES_2002.pdf)
- Paulian R. Observations sur la faune terrestre de l'île Tromelin. *Le Naturaliste Malgache* 1955; 7:1-7.
- Paulian R. Les Insectes des Iles Glorieuses. *Entomologiste* 1989; 45:203-8.
- Vieitte P. L'entomofaune de l'île Europa. *Editions du Muséum* 1966; 91:191-210.
- Staub F. Geography and ecology of Tromelin Island. *Atoll Res Bull* 1970; 136:197-209.
- Tsacas L, Chassagnard MT. Les espèces afrotropicales du sous-genre *Gitonides* Knab du genre *Cacoxenus* Loew à larves prédatrices de cochenilles (Diptera: Drosophilidae). *Ann Soc Entomol Fr* 1999; 35:91-121.
- Bächli G, Vilela CR, McEvey SF. Nine new species of Afrotropical *Leucophenga* (Diptera: Drosophilidae). *Mitt Schweiz Entomol Ges* 2005; 78:23-57.
- Kress WJ, Erickson DL. A two-locus global DNA barcode for land plants: the coding rbcL gene complements the non-coding trnH-psbA spacer region. *PLoS One* 2007; 2:e508; PMID:17551588; <http://dx.doi.org/10.1371/journal.pone.0000508>.
- Panwar J, Tarafdar JC. Distribution of three endangered medicinal plant species and their colonization with arbuscular mycorrhizal fungi. *J Arid Environ* 2006; 65:337-50; <http://dx.doi.org/10.1016/j.jaridenv.2005.07.008>.
- Perrier de la Bâthie H. Note sur la constitution géologique et la flore des îles Chesterfield, Juan de Nova, Europa et Nosy-Trozona. *Bulletin Economique de Madagascar* 1921; 170-176.
- Bosser J. Note sur la végétation des îles Europa et Juan de Nova. *Le Naturaliste Malgache* 1952; 4:41-2.
- McEvey SF, Aulard S, Ralisoa-Randrianasolo O. An Australian drosophilid (Diptera) on *Eucalyptus* and *Eugenia* (Myrtaceae) flowers in Madagascar. *J Aust Entomol Soc* 1989; 28:53-4; <http://dx.doi.org/10.1111/j.1440-6055.1989.tb01193.x>.
- Markow TA, O'Grady PM. Reproductive ecology of *Drosophila*. *Funct Ecol* 2008; 22:747-59; <http://dx.doi.org/10.1111/j.1365-2435.2008.01457.x>.
- David JR, Yassin A, Rasamizafi LA, Raveloson Ravaomanarivo LH, Debat V. Scratching for food: An original feeding behavior in an African flower breeding *Drosophila*. *Fly (Austin)* 2011; 5:285-90; PMID:22041577.
- Sultana F, Hu YG, Toda MJ, Takenaka K, Yafuso M. Phylogeny and classification of *Colocasiomyia* (Diptera, Drosophilidae), and its evolution of pollination mutualism with aroid plants. *Syst Entomol* 2006; 31:684-702; <http://dx.doi.org/10.1111/j.1365-3113.2006.00344.x>.
- Ollerton J, Masinde S, Meve U, Picker M, Whittington A. Fly pollination in *Ceropegia* (Apocynaceae: Asclepiadoideae): biogeographic and phylogenetic perspectives. *Ann Bot* 2009; 103:1501-14; PMID:19339298; <http://dx.doi.org/10.1093/aob/mcp072>.
- Agnew JD. A case of myophily involving Drosophilidae (Diptera). *Journal of South African Botany* 1976; 42:85-95.
- Bhatnagar S. On insect adaptations for pollination in some asclepiads on Central India. In Kapil RP (ed) *Pollination Biology – an Analysis* 1986, pp. 37-57, Inter-India Publs, New Delhi.
- Yamashiro T, Yamashiro A, Yokoyama J, Maki M. Morphological aspects and phylogenetic analyses of pollination systems in the *Tylophora-Vincetoxicum* complex (Apocynaceae-Asclepiadoideae) in Japan. *Biol J Linn Soc Lond* 2008; 93:325-41; <http://dx.doi.org/10.1111/j.1095-8312.2007.00896.x>.
- Lamb CG. The Percy Sladen Trust expedition to the Indian Ocean in 1905. XV. Diptera: Heteroneuridae, Ortalidae, Trypetidae, Sepsidae, Micropezidae, Drosophilidae, Geomyzidae, Milichidae. *Transactions of the Linnean Society of London* 1914; 16:307-72; <http://dx.doi.org/10.1111/j.1096-3642.1913.tb00152.x>.
- Tsacas L, Chassagnard MT, David JR. Un nouveau groupe d'espèces afrotropicales anthophiles dans le sous-genre *Scaptodrosophila* du genre *Drosophila* (Diptera: Drosophilidae). *Ann Soc Entomol Fr* 1988; 24:181-202.
- Tsacas L, Chassagnard MT. Identité de *Drosophila brunnea* de Meijere et description de nouvelles espèces orientales et africaines à pointe du scutellum blanche (Diptera, Drosophilidae). *Bulletin Zoologisch Museum Universiteit van Amsterdam* 1976; 5:89-105.
- Yassin A, David JR. Revision of the afrotropical species of *Zaprionus* (Diptera, Drosophilidae), with descriptions of two new species and notes on internal reproductive structures and immature stages. *ZooKeys* 2010; 51:33-72; PMID:21594121; <http://dx.doi.org/10.3897/zookeys.51.380>.