



RESEARCH

Open Access

DNA Methylation status of Wnt antagonist *SFRP5* can predict the response to the EGFR-tyrosine kinase inhibitor therapy in non-small cell lung cancer

Jian Zhu^{1†}, Yuyan Wang^{1†}, Jianchun Duan¹, Hua Bai¹, Zhijie Wang¹, Lai Wei², Jun Zhao¹, Minglei Zhuo¹, Shuhang Wang¹, Lu Yang¹, Tongtong An¹, Meina Wu¹ and Jie Wang^{1*}

Abstract

Background: It is well known that genetic alternation of epidermal growth factor receptor (*EGFR*) plays critical roles in tumorigenesis of lung cancer and can predict outcome of non-small-cell lung cancer treatment, especially the EGFR tyrosine-kinase inhibitors (EGFR-TKIs) therapy. However, it is unclear whether epigenetic changes such as DNA methylation involve in the response to the EGFR-TKI therapy.

Methods: Tumor samples from 155 patients with stages IIIB to IV NSCLC who received EGFR-TKI therapy were analyzed for DNA methylation status of Wnt antagonist genes, including *SFRP1*, *SFRP2*, *SFRP5*, *DKK3*, *WIF1*, and *APC*, using methylation specific PCR (MSP) method. EGFR mutations detections were performed in the same tissues samples using Denaturing High Performance Liquid Chromatography (DHPLC).

Results: We found that Wnt antagonists tend to methylate simultaneously. Methylation of *sFRP1* and *sFRP5* are reversely correlated with EGFR mutation ($P = 0.005$, $P = 0.011$). However, no correlations of methylations of other Wnt antagonist genes with EGFR mutation were found. The patients with methylated *SFRP5* have a significant shorter progression free survival than those with unmethylated *SFRP5* in response to EGFR-TKI treatment ($P = 0.002$), which is independent of *EGFR* genotype.

Conclusions: Patients with unmethylated *SFRP5* are more likely to benefit from EGFR-TKI therapy.

Keywords: DNA methylation, EGFR-TKI, Wnt antagonists, Non-small cell lung cancer

Background

Lung cancer is the leading cause of cancer death worldwide [1]. NSCLC is the most common form of lung cancer, accounting for approximately 85% of lung cancer cases [2,3]. The efficacy of traditional chemotherapy has reached a plateau [4-6]. Therefore, new approaches are needed to improve the efficacy of lung cancer therapy. A number of targeted anticancer agents have been recently developed and approved for clinical use, among which

the EGFR-TKI has been used as the first-line therapy for lung cancer patients with EGFR mutations [7-11].

EGFR gene product functions as a receptor tyrosine kinase that affects cell proliferation and survival by activating downstream signaling pathways. In 2004, three research groups reported that mutations in the tyrosine kinase domain of *EGFR* can predict the responses to TKIs in NSCLC patients [12-14], which enables the identification of patient populations that are more likely to benefit from TKI therapies and serves as the first step toward personalizing lung cancer therapy. However, according to the theory of "EGFR addition", which refers to the dependency of cancer cells on *EGFR* mutation to maintain their malignant phenotypes [15], lung cancer patients harboring mutations in the tyrosine kinase

* Correspondence: wangjie_cc@yahoo.com

†Equal contributors

¹Department of Thoracic Medical Oncology Peking University School of Oncology, Beijing Cancer Hospital & Institute, Beijing 100036, China
Full list of author information is available at the end of the article

domain of their *EGFR* genes should survive much longer, in response to the EGFR-TKI therapy, than the actual result. This suggested that *EGFR* mutation cannot explain all clinical outcomes of TKI therapy. At least 10~20% of patients with wild-type *EGFR* still significantly benefit from EGFR-TKI treatment, whereas around 10% of patients with mutated *EGFR* are resistant to the TKI therapy [10,16,17]. In addition, previous studies reported that both T790M mutation [18] and c-MET amplification [19] involved in acquired resistance of EGFR-TKI therapy. Therefore, factors in addition to *EGFR* genotype may also contribute to the response to EGFR-TKI therapy.

The Wingless-type (Wnt) signaling cascade is an important regulator of embryonic development [20]. Activation of Wnt signaling pathway leads to elevated expression of β -catenin in cytoplasm, which in turn translocates to the nucleus, interacts with T cell factor/lymphocyte enhancer factor family, induces, downstream target genes that regulate cell proliferation and cancer progression. Aberrant activation of Wnt signaling pathway has been found in a number of tumors [21], which can be categorized into the following three common forms: 1) mutations in *APC* and/or *Axin*; 2) aberrant activation of Wnt signaling induced by activated *EGFR* [22]; 3) methylation of Wnt antagonists. Mutations of *APC* and/or *Axin* are rarely found in lung cancer patients. In addition, EGFR-TKI treatment blocks activation of EGFR in patients. Therefore, we hypothesized that the methylation of Wnt antagonists might significantly affect the responses to the EGFR-TKI therapy in NSCLC patients. Suzuki et al [23] analyzed the synchronous effects and correlations between Wnt antagonists and EGFR mutations and found that EGFR mutation was correlated with a good prognosis in tumors without methylated wnt antagonist genes.

In current study, we analyzed the methylation status of the CpG sites within Wnt antagonist genes, including *SFRP1*, *SFRP2*, *SFRP5*, *WIF1*, *DKK3*, *APC*, and *CDH1*, in 155 Chinese patients who received EGFR-TKI therapy and investigated potential clinical implication of the epigenetic regulation of Wnt antagonists.

Methods

Patients

155 patients were enrolled in current study. They were pathologically diagnosed as stage IIIB or IV NSCLC, with Eastern Cooperative Oncology Group performance (ECOG) status of 0 to 2; and received EGFR-TKI as either first- or second-line therapy at the Peking University Cancer Hospital between June 2006 and December 2009. The study was reviewed and approved by the Institutional Review Board at the Beijing Cancer Hospital. Written informed consent was obtained from all participants.

The smoking status of patients was decided during their first visit. A smoker was defined as the one who smoked more than 100 cigarettes in his/her life time. Patients were treated with either TKI therapy or platinum-based chemotherapy as the first line of treatment until their disease progressed, justified by imaging evidence or aggravated symptoms. The Response Evaluation Criteria in Solid Tumors (RECIST) [24] including progressive disease (PD), stable disease (SD), partial remission (PR) and complete remission (CR) was used to evaluate the drug response after patients received treatment every 6 weeks to 2 months. The objective response rate (ORR) was defined as the sum of PR and CR, while the disease control rate (DCR) was defined as the sum of SD, PR, and CR. Progression-free survival (PFS) was assessed from the beginning of therapy to disease progress or death from any cause. Overall survival (OS) was assessed from the beginning of first-line therapy until death from any cause.

DNA extraction and methylation-specific PCR

Genomic DNA of tumor tissues from patients biopsied before TKI treatment were extracted using QIAamp FFPE DNA kit (Qiagen). The methylation status of the CpG sites within the gene loci of *SFRP1*, *SFRP2*, *SFRP5*, *WIF1*, *DKK3*, *APC*, and *CDH1* was decided by MSP assays as described previously [25-27]. Briefly, genomic DNA was treated with sodium bisulfite, followed by PCR amplifications using the primer pairs that can specific detect either the methylated or the unmethylated CpG sites. Genes were defined as methylated if the PCR products could be detected using the methylated DNA-specific primer pairs, while they were defined as unmethylated if the PCR products could only be detected using the unmethylated DNA-specific primer pairs. DNA from the human adenocarcinomic alveolar basal epithelial cell lines, A549 and A549/DDP, was used as the positive control for methylated DNA, while DNA from lymphocytes of healthy nonsmoking volunteers was used as the negative control. The methylation status results were confirmed by at least one repeat of the methylation-specific PCR assays. The following primers were used:

sFRP1: Methylated F: 5'-GTTTTTCGGAGT TAGTGTGCGCGC-3'; R: 5'-ACGATCGAAAACGACGC GAACG-3'; Unmethylated F: 5'-GTAGTTTTTGGAGT TAGTGTTGTGT-3'; R: 5'-ACCTACAATCAAAAACAA CACAAACA-3'; *sFRP2*: Methylated F: 5'-TCGGAGTTTTTCGGAGTTGCGC-3'; R: 5'-GCTCTCTTCGCTAAATACGACTCG-3'; Unmethylated F: 5'-GGTTGGAGTTTTTGGAGTTGTGT-3'; R: 5'-CCCCTCTCTTCACTAAATACTCA-3'; *sFRP5*: Methylated F: 5'-TGGCGTTGGGCGGGACGTTTC-3'; R: 5'-

AACCCGAACCTCGCCGTACG-3;UnmethylatedF:5'-TGGTGTGGGTGGGATGTTTG-3;R:5'-CAACC
 CAAACCTCACCATACAC-3';**DKK3**MethylatedF:5'-GGGGCGGGCGCGGGGC-3;R:5'-ACATCTCCGCTC
 TACGCCG-3;UnmethylatedF:5'-TTAGGGTGGGTGGTGGGGT-3;R:5'-CTACATCTC
 CACTCTACACCCA-3';**WIF-1**MethylatedF:5'-CGTTTTATTGGGCGTATCGT-3;R:5'-ACTAACGC
 GAACGAAATACGA-3;UnmethylatedF:5'-GGGTGTTTTATTGGGTGTATTGT-3;R:5'-AAAAAC
 TAACACAAACAAAATACAAAC-3';**APC**MethylatedF:5'-TATTGCGGAGTGC GGTC-3;R:5'-TCGAC-
 GAACTCCCGACGA-3;UnmethylatedF:5'-GTGTTTTATTGTGGAGTGTGGGTT-3;R:5'-CCAAT
 CAACAACTCCCAACAA-3';**CDH-1**MethylatedF:5'-TG TAGTTACGTATTTATTTTATGTCGTC-3;R:5'-
 CGAATACGATCGAATCGAACCG-3;UnmethylatedF:5'-TG GTTGTAGTTATGTATTTGTTTTAGTGG-3;R:5'-
 ACACCAAATACAATCAAATCAAACAAA-3'.

Mutation detection

The denaturing high-performance liquid chromatography (DHPLC) was used to detect mutations in the exon 19 and 21 of EGFR tyrosine kinase domains as described previously [28].

Statistical analysis

All data were analyzed using SPSS (version 16.0). Chi-square and Fisher's exact tests were used to assess the association between DNA methylation and EGFR genotypes. Multivariate analysis was performed using Cox proportional hazard regression model. The Kaplan-Meier method was used to determine the overall survival and

progression-free survival curves. P value less than 0.05 was considered statistically significant.

Results

Characteristics of study patients

Table 1 summarized the demographic characteristics of 155 study patients, among which 118 cases were adenocarcinoma and 37 cases were non-adenocarcinoma (29 squamous carcinoma, 5 large cell carcinoma, and 3 adeno-squamous carcinoma cases). 60 of all patients received EGFR-TKI as the first-line therapy, while the rest had EGFR-TKI as the second- or more-line treatment. Among those 95 patients who had EGFR-TKI as the second- or more-line treatment, 63 patients took platinum-based chemotherapy as the first-line treatment. The median follow-up time for all patients was 22.4 months (from 2.4 to 77.2 months).

Epigenotype of Wnt antagonists in NSCLC

Genomic DNA was extracted from tumor tissues of all patients as described in the Method Section. The methylation status of Wnt antagonist genes including *SFRP1*, *SFRP2*, *SFRP5*, *WIF1*, *DKK3*, *APC*, and *CDH1*, defined as their epigenotype, was detected by Methylation Specific PCR Assays (examples were shown in Additional file 1: Figure S1A). The frequency of methylation events in Wnt antagonist genes in patients with different demographic characteristics was listed in Table 1. Interestingly, no significant difference in epigenotype of Wnt antagonist genes was found between male and female, among different age groups, between smokers and non-smokers, or between adenocarcinoma and non-adenocarcinoma cases.

Using DHPLC, we also detected EGFR activating mutations in exon 19 or 21 (the examples of wild type, mutated

Table 1 Methylation and mutation profile of NSCLC

Clinical characteristics (cases)	Methylation (%)							EGFR mutation (%)	
	SFRP1	SFRP2	SFRP5	DKK3	WIF1	APC	CDH1	Any gene	
<i>Gender</i>									
Male (74)	30 (40.5)	20 (27.0)	9 (12.2)	9 (12.2)	3 (4.1)	13 (17.6)	7 (9.5)	44 (59.5)	36 (48.6)
Female (81)	31 (38.3)	20 (24.7)	14 (17.3)	13 (16.0)	3 (3.7)	18 (22.2)	8 (9.9)	48 (59.3)	49 (60.5)
<i>Age</i>									
<65 (89)	33 (37.1)	21 (23.6)	10 (11.2)	12 (13.5)	3 (3.4)	16 (18.0)	7 (7.9)	48 (53.9)	56 (62.9)*
≥65 (66)	28 (42.4)	19 (28.8)	13 (19.7)	10 (15.2)	3 (4.5)	15 (22.7)	8 (12.1)	44 (66.7)	29 (43.9)
<i>Smoking</i>									
Never (93)	35 (37.6)	24 (25.8)	14 (15.1)	15 (16.1)	2 (2.2)	21 (22.6)	8 (8.6)	58 (62.4)	57 (61.3)*
Smokers (62)	26 (41.9)	16 (25.8)	9 (14.5)	7 (11.3)	4 (6.5)	10 (16.1)	7 (11.3)	34 (54.8)	28 (45.2)
<i>Histology</i>									
Adenocarcinoma (118)	46 (38.9)	30 (25.4)	16 (13.6)	16 (13.6)	4 (3.4)	21 (17.8)	14 (11.9)	72 (61.0)	65 (55.1)
Non-adenocarcinoma (37)	15 (40.5)	10 (27.0)	7 (18.9)	6 (16.2)	2 (5.4)	7 (18.9)	1 (2.7)	20 (54.1)	20 (54.1)
Total	61 (39.4)	40 (25.8)	23 (14.8)	22 (14.2)	6 (38.7)	31 (20%)	15 (9.7%)	92 (59.4%)	85 (54.8%)

*The frequency of this group is significantly higher than their counterparts.

Table 2 P value among methylated genes and EGFR mutation

	sFRP1	sFRP2	sFRP5	DKK3	WIF-1	APC	CDH-1	EGFR mutation
sFRP1	NA	0.004	0.005	0.008	0.02	<0.0001	0.266	0.005
sFRP2	0.004	NA	<0.0001	<0.0001	0.007	<0.0001	<0.0001	0.854
sFRP5	0.005	<0.0001	NA	<0.0001	<0.0001	0.06	<0.0001	0.011
DKK3	0.008	<0.0001	<0.0001	NA	0.0001	0.006	<0.0001	0.489
WIF-1	0.02	0.007	<0.0001	<0.0001	NA	0.03	0.02	0.094
APC	<0.0001	<0.0001	0.06	0.006	0.03	NA	0.126	0.546
CDH-1	0.266	<0.0001	<0.0001	<0.0001	0.02	0.126	NA	0.592
EGFR	0.005	0.854	0.011	0.489	0.094	0.546	0.592	NA

exon 19, and mutated exon 21 were shown in Additional file 1: Figure S1B, 1C, and 1D). Among the 155 patients, 85 (55.4%) carried mutations in either exon 19 or 21 of the EGFR genes (Table 1). Similar to the previous studies, we found that EGFR mutation rates were significantly increased among the patients younger than 65 years old ($P = 0.02$, Fisher's exact test) and the patients who are non-smokers ($P = 0.04$, Fisher's exact test). EGFR mutation reversely correlates with sFRP1 methylation ($P = 0.005$) and sFRP5 ($P = 0.011$). We fail to find methylation of other wnt antagonist genes correlated with EGFR mutation (Table 2).

We next investigated whether the epigenotype of any Wnt antagonist genes correlated with the genotype of EGFR. Hierarchical clustering of the epigenotype of *SFRP1*, *SFRP2*, *SFRP5*, *WIF1*, *DKK3*, *APC*, and *CDH1*, as well as the genotype of EGFR (defined as "1" if mutation

was detected in the exon 19 or 21, and as "0" if no mutation was detected) was generated using Partek Genomics Suite 6.5 (Partek Inc., MO). As shown in Figure 1, the epigenotype of Wnt antagonist genes had similar patterns, which were different from the genotype of EGFR. Therefore, our results suggested that the DNA methylation of Wnt antagonist might be independently regulated from the genotype of EGFR.

Epigenotype of Wnt antagonist genes and clinical responses to TKI therapy

The RECIST was used to evaluate the clinical response of all patients to the TKI therapy. By the end of our study, 59 (38.1%), 53 (33.2%), 43 (27.7%) patients were defined with PD, SD, or PR, respectively. We then calculated the ORR and DCR and analyzed the difference




Figure 1 Hierarchical clustering of Wnt antagonist DNA methylation status and EGFR genotype in 155 patients received EGFR-TKI therapy. Red represents methylated gene or mutated EGFR, while blue represents unmethylated gene or wild-type EGFR. The figure of hierarchical clustering showed that the epigenotype of Wnt antagonist genes had similar patterns, which were different from the genotype of EGFR.

Table 3 Multivariate statistic of gender, age, histology, smoking status, treat line, EGFR mutation and SFRP5 methylation for objective response rate (ORR) and disease control rate (DCR)

Variable	Objective response rate (ORR)			Disease control rate (DCR)		
	Univariate	Multivariate		Univariate	Multivariate	
	P value	P value	Hazard ratio (95% CI)	P value	P value	Hazard ratio (95% CI)
Gender (male / female)	0.188	0.881	0.926 (0.337-2.542)	0.001	0.115	2.117 (0.834-5.734)
Age (≤ 65 / >65)	0.351	0.078	2.295 (0.912-5.772)	0.291	0.791	1.110 (0.515-2.393)
Histology (adenocarcinoma / nonadenocarcinoma)	0.002	0.006	6.680 (1.712-26.057)	0.049	0.244	1.663 (0.707-3.915)
Line Treatment (first line / not-first line)	0.016	0.078	2.184 (0.917-5.200)	0.940	0.491	0.756 (0.341-1.678)
Smoking Status (smoker / nonsmoker)	0.016	0.262	0.526 (0.171-1.617)	0.001	0.188	0.524 (0.200-1.371)
EGFR Mutation (wide type / mutation)	<0.0001	<0.0001	7.695 (2.895-20.454)	<0.0001	0.002	3.255 (1.540-6.881)
SFRP5 Methylation (methylated / unmethylated)	0.222	0.650	0.734 (0.193-2.788)	0.04	0.106	0.434 (0.158-1.193)

between patient groups with different demographic characteristics, as well as with different genotypes of *EGFR* and epigenotypes of Wnt antagonist genes. As shown in Table 3, when only single factor was considered, the histology of the cancer (adenocarcinoma/nonadenocarcinoma), line treatment of TKI therapy (first line/not-first line), as well as smoking status (smoker/nonsmoker) significantly affected the ORR to the TKI therapy. Similarly, the gender (male/female), the histology of the cancer (adenocarcinoma/nonadenocarcinoma) as well as smoking status (smoker/nonsmoker) were found to significantly affect the DCR of the TKI therapy. However, when all demographic characteristics were considered, only the histology of the cancer ($P = 0.006$, 95% CI, 1.712-26.057, multivariate logistic regression) was associated with ORR.

Previous studies have indicated that *EGFR* mutation significantly affected the ORR and DCR of the TKI therapy. Consistently, we found that the genotype of *EGFR* significantly affected the ORR ($P < 0.0001$, 95% CI, 2.895-20.454, multivariate logistic regression adjusted by gender, age, histology, line treatment, and smoking status) and the DCR ($P = 0.002$, 95% CI, 1.540-6.881, multivariate logistic regression adjusted by gender, age, histology, line treatment, and smoking status) (Table 3).

Our results confirmed the higher response rate to the TKI therapy among patients with *EGFR* mutations as compared to the patients with wild-type *EGFR*.

Next, we investigated whether epigenotype of Wnt antagonists correlated with the clinical responses rate of the TKI therapy. Our univariate analysis identified the epigenotype of *SFRP5* as the only potential factor significantly affecting DCR but not ORR ($P = 0.04$). However, the positive association of *SFRP5* with DCR was not confirmed in multivariate analysis. When we subgrouped patients based on their demographic characteristics, we found that *SFRP1* methylation significantly reduced DCR in patients older than 65 ($P = 0.038$) and *sFRP5* methylation significantly reduced DCR in patients suffered adenocarcinoma ($P = 0.042$).

Epigenotype of Wnt antagonists and progression-free survival (PFS)

We next analyzed whether the epigenotypes of Wnt antagonists could predict the PFS in response to the TKI therapy. The median PFS time in all patients was 5.1 months (ranging from 0.4 month to 38 months). Interestingly, as shown in Figure 2A, patients with methylated *SFRP5* gene had significantly shorter median




Figure 2 Kaplan-Meier curves are shown comparing the progression free survival of patients with different epigenotypes of *SFRP5* (A), *WIF1* (B), different genotype of *EGFR* (C), or *SFRP5* in adenocarcinoma with *EGFR* mutation group (D).

PFS time (1.2 months, 95% CI, 0.5-1.9) as compared to those with unmethylated *SFRP5* gene (6.1 months, 95% CI, 4.4-7.8) ($P = 0.002$, Logrank Test). Similarly, patients with methylated *WIF1* gene had significantly shorter median PFS time (1.1 months, 95% CI, 1.0-1.2) as compared to those with unmethylated *WIF1* gene (5.4 months, 95% CI, 3.5-7.4) ($P = 0.006$, Logrank Test) (Figure 2B). We did not find association between epigenotype of other Wnt antagonists and PFS in response to the TKI therapy (Additional file 1: Figure S2 A-F). Moreover, after adjusted by age, gender, histology of the cancer, smoking status, and line of treatment, the methylation of *SFRP5* gene was still significantly associated with a shorter PFS ($P = 0.008$; hazard ratio, 2.165, 95% CI, 1.2-3.8; Cox proportional hazards models of survival analysis), while the methylation of *WIF1* gene was no longer associated with a shorter PFS ($P = 0.224$; hazard ratio, 1.804, 95% CI, 0.7-4.7; Cox proportional hazards models of survival analysis) (Table 4). Taken together, our results suggested that the methylation status of *SFRP5* might be able to predict the PFS in response to the TKI therapy.

Similar to the previous discovery [27], we also found that the median PFS time for patients with *EGFR* mutations (8.3 months, 95% CI, 5.5-11.1) was significantly longer than the median PFS for patients with wide-type *EGFR* (2.0 months, 95% CI, 1.5-2.5) ($P = 0.009$, Logrank

test) (Figure 2C). This is still valid when tested by Cox proportional hazards model of survival analysis ($P = 0.024$; hazard ratio, 0.656, 95% CI, 0.5-0.9; adjusted by age, gender, smoking status, histology of the cancer, and line of treatment). More interestingly, we found that in the subgroup of patients with adenocarcinoma and *EGFR* mutation, the ones with methylated *SFRP5* had a significantly shorter PFS (2.0 months), as compared to the ones with unmethylated *SFRP5* (9.0 months) ($P = 0.013$, Logrank Test) (Figure 2D).

Epigenotype of Wnt antagonists and overall survival rate (OS)

To test whether the epigenotype of Wnt antagonists can predict the clinical outcome of the TKI therapy, we first investigated the association of DNA methylation of the Wnt antagonists and overall survival rate in our patient cohort. Nine patients (6.5%) were lost during the follow-up period of our study. The median OS time was 27.4 months (ranging from 3.0 to 93.1 months). Interestingly, patients with methylated *WIF1* genes had significantly reduced overall survival time ($P = 0.006$, Logrank Test) (Figure 3B), while the epigenotypes of *SFRP5* (Figure 3A), *SFRP1*, *SFRP2*, *DKK3*, *APC*, and *CDH1* (Additional file 1: Figure S3 A-E), as well as the genotype of *EGFR* (Figure 3C) were not associated with OS in our patients.

Correlation between Wnt antagonist methylation and Progression-free survival in platinum-based chemotherapy


In order to decide if *WIF-1* and *sFRP5* are TKIs specific biomarkers related to PFS of TKIs treatment, we meanwhile analyzed the association of chemotherapy with the epigenotype of Wnt antagonists in 63 patients out of the whole group, who once took platinum-based chemotherapy as first-line treatment. We failed to find significant differences in PFS between patients with or without *sFRP5* methylation (3.2 ms, 95% CI 2.01-4.5 vs 4.3 ms, 95% CI 2.5-6.2, respectively, $P = 0.487$). We did not find differences in PFS between patients with or without *WIF-1* methylation (3.2 ms, 95% CI 1.89-4.67 vs 2.0 ms, 95% CI 1.71-2.36 $P = 0.798$) either. We accidentally found discrepancy in PFS between patients with or without *sFRP1* methylation (1.8 ms, 95% CI, 1.50-2.09 vs 3.0 ms 95% CI, 1.9-4.0, $P = 0.017$). However, this statistically significant difference in PFS remains limited for patients in clinical practice.

Discussion

Recent studies have demonstrated that cancer is as much an epigenetic disease as it is a genetic disease (Iacobuzio-Donahue). Therefore, in addition to genetic alterations, changes in epigenetic features such as CpG DNA methylation status of specific gene loci also mark

Table 4 Cox proportional hazard regression analysis of gender, age, histology, smoking status, EGFR mutation, WIF1 methylation and SFRP5 methylation for progression-free survival (PFS)

Variable	P value	Hazard ratio (95% CI)
Smoking Status (smokers/nonsmokers)	0.986	1.004 (0.615-1.640)
Histology (adenocarcinoma/Nonadenocarcinoma)	0.689	0.915 (0.592-1.414)
Gender (male/female)	0.006	0.516 (0.322-0.826)
Age (<65/>65)	0.456	0.858 (0.575-1.282)
Lines of Treatment (first line/non-first line)	0.302	0.807 (0.537-1.213)
EGFR Mutation (mutation/wide type)	0.024	0.656 (0.455-0.945)
SFRP5 Methylation (methylated/unmethylated)	0.008	2.165 (1.226-3.823)
WIF1 Methylation (methylated/unmethylated)	0.224	1.804 (0.697-4.674)



the progress of cancers. Our current study showed that methylation of Wnt antagonist *SFRP5* gene before treatment, independent of the genotype of *EGFR* gene, correlated with decreased progression free survival rate in NSCLC patients in response to the *EGFR*-TKI therapy. To our knowledge, this is the first report indicating that DNA methylation at specific gene loci in patient may predict drug response to the *EGFR*-TKI therapy.

Both genetic and epigenetic risk factors for NSCLC have been studied extensively. Suzuki et al [23] has reported that methylation of the Wnt antagonist *DKK3* correlated with low survival rate in NSCLC patients, despite of the different therapies patients received. However, in our study, we did not find significant difference in the *EGFR*-TKI responses between patient groups with or without methylated *DKK3* (Additional file 1: Figure S2 and S3). In contrast, our results suggested epigenotype of *SFRP5* provide better prognostic estimation for the *EGFR*-TKI response, comparing to other Wnt antagonists.

SFRP5 is a member of the SFRP protein family containing a cysteine-rich domain homologous to the putative Wnt-binding site of Frizzled proteins. It acts as soluble antagonist of Wnt signaling and is highly expressed in the retinal pigment epithelium, and moderately expressed in the pancreas ("Entrez Gene: *SFRP5* secreted frizzled-related protein 5"). Previous studies has identified association of *SFRP5* promoter hypermethylation with Acute myeloid leukemia [29], ovarian cancer [30], gastric cancer [31], oral squamous cell carcinoma [32], pancreatic cancer [33] and breast cancer [34].

We found that hypermethylation of *SFRP5* predicted worse outcomes of the *EGFR*-TKI therapy. Therefore,

SFRP5 DNA methylation status may serve as a prognostic molecular marker for appropriately predicting whether NSCLC patients would benefit from the *EGFR*-TKI therapy. Especially, it is interesting that in the subgroup with adenocarcinoma and *EGFR* mutation, patients with *sFRP5* methylation have a significantly shorter PFS than those without *sFRP5* methylation, While in nonsmokers without *EGFR* mutation, patients without *sFRP1* methylation have a longer PFS compared with patients with its methylation(9.7 ms vs 2.0 ms, $p=0.05$). Based on these results, we can make a hypothesis that activation of Wnt signaling by antagonist methylation could confer tumors the characters of stem cell, which consequently causes tumors resistant to *EGFR* TKIs therapy by generating acquired resistance, such as *MET* amplification or changes of *PTEN* tumor suppressor activity and so on. Further study is needed to validate this hypothesis.

Conclusions

In conclusion, our study revealed that *sFRP5* may be an independent factor affecting PFS during long time maintenance of TKIs therapy. Furthermore, the simple, PCR-based detection method of DNA methylation may be more feasible as clinical tests, compared to protein or RNA expression detection in clinics. Both general DNA methylation inhibitors and Wnt-pathway-targeting anticancer drugs are under development [35,36]. Our results that linked Wnt antagonist hypermethylation and *EGFR*-TKI response suggest that the treatment paradigm combining epigenetic drugs and *EGFR*-TKI may be a potential and attractive therapeutic option for patients with NSCLC.

Additional file

Additional file 1: Figure S1. Methylated and unmethylated bands of Wnt antagonist genes and wild/mutant EGFR. S1: The example graphs of methylated and unmethylated bands of Wnt antagonist genes (A) and EGFR wild (B) and mutation types (C, D) by methylation specific PCR and DHPLC respectively. Figure S2 PFS with different epigenotypes of Wnt antagonist genes. Figure S2S A-F. Kaplan-Meier curves of comparing the progression free survival of patients with different epigenotypes of SFRP1 (A), SFRP2 (B), DKK3 (C), APC (D), CDH1 (E) and combination analysis (F). Figure S3 OS with different epigenotypes of Wnt antagonist genes. Figure S3S A-F. Kaplan-Meier curves of comparing the overall survival of patients with different epigenotypes of SFRP1 (A), SFRP2 (B), DKK3 (C), APC (D), CDH1 (E) and combination analysis (F).

Abbreviations

EGFR: Epidermal growth factor receptor; EGFR-TKI: Epidermal growth factor receptor -tyrosine kinase inhibitors; MSP: Methylation specific PCR; Wnt: Wingless-type; ECOG: Eastern cooperative oncology group; ORR: Objective response rate; DCR: Disease control rate; PFS: Progression-free survival; OS: Overall survival; PD: Disease progression; CR: Complete response; PR: Partial response; SD: Stable disease; RECIST: Response evaluation criteria in solid tumors; HR: Hazard ratio.

Competing interests

The authors declare that they have no competing interests.

Authors' contributions

JZ, YW carried out the molecular genetic studies; JD, MZ, ZW, JZ, SW, LY, TA, MW participated in Provision of study materials or patients and collection and assembly of data; LW, JZ, YW, HB and JW analyzed final data and JZ, YW, JW drafted the manuscript. All authors read and approved the final manuscript.

Authors' information

Supported by grants from National Natural Sciences Foundation Distinguished Young Scholars (81025012), National Natural Sciences Foundation General Program (81172235), Beijing Health Systems Academic Leader (2011-2-22).

Acknowledgement

We thank Dr. BM Zhu for her critical review of this manuscript and Dr Ning Wang in the radiological department of Beijing Cancer Hospital for his assessments of the response of treatment. We thank Dr. Guoshuang Feng in (Chaoyang District Center for Disease Control and Prevention) for statistical analysis.

Author details

¹Department of Thoracic Medical Oncology Peking University School of Oncology, Beijing Cancer Hospital & Institute, Beijing 100036, China.

²Laboratory of Immunology, National Eye Institute, National Institutes of Health, 31 Center Drive MSC 2510 Bethesda, MD 20892-2510, USA.

Received: 22 May 2012 Accepted: 16 August 2012

Published: 25 September 2012

References

- Jemal A, Siegel R, Ward E, Hao Y, Xu J, Murray T, et al: **Cancer statistics, 2008.** *CA Cancer J Clin* 2008, **58**(2):71-96.
- Govindan R, Page N, Morgensztern D, Read W, Tierney R, Vlahiotis A, et al: **Changing epidemiology of small-cell lung cancer in the United States over the last 30 years: Analysis of the surveillance, epidemiologic, and end results database.** *J Clin Oncol* 2006, **24**:4539-4544.
- Sekido Y, Fong KM, Minna JD: **Progress in understanding the molecular pathogenesis of human lung cancer.** *Biochim Biophys Acta* 1998, **1378**:F21-F59.
- Fossella F, Pereira JR, Pawel JV, Pluzanska A, Gorbounova V, Kaukel E, et al: **Randomized, multinational, phase III study of docetaxel plus patinnum combinations versus vinorelbine plus cisplatin for advanced NSCLC: the TAX326 Study Group.** *J Clin Oncol* 2003, **21**(16):3016-3024.
- Ramalingam S: **First-line chemotherapy for advanced-stage non-small cell lung cancer: focus on docetaxel.** *Clin Lung Cancer* 2005, **7**:S77-S82.
- Surmont V, Aerts JG, Tan KY, Schramel F, Vernhout R, Hoogsteden HC, et al: **Non-cross resistant sequential single agent chemotherapy in first-line advanced non-small cell lung cancer patients: results of a phase II study.** *J Oncol* 2009. doi:10.1155/2009/457418.
- Mok TS, Wu YL, Thongprasert S, Yang CH, Chu DT, Saijo N, et al: **Gefitinib or carboplatin-paclitaxel in pulmonary adenocarcinoma.** *N Engl J Med* 2009, **361**(10):947-957.
- Maemondo M, Inoue A, Kobayashi K, Sugawara S, Oizumi S, Isobe H, et al: **Gefitinib or Chemotherapy for Non-Small-Cell Lung Cancer with Mutated EGFR.** *N Engl J Med* 2010, **362**(25):2380-2388.
- Mitsudomi T, Morita S, Yatabe Y, Negoro S, Okamoto I, Tsurutani J, et al: **Gefitinib versus cisplatin plus docetaxel in patients with non-small-cell lung cancer harbouring mutations of the epidermal growth factor receptor (WJTOG3405): an open label, randomised phase 3 trial.** *Lancet Oncol* 2010, **11**(2):121-128.
- Zhou CC, Wu YL, Chen GY, Feng JF, Liu XQ, Wang CL, et al: **Erlotinib versus chemotherapy as first-line treatment for patients with advanced EGFR mutation-positive non-small-cell lung cancer (OPTIMAL, CTONG-0802): a multicentre, open-label, randomised, phase 3 study.** *Lancet Oncol* 2011, **8**(12):735-742.
- Rosell R, Gervais R, Vergnenegre A, Massuti B, Felip E, Cardenal F, et al: **Erlotinib vs chemotherapy (CT) in advanced non-small-cell lung cancer (NSCLC) patients (P) with epidermal growth factor receptor (EGFR) activating mutations: interim results of the European Tarceva vs chemotherapy (EURTAC) phase III randomized trial.** *ASCO* 2011, : abs7503.
- Paez JG, Jänne PA, Lee JC, Tracy S, Greulich H, Gabriel S, et al: **EGF mutations in lung cancer: Correlation with clinical response to gefitinib therapy.** *Science* 2004, **304**:1497-1500.
- Lynch TJ, Bell DW, Sordella R, Gurubhagavatula S, Okimoto RA, Brannigan BW, et al: **Activating mutations in the epidermal growth factor receptor underlying responsiveness of non-small-cell lung cancer to gefitinib.** *N Engl J Med* 2004, **350**:2129-2139.
- Pao W, Miller V, Zakowski M, Doherty J, Politi K, Sarkaria I, et al: **EGF receptor gene mutations are common in lung cancers from "never-smokers" and are associated with sensitivity of tumors to gefitinib and erlotinib.** *Proc Natl Acad Sci USA* 2004, **101**:13306-13311.
- Gazdar AF, Shigematsu H, Herz J, Minna JD: **Mutations and addiction to EGFR: the Achilles 'heal' of lung cancers?** *Trends Mol Med* 2004, **10**(10):482-487.
- Mitsudomi T, Kosaka T, Yatabe Y: **Biological and clinical implications of EGFR mutations in lung cancer.** *Int J Clin Oncol* 2006, **11**(3):190-198.
- Sequist LV, Bell DW, Lynch TJ, Haber DA: **Molecular predictors of response to epidermal growth factor receptor antagonists in non-small-cell lung cancer.** *J Clin Oncol* 2007, **25**(5):587-595.
- Kobayashi S, Boggon TJ, Dayaram T, Jänne PA, Kocher O, Meyerson M, et al: **EGFR mutation and resistance of non-small-cell lung cancer to gefitinib.** *N Engl J Med* 2005, **352**:786-792.
- Engelman JA, Zejnullahu K, Mitsudomi T, Song YC, Hyland C, Park JO, et al: **MET Amplification Leads to Gefitinib Resistance in Lung Cancer by Activating ERBB3 Signaling.** *Science* 2007, **316**:1039-1043.
- Lustig B, Behrens J: **The Wnt signaling pathway and its role in tumor development.** *J Cancer Res Clin Oncol* 2003, **129**:199-221.
- Kato H: **WNT/PCP signaling pathway and human cancer [review].** *Oncol Rep* 2005, **14**:1583-1588.
- Hoschuetzky H, Aberle H, Kemler R: **Beta-Catenin mediates the interaction of the cadherin-catenin complex with epidermal growth factor receptor.** *J Cell Biol* 1994, **127**:1375-1380.
- Suzuki M, Shigematsu H, Nakajima T, Kubo R, Motohashi S, Sekine Y, et al: **Synchronous Alterations of Wnt and Epidermal Growth Factor Receptor Signaling Pathways through Aberrant Methylation and Mutation in Non-Small Cell Lung Cancer.** *Clin Cancer Res* 2007, **13**:6087-6092.
- Therasse P, Arbuck SG, Eisenhauer EA, et al: **New guidelines to evaluate the response to treatment in solid tumors.** *J Natl Cancer Inst* 2000, **92**:205-216.
- Mazieres J, He B, You L, Xu ZD, Lee AY, Mikami I, et al: **Wnt inhibitory factor-1 is silenced by promoter hypermethylation in human lung cancer.** *Cancer Res* 2004, **64**:4717-4720.

26. Brabender J, Usadel H, Danenberg KD, Metzger R, Schneider PM, Lord RV, *et al*: Adenomatous polyposis coli gene promoter hypermethylation in non-small cell lung cancer is associated with survival. *Oncogene* 2001, **20**(27):3528–3532.
27. Lee SM, Kim MJ, Lee JY, Park JY, Kim DS: Aberrant methylation of E-cadherin and H-cadherin genes in non-small cell lung cancer and its relation to clinicopathologic features. *Cancer* 2007, **12**:2785–2792.
28. Bai H, Mao L, Wang SH, Zhao J, Yang L, An TT, *et al*: Epidermal Growth Factor Receptor Mutations in Plasma DNA Samples Predict Tumor Response in Chinese Patients With Stages IIIB to IV Non-Small-Cell Lung Cancer. *J Clin Oncol* 2009, **27**:2653–2659.
29. Griffiths EA, Gore SD, Hooker C, McDevitt MA, Karp JE, Smith BD, *et al*: Acute myeloid leukemia is characterized by Wnt pathway inhibitor promoter hypermethylation. *Leuk Lymphoma* 2010, **51**(9):1711–1719.
30. Su HY, Lai HC, Lin YW, Liu VY, Chen CK, Chou YC, *et al*: Epigenetic silencing of SFRP5 is related to malignant phenotype and chemoresistance of ovarian cancer through Wnt signaling pathway. *Int J Cancer* 2010, **127**(3):555–567.
31. Zhao C, Bu X, Zhang N, Wang W: Downregulation of SFRP5 expression and its inverse correlation with those of MMP-7 and MT1-MMP in gastric cancer. *BMC Cancer* 2009, **9**:224.
32. Sogabe Y, Suzuki H, Toyota M, Ogi K, Imai T, Nojima M, *et al*: Epigenetic inactivation of SFRP genes in oral squamous cell carcinoma. *Int J Oncol* 2008, **32**(6):1253–1261.
33. Bu XM, Zhao CH, Zhang N, Gao F, Lin S, Dai XW: Hypermethylation and aberrant expression of secreted frizzled-related protein genes in pancreatic cancer. *World J Gastroenterol* 2008, **14**(21):3421–3424.
34. Veeck J, Geisler C, Noetzel E, Alkaya S, Hartmann A, Knuchel R, *et al*: Epigenetic inactivation of the secreted frizzled-related protein-5 (SFRP5) gene in human breast cancer is associated with unfavorable prognosis. *Carcinogenesis* 2008, **29**(5):991–998.
35. Minke KS, Staib P, Puetter A, Gehrke I, Gandhirajan RK, Schlösser A, *et al*: Small molecule inhibitors of WNT signaling effectively induce apoptosis in acute myeloid leukemia cells. *Eur J Haematol* 2009, **82**(3):165–175.
36. Esteller M: DNA methylation and cancer therapy: new developments and expectations. *Curr Opin Oncol* 2005, **17**(1):55–60.

doi:10.1186/1756-9966-31-80

Cite this article as: Zhu *et al*: DNA Methylation status of Wnt antagonist SFRP5 can predict the response to the EGFR-tyrosine kinase inhibitor therapy in non-small cell lung cancer. *Journal of Experimental & Clinical Cancer Research* 2012 **31**:80.

Submit your next manuscript to BioMed Central and take full advantage of:

- Convenient online submission
- Thorough peer review
- No space constraints or color figure charges
- Immediate publication on acceptance
- Inclusion in PubMed, CAS, Scopus and Google Scholar
- Research which is freely available for redistribution

Submit your manuscript at
www.biomedcentral.com/submit

