

The Incidence of Thyroid Cancer Is Affected by the Characteristics of a Healthcare System

Tae-Jin Lee¹, Sun Kim¹, Hong-Jun Cho²,
and Jae-Ho Lee³

¹Graduate School of Public Health & Institute of Health and Environment, Seoul National University, Seoul; ²Department of Family Medicine, University of Ulsan College of Medicine, Seoul; ³Department of Family Medicine, The Catholic University of Korea College of Medicine, Seoul, Korea

Received: 2 September 2012
Accepted: 22 October 2012

Address for Correspondence:
Jae-Ho Lee, MD

Department of Family Medicine, The Catholic University of Korea College of Medicine, 222 Banpo-daero, Seocho-gu, Seoul 137-701, Korea
Tel: +82.2-2258-6288, Fax: +82.2-2258-2907
E-mail: jaeholee@catholic.ac.kr

The aim of this study was to investigate the associations between the incidence of thyroid cancer and the characteristics of healthcare systems in OECD countries and to demonstrate that the increasing incidence of thyroid cancer is mainly due to overdiagnosis. We used a random effects panel model to regress the incidence of thyroid cancer on the characteristics of healthcare systems (i.e., share of public expenditure on health, mode of health financing, existence of referral system to secondary care, mode of payment to primary care physicians), controlling for macro context variables (i.e., GDP per capita, educational level) on a country level. Data were derived from 34 OECD countries for 2002 and 2008. The share of public expenditure on health was negatively associated with the incidence of thyroid cancer. However, it had no statistically significant effect on the mortality of thyroid cancer and on the incidence of stomach and lung cancer. In the case of colorectal cancer, it had a positive effect on the incidence rate. The upward trend of the incidence of thyroid cancer is closely related to the healthcare system that permits overdiagnosis. Increases in the proportion of public financing may help reduce the overdiagnosis of thyroid cancer.

Key Words: Thyroid Neoplasms; Overdiagnosis; Healthcare System; Public Health Expenditure

INTRODUCTION

During the past several decades, an increasing incidence of thyroid cancer has been reported in many parts of the world, while its mortality has remained stable during this period. The upward trend in the incidence rate started back in the 1940s (1) characterized by the papillary type of thyroid cancer. In the United States, thyroid cancer was diagnosed in 4.9 per 100,000 in 1975, which gradually increased to 13.0 per 100,000 by 2008 (2). European countries reported increases in the incidence rate between 5.3% (Switzerland) and 155.6% (France) between 1973 and 2002, while Southern Australia reported the greatest increases, a 177.8% increase in men and a 252.2% increase in women during the same period (3). The highest female incidence rate ever reported was 62.9 (1995-1999) per 100,000 in New Caledonia, a French territory in the South Pacific (4). However, Korea broke the world record in the incidence rate and its increase: for women, 80.2 (2008) per 100,000 and an annual increase rate of 23.6% (1999-2008) (5).

Earlier studies suggested that the increasing trend was driven by the widespread use of radiation therapy for benign conditions of the head and neck among children and adolescents from the early 1920s to the late 1950s (6). Some studies suggested that the

trend might be associated with atmospheric nuclear fallout (7), or increased exposure to diagnostic radiography (8), particularly among children (9, 10).

Two recent studies concluded that the increasing incidence of thyroid cancer could be a result of improved diagnosis of subclinical cancers due to the increased use of ultrasound-guided fine-needle aspiration biopsy (11, 12). Many of these subclinical cancers are tumors measuring 1 cm or smaller (13) that lie dormant in the general population, never manifesting, and usually left undiagnosed. Besides, it was found that income and gender might affect the stage at the initial disease presentation of thyroid cancer (14), and that there were positive correlations between the incidence of thyroid cancer and socioeconomic indicators of healthcare access including household income, education, and health insurance (15). These are consistent with the hypothesis that the increasing incidence trend of thyroid cancer is attributable to the utilization of new diagnostic practices.

Epidemiologists have labeled this phenomenon "overdiagnosis"—a term perhaps most familiar in the clinical setting for prostate cancer (16). The prerequisites for cancer overdiagnosis are twofold: the existence of a reservoir of detectable cancer and activities leading to the detection (17). In terms of the reservoir,

thyroid cancer is an extreme case. A study that examined slices of thyroid tissue taken every 2.5 mm found at least one papillary carcinoma in 36% of Finnish adults, concluding that the prevalence of histologically verifiable papillary carcinoma would be close to 100% if one could look at thin enough slices of the gland (18). Moreover, the most obvious activities leading to detection is cancer screening, especially use of diagnostic imaging. It is estimated that the probability of overdiagnosis is 99.7%-99.9% in thyroid cancer in which the entire reservoir is detected (17). For this reason, almost all national guidelines do not recommend screening for thyroid cancer in asymptomatic people.

Taking into consideration this overdiagnosis phenomenon, it is possible to postulate that the incidence of thyroid cancer is higher in countries where the healthcare system provides inappropriate incentives for cancer screening for other purposes rather than for the health of the individual (e.g., making profit). For example, overdiagnosis is more likely to occur in less regulated markets, which might be related to a lack of the public sector's role in providing healthcare services. However, there has been no research studying the relationship between the incidence of thyroid cancer and the characteristics of healthcare systems. To support the hypothesis that the increasing incidence of thyroid cancer is mainly caused by overdiagnosis and closely related to the healthcare system, we investigated the associations between the incidence of thyroid cancer and the characteristics of healthcare systems in Organization for Economic Cooperation and Development (OECD) countries.

MATERIALS AND METHODS

Model

To find out the relationship between the incidence of thyroid cancer and the characteristics of healthcare systems, we chose a set of explanatory variables based on already known theories and on empirical studies about individual- and country-level factors that influence health outcomes or healthcare utilization. To begin with, we included two macro context variables in our models to control for their effects. First, GDP per capita is a well-known determinant of health as a measure of income or the socioeconomic environment, which in turn influences healthcare resources (19, 20). Second, education is also a strong determinant of health through lifestyle factors and healthcare utilization since it is associated with knowledge about health and healthcare services (21).

Apart from GDP per capita and education, health outcomes and healthcare utilization can be influenced by other factors that affect the efficiency of healthcare systems (22, 23). First, we included the share of public health expenditure (the proportion of health expenditure covered by the public sector) as a measure of the commercialization of healthcare systems (24). A low share of public health expenditure may mean less regulation in

healthcare delivery, which in turn is likely to lead to commodification of healthcare services. Under the commercialized healthcare system, overdiagnosis is more likely to occur. Besides, public health expenditure was reported as an important factor that affects health outcomes (24, 25). Second, we included mode of health financing as a factor that affects the efficiency of the healthcare system. It was reported that healthcare systems based on tax-financing were more efficient than healthcare systems based on social health insurance in producing better health outcomes and reducing health spending (20). Third, we included the existence of the gatekeeping role in primary care for referral to secondary care since there is a risk of overdiagnosis without gatekeeping. Fourth, we included the mode of payment to primary care physicians as a proxy for how the whole medical system works since primary care usually deals with the majority of health care needs in the first level of contact, and doctors have incentives to generate higher income by increasing the services provided under the fee-for-service system (26).

In terms of the dependent variables, not only the incidence of thyroid cancer, but also the mortality of thyroid cancer and the incidence of stomach, lung, and colorectal cancer were used, respectively, to compare the effects of the characteristics of healthcare systems. Since mass screening for stomach and lung cancers are not usually recommended in developed countries, incidences of those cancers are not likely to be associated with the share of public health expenditure. In contrast, the incidence of colorectal cancer for which well-coordinated screening programs are usually recommended is likely to be affected by the share of public health expenditure.

Given those explanatory and dependent variables, we built two models in this study. In model 1, we focused on the influence of public health expenditure controlling for two macro context variables. In model 2, we added more healthcare system variables, i.e. health financing, payment to primary care physicians, and referral to secondary care, to those variables in model 1. The two models are as follows:

Model 1:

$$\log\text{thyinc}_{it} = \beta_0 + \beta_1 \log\text{GDP}_{it} + \beta_2 \text{educ}_{it} + \beta_3 \text{peh}_{it} + \varepsilon_{it}$$

Model 2:

$$\log\text{thyinc}_{it} = \beta_0 + \beta_1 \log\text{GDP}_{it} + \beta_2 \text{educ}_{it} + \beta_3 \text{peh}_{it} + \beta_4 \text{finance}_{it} + \beta_5 \text{refer}_{it} + \beta_6 \text{primpay}_{it} + \varepsilon_{it}$$

where thyinc_{it} is country i 's incidence of thyroid cancer at time t ; GDP_{it} is the GDP per capita; educ_{it} is the percent of the population with attainment at the tertiary level of education (some college); peh_{it} is the percent of public health expenditure; finance_{it} is a dummy indicating whether country i 's health financing is based on tax or social insurance; refer_{it} is a dummy indicating whether or not country i 's referral to secondary care is

compulsory or financially encouraged, and $primpay_{it}$ is a dummy indicating whether or not country i 's payment to primary care physicians is fee-for-service.

Data sources

This study was done with data from 34 OECD countries for 2002 and 2008. Incidence and mortality data were obtained from the GLOBOCAN project by the International Agency for Research on Cancer (IARC), WHO for 2002 (27) and for 2008 (28). This project provides contemporary estimates of the incidence and mortality from major types of cancers at the national level for all countries (28).

Data about macro context variables, i.e. GDP per capita and education, and share of public health expenditure were obtained

from OECD health data 2010 (29), data on payment to primary care physicians and referral to secondary care from OECD (30), and data on health financing from the World Bank (31). Since data on education for all the target countries did not exist for 2008, we used the data for 2007 instead.

Statistical analysis

Given that some of the explanatory variables are time invariant, this study used a random effects panel model to investigate the effects of the explanatory variables on the dependent variables at the country level. To get correct standard errors despite any pattern of heteroskedasticity or serial correlation, robust-cluster variance estimator was used. Incidence rates and mortality rates as well as the GDP per capita were used after log transformation




Fig. 1. Incidence and mortality of thyroid cancer in OECD countries (2002, 2008). Incidence and mortality of thyroid cancer denote age-standardized rates per 100,000 population in OECD countries in 2002 and 2008. About three fourths of the countries experienced increases in the incidence of thyroid cancer from 2002 to 2008 while most countries maintained a low mortality or even experienced decreases in mortality during this period. Sources: GLOBOCAN 2002 and 2008.

to normalize the distribution.

RESULTS

Fig. 1 shows the incidence and mortality of thyroid cancer for 34 OECD countries in 2002 and 2008. About three fourths of the countries experienced increases in the incidence of thyroid cancer. Especially, the incidence rate for Korea increased remarkably. The incidence rate for Korean women increased nearly ten-fold from 2002 to 2008. Despite the increased incidence, its mortality did not change much and was similar to that of the other countries with a lower incidence rate. Besides Korea, most countries maintained a low mortality and even experienced decreases in mortality during this period.

Table 1 presents the descriptive statistics of the dependent and explanatory variables used in the regression analysis. There are some gender differences in the incidence of cancers. The incidence of thyroid cancer was higher in females while the in-

cidences of stomach, lung, and colorectal cancer were all higher in males. From 2002 to 2008, the GDP per capita and educational level showed increases. The mean share of public health expenditure did not change a lot, which was around 72% of the total health expenditure. In terms of payment to primary care physicians, fee-for-service was the predominant mode of payment in 14 out of 34 countries. Twenty-one countries had a referral system for secondary care, although there were some variations in the forms. Twenty countries had social health insurance or private health insurance while 14 had tax-finance as their major method of health financing.

Table 2 presents the regression results of the random effects panel model for the incidence and mortality of thyroid cancer for males and females, respectively. Both the GDP per capita and education were positively associated with the incidence of thyroid cancer for both genders, but the effect of education was not statistically significant. In contrast, the share of public health expenditure was negatively associated with the incidence of thy-

Table 1. Descriptive statistics of variables used in the regression analyses

Variables	Definition	No.	Mean	S.D.	Min.	Max.
Dependent variables (Age-standardized and expressed as per 100,000)						
Mthyinc	Male incidence of thyroid cancer (2002)	34	1.8	0.8	0.5	4.4
	Male incidence of thyroid cancer (2008)	34	2.7	1.8	0.7	10.9
Fthyinc	Female incidence of thyroid cancer (2002)	34	5.0	2.3	2	11.6
	Female incidence of thyroid cancer (2008)	34	8.8	9.7	2.8	59.5
Mthymort	Male mortality of thyroid cancer (2002)	34	0.4	0.1	0.2	0.7
	Male mortality of thyroid cancer (2008)	34	0.3	0.3	0	1.6
Fthymort	Female mortality of thyroid cancer (2002)	34	0.5	0.2	0.3	1
	Female mortality of thyroid cancer (2008)	34	0.4	0.3	0	1.6
Mstominc	Male incidence of stomach cancer (2002)	34	18.0	14.4	7.2	69.6
	Male incidence of stomach cancer (2008)	34	13.6	11.7	4.6	62.2
Fstominc	Female incidence of stomach cancer (2002)	34	8.3	5.6	3.3	26.8
	Female incidence of stomach cancer (2008)	34	6.2	4.5	2.5	24.6
Mlunginc	Male incidence of lung cancer (2002)	34	49.5	17.3	17	94.6
	Male incidence of lung cancer (2008)	34	43.0	14.1	14	80.9
Flunginc	Female incidence of lung cancer (2002)	34	15.3	7.9	5.3	36.1
	Female incidence of lung cancer (2008)	34	17.8	8.5	5.2	36.2
Mcoloninc	Male incidence of colorectal cancer (2002)	34	38.1	12.1	7.9	58.5
	Male incidence of colorectal cancer (2008)	34	38.2	12.0	7	60.6
Fcoloninc	Female incidence of colorectal cancer (2002)	34	26.4	7.7	7	42.2
	Female incidence of colorectal cancer (2008)	34	25.1	7.4	6.1	37.5
Explanatory variables						
GDP	GDP per capita (US dollars, current prices and PPPs) (2002)	34	25,063.1	9,918.7	8,667	57,546
	GDP per capita (US dollars, current prices and PPPs) (2008)	34	33,065.3	13,458.1	13,952	84,713
Educ	Attainment at the tertiary level	33*	24.0	9.6	9	43
	(% of population with educational level of some college) (2002)					
PEH	Attainment at the tertiary level	34	27.9	9.6	11	48
	(% of population with educational level of some college) (2008)					
Finance	Public health expenditure (% total expenditure on health) (2002)	33 [†]	71.8	11.9	43.9	90.5
	Public health expenditure (% total expenditure on health) (2008)	31 [†]	72.0	10.8	46.5	84.5
Refer	Health financing (tax-finance = 1, social health insurance or private health insurance = 0)	30 [§]	0.4	0.5	0	1
	Referral to secondary care (compulsory or financially-encouraged = 1, no obligation and no incentive = 0)	30	0.7	0.5	0	1
Primpay	Payment to primary care physicians	34	0.4	0.5	0	1
	(fee for service = 1, otherwise = 0)					

Sources: GLOBOCAN 2002 and 2008; OECD Health Data 2010. *Data about educational level of Chile for 2002 was missing in OECD health data; [†]Data about public health expenditure of Belgium for 2002 was missing in OECD health data; [‡]Data about public health expenditure of Belgium, the Netherlands and Portugal for 2008 were missing in OECD health data; [§]Data about health financing of Chile, Estonia, Israel and Slovenia were missing in the reference; ^{||}Data about referral to secondary care of Chile, Estonia, Israel and Slovenia were missing in the reference. S.D., standard deviation; Min., minimum; Max., maximum; GDP, gross domestic product; PPP, purchasing power parity.

Table 2. Factors affecting the incidence and mortality of thyroid cancer

Independent variables	Incidence								Mortality							
	Male				Female				Male				Female			
	Model 1		Model 2		Model 1		Model 2		Model 1		Model 2		Model 1		Model 2	
	Est.	P value	Est.	P value	Est.	P value	Est.	P value	Est.	P value	Est.	P value	Est.	P value	Est.	P value
Intercept	-4.02	0.003	-4.42	0.01	-1.78	0.18	-2.76	0.11	-0.18	0.90	-0.79	0.59	5.09	0.000	5.38	0.000
Log GDP*	0.50	0.001	0.51	0.01	0.41	0.01	0.54	0.006	-0.10	0.47	-0.01	0.94	-0.58	0.000	-0.55	0.000
Educ [†]	0.01	0.28	0.02	0.23	0.02	0.11	0.01	0.47	-0.00	0.58	-0.01	0.14	-0.00	0.84	-0.01	0.03
PEH [‡]	-0.01	0.09	-0.01	0.36	-0.02	0.02	-0.02	0.02	0.00	0.56	0.00	0.73	0.00	0.84	-0.00	0.51
Finance [§]	-	-	-0.30	0.23	-	-	0.00	0.99	-	-	-0.00	1.00	-	-	0.14	0.23
Refer	-	-	0.01	0.97	-	-	-0.17	0.46	-	-	-0.15	0.44	-	-	-0.31	0.048
Primpay [¶]	-	-	-0.20	0.44	-	-	-0.07	0.79	-	-	0.03	0.86	-	-	0.18	0.10
Rho	0.47	-	0.49	-	0.53	-	0.53	-	0.44	-	0.46	-	0.37	-	0.31	-
R ² (overall)	0.19	-	0.19	-	0.20	-	0.19	-	0.03	-	0.08	-	0.30	-	0.42	-
No.	63	-	56	-	63	-	56	-	62	-	55	-	62	-	55	-

*Log transformed GDP per capita (US dollars, current prices and PPPs); [†]Attainment at the tertiary level (% of population with educational level of some college); [‡]Public health expenditure (% total expenditure on health); [§]Health financing (countries which use tax-finance as their major method of health financing were coded to 1 and other countries [use social health insurance or private health insurance, instead] were coded to 0); ^{||}Referral to secondary care (countries in which referral is compulsory or financially encouraged were coded to 1 and other countries [neither compulsory nor financially encouraged] were coded to 0); [¶]Payment to primary care physicians (countries which use fee-for-service as their predominant mode were coded to 1 and other countries [use salary or capitation, or combination of these with fee-for-service] were coded to 0). Est., estimate.

roid cancer for both genders, although its effect became insignificant for males. The other characteristics of healthcare systems had no significant effect on the incidence of thyroid cancer.

The results for mortality were different from those of the incidence. The GDP per capita was negatively associated with the mortality of thyroid cancer, though not significant for males. Interestingly, the effect of public health expenditure on mortality disappeared. When healthcare system variables were added to model 2, educational level became significant and was negatively associated with the mortality of thyroid cancer for females. Countries with no referral system had a higher mortality of thyroid cancer for females.

In Table 3, the regression results of the random effects panel model on the incidence of stomach, lung, and colorectal cancer show different patterns from those on the incidence of thyroid cancer. Specifically, the share of public health expenditure showed no significant effect on the incidence of stomach and lung cancer, while it had a positive effect on the incidence of colorectal cancer for males in model 2. This finding contradicts the finding that a low share of public health expenditure was associated with a high incidence of thyroid cancer.

DISCUSSION

Overdiagnosis of thyroid cancers and subsequent surgeries could not only be an extra burden for patients but also a waste of scarce resources from a social viewpoint. Nevertheless, why is the incidence of thyroid cancer increasing very rapidly in some parts of the world? We examined whether some characteristics of healthcare systems, including the share of public health expenditure, can explain the overdiagnosis of thyroid cancer at the country level.

The results of our analysis show that a low share of public health expenditure is associated with a high incidence of thyroid cancer. Since a low proportion of public health financing means a high proportion of private health financing, it is highly probable that commercialization of the healthcare system occurs in this less regulated market. Thus, a low share of public health expenditure may permit overdiagnosis, which inevitably leads to a high incidence of thyroid cancer. In detail, if the share of public health financing is low, private health financing – private health insurance or patients’ direct payment – would be of great importance. In this context, individuals’ abilities to pay for healthcare services become crucial in the utilization of healthcare, so both demand induced by the providers and overuse by patients could easily occur.

For example, Korea has a low share of public health expenditure (55.3% out of total health expenditure, the third lowest among OECD countries in 2008) and relies heavily on patients’ direct payment (35.0%, the second highest among OECD countries) supplemented with private health insurance (29). Besides, healthcare services are delivered predominantly by private providers. Recently, there has been increasing demand for cancer screening and increasing use of ultrasound. Since ultrasound is excluded from social health insurance benefits, there is literally no regulation about screening with ultrasound. Moreover, there is neither obligation nor incentive for patients to register with primary care physicians who can coordinate cancer screening, and secondary care in which medical imaging equipment including ultrasound is frequently used for screening is easily accessible without referral. In this situation, screening for thyroid cancer can flourish. In practice, it was reported in a survey of 2008 that 60.4% and 90.0% of family physicians and internists working at their community offices performed ultrasound test,

Table 3. Factors affecting the incidence of the stomach (a), lung (b), and colorectal cancer (c)**(a) Stomach cancer**

Independent variables	Male				Female			
	Model 1		Model 2		Model 1		Model 2	
	Est.	P value	Est.	P value	Est.	P value	Est.	P value
Intercept	9.92	0.000	10.00	0.001	8.96	0.000	9.19	0.000
Log GDP*	-0.69	0.001	-0.68	0.02	-0.66	0.000	-0.64	0.009
Educ [†]	-0.01	0.43	-0.01	0.22	-0.01	0.40	-0.02	0.16
PEH [‡]	-0.00	0.81	-0.00	0.67	-0.00	0.60	-0.01	0.32
Finance [§]	-	-	0.01	0.95	-	-	0.09	0.63
Refer	-	-	-0.38	0.16	-	-	-0.44	0.07
Primpay [¶]	-	-	0.46	0.006	-	-	0.40	0.02
Rho	0.91	-	0.90	-	0.87	-	0.86	-
R ² (overall)	0.21	-	0.32	-	0.23	-	0.34	-
No.	63	-	56	-	63	-	56	-

(b) Lung cancer

Independent variables	Male				Female			
	Model 1		Model 2		Model 1		Model 2	
	Est.	P value	Est.	P value	Est.	P value	Est.	P value
Intercept	5.55	0.000	5.80	0.000	-0.54	0.51	-0.63	0.52
Log GDP*	-0.18	0.02	-0.22	0.02	0.28	0.002	0.28	0.008
Educ [†]	-0.01	0.008	-0.01	0.006	0.02	0.004	0.02	0.01
PEH [‡]	0.00	0.31	0.01	0.051	-0.00	0.78	-0.00	0.91
Finance [§]	-	-	-0.19	0.24	-	-	0.03	0.89
Refer	-	-	0.11	0.52	-	-	0.11	0.57
Primpay [¶]	-	-	0.26	0.09	-	-	0.11	0.51
Rho	0.96	-	0.96	-	0.94	-	0.95	-
R ² (overall)	0.08	-	0.19	-	0.34	-	0.38	-
No.	62	-	55	-	63	-	56	-

(c) Colon cancer

Independent variables	Male				Female			
	Model 1		Model 2		Model 1		Model 2	
	Est.	P value	Est.	P value	Est.	P value	Est.	P value
Intercept	0.73	0.59	0.14	0.93	2.22	0.048	1.75	0.19
Log GDP*	0.19	0.10	0.23	0.15	-0.00	0.98	0.03	0.78
Educ [†]	0.00	0.93	-0.01	0.44	0.01	0.12	-0.00	0.69
PEH [‡]	0.01	0.06	0.01	0.03	0.01	0.07	0.01	0.06
Finance [§]	-	-	-0.02	0.85	-	-	0.12	0.28
Refer	-	-	0.23	0.14	-	-	0.18	0.16
Primpay [¶]	-	-	0.24	0.07	-	-	0.31	0.01
Rho	0.87	-	0.85	-	0.84	-	0.81	-
R ² (overall)	0.34	-	0.43	-	0.37	-	0.49	-
No.	63	-	56	-	63	-	56	-

*Log transformed GDP per capita (US dollars, current prices and PPPs); [†]Attainment at the tertiary level (% of population with educational level of some college); [‡]Public health expenditure (% total expenditure on health); [§]Health financing (countries which use tax-finance as their major method of health financing were coded to 1 and other countries [use social health insurance or private health insurance, instead] were coded to 0); ^{||}Referral to secondary care (countries in which referral is compulsory or financially encouraged were coded to 1 and other countries [neither compulsory nor financially encouraged] were coded to 0); [¶]Payment to primary care physicians (countries which use fee-for-service as their predominant mode were coded to 1 and other countries [use salary or capitation, or combination of these with fee-for-service] were coded to 0).

respectively (unpublished observations).

There are some arguments that the rising incidence of thyroid cancer cannot be attributed solely to improved detection because large thyroid cancers have also increased (32, 33) and radiation exposure to CT scans has rapidly increased (34). Nev-

ertheless, these observations do not necessarily rule out the possibility of overdiagnosis which is more likely to occur in less regulated markets. Moreover, the rapid increase in CT use in itself may have to do with the very healthcare system that permits the overdiagnosis of thyroid cancer.

Meanwhile, unlike the incidence of thyroid cancer, incidences of other cancers and mortality of thyroid cancer have no significant associations with the share of public health expenditure in our analysis. Rather, in the case of colorectal cancer, the share of public health expenditure has a significantly positive association with its incidence for males. Since screening for colorectal cancer is generally recommended, it is highly likely that countries with a high share of public health expenditure have well-coordinated screening programs nationwide, which naturally leads to early detection of colorectal cancer.

It is suggested that higher income not only improves access to appropriate care, but also increases the risk of receiving excessive care (35). Our analysis revealed that a high GDP per capita was associated with a high incidence of thyroid cancer, one of good prognosis cancers, while the mortality of thyroid cancer was negatively associated. These results correspond with earlier studies, which reported that the GDP per capita was positively associated with the incidence of good prognosis cancers, such as prostate and colorectal cancer, but negatively associated with ovarian and oral cavity cancer (36, 37).

There are some unexpected findings that should be investigated in further studies. First, countries with no referral system showed a higher mortality of thyroid cancer for females but had no significant differences in the incidences of thyroid, stomach, lung, and colorectal cancer. The role of the physician-gatekeeper can be defined from two perspectives: protecting patients from possible adverse effects of unnecessary care, and ensuring the appropriate use of healthcare services (35). This physician-gatekeeper role does not seem to have been proven in our analysis. Second, countries that use fee-for-service as their predominant mode of payment to primary care physicians showed a higher incidence of stomach cancer for both genders, and a higher incidence of colorectal cancer for females, but no significant association with the incidence of thyroid cancer. It seems that the influences of payment to primary care physicians on health outcomes were not coherent across indicators and models (38, 39).

There are some limitations in this study, which come mainly from the limited availability of data. First, the most important characteristic of healthcare systems that influences the overdiagnosis of thyroid cancer is the number of ultrasounds per capita or the number of check-ups with ultrasound per capita, but they were not available in the published international data. Second, regarding the premise that commercialization of the healthcare system means more dependence on the private sector, it could be reflected in healthcare services as well as in health financing. Thus, we considered including the proportion of beds in publicly owned hospitals among all hospital beds in our model, which was not possible due to a lot of missing data. Third, to investigate the possible relationship between the rising incidence and the healthcare systems, use of aggregate data on the incidence of thyroid cancer at the country level was inevitable.

If available, however, information on how each case of thyroid cancer was diagnosed at the individual level would help figure out the extent of overdiagnosis, which calls for further studies.

To our knowledge, this study is the first to compare the incidence of thyroid cancer across countries with different payment programs and health financing. The findings show that the upward trend of the incidence of thyroid cancer is closely related to the healthcare system that permits overdiagnosis. Therefore, it is suggested that increases in the proportion of public financing may help reduce the overdiagnosis of thyroid cancer to some extent.

REFERENCES

- Zheng TZ, Holford TR, Chen YT, Ma JZ, Flannery J, Liu W, Russi M, Boyle P. *Time trend and age-period-cohort effect on incidence of thyroid cancer in Connecticut, 1935-1992. Int J Cancer* 1996; 67: 504-9.
- Howlander N, Noone AM, Krapcho M, Neyman N, Aminou R, Waldron W, Altekruse SE, Kosary CL, Ruhl J, Tatalovich Z, et al. *SEER Cancer Statistics Review, 1975-2008, National Cancer Institute. Bethesda, MD, http://seer.cancer.gov/csr/1975_2008/, based on November 2010 SEER data submission, posted to the SEER web site, 2011 [accessed on 5 August 2012].*
- Kilfoy BA, Zheng T, Holford TR, Han X, Ward MH, Sjodin A, Ahang Y, Bai Y, Zhu C, Guo GL, et al. *International patterns and trends in thyroid cancer incidence, 1973-2002. Cancer Causes Control* 2009; 20: 525-31.
- Truong T, Rougier Y, Dubourdieu D, Guihenneuc-Jouyaux C, Orsi L, Hémon D, Guénel P. *Time trends and geographic variations for thyroid cancer in New Caledonia, a very high incidence area (1985-1999). Eur J Cancer Prev* 2007; 16: 62-70.
- National Cancer Information Center of Korea. *Statistics about thyroid cancer. Available at http://www.cancer.go.kr/ncic/index.html [accessed on 7 August 2011].*
- Weiss W. *Changing incidence of thyroid cancer. J Natl Cancer Inst* 1979; 62: 1137-42.
- Kazakov VS, Demidchik EP, Astakhova LN. *Thyroid cancer after Chernobyl. Nature* 1992; 359: 21.
- Ron E, Lubin JH, Shore RE, Mabuchi K, Modan B, Pottern LM, Schneider AB, Tucker MA, Boice JD Jr. *Thyroid cancer after exposure to external radiation: a pooled analysis of seven studies. Radiat Res* 1995; 141: 259-77.
- Golding SJ, Shrimpton PC. *Radiation dose in CT: are we meeting the challenge? Br J Radiol* 2002; 75: 1-4.
- Brenner D, Elliston C, Hall E, Berdon W. *Estimated risks of radiation-induced fatal cancer from pediatric CT. AJR Am J Roentgenol* 2001; 176: 289-96.
- Davies L, Welch HG. *Increasing incidence of thyroid cancer in the United States, 1973-2002. JAMA* 2006; 295: 2164-7.
- Kent WD, Hall SF, Isotalo PA, Houlden RL, George RL, Groome PA. *Increased incidence of differentiated thyroid carcinoma and detection of subclinical disease. CMAJ* 2007; 177: 1357-61.
- Lloyd R, De Lellis R, Heitz P, Eng C. *World Health Organization classification of tumours: pathology and genetics of tumours of the endocrine organs. Lyon, France: IARC Press, 2004.*
- Ghori FY, Gutterman-Litofsky DR, Jamal A, Yeung SC, Arem R, Sherman SI. *Socioeconomic factors and the presentation, management, and out-*

- come of patients with differentiated thyroid carcinoma. *Thyroid* 2002; 12: 1009-16.
15. Sprague BL, Warren Andersen S, Trentham-Dietz A. *Thyroid cancer incidence and socioeconomic indicators of health care access. Cancer Causes Control* 2008; 19: 585-93.
 16. Etzioni R, Penson DE, Legler JM, di Tommaso D, Boer R, Gann PH, Feuer EJ. *Overdiagnosis due to prostate-specific antigen screening: lessons from U.S. prostate cancer incidence trends. J Natl Cancer Inst* 2002; 94: 981-90.
 17. Welch HG, Black WC. *Overdiagnosis in cancer. J Natl Cancer Inst* 2010; 102: 605-13.
 18. Harach HR, Franssila KO, Wasenius VM. *Occult papillary carcinoma of the thyroid. A "normal" finding in Finland. A systematic autopsy study. Cancer* 1985; 56: 531-8.
 19. Macinko J, Starfield B, Shi L. *The contribution of primary care systems to health outcomes within Organization for Economic Cooperation and Development (OECD) countries, 1970-1998. Health Serv Res* 2003; 38: 831-65.
 20. Elola J, Daponte A, Navarro V. *Health indicators and the organization of health care systems in western Europe. Am J Public Health* 1995; 85: 1397-401.
 21. Andersen R, Newman JF. *Societal and individual determinants of medical care utilization in the United States. Milbank Mem Fund Q Health Soc* 1973; 51: 95-124.
 22. Quaglia A, Vercelli M, Lillini R, Mugno E, Coebergh JW, Quinn M, Martinez-Garcia C, Capocaccia R, Micheli A; ELDCARE Working Group. *Socio-economic factors and health care system characteristics related to cancer survival in the elderly. A population-based analysis in 16 European countries (ELDCARE project). Crit Rev Oncol Hematol* 2005; 54: 117-28.
 23. Nixon J, Ulmann P. *The relationship between health care expenditure and health outcomes. Eur J Health Econ* 2006; 7: 7-18.
 24. Asiskovitch S. *Gender and health outcomes: the impact of healthcare systems and their financing on life expectancies of women and men. Soc Sci Med* 2010; 70: 886-95.
 25. Babazono A, Hillman AL. *A comparison of international health outcomes and health care spending. Int J Technol Assess Health Care* 1994; 10: 376-81.
 26. Enthoven AC. *The history and principles of managed competition. Health Aff (Millwood)* 1993; 12: 24-48.
 27. Ferlay J, Bray F, Pisani P, Parkin DM. *GLOBOCAN 2002 v2.0. in cancer incidence, mortality and prevalence worldwide. Lyon, France: IARC Press, 2004.*
 28. Ferlay J, Shin HR, Bray F, Forman D, Mathers C, Parkin DM. *GLOBOCAN 2008 v1.2. in cancer incidence, mortality and prevalence worldwide. Lyon, France: IARC Press, 2010.*
 29. Organization for Economic Co-operation and Development. *OECD Health Data 2010 – Frequently Requested Data (online database: <http://www.oecd.org/>, based on Oct 2010 ver). Paris: OECD, 2010.*
 30. Paris V, Devaux M, Wei L. *Health systems institutional characteristics: a survey of 29 OECD countries. Paris: OECD, 2010.*
 31. Wagstaff A. *Social health insurance vs. tax-financed health systems: evidence from the OECD. Washington, DC: World Bank, 2009.*
 32. Chen AY, Jemal A, Ward EM. *Increasing incidence of differentiated thyroid cancer in the United States, 1988-2005. Cancer* 2009; 115: 3801-7.
 33. Morris LG, Myssiorek D. *Improved detection does not fully explain the rising incidence of well-differentiated thyroid cancer: a population-based analysis. Am J Surg* 2010; 200: 454-61.
 34. Baker SR, Bhatti WA. *The thyroid cancer epidemic: is it the dark side of the CT revolution? Eur J Radiol* 2006; 60: 67-9.
 35. Franks P, Clancy CM, Nutting PA. *Gatekeeping revisited: protecting patients from overtreatment. N Engl J Med* 1992; 327: 424-9.
 36. Grant WB. *A multicountry ecological study of risk-modifying factors for prostate cancer: apolipoprotein E epsilon4 as a risk factor and cereals as a risk reduction factor. Anticancer Res* 2010; 30: 189-99.
 37. Chen DT, Chou YF, Wu HP, Hsu LP, Wen IS, Lee CF, Chen PR. *Income and the incidence of oral cavity cancer: cross-national study. J Otolaryngol Head Neck Surg* 2009; 38: 208-11.
 38. Or Z. *Exploring the effects of health care on mortality across OECD countries. Paris: OECD, 2001.*
 39. Or Z, Wang J, Jamison D. *International differences in the impact of doctors on health: a multilevel analysis of OECD countries. J Health Econ* 2005; 24: 531-60.