

Lipid Droplets and Peroxisomes: Key Players in Cellular Lipid Homeostasis or A Matter of Fat—Store 'em Up or Burn 'em Down

Sepp D. Kohlwein,^{*,1} Marten Veenhuis,[†] and Ida J. van der Klei^{†,1}

^{*}Institute of Molecular Biosciences, University of Graz, 8010 Graz, Austria, and [†]Molecular Cell Biology, University of Groningen, 9700CC Groningen, The Netherlands

ABSTRACT Lipid droplets (LDs) and peroxisomes are central players in cellular lipid homeostasis: some of their main functions are to control the metabolic flux and availability of fatty acids (LDs and peroxisomes) as well as of sterols (LDs). Both fatty acids and sterols serve multiple functions in the cell—as membrane stabilizers affecting membrane fluidity, as crucial structural elements of membrane-forming phospholipids and sphingolipids, as protein modifiers and signaling molecules, and last but not least, as a rich carbon and energy source. In addition, peroxisomes harbor enzymes of the malic acid shunt, which is indispensable to regenerate oxaloacetate for gluconeogenesis, thus allowing yeast cells to generate sugars from fatty acids or nonfermentable carbon sources. Therefore, failure of LD and peroxisome biogenesis and function are likely to lead to deregulated lipid fluxes and disrupted energy homeostasis with detrimental consequences for the cell. These pathological consequences of LD and peroxisome failure have indeed sparked great biomedical interest in understanding the biogenesis of these organelles, their functional roles in lipid homeostasis, interaction with cellular metabolism and other organelles, as well as their regulation, turnover, and inheritance. These questions are particularly burning in view of the pandemic development of lipid-associated disorders worldwide.

TABLE OF CONTENTS

Abstract	1
Introduction	2
Lipid Droplets	2
<i>Experimental approaches to studying LD biology</i>	3
<i>Lipid droplet structure</i>	4
<i>Lipid composition of purified LDs</i>	4
<i>Protein composition of LDs</i>	5
<i>Biosynthesis of triacylglycerol and steryl esters</i>	5
<i>Turnover of lipid droplets</i>	10
<i>Regulation of neutral lipid synthesis</i>	11
<i>Regulation of neutral lipid degradation</i>	13

Continued

Copyright © 2013 by the Genetics Society of America

doi: 10.1534/genetics.112.143362

Manuscript received June 29, 2012; accepted for publication September 25, 2012

Available freely online through the author-supported open access option.

¹Corresponding authors: University of Graz, Humboldtstrasse 50/II, A8010 Graz, Austria. E-mail: sepp.kohlwein@uni-graz.at; University of Groningen, 9700CC Groningen, The Netherlands. E-mail: i.j.van.der.klei@rug.nl

CONTENTS, *continued*

<i>Mechanisms of LD biogenesis and inheritance</i>	14
<i>Contribution of acyl transferases to LD biogenesis</i>	14
<i>Targeting of proteins to LDs</i>	15
<i>Lipid droplet morphology and inheritance</i>	15
<i>Physiological role of LDs</i>	15
<i>Physiological interaction of LDs with other organelles</i>	16
<i>Yeast as a model to investigate and understand lipid-associated disorders</i>	17
Peroxisomes	18
<i>Peroxisome composition</i>	18
<i>Peroxisome metabolic functions</i>	18
<i>β-Oxidation:</i>	19
<i>Glyoxylate cycle:</i>	25
<i>Other peroxisomal enzymes:</i>	26
<i>Methods to identify proteins involved in peroxisome biology</i>	27
<i>Identification of peroxisomal enzymes:</i>	27
<i>Discovery of peroxins by genetic approaches:</i>	27
<i>Identification of peroxisomal proteins by organelle proteomics:</i>	28
<i>In silico prediction of peroxisomal proteins:</i>	28
<i>Transcriptome analysis:</i>	28
<i>Microscopic analysis of peroxisomes:</i>	29
Peroxisome Biogenesis	30
<i>Peroxisomal matrix protein import</i>	30
<i>Targeting signals and their receptors:</i>	30
<i>PTS1 and its receptor Pex5:</i>	30
<i>PTS2 and Pex7 with its coreceptors:</i>	31
<i>Piggy-back import:</i>	31
<i>Receptor docking site:</i>	31
<i>RING finger complex:</i>	32
<i>Translocation pore:</i>	33
<i>Receptor recycling and the role of ubiquitin:</i>	33
<i>Sorting and insertion of PMPs</i>	33
<i>Formation of peroxisomes from the ER</i>	35
<i>Peroxisome fission</i>	35
<i>Pex11 and the Fis1/Dnm1 fission machinery:</i>	35
<i>Role of the other Pex11 protein family members:</i>	36
<i>Pex23 protein family:</i>	36
<i>Peroxisome inheritance</i>	37
<i>Peroxisome degradation</i>	37
<i>Peroxisome function and biology in yeast as a paradigm for metabolic disorders in humans</i>	38
<i>Outlook and perspectives</i>	38

WORK for the past five decades on the yeast *Saccharomyces cerevisiae* has contributed fundamental insight into peroxisome biogenesis and function that is also relevant for mammalian cells. While LD research in yeast is still in its infancy and looks back to a much shorter history—the previous edition of *YeastBook* did not even mention LDs as an “organelle”—combined biochemical, cell biological, lipidomic, and proteomic studies in recent years have already

contributed significant insight into LD biogenesis and function.

Lipid Droplets

LDs, also termed “lipid particles,” “lipid bodies,” or “oil bodies,” are ubiquitous subcellular structures that have only in recent years been recognized as metabolically highly dynamic organelles (Daum *et al.* 2007a; Fujimoto *et al.*

2008; Goodman 2008, 2009; Guo *et al.* 2009; Kraemer *et al.* 2009; Murphy *et al.* 2009; Olofsson *et al.* 2009; Walther and Farese 2009, 2012; Athenstaedt and Daum 2011). In the past, LDs were primarily considered as rather inert storage depots for the “neutral lipids,” triacylglycerols (TAG) and steryl esters (SE). However, the increased biomedical interest in understanding neutral lipid homeostasis, fueled by the pandemic increase in lipid-associated disorders, has moved LDs into the spotlight of biomedical research (Farese and Walther 2009; Walther and Farese 2012). Given the significant homology of lipid biosynthetic processes to mammalian cells, yeast LD research has gained a great momentum to address the fundamental mechanisms of LD assembly and the regulation of neutral lipid homeostasis (Athenstaedt and Daum 2006, 2011; Czabany *et al.* 2007; Daum *et al.* 2007a,b; Rajakumari *et al.* 2008; Kohlwein 2010a,b).

Among subcellular organelles, LDs are unique in their structure, as they appear to harbor only a monolayer of phospholipids that surrounds the hydrophobic core consisting of TAG and SE. A second feature standing out is that LDs, like peroxisomes, are organelles that are not essential under standard nutritional conditions, *i.e.*, in the presence of carbon sources other than fatty acids (FA). Unlike other organelles, LD biogenesis and degradation need to be discussed in the context of the synthesis and turnover of their major components, namely neutral lipids: their biogenesis is driven by the availability of precursors for the synthesis of their core compounds, TAG and SE, and cells are devoid of LDs in the absence of the cellular capacity to synthesize these lipids (Garbarino *et al.* 2009; Petschnigg *et al.* 2009). On the other hand, TAG synthesis—and concomitant formation of LDs—is essential for cell survival in the presence of excess FAs (Garbarino *et al.* 2009; Petschnigg *et al.* 2009; Fakas *et al.* 2011b). The LD surface is decorated with numerous proteins that are, in part, also present in the endoplasmic reticulum (ER) membrane, raising the question as to the specific signals that target proteins to the LD surface. The highly dynamic nature of LDs in growing cells reflects the importance of neutral lipids in various stages of cell growth and in response to the nutritional status of the cell; the metabolic role of LDs is highlighted by the recent discoveries that TAG-derived metabolites are required for efficient cell cycle progression (Kurat *et al.* 2009) and that TAG play an essential role in counteracting FA-induced lipotoxicity (Garbarino *et al.* 2009; Petschnigg *et al.* 2009; Fakas *et al.* 2011b).

Experimental approaches to studying LD biology

In vivo, LDs are readily detectable by transmission light microscopy (differential interference contrast (DIC; Nomarski optics) due to their high refractive index (Figure 1). Numerous cell-permeable hydrophobic fluorescence dyes that label LDs with high specificity, including Nile Red, LD540, and BODIPY dyes, exist (Szymanski *et al.* 2007; Fei *et al.* 2008; Wolinski and Kohlwein 2008; Spandl *et al.* 2009; Wolinski *et al.* 2011, 2012). It should be noted, however, that these

dyes are potential substrates of the pleiotropic drug resistance pumps (Ivnitski-Steele *et al.* 2009), and staining efficiency may strongly depend on the activity of these pumps in the respective strain backgrounds. Thus, staining of LDs in growing cultures that contain both young and aged cells may appear quite heterogeneous; fixation of cells with formaldehyde or elimination of Pdr pumps strongly increases labeling efficiency (Wolinski and Kohlwein 2008; Wolinski *et al.* 2009a, 2012). Given the specificity and ease of labeling of both living and fixed cells, several microscopy- or photometry-based screens of yeast mutant collections have been performed to identify mutants with altered LD morphology and content (Szymanski *et al.* 2007; Fei *et al.* 2008; Bozaquel-Morais *et al.* 2010; Adeyo *et al.* 2011; Fei and Yang 2012). In addition, green fluorescent protein-tagged reporter constructs of LD-associated proteins provide an additional tool for studying LD dynamics and inheritance (Kurat *et al.* 2006; Jacquier *et al.* 2011; Wolinski *et al.* 2012). It should be noted that the number and size of LDs vary greatly between various yeast wild-type strains, and it is currently unclear which genetic traits are responsible for this heterogeneity. Microscopy-based screens of the GFP-labeled protein collection (Huh *et al.* 2003) have also led to the identification of numerous novel LD-associated proteins (Natter *et al.* 2005; see below). In addition to the use of hydrophobic fluorescent dyes, recent advances in spectroscopic imaging techniques such as coherent anti-Stokes Raman scattering (CARS) microscopy allow the label-free imaging of yeast LDs (Brackmann *et al.* 2009; Kohlwein 2010b; Wolinski *et al.* 2012). This technology is based on the C-H molecular vibrations in the FA acyl chains that are packed in high density as TAG in the LD, and thus independent of exogenously supplied fluorescent dyes or endogenously expressed fluorescent protein reporter constructs (see also Figure 1).

Higher resolution images of LDs are obtained by electron microscopy and electron tomography (Binns *et al.* 2006; Perktold *et al.* 2007; Czabany *et al.* 2008; Jacquier *et al.* 2011; Wolinski *et al.* 2011) (Figure 1), which also demonstrate their close physical interactions with other intracellular organelles, in particular the ER, mitochondria, and peroxisomes (Binns *et al.* 2006; Pu *et al.* 2011). Biophysical studies on isolated LDs have been performed using X-ray small-angle scattering analyses, dynamic light scattering, and differential scanning calorimetry to unveil LD size distribution and structural organization, depending on lipid composition (Czabany *et al.* 2008; Spanova *et al.* 2012; see below).

LD can be purified from cellular extracts by ultracentrifugation/flotation (Leber *et al.* 1994; Athenstaedt *et al.* 1999; Connerth *et al.* 2009). Since LDs are in close contact with other intracellular organelles, a clean LD preparation requires careful cell lysis (*e.g.*, enzymatic digestion of the cell wall with Zymolyase), followed by differential centrifugation, to obtain a layer of LDs—together with vacuolar membranes—floating on top of the centrifuge tube. Attached vacuolar membranes are separated by an additional centrifugation




Figure 1 Morphological characteristics of yeast lipid droplets. (Rows 1–3, left panels) Fluorescence images of LDs that are labeled with BODIPY 493/503 (Wolinski and Kohlwein 2008; Wolinski *et al.* 2009a, 2012). (Right panels) Corresponding transmission images. All strains, except the *fld1Δ* mutant, were cultivated for 72 h in YPD complete medium; *fld1Δ* mutants were grown in synthetic complete (minimal) medium with 2mg/liter inositol for 12 h. Images were obtained by confocal laser scanning microscopy and represent projections of 8–12 optical sections. wt, wild type; *tgl3Δ tgl4Δ*, mutant lacking the major TAG lipases; *are1Δ are2Δ*, mutant lacking the steryl ester synthases and thus harboring LDs that contain TAG only; *dga1Δ lro1Δ*, mutant lacking acyl-CoA and phospholipid-dependent diacylglycerol (DAG) acyltransferases and thus harboring LDs that contain SE only; *fld1Δ* mutant, lacking the yeast ortholog of seipin. TEM: transmission electron microscopy images of wild type (wt) and the *fld1Δ* mutant. (Row 4) Electron tomography (ET) of LDs in wild type, showing close association of LDs with the ER membrane. CARS: coherent anti-Stokes Raman scattering microscopy of LDs in wild type, *tgl3Δ tgl4Δ* mutant, and *dga1Δ lro1Δ* mutant. CARS is a label-free imaging technique that generates contrast by imaging molecular vibrations at 2840 cm^{-1} . Scale bar: 500 nm in the TEM images, 200 nm in the ET image, and 5 μm in the fluorescence/transmission images. See text for details. Images courtesy of H. Wolinski (fluorescence and CARS microscopy) and D. Kolb (electron microscopy and tomography).

step that requires a pH/buffer change (Athenstaedt *et al.* 1999; Connerth *et al.* 2009; Grillitsch *et al.* 2011). This protocol restricts LD preparations to cells in late log/early stationary phase that are susceptible to Zymolyase lysis of the cell wall.

Lipid droplet structure

LDs isolated from yeast are rather homogeneous in size, ranging from typically 300 nm (in late log phase) to 1 μm (in stationary phase) in diameter. In the late log/early stationary phase of growth, the majority of LDs fall into a rather narrow 350- to 450-nm size range, largely independent of their lipid composition (Czabany *et al.* 2008). As discussed below, LDs are subject to high metabolic turnover and may be almost completely degraded during the early log phase of growth (Kurat *et al.* 2006). Wild-type LDs containing about equal amounts of TAG and SE are typically spherical structures, in which the core of neutral lipids is surrounded by a monolayer of phospholipids, which, according to the current biogenesis models, is derived from the ER membrane

(Mechanisms of LD biogenesis and inheritance). X-ray small-angle scattering experiments have unveiled some level of supramolecular organization of LDs, indicating that SE form a shell surrounding the rather fluid disordered TAG core (Czabany *et al.* 2008). A *hem1* mutant that is defective in sterol synthesis accumulates the sterol intermediate squalene in LDs, which is found in subcellular membranes as well as in LD and leads to a disordering of the shell structure (Spanova *et al.* 2012).

Lipid composition of purified LDs

The major lipid components of LDs are the neutral lipids, TAG and SE (Zinser *et al.* 1991; Leber *et al.* 1994; Connerth *et al.* 2009). In mammalian cells, LD composition may vary, depending on cell type, and contain mostly TAG (as in adipocytes) or TAG and cholesteryl esters, retinylesters, and free cholesterol (as in liver). It should be emphasized that the designation “neutral lipid” of these compounds refers to their uncharged and highly hydrophobic structure, but not to their (active) involvement in cellular metabolism. TAG

and SE are present in about equal amounts in LDs (Daum *et al.* 2007b; Czabany *et al.* 2008; Rajakumari *et al.* 2008; Connerth *et al.* 2009; Grillitsch *et al.* 2011). The TAG molecular species distribution reflects the cellular content of long-chain FAs, namely predominantly C16 and C18 saturated and mono-unsaturated FAs, giving rise to the most prominent 48:2, 50:2, 50:3, 52:2, and 52:3 TAG molecular species¹ in wild-type cells grown on glucose (Connerth *et al.* 2009; Grillitsch *et al.* 2011). Somewhat different TAG profiles were obtained for cells grown on raffinose (Ejsing *et al.* 2009). This finding also reflects the dynamic nature of LDs [and the entire yeast lipidome for that matter (Klose *et al.* 2012)] that respond quickly to growth rate and carbon source. Accordingly, growth of yeast in the presence of oleic acid as the sole carbon source results in TAG species predominantly composed of TAG 54:3 (Grillitsch *et al.* 2011). The SE fraction is mainly composed of ergosterol esterified with oleic acid (C18:1) and palmitoleic acid (Czabany *et al.* 2008), but sterol intermediates, such as zymosterol, episterol, and fecosterol are also found esterified in the SE fraction (Zweytick *et al.* 2000b; Czabany *et al.* 2008).

The phospholipid monolayer of LDs is enriched in the anionic phospholipid, phosphatidylinositol, compared to total cellular phospholipids (Schneider *et al.* 1999; Connerth *et al.* 2009; Grillitsch *et al.* 2011); notably, the molecular species distribution of LD phospholipids is quite distinct from that of the ER membrane, from which it is presumably derived (Connerth *et al.* 2009; Grillitsch *et al.* 2011), and appears to be enriched in double-unsaturated species (Schneider *et al.* 1999). Notably, phosphatidylinositol molecular species with medium-chain fatty acids (C12 and C14), which are quite prominent in subcellular membranes (Ejsing *et al.* 2009; Klose *et al.* 2012), are excluded from the LD phospholipid monolayer (Schneider *et al.* 1999).

Protein composition of LDs

Although LDs are present in almost all cell types, ranging from bacteria to mammals, their protein composition is rather divergent (Murphy 2001; Yang *et al.* 2012). The proteome of highly purified LDs from yeast is composed of a characteristic set of proteins, but the overall protein content is rather low (Table 1). Notably, most of the LD-resident enzymes identified so far play a role in lipid metabolism, emphasizing the active role of this organelle in cellular metabolism (Athenstaedt and Daum 2006; Czabany *et al.* 2007; Daum *et al.* 2007a,b; Rajakumari *et al.* 2008; Kohlwein 2010b; Grillitsch *et al.* 2011). Notably, the set of LD-associated proteins may substantially change during cellular growth, in particular if FAs such as oleic acid are supplied, to induce formation of LDs and peroxisomes (Grillitsch *et al.* 2011). Also, size and phospholipid composition that are dependent on growth conditions and media composition (e.g., presence

or absence of the phospholipid precursor inositol) may influence the LD proteome (Fei *et al.* 2011c). Numerous LD-associated proteins display a dual localization also to the ER (Table 1; see below), and their relative distribution to both organelles may change during various stages of growth. Since LDs closely interact with other subcellular organelles, some of the identified proteins may actually be contaminants during preparation. On the other hand, a transient association of non-LD-resident proteins may also be of physiological significance (see *Physiological role of LDs*): evidence suggests that association of proteins with LDs, at least in mammalian cells, may serve a protective or regulatory role (Hodges and Wu 2010).

Notably, yeast LDs do not contain proteins related to the perilipin family of proteins in mammals (Brasaemle 2007) or oleosins in plants (Chapman *et al.* 2012). Perilipins are prominent LD surface proteins that regulate the access of enzymes to the LD surface during lipogenesis or lipolysis (Brasaemle 2007). Oleosins and related proteins are characteristically shaped proteins that reside on the surface of oil droplets in plant seeds and nonseed tissues and play a role in stress response, hormone signaling, and plant growth and development (Chapman *et al.* 2012). Both types of surface proteins are believed to play important roles in LD biogenesis and structure and lipid mobilization; thus, the question remains how the size of LDs and processes acting on TAG and SE substrates are regulated in yeast in the absence of such LD coat proteins.

Biosynthesis of triacylglycerol and sterol esters

Formation of LD is driven by the synthesis of TAG and SE; in the absence of the biosynthetic capacity to form these lipids, no LD are present and LD-resident proteins may mis-localize to the ER or other intracellular structures and the cytosol (Athenstaedt and Daum 2006, 2011; Daum *et al.* 2007a,b; Rajakumari *et al.* 2008; Garbarino *et al.* 2009; Petschnigg *et al.* 2009; Jacquier *et al.* 2011). The enzymes involved in TAG and SE metabolism are listed in Table 1 (see also Henry *et al.* 2012).

The major substrates for the synthesis of TAG and SE are activated FAs and glycerol-3-phosphate or dihydroxyacetone phosphate (DHAP) and sterols, respectively. The first and rate-limiting step in FA synthesis is catalyzed by acetyl-CoA carboxylase, encoded by *ACC1* (Roggenkamp *et al.* 1980; Al-Feel *et al.* 1992; Hasslacher *et al.* 1993; Tehlivets *et al.* 2007; Henry *et al.* 2012) (Figure 2). *Acc1* converts acetyl-CoA to malonyl-CoA in an ATP, biotin, and CO₂-dependent reaction. Malonyl-CoA is used by FA synthase, which consists of a hexameric $\alpha_6\beta_6$ complex of two subunits encoded by *FAS2* (α -subunit) and *FAS1* (β -subunit) for the step-wise elongation of the growing acyl chain (Tehlivets *et al.* 2007). In contrast to mammalian FA synthase that releases free FAs, the yeast FAS complex generates acyl-CoAs that may be directly channeled into phosphatidic acid (PA), TAG and SE synthesis (Tehlivets *et al.* 2007; Henry *et al.* 2012). FA *de novo* synthesis is a major consumer of acetyl-CoA and

¹The numbers indicate the total number of carbon atoms in the acyl chains and the number of double bonds. Yeast produces only mono-unsaturated FAs; thus TAG 52:3 indicates a species containing C18:1 + C18:1 + C16:1 acyl chains.

Table 1 Neutral lipid metabolism enzymes and LD-associated proteins

Gene	Enzyme	Molecular mass (kDa)	Isoelectric point	Molecules per cell ^a	Location ^b	Transmembrane domains	Phosphorylation sites ^c
Neutral lipid synthesis enzymes							
<i>SCT1 (GAT2)</i>	Glycerol-3-P/dihydroxyacetone-P acyltransferase	85.7	7.27	1,050	ER	4	Few
<i>GPT2 (GAT1)</i>	Glycerol-3-P/dihydroxyacetone-P acyltransferase	83.6	10.3	3,100	ER, lipid droplets	4	Several
<i>AYR1</i>	Acyl DHAP reductase	32.8	9.92	3,670	ER, lipid droplets	None	None
<i>SLC1</i>	LysoPA/Acylglycerol-3-P acyltransferase	33.8	10.41	ND ^d	ER, lipid droplets	1	None
<i>ALE1 (SLC4, LPT1, LCA1)</i>	LysoPA/Acylglycerol-3-P acyltransferase	72.2	10.3	ND	ER	7	Several
<i>PHM8</i>	LysoPA phosphatase	37.7	5.14	195	Cytoplasm, nucleus	None	None
<i>LOA1 (VPS66)</i>	LysoPA acyltransferase	33.8	10.5	6,630	ER, lipid droplets	1	None
<i>PAH1 (SIMP2)</i>	PA phosphatase	95	4.68	3,910	Cytoplasm, ER	None	Several
<i>DGK1 (HSD1)</i>	DAG kinase	32.8	9.48	784	ER	4	Few
<i>DGA1</i>	Acyl-CoA diacylglycerol acyltransferase	47.7	10.39	907	ER, lipid droplets	1	Few
<i>LRO1</i>	Phospholipid diacylglycerol acyltransferase	75.3	6.67	ND	ER	1	Few
<i>ARE1 (SAT2)</i>	Acyl-CoA sterol acyltransferase	71.6	8.27	ND	ER	9	Several
<i>ARE2 (SAT1)</i>	Acyl-CoA sterol acyltransferase	74.0	7.71	279	ER	9	Several
Neutral lipid turnover enzymes							
<i>LDH1</i>	Triacylglycerol lipase, hydrolase	43.3	6.51	ND	Lipid droplets	None	None
<i>TGL1 (YKL5)</i>	Triacylglycerol lipase, steryl ester hydrolase	63.0	6.83	1,470	ER, lipid droplets	1	Several
<i>TGL2</i>	Acylglycerol lipase	37.5	8.41	ND	Mitochondria	None	None
<i>TGL3</i>	Triacylglycerol lipase, lysoPA acyltransferase	73.6	8.50	3,210	Lipid droplets	1	Few
<i>TGL4 (STC1)</i>	Triacylglycerol lipase, Ca ⁺⁺ dependent phospholipase A2, lysoPA acyltransferase	102.7	8.05	195	Lipid droplets	None	Several
<i>TGL5 (STC2)</i>	Triacylglycerol lipase, lysoPA acyltransferase	84.7	9.84	358	Lipid droplets	1	Several
<i>YEH1</i>	Steryl ester hydrolase	66.5	6.33	7,770	Lipid droplets	1-2	None
<i>YEH2</i>	Steryl ester hydrolase	62.4	8.91	1,630	Plasma membrane	1	Few
<i>YU3</i>	Monoacylglycerol lipase	35.6	8.5	2,140	Lipid droplets, ER	None	None
Lipid droplet-associated proteins^d							
<i>ATF1</i>	Alcohol-O-acetyltransferase	61.0	6.94	1,990	Lipid droplets	None	None
<i>AYR1</i>	Acyl DHAP reductase	32.8	9.92	3,670	ER, lipid droplets	None	None
<i>BSC2</i>	Unknown function	26.6	5.10	922	Lipid droplets	1	None
<i>COY1</i>	Unknown function, similarity to mammalian CASP	77.5	5.93	2,650	Lipid droplets, Golgi	1	Several
<i>CPP5 (CYP5)</i>	Peptidyl-prolyl <i>cis-trans</i> isomerase (cyclophilin)	25.3	5.40	ND	Lipid droplets, ER	1	1
<i>CSR1 (SFH2)</i>	Phosphatidylinositol transfer protein	47.5	6.61	9,600	Lipid droplets, cytoplasm	None	1
<i>CST26 (PS1)</i>	Stearyl-CoA acyltransferase, phosphatidylinositol-specific	45.5	10.15	2,010	Lipid droplets, ER	Four	None
<i>EHT1</i>	Acyl-coenzymeA ethanol <i>O</i> -acyltransferase	51.3	7.83	2,550	Lipid droplets, ER	None	None
<i>ENV9</i>	Unknown function, similarity to oxidoreductases	37.5	8.35	967	Lipid droplets	1	None
<i>ERG1</i>	Squalene epoxidase	55.1	6.45	65,400	Lipid droplets, ER	2	None
<i>ERG27</i>	3-Keto sterol reductase	39.7	8.90	ND	Lipid droplets	None	1
<i>ERG6 (ISE1, LIS1, SED6, VID1)</i>	Δ^24 -Sterol C-methyltransferase	43.3	5.60	53,800	Lipid droplets, ER	None	Few
<i>ERG7</i>	Lanosterol synthase	83.5	6.59	2,190	Lipid droplets, ER	None	None

(continued)

Table 1, continued

Gene	Enzyme	Molecular mass (kDa)	Isoelectric point	Molecules per cell ^a	Location ^b	Transmembrane domains	Phosphorylation sites ^c
<i>FAA1</i>	Fatty acyl-CoA synthetase	77.8	7.58	7,470	Lipid droplets, ER	None	None
<i>FAA4</i>	Fatty acyl-CoA synthetase	77.2	6.52	31,200	Lipid droplets, ER	None	None
<i>FAT1</i>	Fatty acid transporter and fatty acyl-CoA synthetase	77.1	8.47	16,900	Lipid droplets, ER	3	None
<i>GTT1</i>	Glutathione S-transferase	26.8	6.65	ND	Lipid droplets, ER	None	None
<i>HFD1</i>	Hexadecenal dehydrogenase	60.0	6.73	2,930	Lipid droplets, mitochondria	1	1
<i>KE51 (LPI3, OSH4, BSR3)</i>	Member of the oxysterol binding protein family	49.5	5.92	32,200	Lipid droplets, cytoplasm	None	None
<i>LDB16</i>	Unknown function	29.0	8.07	149	Lipid droplets	2	1
<i>LOA1 (VPS66)</i>	LysoPA acyltransferase	33.8	10.5	6,630	Lipid droplets, ER	1	None
<i>NUS1</i>	Putative prenyltransferase	42.6	6.94	ND	Lipid droplets, ER ^e	1	None
<i>OSW5</i>	Unknown function	16.4	9.94	922	Lipid droplets	2	1
<i>PDI1 (MFP1, TRG1)</i>	Protein disulfide isomerase	58.2	4.22	ND	Lipid droplets, ER (lumen)	None	None
<i>PDR16 (SFH3)</i>	Phosphatidylinositol transfer protein	40.7	8.16	15,400	Lipid droplets, cytoplasm	None	2
<i>PET10</i>	Unknown function	31.2	8.85	2,160	Lipid droplets	None	None
<i>RRT8</i>	Unknown function	39.6	10.66	ND	Lipid droplets	5	None
<i>SLC1</i>	LysoPA/Acylglycerol-3-P acyltransferase	33.8	10.41	ND	Lipid droplets, ER	1	None
<i>SMA2</i>	Unknown function	9.2	6.50	20,400	Lipid droplets	2	1
<i>SNX41</i>	Sorting nexin	70.7	7.02	1,800	Lipid droplets, cytoplasm	None	None
<i>SRT1</i>	Cis-prenyl transferase	40.2	10.30	ND	Lipid droplets	1	None
<i>SSO1</i>	t-SNARE	33.1	5.06	450	Lipid droplets, vesicles	1	1
<i>TDH1 (GLD3)</i>	Glyceraldehyde-3-phosphate dehydrogenase, isozyme 1	35.8	8.59	120,000	Lipid droplets, cytoplasm	None	Several
<i>TDH2 (GLD2)</i>	Glyceraldehyde-3-phosphate dehydrogenase, isozyme 2	35.8	6.96	121,000	Lipid droplets, cytoplasm	None	Several
<i>TDH3 (GLD1, HSP35, HSP36, SSS2)</i>	Glyceraldehyde-3-phosphate dehydrogenase, isozyme 3	35.7	6.96	169,000	Lipid droplets, nucleus, cytoplasm	None	Several
<i>TGL1 (YKL5)</i>	Triacylglycerol lipase, steryl ester hydrolase	63.0	6.83	1,470	Lipid droplets, ER	1	Several
<i>TGL3</i>	Triacylglycerol lipase, lysoPA acyltransferase	73.6	8.50	3,210	Lipid droplets	1	Few
<i>TGL4</i>	Triacylglycerol lipase, Ca ⁺⁺ dependent phospholipase A2, lysoPA acyltransferase	102.7	8.05	195	Lipid droplets	None	Several
<i>TGL5</i>	Triacylglycerol lipase, lysoPA acyltransferase	84.7	9.84	358	Lipid droplets	1	Several
<i>UBX2 (SEL1)</i>	Bridging factor involved in ER-associated protein degradation (ERAD)	66.8	5.22	12,600	Lipid droplets, ER	None	None
<i>USE1 (SLT1)</i>	SNARE	28.1	5.01	973	Lipid droplets, ER	1	None
<i>YEH1</i>	Steryl ester hydrolase	66.5	6.33	7,770	Lipid droplets	1-2	None
<i>YIM1</i>	Unknown function	41.6	7.98	6,540	Lipid droplets, ER	None	None
<i>YIU3</i>	Monoacylglycerol lipase	35.6	8.5	2,140	Lipid droplets, ER	None	None
<i>YPT7 (AST4, VAM4)</i>	Rab family GTPase	23.0	4.62	5,530	Lipid droplets, cytoplasm	None	None
<i>YOR059c</i>	Putative lipase	51.1	9.81	1,210	Lipid droplets	1	None

Much of the information in this table may be found in the *Saccharomyces* Genome Database. ND, not determined.

^a Ghaemmaghami et al. (2003).

^b Habeler et al. (2002); Grillitsch et al. (2011); Kumar et al. (2002); Huh et al. (2003); Natter et al. (2005); Athenstaedt et al. (1999).

^c Based on PhosphoGrid (<http://www.phosphogrid.org>) and PhosphoPed (<http://www.phosphopep.org>) Databases.

^d Grillitsch et al. (2011); Natter et al. (2005); Athenstaedt et al. (1999).

^e Prein et al. (2000).

NADPH, similar to sterol synthesis. Free FAs that are derived from exogenous supply or from endogenous lipid degradation need to be activated by one of five acyl-CoA synthetases, encoded by *FAA1*, *FAA2*, *FAA3*, *FAA4*, and *FAT1* genes, which differ in their substrate specificities (Black and Dirusso 2007). *Faa2* is required for the activation of FAs that are directed toward β -oxidation (see below). *Faa1*, *Faa4*, and *Fat1* activate exogenously supplied FAs and free FAs that derive from phospholipid, TAG, and SE breakdown. In the absence of these acyl-CoA synthetases, yeast secretes lipolysis-derived FAs (Scharnewski *et al.* 2008), and growth and membrane lipid composition depend solely on the FAs that are generated by *de novo* synthesis, FA desaturation, and elongation (Tehlivets *et al.* 2007).

The central intermediate in glycerolipid metabolism from which TAG and phospholipids are derived is PA (Athenstaedt and Daum 1997, 1999; Kohlwein 2010b) (Figure 2A). PA is synthesized by a two-step acylation reaction: first, glycerol-3-phosphate is acylated by *Sct1* and *Gpt2* acyltransferases to *sn1*-acylglycerol-3-phosphate (also termed lyso-PA) (Zheng and Zou 2001; Zaremborg and McMaster 2002). Alternatively, *Sct1* and *Gpt2* may also acylate dihydroxyacetone phosphate to 1-acyl-DHAP, which is subsequently reduced by the *Ayr1* reductase to *sn1*-acylglycerol-3-phosphate (Athenstaedt and Daum 2000). The acyltransferases and *Ayr* are predominantly localized to the ER membrane, but, notably, *Ayr* and *Gpt2* also partially localize to the LD (Athenstaedt *et al.* 1999; Athenstaedt and Daum 2000; Marr *et al.* 2012), indicating that at least the first steps in PA synthesis are also LD resident. *Gpt2* and *Sct1* acyltransferases exhibit different substrate specificities, giving rise to different populations of phospholipids and TAG molecular species (Zaremborg and McMaster 2002; Marr *et al.* 2012).

Sct1 and *Gpt2* generate lyso-PA, which is further acylated by the ER-resident *Slc1* and *Ale1* acyltransferases to *sn1,2*-diacylglycerol-3-phosphate (PA) (Benghezal *et al.* 2007; Chen *et al.* 2007; Jain *et al.* 2007; Riekhof *et al.* 2007; Henry *et al.* 2012). *Slc1* and *Ale1* are members of the MBOAT, the membrane-bound *O*-acyltransferase family of proteins and also involved in the Lands cycle of phospholipid acyl-chain remodeling (Hofmann 2000; Benghezal *et al.* 2007; Chen *et al.* 2007; Jain *et al.* 2007; Riekhof *et al.* 2007; Pagac *et al.* 2011). PA is the central glycerolipid intermediate that is utilized both for TAG and for phospholipid synthesis (for details see Henry *et al.* 2012). In addition to its role as glycerolipid precursor, PA also plays an important role in regulating cellular lipid metabolism (Henry *et al.* 2012), and its dephosphorylation to diacylglycerol (DAG) is a key step in driving LD formation (Adeyo *et al.* 2011; Fei *et al.* 2011c).

The gatekeeper and major regulator of TAG synthesis—and therefore of LD formation—is the Mg^{++} -dependent PA phosphohydrolase, *Pah1/Smp2* (Carman and Han 2006, 2011; Han *et al.* 2006, 2007; O'Hara *et al.* 2006; Pascual and Carman 2012): in the absence of this enzyme in *pah1* mutants, TAG synthesis is reduced by at least 70%, which also results in a drastically reduced LD formation (Adeyo

et al. 2011; Fei *et al.* 2011c) (Figure 1). Mammals express the *Pah1/Smp2* ortholog, lipin (encoded by *LPIN1-3* genes), mutations of which may cause lipodystrophy in the mouse (Garg 2004; Csaki and Reue 2010). Two additional enzymes, diacylglycerolpyrophosphate phosphatase, encoded by *DPP1* and *LPP1*, may also be involved in DAG formation; however, they serve a regulatory function and their quantitative contribution to TAG formation is unlikely (Henry *et al.* 2012).

Diacylglycerol that is formed by dephosphorylation of PA is converted either by the acyl-CoA-dependent acyltransferase *Dga1* [ortholog of mammalian DGAT (Oelkers *et al.* 2002; Sandager *et al.* 2002; Sorger and Daum 2002, 2003)], or the phospholipid-dependent acyltransferase *Lro1* [ortholog of mammalian lecithin-cholesterol acyltransferase LCAT (Oelkers *et al.* 2000)] to TAG. *Lro1* localizes to the ER whereas *Dga1* localizes both to the ER and LDs (Natter *et al.* 2005; Choudhary *et al.* 2011; Jacquier *et al.* 2011). The primary acyl donors of the *Lro1*-catalyzed reaction are phosphatidylethanolamine and phosphatidylcholine; thus, this reaction not only contributes to the synthesis of TAG but also serves to remodel the acyl chain composition of these phospholipids (Kohlwein 2010b; Horvath *et al.* 2011). Minor contribution to TAG synthesis from DAG stems from the activity of the sterol acyltransferases *Are1* and *Are2* (Yang *et al.* 1996) (see below).

The second major neutral lipid components of LD are the SEs. Sterols are synthesized in the ER membrane, which also harbors the acyl-CoA-dependent acyltransferases *Are1* and *Are2* that are required for SE synthesis (Yang *et al.* 1996; Yu *et al.* 1996; Zweytick *et al.* 2000b) (Table 1). *Are1* and *Are2* share 49% sequence identity with each other, and some 24% identity with mammalian acyl-CoA:cholesterol acyltransferases (ACAT; hence their names ACAT-related enzymes, or Are). Like *Slc1* and *Ale1*, *Are1* and *Are2* are members of the MBOAT family of membrane-bound *O*-acyltransferases (Pagac *et al.* 2011). Notably, both enzymes acylate not only ergosterol, but also intermediates in the ergosterol biosynthetic pathway: whereas the major SE synthase *Are2* prefers ergosterol as the substrate, *Are1* has a preference for the sterol precursor, lanosterol, giving rise to distinct SE compositions in mutants lacking either one of the enzymes (Zweytick *et al.* 2000b; Czabany *et al.* 2007, 2008). *Are1* was also found to contribute most to SE synthesis under anaerobic conditions (Hronska *et al.* 2004).

The localization of the four acyltransferases involved in TAG and SE formation to the ER membrane poses an interesting puzzle as to the transfer mechanism of their products, TAG or SE, to the LD. The close association between the ER and LDs (Figure 1) may be instrumental in supporting this exchange, but the proteins required for this process are presently unknown. Notably, since also intermediates of the ergosterol biosynthesis are stored as SEs in LD, their mobilization and further processing to “mature” ergosterol requires their reshuffling to the ER-resident sterol biosynthetic enzymes (Espenshade and Hughes 2007). The mechanism underlying this transfer and its regulation are unknown.




Figure 2 (A) Metabolic pathways of TAG synthesis and degradation and their subcellular localization (adapted from Kohlwein 2010b and Henry *et al.* 2012). Phospholipids and TAG share DAG and PA as common precursors. In the *de novo* synthesis of phospholipids, PA serves as the immediate precursor of CDP-DAG, precursor to PI, PGP, and PS. PA is dephosphorylated to DAG, which serves as the precursor of PE and PC in the Kennedy pathway. DAG also serves as the precursor for TAG and can be phosphorylated, regenerating PA. The names of the enzymes that are discussed in detail in the text are shown adjacent to the arrows of the metabolic conversions in which they are involved, and the gene-enzyme relationships are listed in Table 1. Lipids and intermediates are boxed, with the most abundant lipid classes boxed by bold lines. Enzyme names are indicated in boldface type. TAG, triacylglycerols; PI, phosphatidylinositol; PA, phosphatidic acid; CDP-DAG, CDP-diacylglycerol; DAG, diacylglycerol; MAG, monoacylglycerol; Gro, glycerol; Gluc-6P, glucose-6 phosphate; DHAP, dihydroxyacetone phosphate, PS, phosphatidylserine; FFA, free fatty acids; Ins, inositol. Nucl, nucleus; ER, endoplasmic reticulum; Mito, mitochondria; LD, lipid droplets; G/E/V, Golgi, endosomes, vacuole; Pex, peroxisomes; Cyt, cytoplasm; PM, plasma membrane. See text for details. (B) Metabolic pathways of fatty acid metabolism. FA *de novo* synthesis and elongation: FA (type I) *de novo* synthesis requires the synthesis of malonyl-CoA by the acetyl-CoA carboxylase Acc1. This cytosolic trifunctional enzyme harbors a covalently bound biotin, an N-terminal biotin carboxylase domain, and a C-terminal transcarboxylase domain (Tehlivets *et al.* 2007). Malonyl-CoA is used by the cytosolic FA synthase complex, consisting of Fas1 (β -subunit) and Fas2 (α -subunit), which are organized in a hexameric $\alpha_6\beta_6$ complex. Fas1 harbors acetyl transferase (AT), enoyl reductase (ER), dehydratase (DH), and malonyl-palmitoyl transferase (MPT) activities; Fas2 contains the acyl carrier protein (ACP), 3-ketoreductase (KR), 3-ketosynthase (KS), and phosphopantetheine transferase activities. The product of FA synthesis in yeast is acyl-CoA, typically C14–C16 carbon atoms in length (Tehlivets *et al.* 2007). Activated FAs may be elongated to VLCFAs by the activity of Elo1, Fen1/Elo2, and Sur4/Elo3 (condensing enzymes); Ybr159w (reductase); Phs1 (dehydratase); and Tsc13 (enoyl-CoA reductase). Yeast also expresses a set of bacterial type II

enzymes (as individual polypeptides) that perform the same reactions in mitochondria, but are encoded by nuclear genes (Tehlivets *et al.* 2007). Mitochondrial FA synthesis presumably generates FA only up to C8, which is a precursor for lipoic acid synthesis. FAs are degraded by β -oxidation. β -oxidation in yeast occurs exclusively in peroxisomes. Medium chain fatty acids enter peroxisomes as free fatty acids (FFA) and are activated by a peroxisomal acyl-CoA synthetase, Faa2. ATP that is required for this activation step is imported into the organelle via Ant1. Long chain fatty acids, such as oleate, are activated outside the organelle by Fat1, Faa1, or Faa4 and taken up as CoA esters (acyl-CoA) via a peroxisomal ABC transporter that consists of the heterodimer Pxa1/Pxa2. Inside peroxisomes, CoA esters undergo dehydrogenation by Pox1, hydration/dehydrogenation by Fox2, and ultimately thiolyl cleavage by Pot1, leading to acetyl-CoA and an acyl-chain shortened by two carbon atoms. Hydrogen peroxide produced by Pox1 is degraded by peroxisomal catalase T, Cta1. NADH is exported to the cytosol via a malate shuttle that involves peroxisomal (Mdh3) and cytosolic (Mdh2) malate dehydrogenases. The transporter for malate and oxaloacetate has not been identified yet. Acetyl-CoA is transported to the cytosol via carnitine-dependent acetyl-CoA transport (involving Cat2) or via the glyoxylate cycle (see Figure 5). Unsaturated FAs, such as oleic acid with the double bond between C9 and C10, can be fully oxidized only in the presence of auxiliary enzymes, but the precise mechanism is controversial. Eci1 is a $\Delta 3, \Delta 2$ -enoyl-CoA isomerase in the so-called isomerase-dependent major pathway, which catalyzes the positional and stereochemical isomerization of *cis*-3-enoyl-CoA to *trans*-2-enoyl-CoA; this reaction is required after oleic acid (as coenzyme A derivative) has been shortened by three rounds of β -oxidation, since only *trans*-2-enoyl-CoA is a β -oxidation substrate. Eci1 also isomerizes a fraction of 2-*trans*, 5-*cis*-dienoyl-CoA to 3,5-dienoyl-CoA, which has two conjugated double bonds in *trans* (3) and *cis* (5) configuration. This compound is presumably degraded by the minor pathway that involves Dci1, Sps19, and Eci1. Alternatively, 3,5-dienoyl-CoA is hydrolyzed by Tes1 thioesterase-dependent pathway) to the free FA and coenzyme A.

Turnover of lipid droplets

A systematic microscopic analysis in growing cells has shown that LDs are readily degraded and their content mobilized by up to 80% within 4–6 hr after transfer of stationary-phase cells into fresh, glucose-containing media (Kurat *et al.* 2006); the LDs are subsequently replenished until cells reach stationary phase. The neutral lipid content of LD is degraded by the activity of TAG lipases and SE hydrolases. Tgl3, Tgl4, and Tgl5 are members of the conserved patatin-domain-containing family of hydrolases (Athenstaedt and Daum 2003, 2005, 2006; Czabany *et al.* 2007; Daum *et al.* 2007a,b) that are characterized by a serine active residue embedded in a G-x-S-x-G motif in a patatin domain (Kienesberger *et al.* 2009); however, in contrast to typical lipases that harbor a Ser-Asp-His catalytic triad (see below), these enzymes harbor only a catalytic dyad, composed of a serine and an aspartic acid residue. Yeast Tgl4 is the functional ortholog of the mammalian adipose triglyceride lipase, ATGL (Zimmermann *et al.* 2004; Kurat *et al.* 2006), which is the major TAG-hydrolyzing enzyme in adipose tissue and in other cell types (Lass *et al.* 2011; Zechner *et al.* 2012). ATGL deficiency in humans is associated with neutral lipid storage disease with myopathy, NLSDM (Schweiger *et al.* 2009; Zechner *et al.* 2012). Tgl3 and Tgl4 are the major TAG lipases in yeast, and deletion of these genes leads to markedly increased LD size and number (Figure 1); Tgl5 only marginally contributes to TAG hydrolysis under standard growth conditions. TAG content is increased by ~15% in the *tgl3* mutant, whereas overexpression reduces TAG content <10% (Athenstaedt and Daum 2003). However, since TAG content is strongly dependent on growth conditions, somewhat different values were obtained in another study from the same lab, showing that TAG levels in the *tgl3* mutant increased to 4.11 $\mu\text{g}/\text{mg}$ dry cells compared to 1.72 $\mu\text{g}/\text{mg}$ TAG/mg dry cells of the wild-type strain (BY4741). Tgl4-deficient cells contained 2.97 $\mu\text{g}/\text{mg}$ TAG/mg cell dry weight. Notably, whereas Tgl5-deficient cells had TAG levels identical to wild type, TAG levels were even further increased in *tgl3 tgl5* double mutants to 5.38 $\mu\text{g}/\text{mg}$ dry weight (Athenstaedt and Daum

2005).² Not only the quantitative contribution to TAG lipolysis between the yeast lipases differs, but also the lipase substrate specificities differ: Tgl3 preferentially hydrolyzes TAG species containing C14, C16, C20, and C26 saturated acyl chains (Athenstaedt and Daum 2003). Similarly, Tgl4 prefers TAG species with C14 and C16 acyl chains. Notably, cells lacking the Tgl5 lipase showed markedly increased levels of C26 acyl chain-containing TAG molecular species, indicating a substrate preference of this lipase for very long chain FAs (Athenstaedt and Daum 2005). In addition to being an efficient TAG lipase, Tgl3 also harbors substantial DAG lipase activity; thus, overexpression of Tgl4 in a *tgl3* mutant background leads to increased accumulation of DAG, which is also accompanied by a slight growth defect (Kurat *et al.* 2006). Tgl4, in addition to being a major TAG lipase, also displays steryl ester hydrolase and phospholipase A₂ activities *in vitro*. Furthermore, this enzyme also catalyzes acyl-CoA dependent re-acylation of lyso-PA to PA (Rajakumari and Daum 2010b). The efficacy of this reaction in contributing to the synthesis of PA *in vivo* is not clear and apparently not sufficient to support growth of an *slc1 ale1* double mutant, lacking the two major yeast lyso-PA acyltransferases (see above). Similarly, Tgl3 and Tgl5 lipases also harbor lyso-PA and lyso-phosphatidylethanolamine acyltransferase activities *in vitro* (Rajakumari and Daum 2010a). Thus, Tgl3, Tgl4, and Tgl5 lipases not only catalyze TAG breakdown to various degrees, but may also be involved in establishing specific acyl-chain compositions to phospholipids.

Despite catalyzing the majority of TAG breakdown, deletion of all three lipases does not result in a significant growth phenotype in logarithmically growing cells (Athenstaedt and Daum 2005; Kurat *et al.* 2006); however, lipase mutants are sensitive to the FA synthesis inhibitor cerulenin, consistent with the role of lipolysis-derived metabolites (FAs, DAG) for the synthesis of membrane phospholipids. Notably, initiation of the cell division cycle upon transfer of stationary phase/quiescent cells into fresh growth media is delayed in mutants

²In the original publication, TAG levels were erroneously printed as “mg TAG/mg cell dry weight” (K. Athenstaedt, personal communication)

lacking Tgl3 and Tgl4 lipases (Kurat *et al.* 2009): in these mutants, G1/S transition is extended by some 30 min, indicating that lipolysis-derived metabolites are required for efficient cell cycle progression (see below). Evidence suggests that lack of lipolysis affects the formation of sphingolipids (Rajakumari *et al.* 2010), which play multiple regulatory and structural roles (Dickson 2010). Also, lipase-deficient mutants are defective in phosphatidylinositol (PI) synthesis: addition of inositol to wild-type cells that were grown in the absence of this lipid precursor results in a rapid burst in PI synthesis, which is significantly attenuated in *tgl3 tgl4 tgl5* lipase mutants. Additional inhibition of *de novo* FA synthesis by cerulenin abolishes the burst in PI synthesis after inositol addition, indicating that both *de novo*-synthesized FAs and metabolites derived from TAG breakdown are required to support PI synthesis (Gaspar *et al.* 2011). PI is also a precursor for the synthesis of complex sphingolipids, which may be the underlying reason for attenuated sphingolipid synthesis in lipase-deficient cells (Rajakumari *et al.* 2010).

Homozygous diploid *tgl3/tgl3* and *tgl5/tgl5* mutants are unable to sporulate, indicating that Tgl3 and Tgl5 provide essential activities that are required for the generation of functional spores (Rajakumari and Daum 2010a). Indeed, it was shown that the Tgl3 acyltransferase activity, rather than the lipase activity, is required for sporulation (Rajakumari and Daum 2010a). However, the specific step in the sporulation program that requires this activity is not known.

The role of the TAG lipases in sustaining viability during stationary phase, in the absence of other carbon sources, is not known. Notably, mutants defective in the DAG kinase Dgk1 also display a delay in growth resumption after transfer of stationary cells into fresh growth medium (Fakas *et al.* 2011a), similar to *tgl3 tgl4* mutants, and TAG degradation is defective, even in the presence of cerulenin. Choline supplementation partially suppresses this defect; it was suggested that lipotoxic lipolysis-derived DAG might accumulate under these conditions, which is drained into the synthesis phospholipids via the cytidine diphosphate (CDP)-choline (Kennedy) pathway if choline is present (Fakas *et al.* 2011a). Dgk1 is localized to the ER, and the transfer of its substrate DAG from the LDs may be facilitated by the close physical interaction between both organelles (Szymanski *et al.* 2007).

Common to Tgl3, Tgl4, and Tgl5 TAG lipases is their exclusive localization to LDs, which is in contrast to other LD proteins—mostly enzymes involved in anabolic processes—that are additionally associated with the ER membrane (Athenstaedt and Daum 2006; Kurat *et al.* 2006; Daum *et al.* 2007a; Rajakumari *et al.* 2008; Kohlwein 2010b). Localization of Tgl3 to LDs may be regulated by the yeast seipin ortholog Fld1 (Wolinski *et al.* 2011).

In addition to the major TAG lipases Tgl3 and Tgl4, which catalyze the majority of TAG and also DAG breakdown, yeast also expresses a monoacylglycerol (MAG) lipase to complete the “lipolytic cascade” analogous to mammalian

cells (Zechner *et al.* 2012). Yeast MAG lipase is encoded by the *YJU3* gene (Heier *et al.* 2010) and localizes to both ER and LDs. Deletion of the *YJU3* gene results in accumulation of MAG, but does not lead to a detectable phenotype under numerous experimental conditions. This is surprising since the specific activity of the Yju3 protein is several orders of magnitude higher than that of the TAG lipases (Heier *et al.* 2010).

The *TGL1*-, *YEH1*-, and *YEH2*-encoded steryl ester hydrolases are involved in SE degradation. These enzymes are related to mammalian acid lipases, and, as “prototypic” hydrolases, they harbor a serine-active site embedded in a G-x-S-x-G motif, a catalytic triad consisting of Ser-Asp-His residues, and an α/β -hydrolase fold (Jandrositz *et al.* 2005; Koffel *et al.* 2005; Mullner *et al.* 2005; Koffel and Schneider 2006; Wagner *et al.* 2009). Whereas Tgl1 and Yeh1 localize predominantly to LDs, Yeh2 is enriched in the plasma membrane (Koffel *et al.* 2005; Mullner *et al.* 2005; Wagner *et al.* 2009), consistent with previous findings derived from cell fractionation experiments that showed significant SE hydrolase activity in the plasma membrane (Zinser *et al.* 1993; Leber *et al.* 1995). In addition to its activity as an SE hydrolase, Tgl1, which shares similarities to mammalian lysosomal acid lipases, also degrades TAG *in vitro*. This activity, however, does not appear to significantly contribute to TAG turnover *in vivo* under standard growth conditions (Jandrositz *et al.* 2005) and does not affect LD abundance and structure.

Regulation of neutral lipid synthesis

Very little is known about the specific regulation of enzymes involved in TAG and SE synthesis and, therefore, in LD biogenesis. Formation of LDs is clearly driven by the availability of lipid precursors, sterols, and FAs and therefore is dependent on the regulatory processes that control the biosynthesis of their lipid constituents. Notably, none of these processes has been specifically investigated in the context of LD formation. Microscopic analysis of LD in growing cells indicated that their degradation and new synthesis may be processes that at least partially overlap (Kurat *et al.* 2006). Similarly, dynamic flux balance analysis also unveiled that degradation and *de novo* formation of LD may occur in parallel to maintain FA and lipid homeostasis (Zanghellini *et al.* 2008). This appears to be a conserved mechanism that also occurs in mammalian cells: FAs taken up from the blood stream into cells may first be incorporated into TAG prior to their release by lipolysis. Failure to degrade TAG in homozygous lipase-deficient *ATGL*^{-/-} mouse mutants leads to a lack of PPAR-agonist release and impaired mitochondrial function (Haemmerle *et al.* 2011; Zechner *et al.* 2012).

A major determinant of TAG synthesis is the availability of FAs and glycolysis-derived glycerol-3-phosphate or DHAP. Exogenously supplied FAs are preferentially stored as TAG, but are also incorporated into membrane phospholipids upon FA supplementation (Grillitsch *et al.* 2011). Very little is known about the regulation of acyl-CoA synthetase

activities that are required to activate free FAs (Black and Dirusso 2007), and it can only be speculated that glycolysis derived glycerol-3-phosphate or DHAP are the limiting compounds that determine cellular TAG levels in the presence of a surplus of exogenous FAs. TAG levels are increased about fivefold when cells are grown in the presence of oleic acid as the sole carbon source (Grillitsch *et al.* 2011). Endogenous FA synthesis is under transcriptional and post-translational control at the level of *Acc1* and the FA synthase complex (Tehlivets *et al.* 2007; Kohlwein 2010b; Henry *et al.* 2012). *Acc1* is phosphorylated and inactivated by *Snf1* kinase, the ortholog of mammalian AMP-activated protein kinase (Woods *et al.* 1994), under conditions of scarce energy. Thus, *Snf1* is an important regulator of TAG homeostasis by regulating the activity of *Acc1* and, thus, FA *de novo* synthesis and TAG accumulation (Tehlivets *et al.* 2007). Recent evidence suggests that TOR and *Snf1*/AMPK pathways are connected to the control of TAG formation through the *Sit4-Sap190* protein phosphatase complex that may control the activity of *Acc1* and/or *Snf1* (Bozaquel-Morais *et al.* 2010). FA desaturation is regulated by the membrane-bound transcription factors *Spt23* and *Mga2*, which are processed in an *Rsp5* ubiquitin ligase-dependent reaction (Hoppe *et al.* 2000; Rape *et al.* 2001); soluble *Spt23* and *Mga2* fragments translocate into the nucleus to regulate the expression of the *OLE1* gene encoding the single FA desaturase in yeast (Stukey *et al.* 1989, 1990; Hoppe *et al.* 2000; Chellappa *et al.* 2001; Rape *et al.* 2001; Martin *et al.* 2002, 2007; Tehlivets *et al.* 2007; Henry *et al.* 2012). Notably, overexpression of constitutively active *Mga2* or *Spt23* fragments stimulates TAG synthesis and leads to altered LD morphology, indicating a regulatory link between *Rsp5*, *Spt23*, and *Mga2* function and lipid homeostasis (Kaliszewski and Żołądek 2008).

The initial steps in glycerolipid synthesis require the activity of *Sct1* and *Gpt2* acyltransferases (Figure 2). *Sct1* localizes to the ER membrane (Bratschi *et al.* 2009), and the *SCT1* gene was originally identified as a suppressor of a choline transport mutant, indicating a functional relationship to phosphatidylcholine synthesis (Matsushita and Nikawa 1995). Indeed, establishment of the acyl-chain composition in phosphatidylcholine requires *Sct1* (Boumann *et al.* 2003). *Sct1* activity, which is regulated by phosphorylation by an as-yet-unknown kinase, competes with the *OLE1*-encoded FA desaturase for their common substrate, palmitoyl-CoA. Thus, overexpression of *Sct1* leads to increased phosphatidylinositol and TAG levels at the expense of phosphatidylethanolamine and a general shift in FA profiles toward more saturated species (De Smet *et al.* 2012). Notably, deletion of the *SCT1* gene has a significant impact on the turnover of phosphatidylcholine that is generated through the CDP-choline (“Kennedy”) pathway (Zaremborg and McMaster 2002), and further evidence suggests that this phospholipid, next to TAG, functions as a reservoir for FAs, in particular for C16:0 (De Smet *et al.* 2012). This is also consistent with the observation that cells become more sensitive to C16:0 sup-

plementation when both TAG synthesis and the phospholipid methylation pathway are blocked (Garbarino *et al.* 2009).

Deletion of the second acyltransferase encoded by the *GPT2* gene has the opposite effect on phosphatidylcholine turnover than a deletion of *SCT1*, namely a highly stimulated turnover of this phospholipid synthesized via the CDP-choline pathway (Zaremborg and McMaster 2002). In contrast to wild-type cells, mutants defective in *Gpt2* acyltransferase are sensitive to oleate supplementation and fail to synthesize TAG and induce LD formation (Marr *et al.* 2012). Oleate may indeed regulate *Gpt2* abundance and its activity by phosphorylation; furthermore, *Gpt2*-containing crescent ER structures that are observed in close vicinity to LDs in the presence of oleate indicate a regulatory crosstalk between LD formation and activity of the initial steps of glycerolipid synthesis (Marr *et al.* 2012).

The redundant lyso-PA acyltransferases encoded by *SLC1* and *ALE1* both contribute to the typical FA spectrum in cellular glycerolipids, whereby *Ale1* may have a somewhat higher preference for C16:1-CoA than *Slc1* (Benghezal *et al.* 2007). Although *Slc1* harbors the majority of cellular *sn1*-acylglycerol-3-phosphate acyltransferase activity, deletion of either *SLC1* or *ALE1* genes does not significantly affect total cellular glycerolipid content (Benghezal *et al.* 2007). This is surprising since these enzymes together execute an essential reaction, indicated by the synthetic lethal phenotype of *slc1 ale1* double mutants, which also suggests the absence of significant additional lyso-PA acyltransferase activities in yeast (Jain *et al.* 2007).

The dephosphorylation of PA to DAG is considered the rate-limiting step in TAG formation, and mutants lacking the PA phosphatase *Pah1* are characterized by drastically reduced TAG levels (Carman and Han 2006, 2009; Han *et al.* 2006; Fakas *et al.* 2011b; Henry *et al.* 2012; Pascual and Carman 2012). *Pah1* is under multiple levels of regulation by phosphorylation, which controls its localization to the cytosol (phosphorylated) or its association with the ER membrane (dephosphorylated) (Carman and Han 2009; Karanasios *et al.* 2010; Choi *et al.* 2011). Dephosphorylation of *Pah1* by the *Nem1-Spo7* phosphatase complex favors its association with the ER membrane and facilitates generation of the TAG precursor DAG (Siniosoglou *et al.* 1998; Santos-Rosa *et al.* 2005).

Notably, *Pah1* is phosphorylated by the cyclin-dependent protein kinases *Cdc28/Cdk1* and *Pho85* (Karanasios *et al.* 2010; Choi *et al.* 2011), indicating that its membrane association and activity are regulated in a cell cycle-dependent manner. Since TAG degradation also is regulated in a cell cycle-dependent manner (Kurat *et al.* 2009) (see below), the picture emerges that TAG synthesis and degradation may indeed oscillate during the cell cycle (Kurat *et al.* 2009; Kohlwein 2010b).

The activity of *Pah1* is counteracted by the CTP-dependent diacylglycerol kinase *Dgk1* and may thus contribute to the regulation of TAG homeostasis. Overexpression of the *DGKI*

gene results in proliferation of ER membranes, consistent with an overproduction of PA that is preferentially channeled into phospholipid synthesis (Han *et al.* 2008a,b). Whereas the impact of *DGK1* overexpression on cellular TAG levels is unclear, deletion of this gene hardly has any effect on cellular TAG content in growing cells. However, *Dgk1* activity is important during periods of growth resumption, *i.e.*, after transfer of stationary-phase cells to fresh media, presumably to convert lipolysis-derived DAG to PA and subsequently to phospholipids. Absence of *Dgk1* activity leads to the accumulation of DAG with potentially detrimental effects on the cells, which can be attenuated by utilizing DAG for the CDP-choline pathway in the presence of choline (Fakas *et al.* 2011a).

The specific regulatory mechanisms that control the activity of *Dgal*, *Lro1*, *Are1*, and *Are2* acyltransferases are unknown (Yang *et al.* 1996; Oelkers *et al.* 2000, 2002; Zweytick *et al.* 2000b; Sorger and Daum 2002). Mutant analysis indicates that *Dgal* contributes more significantly to TAG synthesis in the stationary phase, whereas *Lro1* apparently is more active during logarithmic growth (Oelkers *et al.* 2002). Notably, supplementation of wild-type cells with oleic acid, which stimulates TAG synthesis, simultaneously reduces cellular SE levels (Connerth *et al.* 2010). This observation indicates a regulatory crosstalk between TAG synthesis and SE synthesis, the molecular basis of which, however, has not been uncovered yet. These findings also raise the question of whether distinct types of yeast LDs that harbor either SE or TAG may exist (see below). Clearly, biophysical properties differ between SE- or TAG-only LDs, despite similar size distribution (Czabany *et al.* 2008). For example, tri-oleoyl glycerol and tri-palmitoleoyl glycerol, which are the major TAG species, have a melting point below -4° , whereas cholesteryl oleate (related to the yeast ergosteryl oleate) has a melting point above $+40^{\circ}$ (PubChem Substance database).

Acylation of sterols may regulate the flux through the ergosterol biosynthetic pathway by sequestering and storing intermediate products as SEs in the LD. Thus, acylation may prevent buildup of potentially harmful sterol intermediates. Sterol synthesis is under a tight feedback regulatory loop that controls the expression of HMG-CoA reductase, the key enzyme of sterol synthesis both in yeast and in mammals (Espenshade and Hughes 2007; Burg and Espenshade 2011; Raychaudhuri *et al.* 2012). Mutants lacking *Dgal*, *Lro1*, *Are1*, and *Are2* acyltransferases altogether display a defect in sterol synthesis, which is due to the reduced amount of squalene epoxidase, *Erg1* (Sorger *et al.* 2004). This reduction in *Erg1* abundance is not due to attenuated expression but rather is a result of decreased protein stability in the *dgal lro1 are1 are2* quadruple mutant. *Erg1* typically localizes both to the ER and LDs (Leber *et al.* 1998), which, however, are absent in the quadruple mutant. Thus, the decreased *Erg1* stability indicates a tight regulation of the amount of ER-resident *Erg1* protein. It furthermore suggests that localization of *Erg1* to the LD provides a mechanism to store (catalytically inactive) enzyme that is not subject to

this regulation (Leber *et al.* 1998), but may be relocalized to the ER upon metabolic requirements. The mechanisms that govern sterol lipid exchange between the ER and LDs, and the regulation of these processes, remain obscure.

TAG accumulation is also influenced by the cellular capacity to synthesize phospholipids: attenuated phosphatidylcholine synthesis, *i.e.*, in mutants lacking the *CHO2*- and *OPI3*-encoded phospholipid methyltransferases or defective in *S*-adenosylhomocysteine hydrolase (*Sah1*), which affects the methylation activity, leads to an increased synthesis of TAG and LD proliferation (Malanovic *et al.* 2008). The reduced flow of FAs into phosphatidylcholine synthesis presumably leads to the accumulation of PA, which is preferentially channeled into the synthesis of TAG. However, these observations also support the notion that PC synthesis, in addition to TAG, may also provide some (limited) buffering capacity for accommodating excess FA.

Regulation of neutral lipid degradation

Lipolysis is most active during growth resumption of stationary-phase cells that are transferred into fresh, glucose-containing medium (Kurat *et al.* 2006; Zanghellini *et al.* 2008). TAG is degraded by *Tgl3* and *Tgl4* lipases that both reside on the lipid droplet (Czabany *et al.* 2007; Daum *et al.* 2007a; Rajakumari *et al.* 2008; Kohlwein 2010b; Henry *et al.* 2012). Neither protein abundance nor localization appear to change during the phase of lipolysis. *Tgl5* and *Tgl1*, which are also TAG lipases *in vitro*, do not appear to contribute significantly to TAG degradation. Indeed, *Tgl1* is more active as a SE hydrolase (Koffel *et al.* 2005). *Tgl4* is phosphorylated and activated by the cyclin-dependent kinase *Cdk1/Cdc28* (Kurat *et al.* 2009) at the G1/S transition of the cell cycle, suggesting that lipolysis-derived products (*i.e.*, FAs or DAG) are required to drive cell cycle progression. The specific checkpoint-monitoring availability of lipolysis products is unknown. Similarly, *Tgl5* may also be a substrate of *Cdk1/Cdc28* (Ubersax *et al.* 2003), and *Tgl3* lipase is a potential target of the second, nonessential cyclin-dependent kinase *Pho85*, according to large-scale studies (Ptacek *et al.* 2005). Whether *Tgl3* and *Tgl5* activities are indeed regulated during the cell cycle is not known.

The observation that lipolysis in yeast is linked to cell cycle progression is unexpected; indeed, *Tgl4* and *Pah1* are among the very few direct enzymatic targets of the cyclin-dependent kinase *Cdk1/Cdc28*. Since both *de novo* TAG synthesis, driven by the activity of PA phosphatase *Pah1*, and lipolysis are regulated in a cell cycle-dependent manner, it becomes obvious that maintenance of lipid homeostasis during the cell cycle is critical (Kurat *et al.* 2009; Kohlwein 2010b). Neither the TAG degradation products nor the checkpoint regulator that senses their availability are known. Notably, *Tgl4* phosphorylation—and activation—occurs at the G1/S transition of the cell cycle, at bud emergence, whereas *Pah1* phosphorylation—and inactivation—occurs at the G2/M transition. This leaves both enzymes active during a large part of the cell cycle, consistent with a model that lipogenesis

and lipolysis may occur in parallel to sustain cellular lipid homeostasis (Zanghellini *et al.* 2008; Kohlwein 2010b).

The mechanisms of the regulation of steryl ester hydrolysis by *Yeh1*, *Yeh2*, and *Tgl1* are currently unknown. Since LDs that are composed of about equal amounts of TAG and SEs are mobilized by 80% during the initial phase (~6 hr) of growth resumption (Kurat *et al.* 2006), one can assume the highest activity of these enzymes during this period of growth. The LD-resident enzymes *Yeh1* and *Tgl1*, but not the plasma membrane-resident enzyme *Yeh2*, harbor potential cAMP-dependent protein kinase A phosphorylation sites, which may be responsible for stimulation of activity (Koffel *et al.* 2005). Also, *Yeh1* is the major SE hydrolase in *hem1*-deficient mutant cells that lack *de novo* sterol synthesis and require ergosterol supplementation (Koffel and Schneiter 2006).

Mechanisms of LD biogenesis and inheritance

No clear picture currently exists of how LDs are actually assembled, neither in yeast nor in other cell types. Current models of LD formation are summarized in Figure 3 (Zweytick *et al.* 2000a; Mullner and Daum 2004; Czabany *et al.* 2007; Daum *et al.* 2007a; Jacquier *et al.* 2011) (see below). The “lensing” model (Figure 3A) and the “bicelle” model (Figure 3B) share the idea that TAG accumulates between the leaflets of the ER membrane; after reaching a critical size, LD may bud off toward the cytosol (lensing model) or are excised from the ER, leaving behind a gap in the membrane, which, however, may be quickly filled up again. In the former model, the monolayer surrounding the LD is solely derived from the cytosolic leaflet of the ER membrane, whereas in the bicelle model both ER membrane leaflets contribute to the LD surface monolayer. The “vesicle budding” model (Figure 3C) suggests the formation of TAG-filled secretory vesicles that undergo remodeling of the ER-derived phospholipid bilayer to yield the observed phospholipid monolayer covering the LDs. Common to these models is a tight functional interaction between emerging LDs and the endoplasmic reticulum from which they presumably derive. Indeed, LDs may be in continuous ER contact throughout their life cycle (Wolinski *et al.* 2011). Recently, first attempts have been made to understand LD formation from a theoretical point of view, based on biophysical models (Zanghellini *et al.* 2010a,b). According to these models, LDs bud off the ER membrane in a process that is driven by lipid de-mixing in the membrane, when a critical size of some 12 nm is reached. Since this size is more than an order of magnitude below the observed LD size *in vivo*, the authors concluded that LD formation is a two-step process in which initial LD formation is followed by fusion events, giving rise to native “ripe” LDs (Zanghellini *et al.* 2010a,b).

Contribution of acyl transferases to LD biogenesis

In the absence of both DAG acyltransferases, *Dga1* and *Lro1*, LDs are solely composed of SE. Notably, despite the fact that SE make ~50% of the total neutral lipid content in wild-type cells, LD numbers are drastically reduced in *dga1 lro1* double mutants to one or two LDs (Oelkers *et al.* 2002; Sorger




Figure 3 Models of lipid droplet biogenesis (adapted from Guo *et al.* (2009). (A) According to the “lensing model,” neutral lipids are deposited between the leaflets of the ER membrane: after reaching a critical size, the neutral lipid core bulges out and the LD is formed; the LD surface monolayer is derived solely from the cytosolic leaflet of the ER membrane. Subsequently, the LD may completely separate from the ER membrane, or remain attached, with the surface layer forming a continuum with the ER. (B) Bicelle formation: LD formation similar to model in A, but the LD is excised from the ER membrane, and both ER membrane leaflets contribute to the LD surface monolayer. (C) Vesicle formation: Inclusion of the neutral lipid core in the membrane vesicle requires rearrangement of the inner leaflet of the bilayer. These models explain the origin of the phospholipid membrane, which stems either from the cytoplasmic leaflet or from both leaflets of the ER membrane, respectively. Unclear is what limits the expansion of the neutral lipid core between the leaflets, what determines the orientation of LD extrusion toward the cytosol, and how the integrity of the ER membrane is maintained. Notably, none of the intermediate stages representing neutral lipid deposits between the ER membrane leaflets, nascent lipid droplets in the ER, or lipid-filled vesicular structures have been experimentally observed in wild-type cells.

and Daum 2002; Athenstaedt and Daum 2006; Czabany *et al.* 2007; Daum *et al.* 2007b; Rajakumari *et al.* 2008; Walther and Farese 2009; Kohlwein 2010b) (Figure 1). In contrast, simultaneous deletion of *Are1* and *Are2* sterol acyltransferases has only a marginal effect on LD content. No LDs are present in mutants lacking *Dga1 Lro1 Are1* and *Are2* acyltransferases (Oelkers *et al.* 2002; Sandager *et al.* 2002; Garbarino *et al.* 2009; Petschnigg *et al.* 2009; Kohlwein 2010b). Thus, LD formation is clearly correlated with the activity of these acyltransferases. This has also led to the establishment of a test system to study LD biogenesis by expressing the major DAG acyltransferases, *Dga1* or *Lro1*, under control of the galactose-inducible *GAL1^P* promoter, in cells lacking other acyltransferases (Jacquier *et al.* 2011). In this system, LDs are absent from cells grown on glucose, but LD formation is induced upon shift of cells to galactose medium; LD formation could be observed within 2 hr of

induction of the acyltransferases. Formation of new LDs occurs close to the nuclear ER, consistent with the current biogenesis model that LD may derive from the ER. Furthermore, LD proteins that relocalize to the ER in the absence of LDs translocate to the newly formed, nascent LD; this protein relocalization is independent of *de novo* protein synthesis or energy (Jacquier *et al.* 2011). Fluorescence recovery after photobleaching and fluorescence loss in photobleaching experiments suggest that ER and LD membrane may indeed form a continuum that allows the free diffusion of LD-resident proteins from the ER to the growing LD, and back to the ER, upon stimulation of TAG breakdown (Jacquier *et al.* 2011).

The topology of acyltransferases involved in TAG formation may provide some clues as to the origin of the LD core lipids and thus the mechanism of LD formation (Choudhary *et al.* 2011; Jacquier *et al.* 2011; Pagac *et al.* 2011). *Dgal* harbors a stretch of hydrophobic amino acids compatible with two membrane-spanning domains; since the enzyme is active both in the ER and on LDs, which contain only a phospholipid monolayer, any potential rearrangement of the enzyme does not appear to affect its activity (Jacquier *et al.* 2011). In contrast, *Lro1*, the phospholipid-dependent acyltransferase, which is exclusively localized to the ER, harbors only one membrane-spanning domain. Its presumed active site residing in the lumen of the ER suggests that TAG may indeed be formed in the lumen of the ER, rather than between ER membrane leaflets (Choudhary *et al.* 2011). Similarly, evidence suggests that in the MBOAT enzymes *Are1* and *Are2*, the conserved histidine residue involved in catalysis is also exposed to the luminal side of the ER (Pagac *et al.* 2011).

Targeting of proteins to LDs

Unlike other proteins targeted to organelles, LD-associated proteins apparently do not harbor targeting consensus sequences as determined by primary structure comparison of LD-associated proteins. However, a common feature appears to be the presence of hydrophobic domains, although exceptions exist (Leber *et al.* 1998; Mullner *et al.* 2004; Grillitsch *et al.* 2011). As shown in Table 1, several of the LD-associated proteins contain even one or two (predicted) transmembrane domains, which appear to be incompatible with the generally accepted view that the LD surface is covered by a phospholipid monolayer. Thus it is unclear how these extended stretches of hydrophobic amino acids are accommodated in the LD surface layer. Also, numerous LD proteins lack hydrophobic stretches indicative of membrane-anchoring sequences altogether (Table 1), suggesting that their interaction with LDs may be indirect and through the interaction with LD-anchored proteins.

Notably, numerous LD-associated proteins are dually localized also to the ER membrane (Table 1), including the enzymes involved in sterol synthesis *Erg1*, *Erg6*, and *Erg7* (Mullner *et al.* 2004). The physiological relevance of this dual localization is unclear, since the other enzymes of ergosterol biosynthesis are ER-resident; However, as shown

for *Erg1*, localization to the LD may serve a regulatory function—to provide a pool of enzyme that is inactive on the LD, but which may readily relocalize to the ER upon demand (Sorger *et al.* 2004). Truncated versions of the *Erg1* lacking a single C-terminal hydrophobic stretch of 55 amino acids lost their affinity to the LDs and relocalized predominantly to the ER. Deletion of 87–139 C-terminal amino acids of the *Erg7* protein also led to significant retention of protein in the ER and reduced association to LDs. A C-terminal deletion of 26 hydrophobic amino acids in *Erg6* did not significantly alter its localization, whereas a deletion of 66 C-terminal amino acids abolished LD association and led to full translocation of the truncated *Erg6* protein to the ER (Mullner *et al.* 2004). It should be noted, however, that the relative distribution of proteins to the ER and LDs might depend on protein abundance, which somewhat limits the use of episomal overexpression clones for LD localization studies. Apparently, hydrophobic stretches are required for LD association, but the factors that discriminate relative distribution to LDs and the ER are currently unknown. Notably, heterologous LD proteins expressed in yeast also localize faithfully to LDs, such as mammalian adipose triglyceride lipase, ATGL (Kurat *et al.* 2006), or methyltransferase like 7B (AAMB) (Zehmer *et al.* 2008), despite the absence in yeast of perilipins that play an important role in regulating the access of proteins to the LDs in mammals (Brasaemle 2007).

Lipid droplet morphology and inheritance

Notably, in a given population of cells, LD size distribution is quite homogeneous, and it is currently unclear which factors regulate LD size independently of neutral lipid composition (Czabany *et al.* 2008). Yeast expresses the protein *Fld1* that is distantly homologous to mammalian seipin, implicated in the serious inheritable Berardinelli–Seip congenital lipodystrophy type 2 that results from defects in the *BSC1* gene (Szymanski *et al.* 2007; Fei *et al.* 2008, 2011a,b). Mutations in the *FLD1* gene lead to “supersized” LDs under inositol-limiting conditions, indicating that *Fld1* may play a role in LD biogenesis and organization. Indeed, morphological analysis in growing yeast cells indicates that *Fld1* plays a role in LD subcellular distribution and inheritance (Wolinski *et al.* 2011). Furthermore, access of the TAG lipase *Tgl3* to LDs seems to be impaired in *fld1* mutants. Its role in LD formation is derived from observations that *Fld1* may form homo-oligomers and localizes at the interface between the ER membrane and LDs (Szymanski *et al.* 2007; Binns *et al.* 2010). Thus, although not itself an LD-resident protein, *Fld1* is a potential regulator of LD assembly.

Physiological role of LDs

LDs function as the storage depot for TAG and SE. Thus, processes that depend on TAG and SE formation, or metabolites derived from TAG or SE, are affected by the cell's capacity to generate LDs. LDs lacking SE are more sensitive to sterol synthesis inhibitors, such as terbinafine (Zweytick *et al.* 2000b), which is in line with the function

of SE as storage molecules. In the presence of sterol synthesis inhibitors, SE are degraded and sterols incorporated into membranes until the SE content of the cell is exhausted and growth ceases (Zweytick *et al.* 2000b). Similarly, inhibition of FA *de novo* synthesis by cerulenin results in rapid mobilization of TAG and the utilization of released FAs or DAG for membrane lipid synthesis (Kurat *et al.* 2006; Fakas *et al.* 2011a).

A second major function of TAG (and LDs) is to serve as a buffer to “neutralize” excess FA. Cells lacking the capacity to synthesize TAG, *i.e.*, *dgal lro1 are1 are2* quadruple mutants, are highly sensitive to supplementation with unsaturated FAs: in the absence of TAG formation, oleic acid is preferentially incorporated into phospholipids, which leads to massive membrane proliferation and rapid loss of viability (Kohlwein and Petschnigg 2007; Garbarino *et al.* 2009; Petschnigg *et al.* 2009). This also reflects the sensitivity of mammalian cells to FA overload (Listenberger *et al.* 2003; Schaffer 2003). Notably, quadruple mutants exposed to oleic acid appear to “adapt” to this challenge and recover after an extended lag period (Connerth *et al.* 2010). This adaptation, however, seems to be a stable trait and suggests the appearance of suppressor mutations that allow cells devoid in TAG synthesis to sustain oleic acid challenge. Indeed, mutations in mitochondrial DNA confer resistance of the quadruple mutant to oleic acid-induced cell death (Rockenfeller *et al.* 2010). Similarly to the quadruple mutant that lacks TAG altogether, *pah1* mutants lacking PA phosphatase and containing drastically reduced levels of TAG are also highly sensitive to unsaturated FA supplementation (Fakas *et al.* 2011b). Thus, the picture emerges that FA overload leads to a critical imbalance in cellular phospholipid composition in the absence of TAG synthesis (Kohlwein and Petschnigg 2007; Garbarino *et al.* 2009; Kohlwein 2010a).

As mentioned above, attenuation of phosphatidylcholine synthesis in *cho2 opi3* mutants or in mutants defective in *S*-adenosylhomocysteine hydrolase, which regulates the cellular methylation potential (Malanovic *et al.* 2008; Tehlivets 2011), leads to an increased flux of FAs into TAG and subsequent LD accumulation. Notably, levels of phosphatidylcholine can be substantially reduced in yeast cells without leading to a significant growth phenotype (Henry *et al.* 2012), indicating that this phospholipid may also serve, at least in part, as a buffer for FAs. Similarly, a block of the early secretory pathway that can also be considered as a process to regulate the metabolic flux of phospholipids out of the ER leads to elevated TAG levels (Gaspar *et al.* 2008). These observations support the notion that phospholipid and TAG metabolism are metabolically tightly interconnected and that inactivation of either biosynthetic branch forces the channeling of FAs into the other, with potentially detrimental consequences for the cell. Furthermore, TAG and LD formation appear to play a crucial role in modulating ER stress that is induced by altered phospholipid composition or turnover (Hapala *et al.* 2011).

In addition to serving as an overflow reservoir for excess FAs, TAG—and LD altogether—may also serve as an over-

flow storage compartment for proteins. For example, *Erg1* is a prominent protein residing on LDs, but inactive in the absence of the ER-resident reductase (Leber *et al.* 1998). Thus, localization of *Erg1* to LD may serve as a reservoir to control the catalytic capacity of the ER-resident sterol biosynthetic pathway. Absence of LDs leads to relocalization of *Erg1* to the ER membrane and its partial degradation to regulate the concentration of *Erg1* in the ER (Sorger *et al.* 2004).

Notably, induction of LDs was observed in cells expressing mammalian α -synuclein (Outeiro and Lindquist 2003). α -Synuclein is implicated in neurodegenerative diseases, such as Parkinson’s and Alzheimer’s, and its expression in yeast is toxic and leads to impaired vesicular trafficking and inhibits phospholipase D expression (Outeiro and Lindquist 2003). Notably, yeast quadruple mutants lacking *Dgal Lro1 Are1* and *Are2* acyltransferases and, therefore, LDs altogether (see above) are more resistant to α -synuclein expression (Sere *et al.* 2010). In the quadruple mutant, the basal levels of reactive oxygen species (Sere *et al.* 2010) as well as unfolded protein response (Petschnigg *et al.* 2009) are elevated in the absence of neutral lipid synthesis; thus it was suggested that upregulated oxidative defense mechanisms may protect LD-deficient cells from α -synuclein toxicity. Notably, the sterol precursor, squalene, may play an important role in oxidative stress defense (Sere *et al.* 2010). Squalene accumulates in LDs, but also in subcellular membranes, when LDs are absent (Spanova *et al.* 2010, 2012).

Physiological interaction of LDs with other organelles

It is currently unclear whether and how TAG homeostasis and peroxisome (PEX) function (see *Peroxisomes*) are coupled in yeast, despite their apparent close physical interaction *in vivo* (Binns *et al.* 2006). A physiological interaction between these organelles may be restricted to FA β -oxidation, which, in *S. cerevisiae*, occurs exclusively in peroxisomes and is absent from mitochondria, which are the major site of β -oxidation in mammalian cells. *Tgl3* and *Tgl4* lipases are not required for induction of peroxisome formation after glucose depletion in the absence of exogenous FA supplementation (Petschnigg *et al.* 2009). By using bi-molecular fluorescence complementation, Pu *et al.* (2011) identified several interactions of LD-resident proteins with peroxisomal and mitochondrial proteins, indicating their direct physical interaction. This analysis technique is based on the reconstitution of a fluorescent protein (Venus) from two nonfluorescent fragments that is driven by the interaction of two proteins fused to these fragments. Most significant interactions were observed for the LD-resident *Erg6* and *Pet10* with other LD proteins, but also with mitochondrial and peroxisomal proteins (Pu *et al.* 2011). According to this analysis, *Tgl3* lipase interacts with the *Ayr1* protein that catalyzes the reduction of *1*-acyl-DHAP to lyso-PA (Figure 2), thus indicating a feedback loop between lipolysis and *de novo* glycerolipid synthesis. Furthermore, the interaction of the TAG lipase *Tgl3* with the peroxisomal protein *Pex11* is

consistent with the concept of metabolic channeling of TAG-derived FAs to peroxisome biogenesis and β -oxidation.

Yeast as a model to investigate and understand lipid-associated disorders

The pandemic development of obesity and related disorders, such as cardiovascular disease and type 2 diabetes, has led to significant efforts to study the molecular basis of lipid-associated disorders in various experimental model systems, including mice, worms, flies, and yeast. The similarities in the metabolic pathways involved in TAG and SE metabolism between yeast and “larger” cells make yeast an attractive experimental system to study lipid function and malfunction at the molecular and cellular levels (Kohlwein and Petschnigg 2007; Kohlwein 2010a,b; Zechner *et al.* 2012). Although lipid-associated disorders typically affect multiple organs and cell types, fundamental insights into key factors of mammalian TAG synthesis have been generated in yeast, *e.g.*, by the discovery of lipin function as a phosphatidic acid hydrolyase. Lipin is a protein known for a long time to be implicated in lipodystrophies in mouse model systems (Peterfy *et al.* 2001; Carman and Han 2006; Reue and Zhang 2008; Csaki and Reue 2010; Han and Carman 2010), yet its enzymatic function was identified by studies in yeast (Han *et al.* 2006, 2007; O'Hara *et al.* 2006; Carman and Wu 2007).

Sah1, a key enzyme in methylation metabolism, is one of the most highly conserved proteins, sharing some 60% sequence identity between yeast and the mammalian enzymes (Tehlivets *et al.* 2004; Malanovic *et al.* 2008; Tehlivets 2011). Notably, the *SAH1* gene in yeast was found to be transcriptionally coregulated with phospholipid biosynthetic genes, indicating a functional link between methylation and lipid metabolism. Indeed, phospholipid methylation is a major consumer of *S*-adenosylmethionine (AdoMet), leading to the accumulation of *S*-adenosylhomocysteine (AdoHcy) as a by-product, which also acts as a potent product inhibitor. *Sah1* is responsible for the degradation of AdoHcy to adenosine and homocysteine, thus regulating the cellular methylation potential (Tehlivets 2011); however, this reaction is reversible, and an accumulation of homocysteine may, in fact, drive the formation of the methylation inhibitor, AdoHcy. This puts the risk factor for atherosclerosis, homocysteine, into a new perspective, as a regulator of cellular methylation by its *Sah1*-dependent conversion to AdoHcy (Tehlivets 2011). Since *Sah1* is an essential enzyme, studies in mouse model systems are very limited due to embryonic lethality of the *ko* mutation. On the other hand, mammalian *Sah1* complements a yeast *sah1* deletion, providing an attractive experimental system for studies on structure–function relationships (Malanovic *et al.* 2008; Tehlivets 2011).

A high level of functional or structural conservation was also observed for the “lipolytic cascade,” a sequence of enzymatic steps that results in complete TAG degradation via DAG and MAG to glycerol and free FAs (Kurat *et al.* 2006; Zechner *et al.* 2012). These reactions are governed in yeast by the *Tgl3* and *Tgl4* TAG lipases, of which *Tgl3* also harbors

DAG lipase activity (Kurat *et al.* 2006); the final step of MAG hydrolysis is catalyzed by *Yju3* in yeast (Heier *et al.* 2010). In mammals, ATGL is responsible for TAG breakdown to DAG, which is further subject to degradation by hormone-sensitive lipase and monoacylglycerol lipase. Notably, *Tgl3*, *Tgl4*, and ATGL are members of the patatin-domain-containing family of enzymes (Kienesberger *et al.* 2009), and *Tgl4* deficiency in yeast can be functionally complemented by mouse or human ATGL, which also correctly localizes to the LD, despite the absence of perilipins in yeast that regulate access of ATGL to LDs in mammalian cells (Kurat *et al.* 2006). Lypolysis clearly not only provides FAs for β -oxidation and energy production but also generates TAG-derived signaling molecules important for mitochondrial function in mammals (Zechner *et al.* 2012) or cell cycle progression in yeast (Kurat *et al.* 2009).

Many of the enzymes involved in TAG synthesis have orthologs in mammals and are now extensively studied with respect to their topology in the ER membrane and with respect to their contribution to LD formation (Choudhary *et al.* 2011; Jacquier *et al.* 2011; Pagac *et al.* 2011). A class of proteins implicated in TAG storage and LD biogenesis are the mammalian FIT proteins [fat storage-inducing transmembrane proteins (Kadereit *et al.* 2008; Gross *et al.* 2010, 2011; Moir *et al.* 2012)], of which two orthologs exist in yeast, encoded by *SCS3* and *YFT2*. Yeast mutants lacking *Scs3* are inositol auxotrophs, indicating a functional link to the transcriptional regulation of phospholipid synthesis (Henry *et al.* 2012); however, their specific roles in TAG metabolism and LD formation in yeast are unknown. Large-scale interaction studies indicate that both *SCS3* and *YFT2* have shared and unique functions and may be required for ER membrane biosynthesis in response to perturbations in lipid metabolism and ER stress (Moir *et al.* 2012). In two independent imaging-based screens, Fei *et al.* (2008, 2011b) and Szymanski *et al.* (2007) identified yeast mutants with aberrant LD morphology (“supersized LDs”) that are defective in the *FLD1* gene. *FLD1* is an ortholog of the mammalian *BSC2* gene encoding seipin that is associated with severe inherited Berardinelli–Seip congenital lipodystrophy type 2 (Agarwal and Garg 2003; Agarwal *et al.* 2004; Garg and Agarwal 2009). Notably, the supersized LD phenotype in yeast *fld1* mutants can be complemented by mammalian wild-type *BSC2*, supporting the high level of functional conservation in mammals and yeast (Fei *et al.* 2008, 2011b).

In addition to the lipogenic and lipolytic pathways, the major regulatory and signaling processes, such as TOR and AMPK/*Snf1*, are also conserved in yeast (Zaman *et al.* 2008; De Virgilio 2012) and apparently are connected to lipid homeostasis (Bozaquel-Morais *et al.* 2010); however, this “lipid connection” clearly needs further exploration. Notably, many yeast wild-type strains significantly differ in their LD content, but the genetic basis for these diverse lipid phenotypes is not known. This observation clearly reflects the polygenic nature of lipid and energy metabolism in mammals and requires QTL analysis to obtain further insight into

the specific contributions of genes and pathways to cellular lipid homeostasis.

Peroxisomes

Microbodies were first described on the basis of their simple morphology in mouse renal tubule cells by Rhodin (1954). Later, biochemical functions could be attributed to these organelles, and they were functionally subdivided into peroxisomes (containing at least one hydrogen peroxide-producing oxidase and catalase), glyoxysomes (containing enzymes of the glyoxylate cycle) (Tolbert and Essner 1981), glycosomes (harboring glycolytic enzymes, observed only in trypanosomes) (Oppenheimer and Borst 1977), Woronin bodies that are involved in plugging of septal pores in filamentous ascomycetes (Dhavale and Jedd 2007), and hydrogenosomes (producing hydrogen, observed only in anaerobic fungi) (Martin and Muller 1998). Of these, peroxisomes were first characterized as organelles implicated in hydrogen peroxide metabolism. However, today various other peroxisomal functions are known (for reviews see Nyathi and Baker 2006; van der Klei and Veenhuis 2006b; Wanders and Waterham 2006), including biosynthetic [e.g., in secondary metabolite biosynthesis (Bartoszewska *et al.* 2011)] and non-metabolic ones [e.g., in the innate immune response (Lazarow 2011)].

In yeast, the morphology of microbodies was described for the first time in *S. cerevisiae* by Avers and Federman (1968). However, it took almost 20 years before collaborative efforts of the groups of Veenhuis and Kunau demonstrated that peroxisomes play a crucial role in oleate metabolism in *S. cerevisiae* and that, consequently, these organelles are massively induced during growth of yeast on oleate as the sole carbon and energy source (Veenhuis *et al.* 1987). These findings opened a new era in peroxisome research, which contributed to the identification of the first *PEX* genes (Erdmann *et al.* 1989, 1991) involved in peroxisome development. This information subsequently allowed unraveling of the principles of human peroxisome biogenesis disorders because of the strong conservation of the molecular mechanisms of peroxisome development between lower and higher eukaryotes.

Peroxisome composition

Peroxisomes consist of a single membrane encompassing a protein-rich matrix. A typical feature of peroxisomal membranes is their low protein content, which is supported by their very smooth fracture faces in freeze-etch replicas (Figure 4A). The peroxisomal matrix is generally considered to represent the site of the highest protein concentration in eukaryotic cells. Indeed, peroxisomes are the highest density organelles after density gradient centrifugation of post-nuclear cell homogenates. The high protein concentration in these organelles often results in the formation of electron dense inclusions or protein crystalloids in the peroxisomal matrix, e.g., in peroxisomes of plants or methylotrophic yeasts. However, in wild-type *S. cerevisiae* this is never observed (Figure 4D): in this organism, peroxisome proliferation is induced

by oleate and strongly repressed by glucose (Figure 4, C and D). Repression of peroxisome proliferation is especially evident in certain *S. cerevisiae* strains such as G910 (Veenhuis *et al.* 1987). These cells contain only one or very few peroxisomes during exponential growth on glucose (Figure 4C), in contrast to oleate-grown cells in which the abundance and size of peroxisomes is strongly increased (Figure 4D). However, in various currently used strains, peroxisome numbers in glucose-grown cells are only slightly lower compared to cells grown on oleate (Figure 4, E and F; *S. cerevisiae* BY4742).

The peroxisomal matrix almost exclusively contains enzymes, which generally harbor cofactors and are mostly oligomeric. Peroxisomal membrane-bound enzymes are rare. This may explain the relatively low membrane surface/volume ratio and the very low protein/phospholipid ratio of peroxisomal membranes. All characterized peroxisomal membrane proteins (PMPs) are involved in peroxisome biogenesis and dynamics or in solute transport processes; protein modifications, such as glycosylation or phosphorylation, have been reported for only a few PMPs and are absent from matrix proteins.

The *Saccharomyces cerevisiae* Genome Database currently contains 66 proteins that have been demonstrated to reside at peroxisomes (listed in Table 2). Of these, 24 encode enzymes, whereas only 3 represent membrane transporters for solutes. The remaining proteins are peroxins or proteins involved in various other peroxisomal processes, such as fission or inheritance (Table 2).

The lipid composition of oleate-grown *S. cerevisiae* peroxisomes has been determined by Zinser *et al.* (1991). The peroxisomal membrane contains the major cellular phospholipids—phosphatidylcholine (48.2%), phosphatidylethanolamine (22.9%), and phosphatidylinositol (15.8%)—but also has a remarkably high cardiolipin content (7%). The relative abundance of cardiolipin is noteworthy since this lipid is synthesized in mitochondria (Henry *et al.* 2012). The other lipids are derived from the ER; however, the mechanisms by which these lipids reach the peroxisomes are not yet firmly established; some evidence suggests that this process involves vesicular transport both from the ER and from mitochondria (Braschi *et al.* 2010).

Peroxisome metabolic functions

By definition, peroxisomes contain at least one hydrogen peroxide-producing oxidase together with catalase, which decomposes the hydrogen peroxide by-product of the oxidation reaction. *S. cerevisiae* contains only one oxidase, namely the flavo-enzyme *Pox1* (acyl-CoA oxidase), an enzyme of the β -oxidation pathway. Unlike most other species, *S. cerevisiae* contains a second, cytosolic catalase T, *Ctt1*, in addition to the peroxisomal catalase A isoenzyme, *Cta1* (Skoneczny *et al.* 1988).

The two best-characterized peroxisomal metabolic pathways in *S. cerevisiae* are the β -oxidation of fatty acids (Poirier *et al.* 2006) (Figure 2B, Figure 5) and the glyoxylate




Figure 4 Morphological characteristics of yeast peroxisomes. (A) Freeze-etch replica of oleic acid-grown *S. cerevisiae* cells showing the fraction faces of the different organelles. Peroxisomes contain very smooth fracture faces, indicative of a low abundance of integral membrane proteins. (B) Thin section of a cell grown on oleic acid cytochemically stained for catalase activity, using diaminobenzidine and hydrogen peroxide. In these cells, numerous stained, electron-dense peroxisomes are present. Staining of the mitochondrial cristae is due to cytochrome c peroxidase activity. Ultrathin sections of (C) glucose-grown and (D) oleic acid-grown and KMnO_4 -fixed *S. cerevisiae* G910 cells (Veenhuis *et al.* 1987). The cell grown on glucose displays only a few very small peroxisomes (arrows), whereas strong peroxisome proliferation is evident in the oleic acid-grown cell. (E and F) Fluorescence microscopy of (E) glucose-grown and (F) oleic acid-grown cells expressing GFP-SKL to label peroxisomes in wild-type strain BY4742. Notably, in this strain background, the difference in peroxisome number on glucose and oleic acid media is far less pronounced compared to strain G910. N, nucleus; M, mitochondrion; P, peroxisome; V, vacuole; LD, lipid droplet. Scale bar: 200 nm in A, 1 μm in B–D, and 3 μm in E and F.

cycle (Kunze *et al.* 2006) (Figure 5). In most other yeast species (*e.g.* *Candida tropicalis*, *Hansenula polymorpha*, *Pichia pastoris*), several other important peroxisome-bound pathways occur, such as the metabolism of alkanes and methanol and the oxidation of various organic nitrogen sources such as primary amines, purines, and *D*-amino acids (for a review see van der Klei and Veenhuis 2006b).

β -Oxidation: *S. cerevisiae* can utilize a range of saturated and unsaturated fatty acids of different chain lengths, which, in the absence of glucose, are oxidized by peroxisomal β -oxidation. β -Oxidation involves four steps, namely CoA activation, oxidation, hydratation/dehydrogenation, and thiolyl cleavage to generate acetyl-CoA and an acyl chain that is shortened by two carbon atoms (Figure 2B, Figure 5). Activation of medium chain fatty acids occurs in the organelle matrix via acyl-CoA synthetase, *Faa2* (Figure 5). This process requires ATP, which is imported into the organelle by the adenine nucleotide transporter *Ant1* (Table 2) (Palmieri *et al.* 2001). It has been suggested that *Faa2* produces AMP and pyrophosphate; hence, most likely *Ant1* exchanges AMP for ATP across the peroxisomal membrane.

Long-chain fatty acids are activated outside the organelle by *Fat1* and taken up as CoA esters via the heterodimeric ABC transporter consisting of *Pxa1* and *Pxa2* (Hettema *et al.* 1996) (Figure 5). A portion of the cellular *Fat1* activity is associated with peroxisomes (Watkins *et al.* 1998). The acetyl-CoA product of the β -oxidation is transported to mitochondria for further oxidation by the citric acid cycle. Export from peroxisomes occurs via two different pathways, namely via carnitine-dependent acetyl-CoA transport that involves *Cat2* or via the glyoxylate cycle (Figure 5). NADH is transferred to the cytosol via the malate shuttle (Figure 5) (for an excellent review see van Roermund *et al.* 2003).

For β -oxidation of unsaturated fatty acids with *trans* and *cis* double bonds at odd-numbered positions or *cis* double bonds at even positions, auxiliary peroxisomal enzymes are required, *i.e.*, *Eci1*, *Dci1*, *Tes1*, and *Sps19* (van Roermund *et al.* 2003) (Table 2, Figure 2). For example, oleic acid metabolism requires the auxiliary enzyme *Eci1*, which catalyzes the isomerization of the *cis* double bond of oleate after shortening of the oleoyl-CoA chain by three rounds of β -oxidation (Kunau *et al.* 1995). *Eci1* also isomerizes a fraction of 2-*trans*, 5-*cis*-tetradecadienoyl-CoA, an

Table 2 Peroxisomal proteins

Gene	Required for growth on oleate	Expression induced by oleate	Enzyme/activity	Molecular mass (kDa)	Isoelectric point	Molecules per cell ^a	Localization ^b	Function
β-Oxidation enzymes								
<i>PCS60 (FAT2)</i>	No	Yes	Medium chain fatty acyl-CoA synthetase	60.5	9.98	8,770	Peripheral peroxisomal membrane and matrix	Activates fatty acids with a preference for medium chain lengths, C9–C13
<i>FAT1</i>	No		Very long chain fatty acyl-CoA synthetase and long chain fatty acid transporter	77.1	8.47	16,900	Lipid droplet, ER, peroxisome Three predicted TM	Activates fatty acids with a preference for very long chain lengths, C20–C26
<i>POX1</i>	Yes	Yes	Acyl-CoA oxidase	84.0	8.73	ND	Peroxisomal matrix	Oxidation of acyl-CoA
<i>CTA1</i>	No	Yes	Catalase	58.6	7.46	623	Peroxisomal matrix	Degrades hydrogen peroxide produced by Pox1
<i>FOX2 (POX2)</i>	Yes	Yes	Multifunctional enzyme; 3-hydroxyacyl-CoA dehydrogenase and enoyl-CoA hydratase	98.7	9.75	ND	Peroxisomal matrix	
<i>POT1 (FOX3, POX3)</i>	Yes	Yes	3-Ketoacyl-CoA thiolase	44.7	7.56	ND	Peroxisomal matrix	Cleaves 3-ketoacyl-CoA into acyl-CoA and acetyl-CoAs
<i>DCI1 (ECI2)</i>			$\Delta(3,5)\text{-}\Delta(2,4)\text{-dienoyl-CoA}$ isomerase (putative)	30.1	8.83	ND	Peroxisomal matrix	Auxiliary enzyme of fatty acid β -oxidation; role in β -oxidation debated
<i>SPS19 (SPX19)</i>	Yes	Yes	2,4-Dienoyl-CoA reductase	31.1	9.67	ND	Peroxisomal matrix	Auxiliary enzyme of fatty acid β -oxidation
<i>ECI1</i>	Yes	Yes	$\Delta 3, \Delta 2$ -enoyl-CoA isomerase	31.7	8.21	ND	Peroxisomal matrix	Auxiliary enzyme of fatty acid β -oxidation
<i>TES1 (PTE1)</i>	Yes	Yes	Acyl-CoA thioesterase	40.3	9.58	ND	Peroxisomal matrix	Auxiliary enzyme of fatty acid β -oxidation
<i>MDH3</i>	Yes	Yes	Malate dehydrogenase	37.3	10.00	3,300	Peroxisomal matrix	Required for the malate-oxaloacetate shuttle, to exchange peroxisomal NADH for cytosolic NAD ⁺ ; part of the glyoxylate cycle

(continued)

Table 2, continued

Gene	Required for growth on oleate	Expression induced by oleate	Enzyme/activity	Molecular mass (kDa)	Isoelectric point	Molecules per cell ^a	Localization ^b	Function
<i>IDP3</i>	Yes	Yes	NADP ⁺ dependent isocitrate dehydrogenase	47.91	10.02	ND	Peroxisomal matrix	Required for the 2-ketoglutarate/isocitrate shuttle, exchanging peroxisomal NADP ⁺ for cytosolic NADPH
<i>CAT2</i>	No	No	Carnitine acetyl-CoA transferase	77.2	8.34	470	Peroxisome, mitochondria	Transfers activated acetyl groups to carnitine to form acetyl/carnitine which can be shuttled across membranes
<i>CIT2</i>	No			Glyoxylate cycle				
			Citrate synthase	51.4	6.34	2,310	Peroxisomal matrix	Condensation of acetyl CoA and oxaloacetate to form citrate
<i>MDH3</i>	Yes	Yes	Malate dehydrogenase	37.3	10.00	3,300	Peroxisomal matrix	Required for the malate-oxaloacetate shuttle, to exchange peroxisomal NADH for cytosolic NAD ⁺
<i>MLS1</i>	Yes	Yes	Malate synthase	62.8	7.18	ND	Peroxisomal matrix	Required for utilization of nonfermentable carbon sources
<i>GPD1 (DART1, HOR1, OSG1, OSR5)</i>			Other peroxisome-associated enzyme activities					
			NAD-dependent glycerol-3-phosphate dehydrogenase	42.9	5.26	807	Peroxisome, cytosol, nucleus	Key enzyme of glycerol synthesis, essential for growth under osmotic stress
<i>PNC1</i>			Nicotinamidase	25.0	6.23	7,720	Peroxisome, cytosol	Converts nicotinamide to nicotinic acid as part of the NAD(+) salvage pathway
<i>NPY1</i>			NADH diphosphatase	43.5	6.26	846	Peroxisome, cytosol	Hydrolyzes the pyrophosphate linkage in NADH and related nucleotides
<i>STR3</i>			Cystathionine β-lyase	51.8	7.96	ND	Peroxisome	Converts cystathionine into homocysteine
<i>GTO1</i>			ω-Class glutathione transferase	41.3	9.53		Peroxisome	Induced under oxidative stress

(continued)

Table 2, continued

Gene	Required for growth on oleate	Expression induced by oleate	Enzyme/activity	Molecular mass (kDa)	Isoelectric point	Molecules per cell ^a	Localization ^b	Function
AAT2 (ASP5)	Yes	Yes	Aspartate aminotransferase	46.1	8.50	7,700	Cytosol, peroxisome	Involved in nitrogen metabolism
PCD1			Nudix pyrophosphatase with specificity for coenzyme A and CoA derivatives	39.8	6.59	238	Peroxisome	May function to remove potentially toxic oxidized CoA disulfide from peroxisomes
LPX1	Yes	Yes	Triacylglycerol lipase	43.7	8.16	2,350	Peroxisomal matrix	
Peroxisomal transporters								
PXA1 (LPI1, PAL1, PAT2, SSH2)			Subunit of a heterodimeric ATP-binding cassette transporter complex	100.0	10.34	ND	Peroxisomal membrane	Import of long-chain fatty acids into peroxisomes
PXA2 (PAT1)			Subunit of a heterodimeric ATP-binding cassette transporter complex	97.1	9.47	ND	Peroxisomal membrane	Import of long-chain fatty acids into peroxisomes
ANT1 (YPR128C)			Adenine nucleotide transporter	36.4	10.6	2,250	Peroxisomal membrane	Involved in β -oxidation of medium-chain fatty acids
Peroxisins								
PEX1 (PAS1)			AAA ATPase	117.3	6.93	2,100	Peroxisomal membrane	Involved in recycling of Pex5, forms heterodimer with Pex6
PEX2 (CRT1, PAS5)			E3 ubiquitin ligase	30.8	9.02	339	Peroxisomal membrane	RING finger protein, forms complex with Pex10 and Pex12. Involved in matrix protein import
PEX3 (PAS3)				50.7	6.29	1,400	Peroxisomal membrane	Required for proper localization of PMPs
PEX4 (PAS2, UBC10)			Ubiquitin-conjugating enzyme	21.1	5.36	ND	Peroxisomal membrane	Involved in matrix protein import
PEX5 (PAS10)			Soluble PTS1 receptor	69.3	4.79	2,070	Cytosol and peroxisomal matrix	Required for import of PTS1-containing peroxisomal proteins, contains TPR domains
PEX6 (PAS8)			AAA ATPase	115.6	5.44	1,630	Peroxisomal membrane	Involved in recycling of Pex5, forms heterodimer with Pex1
PEX7 (PAS7, PEB1)			Soluble PTS2 receptor	42.3	8.34	589	Cytosol and peroxisomal matrix	Requires Pex18 and Pex21 for association to the receptor docking site, contains WD40 repeat

(continued)

Table 2, continued

Gene	Required for growth on oleate	Expression induced by oleate	Enzyme/activity	Molecular mass (kDa)	Isoelectric point	Molecules per cell ^a	Localization ^b	Function
PEX8 (PAS6)			Intraperoxisomal organizer of the peroxisomal import machinery E3 ubiquitin ligase	68.2	7.62	538	Peroxisomal matrix and luminal membrane face	Pex5-cargo dissociation
PEX10 (PAS4)			E3 ubiquitin ligase	39.1	9.88	ND	Peroxisomal membrane	RING finger protein involved in Ubc4-dependent Pex5 ubiquitination. Forms complex with Pex2 and Pex12
PEX11 (PMP24, PMP27)				26.9	10.65	1,630	Peroxisomal membrane	Involved in peroxisome fission, required for medium-chain fatty acid oxidation
PEX12 (PAS11)			E3 ubiquitin ligase	46.0	9.86	907		RING finger protein, forms complex with Pex2 and Pex10
PEX13 (PAS20)			Component of docking complex for Pex5 and Pex7	42.7	9.83	7,900	Peroxisomal membrane	Forms complex with Pex14 and Pex17
PEX14			Central component of the receptor docking complex	38.4	4.61	2,570	Peroxisomal membrane	Interacts with Pex13
PEX15 (PAS21)				43.7	8.42	1,070	Peroxisomal membrane	Recruits Pex6 to the peroxisomal membrane, tail anchored PMP
PEX17 (PAS9)			Component of docking complex for Pex5 and Pex7	23.2	10.24	656	Peroxisomal membrane	Forms complex with Pex13 and Pex14
PEX18			Required for PTS2 import	32.0	4.78	ND		Interacts with Pex7; partially redundant with Pex21
PEX19 (PAS12)			Chaperone and import receptor for newly synthesized PMPs	38.7	4.08	5,350	Cytosol, peroxisome; farnesylated	Interacts with PMPs, involved in PMP sorting. Also interacts with Myo2 and contributes to peroxisome partitioning
PEX21			Required for PTS2 protein import	33.0	6.67	ND	Cytosol	Interacts with Pex7, partially redundant with Pex18
PEX22 (YAF5)			Required for import of peroxisomal proteins	19.9	8.33	259	Peroxisomal membrane	Recruits Pex4 to the peroxisomal membrane
PEX25			Involved in the regulation of peroxisome size and maintenance, required for re-introduction of peroxisomes in peroxisome deficient cells	44.9	9.77	2,420	Peripheral peroxisomal membrane	Recruits GTPase Rho1 to peroxisomes, interacts with homologous protein Pex27

(continued)

Table 2, continued

Gene	Required for growth on oleate	Expression induced by oleate	Enzyme/activity	Molecular mass (kDa)	Isoelectric point	Molecules per cell ^a	Localization ^b	Function
<i>PEX27</i>			Involved in the regulation of peroxisome size and number	44.1	10.49	382	Peripheral peroxisomal membrane	Interacts with homologous protein Pex25
<i>PEX28</i>			Involved in the regulation of peroxisome size, number and distribution	66.1	7.09	ND	Peroxisomal membrane	May act upstream of Pex30, Pex31 and Pex32
<i>PEX29</i>			Involved in the regulation of peroxisome size, number and distribution	63.5	6.8	5,040	Peroxisomal membrane	May act upstream of Pex30, Pex31 and Pex32
<i>PEX30</i>			Involved in the regulation of peroxisome number	59.5	5.59	4,570	Peroxisomal membrane	Negative regulator, partially functionally redundant with Pex31 and Pex32
<i>PEX31</i>			Involved in the regulation of peroxisome number	52.9	10.15	238	Peroxisomal membrane	Negative regulator, partially functionally redundant with Pex30 and Pex32
<i>PEX32</i>			Involved in the regulation of peroxisome number	48.6	9.14	ND	Peroxisomal membrane	Negative regulator, partially functionally redundant with Pex30 and Pex31
<i>PEX34</i>			Involved in the regulation of peroxisome number	16.6	10.30	ND	Peroxisomal membrane	
<i>DYN2 (SLC1)</i>								
<i>SEC20</i>								
Peroxisome fission and inheritance								
			Light chain dynein v-SNARE	10.4 43.9	9.03 5.92	1,310 4,910	Cytosol Golgi, ER	Microtubule motor protein Involved in retrograde transport from the Golgi to the ER; interacts with the Dsl1 complex through Tip20
<i>SEC39 (DSL3)</i>			Component of the Dsl1p-tethering complex	82.4	4.65	1,840	ER, nuclear envelope	Proposed to be involved in protein secretion
<i>DSL1 (RNS1)</i>			Component of the ER target site that interacts with coatomer	88.1	4.69	8,970	Peripheral ER, Golgi membrane	Forms a complex with Sec39 and Tip20 that interacts with ER SNAREs Sec20 and Use1
<i>FIS1 (MDV2)</i>			Required for peroxisome fission	17.7	9.87	2,410	Peroxisomal membrane, mitochondria	Tail-anchored protein; recruits Dnm1 via Mdv1/Caf4; also involved in mitochondrial fission
<i>DNM1</i>			GTPase, dynamin-like protein involved in peroxisome fission	85.0	5.25	9,620		Also involved in mitochondrial fission

(continued)

Table 2, continued

Gene	Required Expression induced on oleate	Enzyme/activity	Molecular mass (kDa)	Isoelectric point	Molecules per cell ^a	Localization ^b	Function
VPS1 (GRD1, LAM1, SPO15, VPL1, VPT26) VPS34 (END12, PEP15, VPL7, VPT29, STT8, VPS7) INP1		GTPase, dynamin-like protein involved in peroxisome fission Phosphatidylinositol 3-kinase Involved in retention of peroxisomes in mother cells	78.7 100.9 47.3	8.15 7.79 8.34	5,960 1,080 639	Peroxisomal membrane	Also involved in vacuolar protein sorting Forms complex with Vps15 Recruited to the peroxisome by binding to Pex3
INP2		Myo2 receptor, involved in peroxisome inheritance	81.5	9.41	736	Peroxisomal membrane	
RHO1		GTP-binding protein of the Rho subfamily of Ras-like proteins; involved in actin assembly at the peroxisome	23.2	6.07	ND		Involved in <i>de novo</i> peroxisome formation, recruited to peroxisomes by Pex25

Much of the information in the table may be found in the *Saccharomyces* Genome Database. ND, not determined.

^a Ghaemmaghami *et al.* (2003).

^b Kumar *et al.* (2002); Huh *et al.* (2003); Kals *et al.* (2005); Natter *et al.* (2005); Wolinski *et al.* (2009b).

intermediate of oleic acid β -oxidation, to 3,5-*cis*-tetradecadienoyl-CoA, which has two conjugated double bonds that prevent further β -oxidation. Gurvitz *et al.* (1999) proposed that 3,5-*cis*-tetradecadienoyl-CoA is oxidized by the reductase-dependent pathway and involves the dienoyl isomerase *Dci1*. However, recent data also indicate that the thioesterase-dependent pathway, which involves the acyl-CoA thioesterase *Tes1*, is operative in *S. cerevisiae* (Ntamack *et al.* 2009).

The β -oxidation of unsaturated fatty acids with a double bond at an even position requires the function of the NADPH-dependent 2,4-dienoyl-CoA reductase *Sps19*. The NADP⁺ generated by this reaction is reduced by the peroxisomal isocitrate dehydrogenase *Idp3*, and, for the regeneration of NADPH, an isocitrate/2-oxoglutarate shuttle exists (reviewed in van Roermund *et al.* 2003).

The role of *Pcs60/Fat2*, which is associated with the inside of the peroxisomal membrane, is still unclear. *Pcs60/Fat2* belongs to the family of proteins that act via an ATP-dependent covalent binding of AMP to their substrates and shows high similarity to *Escherichia coli* long-chain acyl-CoA synthetases (Blobel and Erdmann 1996). However, *Pcs60/Fat2* is not required for growth of *S. cerevisiae* on oleate. Interestingly, this protein binds mRNAs encoding proteins involved in triglyceride metabolism (Tsvetanova *et al.* 2010).

Glyoxylate cycle: The glyoxylate cycle allows cells to convert two acetyl-CoA molecules into succinate, which can be used to replenish the citric acid cycle or to function as precursors for amino acid or carbohydrate biosynthesis (Figure 5). In yeast, this cycle is essential for growth on oleate or C2 substrates such as ethanol or acetate. In *S. cerevisiae*, three glyoxylate cycle enzymes are cytosolic, namely the malate dehydrogenase *Mdh2*, the aconitase *Aco1*, and the isocitrate lyase *Icl1*, whereas two are peroxisomal: namely citrate synthase *Cit2* and malate synthase *Mls1* (McCammon *et al.* 1990; Taylor *et al.* 1996) (Figure 5). Remarkably, in most other yeast species, in plants, and in filamentous fungi, isocitrate lyase also is a peroxisomal enzyme (for a review see Kunze *et al.* 2006). The presence of *Icl1* in the cytosol of *S. cerevisiae* is surprising as it catalyzes the production of the reactive compound glyoxylate. Thus, the compartmentalization of the enzymes of the glyoxylate cycle predicts the presence of solute transporters for glyoxylate cycle intermediates in the peroxisomal membrane (Figure 5). It is, however, also possible that no specific transport proteins are required since yeast peroxisomal membranes contain pore-forming proteins that allow passage of small molecules with a molecular mass up to 400 Da, which is sufficient to allow passage of these intermediates (Antonenkov *et al.* 2009). However, the genes encoding these pore-forming proteins have not yet been identified.




Figure 5 Compartmentalization of the β -oxidation pathway and the glyoxylate cycle. The enzymes of the β -oxidation (see Figure 2B) as well as key enzymes of the glyoxylate cycle are localized to peroxisomes in *S. cerevisiae*. In addition to the β -oxidation enzymes, three peroxisomal membrane-associated proteins are required for fatty acid oxidation in peroxisomes, namely the ABC transporter Pxa1/Pxa2 for the import of long-chain acyl-CoA, Faa2 for the activation of medium chain FAs, and the transporter Ant1 for import of ATP. The glyoxylate cycle converts two acetyl-CoA molecules into succinate and contributes to the export of acetyl-CoA that is produced in the β -oxidation cycle. Glyoxylate and the first acetyl-CoA molecule are condensed by malate synthase (Mls) to malate; malate dehydrogenase (Mdh) converts malate to oxaloacetate (OAA); and isocitrate synthase (Cit) condenses OAA and a second acetyl-CoA molecule to form citrate. Aconitase (Aco) catalyzes the isomerization of citrate into isocitrate, which is cleaved by isocitrate lyase (Icl) into succinate and glyoxylate. Glyoxylate can be used for the next round of the glyoxylate cycle, whereas succinate is used to replenish the citric acid cycle or to function as a precursor for amino acid or carbohydrate biosynthesis. In *S. cerevisiae*, citrate synthase, Cit2, and malate synthase Mls1 are peroxisomal enzymes, whereas Icl1 is cytosolic. This is in contrast to plants, filamentous fungi, and other yeast species, in which Icl is peroxisomal as well. Acetyl-CoA can also be exported via the carnitine shuttle, which involves peroxisomal Cat2. The malate shuttle is responsible for NADH export. The predicted small molecule transporters involved in both shuttles or in the glyoxylate cycle have not been identified yet.

Other peroxisomal enzymes: Except for the key enzymes involved in fatty acid utilization and the glyoxylate cycle, additional enzymes are (perhaps only transiently) associated with peroxisomes in *S. cerevisiae*. However, for many of these proteins the physiological relevance is unclear (Table 2). Moreover, in some cases their peroxisomal localization has not yet been firmly established. Some examples are detailed below. Lpx1 is a peroxisomal protein that has acyl hydrolase and phospholipase A activity *in vitro*. Deletion of *L PX1* results in peroxisomes with aberrant morphology that is characterized by intraperoxisomal vesicles or invaginations, which may point to a role of vesicle fusions in peroxisome development (Thoms *et al.* 2008). Several peroxisomal matrix proteins have been implicated to play a role in (oxidative) stress response or aging. Glycerol-3-phosphate dehydrogenase 1 (Gpd1) is a key enzyme in glycerol biosynthesis and essential for *S. cerevisiae* to cope with osmotic stress (Albertyn *et al.*

1994); its expression is regulated by the high-osmolarity glycerol response pathway. Fluorescence microscopy (Huh *et al.* 2003) and proteomics analysis of peroxisomal fractions (Marelli *et al.* 2004) indicated a (partial) peroxisomal localization of Gpd1. This was recently confirmed by Jung *et al.* (2010) who showed that Gpd1 is targeted to peroxisomes via the PTS2 pathway (see below). However, under stress conditions, Gpd1 is localized to the cytosol and the nucleus, a process that is regulated by phosphorylation of a residue adjacent to the PTS2 in Gpd1. The physiological relevance of the different subcellular locations remains to be established.

Interestingly, Pnc1, whose import into peroxisomes also depends on the PTS2 pathway, shows a similar shift in localization in response to stress (Anderson *et al.* 2003). Moreover, the expression of *GPD1* and *PNC1* is strongly correlated (Jung *et al.* 2010). Pnc1 converts nicotinamide to nicotinic acid as part of the NAD⁺ salvage pathway. Nicotinamide

strongly inhibits Sir2, a protein important for life-span extension by calorie restriction (Kaeberlein *et al.* 1999), and, like Sir2, Pnc1 also functions in life-span extension (Anderson *et al.* 2003). Most likely, Pnc1 regulates longevity by reducing nicotinamide levels, which activates Sir2 (Gallo *et al.* 2004). Whether the localization of Pnc1 to peroxisomes is important for its function in life-span extension is not yet known.

GTO1 encodes a PTS1 containing ω -class peroxisomal glutathione transferase (Barreto *et al.* 2006), whose expression is induced by oxidative stress. The role of Gto1 may be related to the redox regulation of cystathionine β -lyase, Str3, another putative peroxisomal protein (Schafer *et al.* 2001; Yi *et al.* 2002). Str3 is involved in transulfuration of cysteine to homocysteine (Table 2).

Finally, two phosphatases, Npy1 and Pcd1, were found to be localized to peroxisomes. Npy1 is a PTS1-containing diphosphatase, which utilizes NADH as its preferred substrate. This enzyme may function in the regulation of nicotinamide coenzyme concentrations or in the elimination of oxidized nucleotide derivatives (Abdelraheim *et al.* 2001). Pcd1 is a PTS2-containing diphosphatase, which is active toward coenzyme A and its derivatives (Cartwright *et al.* 2000). A proposed role for this enzyme is the removal of potentially toxic oxidized CoA disulfide in peroxisomes (Cartwright *et al.* 2000).

Methods to identify proteins involved in peroxisome biology

Identification of peroxisomal enzymes: Peroxisome-borne proteins were first identified by biochemical analysis of fractions enriched in peroxisomes obtained by subcellular fractionation. This approach became feasible because of the pioneering work of the Nobel Laureate Christian de Duve, who was the first to isolate peroxisomes from rat liver (De Duve 1965). However, it took a decade before the first report appeared on the isolation of *S. cerevisiae* peroxisomes from derepressed cells (Parish 1975), and controversies regarding the yeast peroxisomal protein content remained over several years. Major improvements in the preparation techniques were achieved after the finding that peroxisome proliferation in *S. cerevisiae* can be induced by oleate (Veenhuis *et al.* 1987). Together with the optimization of cell fractionation protocols for *S. cerevisiae*, McCammon *et al.* (1990) were able to convincingly show the localization of enzymes of the glyoxylate and β -oxidation in highly purified peroxisomal fractions of oleate-grown yeast. Today, these procedures are classic in peroxisome research and still in use (among others) to determine the organelle proteome and to characterize defects in peroxisome assembly in mutant strains (for detailed protocols see Distel and Kragt 2006).

Discovery of peroxins by genetic approaches: Growth of *S. cerevisiae* on oleate requires intact peroxisomes; hence peroxisome-deficient mutants were readily selected from collections of oleate utilization-deficient mutants (Erdmann *et al.* 1989). The corresponding PEX genes were cloned by

functional complementation upon transformation of the selected mutant with a genomic library. In this way, the first PEX gene was identified in baker's yeast (Erdmann *et al.* 1991). Subsequently, similar approaches were used for other yeast species, including *Yarrowia lipolytica* and *Pichia pastoris*, two yeast species that are also capable of growing on oleate, as well as *Hansenula polymorpha*, which, like *P. pastoris*, can grow on methanol as the sole source of carbon and energy (van der Klei *et al.* 1991; Liu *et al.* 1992; Nuttley *et al.* 1994). In these latter two species, *H. polymorpha* and *P. pastoris*, peroxisomes are essential for growth on methanol, a property used for the identification of PEX genes. The identification of novel PEX genes in the other yeast species facilitated the work in *S. cerevisiae* because homologous genes were readily cloned on the basis of sequence homology, especially after completion of the sequencing of the entire *S. cerevisiae* genome (Goffeau *et al.* 1996). This was particularly advantageous because of the functional redundancy of some *S. cerevisiae* PEX genes and the lack of clear phenotypes of certain *pex* mutants.

Most of the *S. cerevisiae* PEX genes identified by the above approaches appear to be involved in matrix protein import—namely PEX1, -2, -4, -5, -6, -7, -8, -10, -12, -13, -14, -15, -17, -18, -21, -22 (Table 2)—whereas two genes, PEX3 and PEX19, are proposed to be essential for the formation of functional peroxisomal membranes. In the first group of mutants, remnant peroxisomal membrane structures (also termed “ghosts”) are still present, indicating that the formation of peroxisomal membranes is independent of the matrix protein import process. In *pex3* and *pex19* mutants, however, peroxisome membrane structures are not detectable. In both groups of mutants, most of the peroxisomal matrix proteins are mislocalized to the cytosol, where they are relatively stable. PMPs are also stable in mutants of the first group and localize to the peroxisomal ghosts. In *pex3* or *pex19* mutants, however, PMPs are mislocalized to the cytosol or the ER, or are rapidly degraded (Hettema *et al.* 2000; Otzen *et al.* 2004).

Most of the recently identified PEX genes (Table 2) have been identified by alternative approaches (see section below). In general, deletion of these genes results in alterations of peroxisome numbers and/or size and is not accompanied by mislocalization of peroxisomal protein or major defects in growth on oleate media.

As shown in Table 2, not all PEX genes that have been identified so far are also present in *S. cerevisiae*. PEX9 was solely identified in the yeast *Y. lipolytica*, but later studies revealed that it was incorrectly annotated and in fact encodes PEX26 (Kiel *et al.* 2006). PEX16 occurs only in higher eukaryotes and in filamentous fungi, but is absent from most yeast species, with the exception of *Y. lipolytica* (Kiel *et al.* 2006). PEX20 encodes a peroxin in filamentous fungi, which is the functional ortholog of *S. cerevisiae* Pex18 and the partially redundant protein Pex21, coreceptors in the PTS2 matrix protein import pathway (for details see below). Similarly, Pex26 occurs only in higher eukaryotes

and filamentous fungi, but can be regarded as the functional homolog of *S. cerevisiae* Pex15, which is a PMP that recruits the two ATPases and members of the AAA-protein family, Pex1 and Pex6, to the peroxisomal membrane.

Pex33 was identified in the filamentous fungus *Neurospora crassa* and shows homology to a short N-terminal domain of Pex14 (Managadze *et al.* 2010). A similar protein was identified in the fungus *Penicillium chrysogenum*, where it was designated Pex14/17 (Opalinski *et al.* 2010).

Y. lipolytica Pex23 has three homologs in *S. cerevisiae* that are designated Pex30, Pex31, and Pex32 in this organism; *Y. lipolytica* Pex24 is homologous to *S. cerevisiae* Pex28 and Pex29 (Table 2).

Identification of peroxisomal proteins by organelle proteomics: PEX11 is the first peroxin that has not been cloned by classic genetic approaches. Instead, it was identified by sequencing of a protein present in purified peroxisomal membranes followed by reversed genetics (Erdmann and Blobel 1995; Marshall *et al.* 1995). The same approach resulted in the identification of other novel PMPs, such as Psc60, Pex13, and Ant1. The first mass spectrometry analyses of purified peroxisomal membrane fractions were reported in 2001 (Schafer *et al.* 2001) and in 2002 (Yi *et al.* 2002) and resulted in the identification of several novel candidate peroxisomal proteins. An extensive analysis of the proteome of isolated *S. cerevisiae* peroxisomes was reported by Marelli *et al.* (2004). In this study, classic subcellular fractionation procedures were combined with immuno-isolation to obtain fractions enriched in peroxisomes from oleate-grown cells. This quantitative proteomics study resulted in a list of 71 candidate proteins that had a high likelihood of being peroxisomal because 28 of them were already annotated as being peroxisomal. Interestingly, the list contained several candidate proteins that were previously localized to other cell compartments, such as the LD protein Faa1 and six proteins linked to the secretory pathway, namely Dpm1, Ybr159w, Yor086c, Ygr266w, Rho1, and Cdc42. Biochemical and microscopic studies of a selection of eight of the candidate proteins confirmed that most of these proteins were, at least transiently, located to peroxisomes such as Rho1, Emp24, Faa1, and Erg6 (Marelli *et al.* 2004).

Rho1 is a small GTPase typically localizing to the plasma membranes and endomembranes in glucose-grown cells. However, Rho1 also localizes to the peroxisomal surface where it is recruited by the PMP Pex25 (Marelli *et al.* 2004); it is suggested that it regulates the assembly state of actin and thereby controls peroxisome membrane dynamics. Recent data in *H. polymorpha* revealed that in that yeast species a significant portion of Rho1 also colocalizes with peroxisomes (Saraya *et al.* 2011); Rho1 as well as Pex25 are essential for the generation of peroxisomes in *H. polymorpha* pex3 cells upon transformation and expression of PEX3 (for details see below).

The role of *S. cerevisiae* Emp24 in peroxisome biology has not yet been analyzed in greater detail. Emp24 is a member

of the family of p24 proteins, which are membrane proteins that are abundantly present in the membranes of the early secretory pathway (for a review see Strating *et al.* 2009). Studies in *H. polymorpha* showed that a (minor) portion of *H. polymorpha* Emp24 colocalizes to peroxisomes, where it is important for peroxisome fission by an as-yet-unknown mechanism (Kurbatova *et al.* 2009).

Erg6 and Faa1 are typically localized to the ER and LDs (see above; Table 1), but were also found to colocalize with peroxisomes, which may reflect the close physical and physiological interaction between these organelles. Alternatively, however, the peroxisomal fractions analyzed may have been contaminated with LDs, which are also induced in the presence of oleate (Binns *et al.* 2006; Pu *et al.* 2011). A large number of candidate peroxisomal proteins identified in the study by Marelli *et al.* (2004) have not yet been further analyzed, but are of high interest for future peroxisome research.

In silico prediction of peroxisomal proteins: Comparative genomics approaches have been used to identify putative peroxisomal matrix proteins based on consensus sequences of the peroxisomal targeting signals (PTSs) PTS1 and PTS2. The first report on *in silico* prediction of *S. cerevisiae* peroxisomal matrix proteins was by Kal *et al.* (2000). Emanuelsson *et al.* (2003) designed an improved predictor (PeroxiP, <http://www.bioinfo.se/PeroxiP/>) in which the sequence of the nine amino acids that immediately precede the PTS1 consensus sequence also was included as a criterion. Another predictor specific for PTS1 proteins has been generated by Neuberger *et al.* (2003a,b; mendel.imp.ac.at/mendeljsp/sat/pts1/PTS1predictor.jsp). Although very successful for metazoans and plants, these methods have not been very effective in identifying novel peroxisomal yeast proteins.

The limited success of using peroxisomal targeting sequences for the prediction of yeast peroxisomal proteins may be due to the fact that several peroxisomal matrix proteins exist that lack a consensus PTS, e.g., Pox1 and Cat2 (van der Klei and Veenhuis 2006a). On the other hand, proteins that do have a PTS based on computer predictions may not be peroxisomal because these signals are masked by protein conformation or altered/removed by post-translational processes. Moreover, it was recently established that in certain fungi some glycolytic enzymes obtain a PTS only upon ribosomal readthrough or by differential splicing, resulting in partial peroxisomal localization of these predominantly cytosolic enzymes (Freitag *et al.* 2012). In addition, *in silico* prediction is not yet possible for PMPs because no specific signature is currently known for these proteins.

Transcriptome analysis: Another approach that has been used to identify peroxisomal proteins is the analysis of transcripts that are induced upon stimulation of peroxisome proliferation. Serial Analysis of Gene Expression analysis (Kal *et al.* 1999) revealed that predominantly genes encoding peroxisomal enzymes of the β -oxidation pathway and

other proteins participating in oleate metabolism were upregulated upon shifting *S. cerevisiae* from glucose to oleate media. This study resulted in the first identification of the acyl-CoA thioesterase *Tes1*. The rather surprising outcome of this study was that, with the exception of *PEX11*, the messenger RNA levels of *PEX* genes remained unaltered after the shift from glucose to oleate medium (Kal *et al.* 1999).

Smith *et al.* (2002) were the first to perform microarray analysis to identify novel oleate-inducible peroxisomal proteins. In this study, candidate genes were identified by pattern matching of profiles of genes known to be involved in peroxisome biogenesis or function by using three complementary clustering algorithms. Thus, in addition to highly induced genes, genes that have lower levels of induction but similar induction patterns were identified. The screen resulted in a list of 225 candidate genes of which 2 were further analyzed to validate the approach. Indeed, both genes, *LPX1* and *PEX25* (Table 2), encoded novel peroxisomal proteins. Microarray studies using the methylotrophic yeast *H. polymorpha* also showed that predominantly the peroxisomal enzymes of methanol metabolism are strongly induced upon a shift of cells from glucose to methanol medium (van Zutphen *et al.* 2010). Again, *PEX* genes were not or were only slightly induced under these conditions; the highest increase in gene expression (four- to fivefold) was observed for *PEX11* and *PEX32* upon induction of peroxisome proliferation.

Microscopic analysis of peroxisomes: A direct way to identify peroxisomal proteins is by fluorescence microscopy. Today several genome-wide studies to localize the entire *S. cerevisiae* proteome have been reported. The first study was performed by Kumar *et al.* (2002) who used indirect, high-throughput immunofluorescence. These authors epitope-tagged 2085 *S. cerevisiae* ORFs and also randomly tagged genes by transposon mutagenesis. However, novel peroxisomal proteins were not identified by this approach.

In 2003, Huh *et al.* performed fluorescence microscopy of glucose-grown cells of an *S. cerevisiae* collection of strains that express C-terminal GFP fusion proteins. This study resulted in 21 proteins with a putative peroxisomal localization. This rather low number may be related to the fact that the proteins were C-terminally tagged, which masks the function of the C-terminal PTS1 sequences. Moreover, cells were grown on glucose in this study, and thus peroxisome proliferation was not strongly induced (Huh *et al.* 2003). This study, however, resulted in the detection of novel peroxisomal proteins and showed the association of *Gpd1* with peroxisomes (Table 1). Natter and colleagues (Kals *et al.* 2005; Natter *et al.* 2005) carried out a similar study with plasmid-encoded C-terminal GFP fusion proteins. This study by Natter *et al.* was limited to 493 proteins, which were selected on the basis of their potential participation in lipid metabolism and membrane assembly; the localization of 16 known yeast peroxisomal proteins was also confirmed by that study.

Wolinski *et al.* (2009b) used GFP fused to a PTS1 under the control of the *ADH1* promoter to analyze all 4740 viable yeast deletion strains by confocal imaging and automated quantitative analysis for peroxisome deficiency or morphological alterations. In addition to all previously known *pex* mutants defective in PTS1 protein import, *mdh2* and *afg1* deletion mutants showed a significant amount of cytosolic GFP-PTS1 as well. *MDH2* encodes a cytosolic malate dehydrogenase, whose function is important in peroxisomal β -oxidation (Figure 5). *AFG1* encodes a mitochondrial member of the AAA family of proteins. Analysis of the localization of additional peroxisomal membrane and matrix markers in *mdh2* and *afg1* cells indicated that both strains are defective in PTS1 and PTS2 matrix protein import, but not in the formation of peroxisomal membranes. How these proteins specifically function in matrix protein import is not yet known.

Saleem *et al.* (2008) analyzed a collection of 249 mutant strains lacking nonessential kinases, phosphatases and cyclins for alterations in peroxisome biology. In this study, GFP was fused to the C terminus of the PTS2 protein *Pot1*. The *POT1* promoter contains an oleate-responsive element, which is typical for oleate-induced peroxisomal β -oxidation enzymes (for a review see Gurvitz and Rottensteiner 2006). Cells were analyzed using a combination of confocal laser scanning microscopy to monitor peroxisome abundance and morphology and fluorescence activated cell sorting (FACS) to measure *Pot1*-GFP fluorescence levels. Different classes of regulatory proteins were found to regulate *POT1* expression and peroxisome number and size; interestingly, the nonessential cyclin-dependent kinase *Pho85* was shown to be involved in both. Prominent effectors in peroxisome biogenesis identified in that study include actin-regulating proteins, which function through the action of Rho regulators (e.g., *Rho1*), as well as proteins involved in phosphatidylinositol metabolism (e.g., *Vps34*). The latter group of proteins may mediate so far unknown membrane fusion processes in peroxisome biogenesis (Saleem *et al.* 2008). In a recent update to this study, Saleem *et al.* (2010) analyzed some 4000 *S. cerevisiae* deletion strains expressing chromosomally integrated *Pox1*-GFP. In this screen, most of the known *PEX* genes required for PTS2 protein import were identified. In addition, deletion of *CBS1* resulted in mislocalization of *Pot1*-GFP. This was a very unexpected finding because *Cbs1* is a mitochondrial translational activator. Interestingly, *Pot1*-GFP import into peroxisomes partially recovered when *cbs1* cells were maintained for several days on solid growth media. In addition, the screen revealed three novel proteins involved in peroxisome inheritance: *Vps52*, a regulator of actin; *Pir3*, involved in cell-wall organization; and *Ykr015c*, a protein of unknown function. Several additional mutants that displayed alterations in peroxisome size and/or number were identified (see the Yeast Peroxisome Cellular Imaging Resource at <http://pbeid.systemsbiology.net>). These include the known genes involved in the regulation of peroxisome size, *PEX11*, *VPS1*, and *DNM1*, as well as two novel genes, *MNN11*, encoding a subunit of a Golgi-resident mannosyltransferase, and *HSL7*,

encoding the bud-neck-localized protein arginine *N*-methyltransferase. The role of these proteins in regulating peroxisome proliferation still needs to be established.

Many of the proteins identified in the various imaging-based screens do not localize to peroxisomes or are rather known to be involved in other cellular processes. Deletion of genes encoding mitochondrial proteins leads to a reduction in peroxisome numbers, e.g., *Cox6*, *Cox9*, and *Cox10*, which are involved in mitochondrial respiration, further stressing the functional and metabolic links between peroxisomes and mitochondria. Indeed, defects in mitochondrial function have previously been indicated to stimulate peroxisome proliferation—the so-called retrograde response. Also, deletion of two genes encoding vacuolar proteins results in reduced numbers of peroxisomes, i.e., the vacuolar t-SNARE *Vam3* or the subunit of the vacuolar ATPase *Vph1*. Similarly, deletion of the *VAM3* and *VAM7* genes encoding two vacuolar t-SNARES in *H. polymorpha* also results in unusual peroxisomal structures with multiple membrane-enclosed compartments (Stevens *et al.* 2005).

Surprisingly, given the important proposed role for the ER in peroxisome biogenesis (see below), no ER proteins were identified in the screen of Saleem *et al.* (2010) that were important for peroxisome biogenesis, inheritance, or regulation of organelle number. Only deletion of *FEN1*, a gene required for fatty acid elongation (Henry *et al.* 2012), showed an effect on peroxisome size and resulted in enlarged organelles (Saleem *et al.* 2010).

Peroxisome Biogenesis

Peroxisomal matrix protein import

Targeting signals and their receptors: The peroxisomal matrix protein import machinery has unique properties and differs fundamentally from import systems of other eukaryotic organelles in that it can translocate *folded* proteins and even protein complexes across the membrane. Peroxisomal matrix proteins are nuclear encoded, synthesized in the cytosol, and subsequently post-translationally imported into the organelle. These proteins harbor PTSs that are recognized by the soluble receptor proteins *Pex5p* (PTS1) and *Pex7p* (PTS2), respectively. It is generally assumed that PTS1 signals are not cleaved upon import, whereas a few PTS2 sequences were indeed shown to be processed upon import in higher eukaryotes.

PTS1 and its receptor *Pex5*: The first PTS1 that has been described is the C-terminal tripeptide SKL in firefly luciferase (Gould *et al.* 1987). This PTS1 tripeptide, as well as the PTS1 peroxisomal import machinery, is highly conserved in yeast, plants, insects, and mammals (Gould *et al.* 1990). Detailed sequence analysis and mutagenesis studies revealed that the PTS1 tripeptide invariably consists of a small neutral amino acid at the first position, followed by an amino acid residue capable of hydrogen bonding at the penultimate position and a hydrophobic residue at the extreme C terminus. Also, the

tripeptide must be present at the extreme C terminus of the protein. The most commonly used consensus of the PTS1 tripeptide is [(S/A/C)(K/R/H)(L/M)]. Studies by Lametschwandtner *et al.* (1998) have suggested a broader degeneracy in yeast and human PTS1 sequences. Also, not all PTS1 peptides that bind human *Pex5* can bind yeast *Pex5* as well, demonstrating some species specificity.

Detailed studies on the interaction of the PTS1-binding domain of *Pex5* with various proteins/peptides indicate that residues upstream of the PTS1 tripeptide also are important for cargo recognition. Therefore, the PTS1 is now defined as a C-terminal 12-amino-acid sequence, which consists of the C-terminal tripeptide that interacts with the PTS1-binding site in *Pex5*, a tetrapeptide immediately upstream of this tripeptide, which may interact with the surface of *Pex5*, and a flexible hinge region of five residues (reviewed by Brocard and Hartig 2006).

The PTS1 receptor *Pex5* consists of a relatively poorly conserved N-terminal domain and a C-terminal domain, which contains six tetratricopeptide repeats (TPR). A typical feature of the N terminus of *Pex5* is the presence of several WxxxF/Y motifs. In human *Pex5*, these motifs are required for binding of the receptor to *Pex13* and *Pex14*, but in yeast the WxxxF/Y motif is important only for *Pex13* binding (Williams *et al.* 2005). The N terminus of *Pex5* also is the region where the protein is modified by mono- or multi-ubiquitination, which is important for receptor recycling (see below).

The *Pex5* TPR domain is the actual PTS1-binding site (Brocard *et al.* 1994). According to a crystal structure of the TPR domain of human *Pex5*, the PTS1 is locked in a groove allowing various interactions with two sets of three TPR motifs linked together by a helical hinge (Gatto *et al.* 2000). Structural studies by Stanley *et al.* (2006) using the human *Pex5* TPR domain and a full-length peroxisomal matrix protein, the sterol carrier protein 2 (SCP2), indicated that the cargo is bound to the receptor by two separate binding sites: a C-terminal PTS1 motif and a topologically separated secondary site. These studies also revealed major conformational changes of the receptor that occur upon cargo loading. Unfortunately, so far no structural data on a full-length *Pex5* protein have been reported. For further details on the structural properties of *Pex5*, the reader is referred to the excellent review by Stanley *et al.* (2007).

As discussed above for the SCP2 protein, the PTS1-binding site in the C-terminal TPR domain of the *Pex5* receptor is not the only site of interaction with cargo. Some matrix proteins without (Klein *et al.* 2002) or with a redundant PTS1 (Gunkel *et al.* 2004) also bind to the N terminus of *Pex5*, such as *Pox1* and *Cat2* in *S. cerevisiae* (Schafer *et al.* 2004) and alcohol oxidase in *H. polymorpha* (Gunkel *et al.* 2004). Mutational analysis of the *S. cerevisiae* *Pex5* N terminus indicates that the domains that are required for *Cat2* and *Pox1* binding are overlapping but not identical. Hence, most likely multiple residues in the N terminus of *Pex5* are involved in the recognition of non-PTS1 proteins.

The occurrence of different matrix protein-binding sites in Pex5 variants was nicely illustrated by studies of Ozimek *et al.* (2006), who showed that *S. cerevisiae* Pex5 (via its TPR domain) and *H. polymorpha* Pex5 (via its N-terminal domain) recognize different, independent binding sites in the same peroxisomal matrix protein. Also, *S. cerevisiae* peroxisomal catalase A has a redundant PTS1, which is the very unusual hexapeptide SSNSKF. Catalase A has a second internal PTS, but it is unknown to which region (N or C terminus) of Pex5 this sequence actually binds (Kragler *et al.* 1993).

PTS2 and Pex7 with its coreceptors: PTS2 was first identified in rat liver thiolase, where it is present as an N-terminal presequence that is processed upon import. Only a few proteins have a PTS2 for which the consensus sequence is (R/K)(L/V/I)X₅(Q/H)(L/A/I). *S. cerevisiae* expresses only three peroxisomal proteins with a firmly established PTS2, namely Pot1, Pcd1, and Gpd1.

The PTS2 is recognized by the soluble receptor Pex7, which is characterized by the presence of six WD repeats. On the basis of its sequence, the receptor is predicted to fold as a seven-bladed β -propeller domain, in which each “blade” comprises a so-called WD repeat (Li and Roberts 2001; Stanley *et al.* 2007); however, no Pex7 crystal structures have been resolved yet.

Pex7 requires additional proteins that function as coreceptors. In *S. cerevisiae*, these are the partially redundant, homologous proteins Pex18 and Pex21 (for a review see Schliebs and Kunau 2006). In other yeast species and filamentous fungi, this function is fulfilled by Pex20 (Einwachter *et al.* 2001). The coreceptors share a characteristic N-terminal region that contains WxxxF/Y sequence motifs, such as the N terminus of Pex5 and a Pex7-binding region. Also, like the Pex5 N terminus, these proteins are ubiquitinated, a process that is required for recycling of the proteins (Leon and Subramani 2007). Additional functions have also been attributed to the coreceptor proteins, *e.g.*, in oligomerization of the cargo protein (Titorenko *et al.* 1998) or in PTS2 binding (Otzen *et al.* 2005).

Piggy-back import: Some peroxisomal proteins without a recognizable PTS still can be imported into peroxisomes in complex with a PTS-containing protein. Recently, the first example of piggy-back import was reported for the mammalian peroxisomal protein, superoxide dismutase (Islinger *et al.* 2009). The examples that have been reported in yeast (Glover *et al.* 1994; McNew and Goodman 1994) all comprise artificial co-expression of oligomeric proteins with and without PTS sequences, and it is currently unclear whether piggy-back import indeed occurs for endogenous yeast proteins *in vivo*.

Receptor docking site: The receptor-cargo complex docks at the outer surface of the peroxisomal membrane to a proteinaceous receptor docking site that consists of the three peroxisomal membrane proteins Pex13, Pex14, and Pex17. Pex17, which interacts with Pex14, occurs only in *S. cerevi-*

siae, and its function is still enigmatic. Pex13 and Pex14 are highly conserved, and they interact with each other as well as with both Pex5 and Pex7 import receptors (Figure 6A). Pex13 is an integral membrane protein that contains two transmembrane regions and a Src homology 3 (SH3) domain at the extreme C terminus, which is exposed to the cytosol. The extreme N terminus of Pex13 binds Pex7 (Stein *et al.* 2002), whereas the SH3 domain has distinct binding sites for Pex5 and Pex14 (Pires *et al.* 2003). In Pex14, a PxxP domain, as well as other conserved residues, is involved in the interaction with the Pex13 SH3 domain (Girzalsky *et al.* 1999; Bottger *et al.* 2000). Pex14 also binds to a region in between the two transmembrane domains of Pex13 (Schell-Steven *et al.* 2005). In addition, indirect interactions may occur between both peroxins, which could explain why disruption of individual interaction sites often does not significantly affect protein import *in vivo* (Williams and Distel 2006).

Although Pex13 can directly interact with Pex5, Pex14 is most likely the initial docking protein in the peroxisomal membrane in the course of PTS1 protein import (Urquhart *et al.* 2000); Pex13 may have a similar function in docking of Pex7. Recently, Grunau *et al.* (2009) showed that the PTS2 protein Pot1 first binds Pex7 in a process that is independent of the coreceptors Pex18 and Pex21. Next, a cytosolic Pot1-Pex7-Pex18 complex that docks to the peroxisomal membrane via the interaction between Pex7 and Pex13 is formed. It was proposed that Pex7 and the cargo dissociate during or after assembly of a large complex with Pex14 and Pex13 (for details see Figure 6B).

The first PEX14 gene was cloned in *H. polymorpha* by functional complementation of a mutant defective in growth on methanol (Komori *et al.* 1997). The *S. cerevisiae* homolog was cloned by the Kunau group, based on the sequence homology with *H. polymorpha* PEX14, and was shown to represent the Pex5 docking site (Albertini *et al.* 1997). At the same time, Brocard *et al.* (1997) isolated the yeast homolog by a two-hybrid screen using Pex5 as a bait. Pex14 is characterized by a coiled-coil region in the middle part of the protein that is involved in homo-dimerization; however, its topology in the membrane is still debated (Azevedo and Schliebs 2006).

The conserved extreme N-terminal region adopts an α -helical structure (de Vries *et al.* 2007). The N terminus of *S. cerevisiae* Pex14 has been implicated in Pex5 binding, whereas its C terminus contains overlapping Pex5- and Pex7-binding sites (Niederhoff *et al.* 2005; Williams *et al.* 2005). In *H. polymorpha*, however, no receptor binding sites have been identified in the Pex14 C terminus (Bellu *et al.* 2001); instead, in *H. polymorpha* the Pex14 N terminus most likely binds both Pex5 and Pex7, as it is involved in both PTS1 and PTS2 protein import (de Vries *et al.* 2007). Structural studies using purified proteins showed that the N terminus of human Pex14 binds Pex19 in addition to Pex5 (Neufeld *et al.* 2009). Interestingly, an F/YFxxxF motif in the N terminus of Pex19 associates with the same site in




Figure 6 Peroxisomal protein import. (A) Hypothetical model of PTS1 protein import. First, cytosolic Pex5 binds a newly synthesized PTS1-containing cargo protein (“cargo”). The PTS1 binds to the C-terminal TPR domain of Pex5. Next, Pex5 docks to the receptor-docking complex at the peroxisomal membrane, which is composed of Pex13, Pex14, and Pex17. Docking involves the N-terminal domain of Pex5, indicated as a spiral. Subsequently, the Pex5-cargo complex is imported into the organellar matrix. Pex5 most likely forms a transient pore in the peroxisomal membrane. The Pex5-cargo complex then dissociates in a process that involves Pex8, a peripheral membrane protein in the peroxisomal matrix. Finally, Pex5 is recycled back to the cytosol, a process that enables it to bind the next PTS1 cargo protein. Recycling involves mono-ubiquitination of Pex5 by the UBC protein Pex4, which is recruited to the peroxisomal membrane by Pex22. The three RING finger proteins Pex2, Pex10, and Pex12 are proposed to serve as E3 ligases. The ubiquitinated Pex5 is pulled out of the membrane by the AAA proteins Pex1 and Pex6, which are associated with the peroxisomal membrane via Pex15. When Pex5 recycling fails, it becomes polyubiquitinated by Ubc4/5 and degraded by the proteasome. (B) Hypothetical model of PTS2 protein import. Dimeric Pot1 is shown as an example of a typical PTS2 protein. Dimeric Pot1 first binds to the PTS2 receptor Pex7. Subsequently, the coreceptor Pex18 (and possibly also Pex21) binds to the receptor/cargo complex. Pex7 associates with Pex13 of the docking complex. After import of the Pot1 cargo into peroxisomes by an as-yet-unknown mechanism Pex7 recycles back to the cytosol. Pex18, however, first forms a complex with Pex14. Whether and how Pex18 recycles for another round of import is unknown.

Pex14 as the Pex5 WxxxY motif. Although the key role of Pex19 is in sorting of PMPs (see below), in this case it possibly plays a role in the assembly of the docking complex (Fransen *et al.* 2004).

Although it is generally accepted that Pex14 is the initial Pex5 receptor in the peroxisomal membrane, the molecular mechanisms involved in its function are still unknown. Moreover, *H. polymorpha* Pex14 has additional functions, for example, in peroxisome degradation by autophagy (Bellu *et al.* 2001; Zutphen *et al.* 2008). Interestingly, data in *H. polymorpha* also indicate that PTS1 protein import can proceed in a *PEX14* deletion strain upon overexpression of Pex5 (Salomons

et al. 2000). This suggests that Pex14 is not essential but rather important for the efficacy of matrix import and that Pex5 also takes part in the protein translocation process.

RING finger complex: The “really interesting new gene” (RING) complex involved in peroxisome biogenesis and function consists of three proteins—Pex2, Pex10, and Pex12—that interact with each other. They contain RING domains at their cytosolic C termini, which bind zinc ions through characteristic conserved cysteine and histidine residues. As these proteins are integral membrane proteins, it was proposed that they form the actual translocation pore

through the membrane. Recent findings, however, indicate that these proteins have E3 ubiquitin ligase activity and function together with the ubiquitin-conjugating E2 enzymes *Pex4* or *Ubc4*. Indeed, *Pex4* plays an important role in matrix protein import (Wiebel and Kunau 1992), whereas *Ubc4* is not required for this process. Instead, *Ubc4* is responsible for a quality control mechanism that results in degradation of nonfunctional *Pex5* molecules by polyubiquitination and their subsequent degradation by the proteasome (Figure 6A) (Kiel *et al.* 2005b).

Williams *et al.* (2008) were the first to show that the RING domain of *Pex10* exhibits E3 ligase activity and acts as E3 ligase for *Ubc4*-dependent polyubiquitination of *Pex5*. Platta *et al.* (2009) subsequently demonstrated that all three RING finger peroxins exhibit ubiquitin-ligase activity. These authors, however, reported that *Pex2*, but not *Pex10*, is essential for *Ubc4*-dependent polyubiquitination, whereas *Pex12* catalyzes *Pex4*-dependent monoubiquitination of *Pex5*. Monoubiquitination is important for recycling of functional *Pex5* molecules to the cytosol to mediate another round of PTS1 protein import (Figure 6A). *pex10* mutants show a severe defect in matrix protein import, which could be due to the stabilizing function of *Pex10* on the other RING finger proteins. However, as *Pex10* also has E3 ligase activity, the important current question concerns what the substrate of *Pex10* actually is.

Translocation pore: The general view on the peroxisomal translocon is that it is composed of two subcomplexes, the receptor docking complex and the RING finger complex, which are organized into the so-called importomer by *Pex8* (Agne *et al.* 2003). Studies in *H. polymorpha* indicated that *Pex8* is also required for cargo release from *Pex5* (Wang *et al.* 2003), indicating a dual role for *Pex8*. A recent proteomics study confirmed the presence of proteins of the docking complex and the RING finger complex, together with *Pex8*, in the core complex of the peroxisomal translocon. In addition, *Pex11* and *Dyn2*, the microtubule motor protein, also were observed to be closely associated with the peroxisomal core translocon (Oeljeklaus *et al.* 2012).

In *Pichia pastoris* Hazra *et al.* (2002) proposed that *Pex3*, rather than *Pex8*, is the protein that sequesters both subcomplexes into the translocon. However, this protein was identified as a component of neither the 9 core nor the 12 transient components of the importomer in *S. cerevisiae* (Oeljeklaus *et al.* 2012). Rather, *Pex3* is a peroxin that is implicated in peroxisomal membrane biogenesis (see below).

The actual mode of matrix protein import is still elusive. Recently, the group of Erdmann (Meinecke *et al.* 2010) provided evidence for the concept of a transient pore, composed of *Pex5* and *Pex14* oligomers. This transiently gated ion-conducting channel may form pores of variable sizes up to 9 nm, which are induced by the interaction with the receptor cargo complex. This model also explains that the peroxisomal translocon pore may accommodate 9-nm gold particles (Walton *et al.* 1995). The occurrence of *Pex5*

oligomers would be in line with electron microscopy data obtained from isolated *H. polymorpha* *Pex5* (Moscicka *et al.* 2007). On the other hand, by using small-angle X-ray scattering, Shiozawa *et al.* (2009) suggested that human *Pex5* is rather monomeric in solution. Nevertheless, the option of a transient pore consisting of *Pex14* and *Pex5* oligomers to mediate matrix protein import is quite attractive. As it stands now, however, this model still leaves many questions open, *i.e.*, how the import process is regulated and how the translocon is able to accommodate large receptor-bound oligomeric matrix proteins.

Receptor recycling and the role of ubiquitin: The final step in PTS1 matrix protein import is the recycling of *Pex5* to the cytosol. The first indication that *Pex5* is indeed a cycling receptor came from the observation that in wild-type *H. polymorpha* *Pex5* shows a dual localization in the cytosol and the peroxisomal matrix (van der Klei *et al.* 1995). Subsequent studies in the same organism indicated that the absence of *Pex4* resulted in a specific block in PTS1-protein import that could be suppressed by overexpression of *PEX5*. Whereas in wild-type cells overproduced *Pex5* was localized mainly to the cytosol, it accumulated in *pex4* cells at the inner surface of the peroxisomal membrane. This observation led to the proposal that *Pex4*, and hence ubiquitination, is involved in receptor recycling (van der Klei *et al.* 1998). Erdmann's group has significantly contributed to unraveling the principles of the *Pex5* recycling process (for a review see Schliebs *et al.* 2010). In the normal receptor recycling process, *Pex5* is mono-ubiquitinated at a conserved cysteine by *Pex4* and *Pex12*. Poly-ubiquitination of *Pex5* occurs on a lysine residue (Kiel *et al.* 2005a). Release of the receptor from the membrane is an ATP-dependent process that is facilitated by the two AAA ATPases, *Pex1* and *Pex6*, which are recruited to the membrane by *Pex15*. Debelyy *et al.* (2011) recently identified a novel component, *Ubp15*, which is capable of removing ubiquitin from *Pex5*. Although the model of a cycling *Pex5* receptor is today generally accepted, it is still debated whether in yeast *Pex5* associates to the outer surface of the membrane, inserts into the membrane, or even transiently enters the peroxisomal membrane, as suggested by the data obtained in *H. polymorpha*. The current knowledge of protein import and receptor recycling is summarized in Figure 6A.

On this point only the PTS1 peroxisomal protein import pathway has been studied in greater detail, but evidence suggests that the PTS2 receptor *Pex7* may also cycle and enter the matrix, based on the observed localization of this protein in the peroxisomal matrix (for a review see Lazarow 2006) (Figure 6B).

Sorting and insertion of PMPs

Like matrix proteins, PMPs are synthesized in the cytosol. However, whether they immediately insert into the peroxisomal membrane or traffic first to the ER is currently still strongly debated. PMP-targeting signals (mPTSs) may

involve different regions of the PMPs and are not characterized by a simple consensus sequence. Two classes of PMPs have been defined: class I PMPs contain an mPTS that is recognized by the soluble farnesylated protein Pex19. Class II PMPs, such as Pex3, Pex16, and Pex22, contain membrane-targeting information that is not recognized by Pex19 (Jones *et al.* 2004). The prevailing model for class I PMPs sorting suggests that cytosolic Pex19 serves as a cycling mPTS receptor, which docks to Pex3 on the peroxisomal surface, after which the cargo PMP is inserted into the membrane via a still-unknown pathway (reviewed in Schliebs and Kunau 2004). In line with this model is the observation that Pex19 binds mPTSs that are required for targeting (Jones *et al.* 2004). Recently, the crystal structure of the C-terminal part of human Pex19 was solved, which revealed that mPTSs bind to a globular α -helical domain in the C terminus of the protein (Schueller *et al.* 2010). However, there are conflicting data on the exact role of Pex19 in the biogenesis of the peroxisomal membrane. For example, in *H. polymorpha* and *S. cerevisiae*, *pex19* mutant cells that overproduce Pex3 peroxisomal membrane structures that contain PMPs exist, suggesting that Pex19 is not essential for the insertion of PMPs into peroxisomal membranes (Otzen *et al.* 2004). Similarly, in *Y. lipolytica* cells lacking Pex19 structures are present that resemble normal peroxisomes even without Pex3 overproduction (Lambkin and Rachubinski 2001). On the basis of these and other observations it has been suggested that Pex19 is required mainly to assemble PMP complexes at the peroxisomal membrane and, hence, may function as a chaperone instead of being an mPTS receptor protein.

Data by van der Zand *et al.* (2010) suggest, on the basis of live-cell imaging experiments, that in *S. cerevisiae* all PMPs travel via the ER to peroxisomes. The authors propose novel functions for Pex3 and Pex19 in the budding of peroxisomal vesicles from the ER. According to these studies, insertion of PMPs into the ER depends on the Sec61 translocon, except for the tail-anchored protein Pex15, which requires Get3. Notably, in previous studies by South *et al.* (2001) no effect of the inactivation of Sec61 on peroxisome formation could be demonstrated.

The process of budding of PMP-containing vesicles from the ER proposed by van der Zand *et al.* (2010) is supported by data generated by Lam *et al.* (2010), who used a cell-free *in vitro* vesicle-budding assay. Using this assay, *S. cerevisiae* Pex15 and Pex3 were shown to be packaged into small vesicles that derived from the ER in a process that was dependent on Pex19. These vesicles were formed by a novel budding mechanism and independently of the COPII machinery that is responsible for secretory cargo packaging at the ER. Possibly, these vesicles fuse homotypically to create a new peroxisome, the so-called *de novo* peroxisome formation process (see below), or they fuse with preexisting peroxisomes as a mode to transfer PMPs and lipids from the ER to these organelles. In a similar study using permeabilized *P. pastoris* cells, Pex3 and Pex11 were shown to be copackaged

into ER-derived vesicles in a Pex19-dependent process (Agrawal *et al.* 2011). Interestingly, this study revealed that the vesicles were also formed in the absence of Pex3, suggesting that this peroxin is not essential for vesicle formation. This finding also implies that Pex19 can dock to the ER membrane without its presumed membrane docking protein Pex3. Notably, this observation is in line with earlier data, which indicate that, in *P. pastoris pex3* mutant cells, small structures that contain PMPs are present (Hazra *et al.* 2002).

Although the above studies indicate that some PMPs initially may insert into the ER membrane, data have been reported that indicate that targeting of several PMPs (including Pex3) to the ER is rather inefficient, especially when endogenous promoters are used to drive gene expression of the reporter constructs. For example, a fusion protein of full-length Pex3 and GFP, expressed under control of the *PEX3* promoter in *H. polymorpha pex19* cells, localizes to the cytosol (Otzen *et al.* 2004). Similarly, tagged Pex11 produced under control of the endogenous promoter is localized to the cytosol in *S. cerevisiae pex3* and *pex19* mutant strains (Hettema *et al.* 2000). Also, in *H. polymorpha pex11 pex25* double-mutant cells, which are fully devoid of peroxisomal membrane structures, Pex3-GFP produced under control of the endogenous *PEX3* promoter is cytosolic (Saraya *et al.* 2011).

Although in most studies PMPs were shown to colocalize with ER markers even when expressed at wild-type levels, the kinetics of their production (e.g., pulsed induction of GFP-tagged proteins using strong inducible promoters) may be quite different from the wild-type situation. Hence, it cannot be excluded that deregulated expression contributes to enhanced targeting of these proteins to the ER. The apparently contradictory data on PMP sorting may also depend on the model and reporter proteins used. This may be concluded from data of the Fujiki laboratory (Matsuzaki and Fujiki 2008), who demonstrated that in mammals Pex3 is directly sorted to peroxisomes. For this sorting, Pex19 serves as a chaperone for full-length Pex3 to form a soluble complex in the cytosol, which docks to Pex16 at the peroxisomal membrane.

A number of PMPs in various cell types show a dual localization to the ER and peroxisomes under steady-state conditions, for example, plant Pex16 (Karnik and Trelease 2005), *P. pastoris* Pex30 and Pex31 (Yan *et al.* 2008), and *S. cerevisiae* Pex11. In the case of *S. cerevisiae* Pex11, the localization varies with the phosphorylation state of the protein (Knoblach and Rachubinski 2010). Whether these ER-localized proteins are on their way to peroxisomes is, however, not firmly established. The key questions to date are whether in wild-type cells all or a subset of the PMPs travel via the ER to peroxisomes under normal physiological conditions and whether the vesicles formed in the *in vitro* assays represent transport vesicles or preperoxisomes of a *de novo* synthesis pathway. The current availability of *in vitro* assays will significantly help in further understanding these important questions in PMP sorting.

Formation of peroxisomes from the ER

Several observations based on electron microscopy have suggested that new peroxisomes can be formed from the ER. Membrane connections appear to exist between peroxisomes and the ER in mouse dendritic cells, allowing new peroxisomes to pinch off from specialized regions of the ER (Geuze *et al.* 2003). Also, evidence obtained in yeast *pex3* and *pex19* mutants that lack peroxisomal membrane structures suggests that peroxisomes can be formed from the ER upon reintroduction of the corresponding *PEX3* or *PEX19* wild-type genes (Hoepfner *et al.* 2005; Kragt *et al.* 2005; Tam *et al.* 2005; Haan *et al.* 2006). *In vivo* pulse-chase experiments using *Pex3*-GFP in *S. cerevisiae pex3* cells (Hoepfner *et al.* 2005) revealed that *Pex3*-GFP is first targeted to the ER, but later present in new, small peroxisomes. Very similar observations have been made in other laboratories using *S. cerevisiae* and in *H. polymorpha* (for a review see Tabak *et al.* 2008).

An alternative explanation for the ER-localized *Pex3* in these experiments, however, may be that peroxisomes are formed by an alternative, relatively slow process, thus promoting excess *Pex3* to sort to the ER, when insufficient peroxisomal membrane is present. Likely, *Pex3* and other PMPs may contain ER-targeting signals, but these are not yet resolved. Therefore, during *de novo* synthesis, these proteins, including *Pex3*, may initially sort to the ER but later on be targeted directly from the cytosol to the newly formed organelle without a requirement for translocation via the ER. In line with this suggestion is the general observation that these PMPs are never observed at the ER in wild-type cells grown under normal conditions. However, this could also be explained by an extremely short residence time of these proteins at the ER.

A role for the ER in peroxisome biogenesis is also indicated by observations of Perry *et al.* (2009), who showed that *Pex3*-GFP localizes to tubular-vesicular structures in cells suppressed for *Sec20*, *Sec39*, and *Dsl1*, which form a complex in the ER. Also, deletion of *ARF1* and *ARF3* in *S. cerevisiae* affects peroxisome proliferation (Anthonio *et al.* 2009). However, details on the exact function of these proteins in peroxisome biogenesis are still lacking.

Several attempts have been made to identify genes that are essential for the *de novo* peroxisome formation process. In *H. polymorpha*, several gene deletions have been tested for their effect on the reintroduction of peroxisomes in *pex3* mutant cells upon induction of *Pex3*-GFP synthesis. These studies revealed that *Dnm1*, *Vps1*, *Emp24*, *Pex11*, and *Pex11C* are not required (Nagotu *et al.* 2008b; Kurbatova *et al.* 2009; Saraya *et al.* 2011). However, Saraya *et al.* (2011) showed that *Pex25* as well as the GTPase *Rho1* are necessary to allow the formation of new peroxisomes in *pex3* cells. Huber *et al.* (2012) showed that in *S. cerevisiae* *Pex25* also is essential for reintroduction of peroxisomes in *Pex3*-deficient cells. Since *H. polymorpha pex25* and *S. cerevisiae pex25* cells contain normal peroxisomes, the above process

apparently is not essential in wild-type cells to maintain peroxisomes.

Recent findings by van der Zand *et al.* (2012) suggest that in *S. cerevisiae* the process of peroxisome formation from the ER involves at least two biochemically distinct types of preperoxisomal vesicles that are initially formed from the ER, which each carry half a peroxisomal translocon complex and hence are unable to import peroxisomal matrix proteins. Upon their fusion, mediated by the AAA proteins *Pex1* and *Pex6*, a functional translocon is formed allowing uptake of peroxisomal matrix proteins from the cytosol. This model suggests a novel role for *Pex1* and *Pex6* in vesicle fusion, in addition to their established function in *Pex5* recycling. Notably, earlier data obtained in the yeast *Y. lipolytica* also suggested the formation of mature peroxisomes upon fusion of different subtypes of small preperoxisomal vesicles (Titorenko *et al.* 2000). Fusion of various types of vesicles was mediated by *Pex1* and *Pex6*; however, the origin of the vesicles remains unknown. All vesicles were matrix protein import competent, although the vesicle subtypes contained different combinations of peroxisomal proteins.

In summary, peroxisome formation can occur *de novo* in cells lacking pre-existing peroxisomes. However, the significance of *de novo* peroxisome formation in cells that already have peroxisomes is still debated.

Peroxisome fission

Pex11 and the Fis1/Dnm1 fission machinery: For decades, peroxisomes have been considered to be autonomous organelles that multiply by growth and division (Lazarow and Fujiki 1985). Various proteins are known to function in peroxisome fission. The first step in fission is organelle elongation, a process that is mediated by *Pex11* (Opalinski *et al.* 2011). In all studies performed in various species so far artificial modulation of *Pex11* levels has resulted in variations in peroxisome size and abundance; deletion of the *PEX11* gene invariably leads to fewer enlarged organelles, whereas *PEX11* overexpression leads to an increased number of smaller and often tubulated peroxisomes. Studies in *H. polymorpha* (Opalinski *et al.* 2011) revealed that the insertion of the N-terminal amphipathic α -helix of *Pex11* into the membrane causes the initial membrane curvature, which initiates organelle elongation. The activation of *Pex11* in organelle fission may be regulated by phosphorylation/dephosphorylation (Saleem *et al.* 2008; Knoblach and Rachubinski 2010). Mutant studies indicate that strains producing constitutively dephosphorylated *Pex11* show a phenotype similar to *pex11* cells, whereas strains expressing a phosphomimetic *Pex11* mutant allele show enhanced peroxisome proliferation, similar to cells overexpressing *PEX11*.

Recent observations indicate that in the yeasts *S. cerevisiae* and *H. polymorpha* most of the organelles are formed by fission of existing peroxisomes by the activity of the dynamin-related proteins (DRP) *Dnm1* and *Vps1* (Hoepfner *et al.* 2001; Motley and Hettema 2007; Nagotu *et al.* 2008b). This is, among other observations, suggested by the finding

that, in cells of a *dnm1 vps1* double-deletion strain, peroxisome fission is completely blocked, resulting in the presence of a single enlarged peroxisome per cell, even after prolonged cultivation under peroxisome-inducing conditions. In these cells, peroxisome formation from the ER is not affected, but generation of additional organelles was never observed (Motley and Hettema 2007; Nagotu *et al.* 2008b).

DRPs are large GTPases that are involved in multiple membrane fission and fusion events. *Vps1* was initially found to be involved in vacuolar protein sorting, whereas *Dnm1* was first identified as a protein required for mitochondrial fission. The peroxisomal fission machinery is therefore not unique for this organelle, but shares components with other membrane fission/fusion processes. *Dnm1* is essential for peroxisome fission under conditions of peroxisome induction by oleate, whereas *Vps1* functions under glucose-repressing conditions (Hoepfner *et al.* 2001; Kuravi *et al.* 2006). In *S. cerevisiae*, *Dnm1* is recruited to peroxisomes by *Mdv1* or its paralog *Caf4*, which are both associated with the peroxisomal membrane via the tail-anchored protein *Fis1* (Motley and Hettema 2007; Motley *et al.* 2008). *Mdv1* and *Caf4* are WD repeat proteins, which are absent in higher eukaryotes. How *Vps1* is recruited to peroxisomes is not yet known. Notably, in *H. polymorpha*, *Vps1* does not play a role in peroxisome fission, and, thus, this yeast species seems to be more similar to higher eukaryotes in this respect, in which a single DRP is involved in mitochondrial and peroxisome fission, as well as in chloroplast fission, in plants.

It is an intriguing question how the *Fis1*/Drp fission machinery is properly distributed over the individual organelles. In higher eukaryotes, *Pex11* has been implicated in the recruitment of *Fis1* to peroxisomes (Kobayashi *et al.* 2007; Lingard *et al.* 2008); in mammals, a role of *Pex19* in *Fis1* targeting was established (Delille and Schrader 2008), but these processes have not yet been confirmed in yeast. Fluorescence microscopy studies in *H. polymorpha* revealed that GFP-tagged *Dnm1* is not evenly distributed in the cell, but rather present in multiple spots to which *Mdv1* colocalizes. These spots dynamically associate and disassociate from mitochondria and peroxisomes, demonstrating that the same protein molecules may be involved in the fission of either one of these organelles (Nagotu *et al.* 2008a).

Peroxisome fission in *H. polymorpha* is fully blocked in *dnm1* cells. Growing *H. polymorpha dnm1* mutant cells contain a single enlarged peroxisome, which forms a long extension that protrudes into the developing bud. These extensions are not observed in *dnm1 pex11* cells, which is in agreement with the notion that *Pex11* plays a role in peroxisome elongation (Nagotu *et al.* 2008b). Remarkably, *Pex11* protein concentrates at the base of these peroxisome extensions, indicating that during fission of the organelles *Pex11* segregates into *Pex11*-enriched patches at the membrane. Recent findings show that other peroxins including *Pex10*, *Pex14*, but also the *Pex11*-family protein *Pex25*, do

not segregate to the same patches; instead, these proteins move to the developing new organelles or extensions in *dnm1* cells (Cepinska *et al.* 2011).

Role of the other *Pex11* protein family members: Most organisms contain at least three *Pex11* protein family members (see Table 2). The human genome encodes three family members, namely *Pex11* α , *Pex11* β , and *Pex11* γ , all of which have high similarity to *S. cerevisiae Pex11*. *S. cerevisiae* expresses, in addition to *Pex11*, the weakly homologous *Pex25* protein and its partially redundant paralogue, *Pex27* (Rottensteiner *et al.* 2003), which have also been implicated in peroxisome proliferation. Quantitative analysis of electron microscopy images revealed that peroxisomes are enlarged in *pex11*, *pex25*, or *pex27* mutant cells (Tam *et al.* 2003). The molecular function of *Pex27* is still unclear, whereas *Pex25* plays a role in recruiting *Rho1* to the peroxisomal membrane (Marelli *et al.* 2004). In *H. polymorpha*, *Pex25*, but not *Pex11* or *Pex11c*, has been shown to be important for the formation of peroxisomes from the ER (Saraya *et al.* 2011). Large-scale protein interaction studies by two-hybrid analysis (Yu *et al.* 2008), together with information from a global analysis of protein localization (Huh *et al.* 2003), led to the identification of a novel peroxisomal interaction partner of *Pex11*, *Pex25*, and *Pex27*, named *Pex34*; *Pex34* was proposed to act as a positive effector of peroxisome division (Tower *et al.* 2011).

***Pex23* protein family:** Based on sequence homology, Kiel *et al.* (2006) defined the *Pex23* protein family, which contains the membrane proteins *Pex23*, *Pex24*, *Pex28*, *Pex29*, *Pex30*, *Pex31*, and *Pex32*. This family of peroxisomal proteins can be divided into two groups of proteins with weak similarity (see Table 2; Kiel *et al.* 2006): The first group consists of *Y. lipolytica Pex23* (Brown *et al.* 2000) and the related *S. cerevisiae* proteins *Pex30*, *Pex31*, and *Pex32* (Vizeacoumar *et al.* 2004) and *P. pastoris Pex30* and *Pex31* (Yan *et al.* 2008). These proteins contain a DysF motif with an unknown function that was first observed in human dysferlin. The second group contains *Y. lipolytica Pex24* (Tam and Rachubinski 2002) and *S. cerevisiae Pex28* and *Pex29* (Vizeacoumar *et al.* 2003).

Y. lipolytica Pex23 is an integral peroxisomal membrane protein. *S. cerevisiae Pex30*, *Pex31*, and *Pex32* are localized to peroxisomes; however, a significant portion of *Pex30* is present in lighter fractions in sucrose gradients. These fractions may represent the ER, as *P. pastoris Pex30* and *Pex31* show a dual localization to the ER and peroxisomes. In *Y. lipolytica pex23* mutant cells, the majority of the peroxisomal matrix proteins is mislocalized to the cytosol, but the cells still contain small vesicular structures that contain *PTS1* and *PTS2* proteins. The phenotypes of *S. cerevisiae pex30*, *pex31*, and *pex32* cells are quite different because they contain peroxisomes in which matrix proteins are normally imported. In *pex30* mutant cells, the number of peroxisomes is increased, and *pex31* and *pex32* cells have enlarged

peroxisomes. It was proposed that, in *S. cerevisiae*, *Pex30* is a negative regulator of peroxisome number, whereas *Pex31* and *Pex32* are negative regulators of peroxisome size (Vizeacoumar *et al.* 2004). Notably, in *P. pastoris*, deletion of *PEX30* or *PEX31* results in the opposite effect, namely a reduction in peroxisome number (Yan *et al.* 2008).

Y. lipolytica *PEX24* encodes for a peroxisomal membrane protein, and *PEX24*-deficient mutants lack morphologically recognizable peroxisomes, but instead contain unusual extended membrane structures (Tam and Rachubinski 2002). Immunofluorescence microscopy suggested a cytosolic localization for PMPs and matrix proteins in these mutants; however, using biochemical approaches, membrane fractions could be detected that contained minor amounts of matrix marker proteins. Hence, *pex24* mutant cells apparently fail to assemble functional peroxisomes, but still contain membrane structures that exhibit some peroxisomal characteristics.

In *S. cerevisiae* cells in which either one or both *PEX28* and/or *PEX29* are deleted, peroxisome assembly is not affected. These mutant cells contain organelles that have a lower density, are smaller, are more abundant, and tend to cluster (Vizeacoumar *et al.* 2003). Two-hybrid studies revealed that *Pex28* and *Pex29* interact with *Pex30*, *Pex31*, and *Pex32*. Systematic deletion of genes demonstrated that *PEX28* and *PEX29* function upstream of *Pex30*, *Pex31*, and *Pex32* and function together with these proteins in the regulation of peroxisome proliferation (Vizeacoumar *et al.* 2004).

In summary, the *Y. lipolytica* members of the Pex23 family seem to play a key role in peroxisome assembly, whereas the *S. cerevisiae* and *P. pastoris* members are important in the regulation of peroxisome proliferation. Detailed knowledge about the function of these proteins is required to understand the major species-specific differences in the phenotypes of the respective deletion strains.

Peroxisome inheritance

During budding of yeast cells, one or a few organelles are actively transported to the developing bud, a process that is mediated by the class V myosin motor protein *Myo2* and *actin* filaments (Hoepfner *et al.* 2001). The remaining peroxisomes are retained in the mother cell. *Inp1* has been identified as the peroxisome-specific retention factor, connecting peroxisomes to an as-yet-unknown anchoring structure in the mother cell. The Hettema laboratory demonstrated that *Pex3* recruits *Inp1* to the peroxisomal membrane (Munck *et al.* 2009). Importantly, the *Inp1*-binding region in the *Pex3* protein is distinct from the regions involved in membrane formation. Unexpectedly, in the absence of *Pex11*, peroxisome retention is also defective in *H. polymorpha*, despite the fact that *Inp1* is properly localized to peroxisomes (Krikken *et al.* 2009). Hence, *Pex11* appears to have a second function in organelle retention, in addition to its role in peroxisome fission.

Inp2 is a PMP that acts as the peroxisomal receptor for *Myo2* and attaches the globular tail of *Myo2* to the peroxisome, thus allowing transport of the organelle to the bud.

The region of *Myo2* involved in *Inp2* binding was identified using mutant variants of *Myo2*. These studies also showed that *Inp2* is a phosphoprotein whose level of phosphorylation is coupled to the cell cycle (Fagarasanu *et al.* 2006; Fagarasanu *et al.* 2009). Recently, Otzen *et al.* (2012) provided evidence that *Pex19* also plays a role in peroxisome inheritance by associating peroxisomes to *Myo2*. Interestingly, mutations that affect the interaction between *Myo2* and *Pex19* do not abolish the *Inp2*–*Myo2* interaction.

Chang *et al.* (2009) suggested that *Inp2* is unique for *S. cerevisiae* and related species because in *Y. lipolytica* *Pex3* and its paralog *Pex3B* function as peroxisome-specific receptors of *Myo2*. However, *Inp2* is also present in other yeast species. The finding that *H. polymorpha* *Inp2* interacts with *Myo2* points to a conserved function of *Inp2* as a binding protein for *Myo2* (Saraya *et al.* 2010). Remarkably, in *H. polymorpha*, *Myo2*–*Inp2* binding was dependent on *Pex19*. This is consistent with the view that *Pex19* may have a stabilizing role in the interaction between *Inp2* and *Myo2* and is also in line with the previously observed defect in peroxisome inheritance in *H. polymorpha pex19* cells (Otzen *et al.* 2006).

Peroxisome degradation

The actual peroxisome population per cell is largely prescribed by physiological needs and determined by the machineries of organelle proliferation, inheritance, and degradation. Peroxisome inactivation can be achieved by degradation of (part of) their constituents or by turnover of the whole organelle by autophagy (Zwart *et al.* 1979). Organelle degradation by autophagy can serve three main cellular functions, namely nonselective degradation, *e.g.*, under nutrient depletion conditions to recycle cellular material; selective degradation of redundant organelles; and constitutive degradation to remove exhausted organelles as a mode to continuously rejuvenate the organelle population (Aksam *et al.* 2007). Selective peroxisome degradation is also designated as “pexophagy” (Klionsky *et al.* 2007). Two distinct mechanisms of pexophagy, termed “macropexophagy” and “micropexophagy,” which can be morphologically distinguished, may occur. During macropexophagy, individual organelles are sequestered by a double-membrane structure, the autophagosome, which fuses with the vacuole and releases the organelle into the lytic environment of the vacuole. During micropexophagy an organelle or a cluster of organelles is engulfed by vacuolar extensions, followed by incorporation of the organelle into the vacuole.

Most studies of the molecular mechanisms involved in pexophagy have been performed with methylotrophic yeast species such as *H. polymorpha* or *P. pastoris*. When these organisms are grown on methanol, peroxisomes are massively induced. However, upon transfer of methanol-grown cells to glucose media, these organelles become redundant for growth and are therefore rapidly degraded by pexophagy. Pexophagy involves a core set of *ATG* genes that also plays a role in other autophagy processes (Meijer *et al.* 2007). In addition, a few genes that are specifically involved in

pexophagy are known (for recent reviews see Sakai *et al.* 2006 and Manjithaya *et al.* 2010). Although homologous genes exist in *S. cerevisiae*, their specific roles in pexophagy in this organism have not been elucidated yet.

Peroxisome function and biology in yeast as a paradigm for metabolic disorders in humans

Yeast is a very useful model (“reference organism”) to understand basic processes of peroxisome biology in humans. As in yeast, human peroxisomes harbor enzymes of the β -oxidation pathway. However, in addition, they are involved in other processes that do not occur in *S. cerevisiae*, such as ether phospholipid biosynthesis, fatty acid α -oxidation, and the oxidation of *D*-amino acids and of polyamines (for reviews see Wanders and Waterham 2006 and Wanders *et al.* 2010). Also, unlike in yeast, mammalian peroxisomal β -oxidation is restricted to very long chain fatty acids (VLCFAs); short- and medium-chain FAs are oxidized by mitochondrial enzymes in mammals, whereas mitochondrial β -oxidation does not exist in *S. cerevisiae*, and short-, medium-, and long-chain fatty acids are solely degraded in peroxisomes in that organism. Since the metabolic pathways in yeast and human peroxisomes are quite distinct, yeast research has only poorly contributed to the understanding of the metabolism of human peroxisomes. In marked contrast, however, studies of yeast peroxisome biogenesis have been instrumental in identifying human counterparts of yeast peroxins and in understanding the molecular basis of peroxisome biogenesis disorders (PBDs). In PBD patients, normal peroxisomal structures are absent, causing dramatic defects in peroxisomal metabolism. Among other deficiencies, PBDs lead to the accumulation of very long chain fatty acids or defective plasmalogen synthesis, and PBD patients develop liver diseases, variable neurodevelopmental delay, retinopathy, and perceptive deafness with onset in the first months of life. The most severe PBD is Zellweger syndrome, and patients suffering from this disease typically die before one year of age (Steinberg *et al.* 2006).

In 1973 Goldfischer *et al.* (1973) described that cells of Zellweger syndrome patients were fully devoid of peroxisomal structures. However, it was not until 1992 that the first gene associated with a peroxisome biogenesis disorder was identified by complementation of fibroblasts isolated from a patient with Zellweger syndrome (Shimozawa *et al.* 1992). In later studies, the cloning and sequencing of yeast *PEX* genes has greatly facilitated the identification of the corresponding human genes by homology probing and searches in human genome databases. The genes that have been shown to be defective in PBDs are so far limited to those that play a role in matrix protein import and in the formation of the peroxisomal membrane, *i.e.*, *Pex3* and *Pex19*. In addition, *Pex16* is required for membrane biogenesis in humans. However, several of the other *PEX* genes appear to be yeast specific and do not have a clearly defined structural ortholog in humans (for a review see Steinberg *et al.* 2006).

Most of the research on yeast peroxisomes is performed using *S. cerevisiae* due to the extensive set of experimental tools that was first available for this species; however, other yeast species have specific advantages for peroxisome research, as their peroxisomes are more similar to their human counterparts in certain aspects. This has become apparent from the presence of orthologs of typical human peroxisomal enzymes in several yeast species, such as *D*-amino acid oxidases and polyamine oxidases, which are absent from *S. cerevisiae*. As outlined above, this is also true for peroxisomal processes, such as organelle fission. All eukaryotes, with the exception of *S. cerevisiae*, contain a peroxisomal Lon protease (Aksam *et al.* 2007), and studies in *H. polymorpha* and *Penicillium chysogenum* revealed that this protease is very important for quality control processes in the organelle (Aksam *et al.* 2007; Bartoszewska *et al.* 2012). Hence, studies on peroxisome biology in other fungi may have specific advantages over *S. cerevisiae* in understanding peroxisome-related processes in humans.

Outlook and perspectives

For many years the peroxisome field has struggled, and in fact still struggles, with major controversies regarding the mode of peroxisome biogenesis and development. One topic currently under heavy debate are the mechanisms by which peroxisomes are formed. Data in several yeast species indicate that, following their induction, peroxisomes predominantly proliferate by fission, whereas other data suggest that all organelles are derived from the ER, as suggested for mammalian peroxisomes. Potentially, ER-derived peroxisomes are capable of one or only a few fission events. Clearly, this point needs urgent elucidation and requires the identification and analysis of novel components that are essential for this process. Similarly, novel approaches will help to resolve the question of how the various PMPs reach the peroxisome membrane. It is attractive to suggest that sorting of at least some PMPs may be associated with lipid transfer from the ER, since the organelles increase in size during maturation; on the other hand, PMP and lipid import may also be separate processes. Notably, a major constituent of the peroxisomal membrane is cardiolipin, which is synthesized in the inner mitochondrial membrane, adding another level of complexity to the peroxisome assembly process.

The question of how matrix proteins enter peroxisomes has long been enigmatic but seems now to be have been cracked by the observations of the Erdmann group that these proteins enter via a transient pore, formed by *Pex5* and *Pex14* molecules (Meinecke *et al.* 2010). However, many questions remain, for example, regarding the composition and regulation of this pore. Particularly interesting is the excellent suggestion by Gould and Collins (2002) who proposed that receptor and matrix proteins may form large complexes (pre-import complex) prior to the membrane translocation step. An estimated pore diameter of up to 90 Å would indeed allow the import of such complexes. The formation of such complexes at the organelle membrane

may also explain why the pool of matrix precursor proteins in the cytosol is invariably extremely low. In the end, detailed protein structure information will be required to end the debate over the function of this fascinating and unique protein translocation machinery.

For the identification of novel peroxisome components, transcriptome studies have been proven to be of low value. Also, *in silico* prediction methods need improved programs. A major limitation in proteomics studies to identify novel components of the peroxisomal import and assembly machinery is the insufficient purity of the organelle fractions that are obtained by currently available fractionation protocols. Moreover, because of the occurrence of dual localizations of proteins, *e.g.*, of *Fis1* and *Rho1*, which are dually localized to mitochondria and peroxisomes, or *Pex30*, which is present in peroxisomes and the ER, certain proteins can erroneously be regarded as contaminants. Also, current procedures to isolate peroxisomal fractions are time-consuming and involve many purification steps, which may result in the dissociation of proteins that only weakly or transiently interact with the peroxisomal surface. Moreover, it is difficult to isolate intact organelles, and leakage of matrix proteins during the isolation procedure is invariably observed. An additional potential drawback may be the use of density centrifugation in current protocols since only organelles of high density are isolated; several data, however, indicate the presence of organelle subpopulations (Veenhuis *et al.* 1989), and immature developing organelles that do not cofractionate with the dense mature organelle fractions exist. In fact, the lighter, nascent organelles may carry the bulk of proteins important for peroxisome biogenesis and dynamics (Erdmann and Blobel 1995; Cepinska *et al.* 2011) and therefore are missed in the current isolation procedures. Procedures that are independent of peroxisome size or density would help in isolating the whole peroxisome population of cells. Immunopurification and FACS-based methods have been attempted, but need further improvement.

In general, the importance of peroxisomes has long been greatly underestimated. We anticipate that due to the technical drawbacks in identifying all relevant peroxisomal components the atlas of peroxisome functions is still far from being complete. This assumption is supported by the recent identification of several novel crucial functions, *i.e.*, non-metabolic activities that are required to cope with stress conditions other than oxidative stress, especially in plants. Among these are roles in reactive nitrogen species signaling, aging, antiviral innate immunity, and plant defense against pathogens (Dixit *et al.* 2010; Lazarow 2011). Of particular interest also are data indicating that peroxisomes play a role in aging in various species, including humans (Aksam *et al.* 2009; Bonekamp *et al.* 2009; Titorenko and Terlecky 2011). It is therefore important to analyze cells that have been grown under various conditions, including stress scenarios, to identify novel potential peroxisome functions in stress and aging.

Despite technical limitations, peroxisomes are among the best-characterized organelles in terms of composition, biogenesis, inheritance, and turnover in yeast. The field of lipid droplet research, on the other hand, is still in its infancy, despite the important role of this organelle in lipid homeostasis and its implication in prevalent metabolic diseases in mammals. Only models of how the neutral lipid of the LD core is formed exist, and although increasing evidence suggests a close functional interaction of LDs and the ER, the specific molecular mechanisms of LD formation are obscure (Farese and Walther 2009; Walther and Farese 2009, 2012). As discussed above for peroxisomes, the biochemical characterization of LD composition is limited since current LD isolation methods based on cell fractionation are time-consuming and may lead to cross-contamination with other cellular compartments or to loss of transiently associated components. Since LDs are typically isolated by flotation due to the low density of the TAG and SE core components, nascent LDs that contain less TAG and SE cannot be isolated by that method. As a consequence, the “time window” for LD isolation during cell cultivation is rather narrow and limited to the late-log/early stationary phase of growth. Thus, in addition to the mechanisms that drive LD formation, such fundamental concerns as determining the mechanism of protein targeting to LDs are largely unresolved. Anabolic LD enzymes are frequently also associated with the ER, whereas catabolic LD enzymes are exclusively LD resident: Which signals regulate the distribution of proteins between the ER and LDs? How are ER-resident proteins excluded from LDs? Evidence suggests that hydrophobic stretches of amino acids are required to drive LD association, but the topology of these potential transmembrane domains in the monolayer leaflet of the LD surface remains obscure. Since LDs may also serve as an overflow compartment for (misfolded) hydrophobic proteins, experimental use of episomally expressed fusion constructs to investigate targeting sequences may be misleading. Wild-type cells are characterized by a remarkably homogeneous size distribution of LDs (Czabany *et al.* 2008). What limits the size of LDs? Do LDs fuse *in vivo*? Are there specific subpopulations of LDs, harboring either SEs or TAGs, and are there also differences in the protein content between LDs? Yeast provides the unique opportunity to deplete LDs and induce their formation by regulated expression of acyltransferases that drive TAG synthesis (Jacquier *et al.* 2011), thus enabling studies on the early events of LD biogenesis and their interaction with the ER. Further refinement of isolation procedures and proteomic and lipidomic analysis will provide better insight into composition under various nutritional conditions (Connerth *et al.* 2009; Grillitsch *et al.* 2011). The exploitation of components involved in the physical interaction between LDs and mitochondria or peroxisomes (Pu *et al.* 2011) may also contribute to a better understanding of the interconnection between lipid storage and cellular physiology. In conclusion: many challenges remain and are open to further exploration

of the biology of these fascinating organelles, the peroxisome and lipid droplets.

Acknowledgments

We thank the members of our laboratories for critical reading of the manuscript and helpful suggestions. We gratefully acknowledge Khaw Aik Kia for the artwork shown in Figures 3 and 6 and Heimo Wolinski and Dagmar Kolb for providing fluorescence and electron micrographs shown in Figure 1. Work in the authors' laboratories is supported by funds from the Netherlands Organization for Scientific Research (NWO) and the Kluyver Centre for Genomics of Industrial Fermentation (to L.J.v.d.K.), which is part of the Netherlands Genomics Initiative/NWO, and by the Fonds zur Förderung der wissenschaftlichen Forschung in Österreich, FWF (to S.D.K.) (projects F3005-B12 SFB LIPOTOX and Ph.D. program "Molecular Enzymology").

Literature Cited

- AbdelRaheim, S. R., J. L. Cartwright, L. Gasmi, and A. G. McLennan, 2001 The NADH diphosphatase encoded by the *Saccharomyces cerevisiae* NPY1 nudix hydrolase gene is located in peroxisomes. *Arch. Biochem. Biophys.* 388: 18–24.
- Adeyo, O., P. J. Horn, S. Lee, D. D. Binns, A. Chandrashekar *et al.*, 2011 The yeast lipin orthologue Pah1p is important for biogenesis of lipid droplets. *J. Cell Biol.* 192: 1043–1055.
- Agarwal, A. K., and A. Garg, 2003 Congenital generalized lipodystrophy: significance of triglyceride biosynthetic pathways. *Trends Endocrinol. Metab.* 14: 214–221.
- Agarwal, A. K., R. I. Barnes, and A. Garg, 2004 Genetic basis of congenital generalized lipodystrophy. *Int. J. Obes. Relat. Metab. Disord.* 28: 336–339.
- Agne, B., N. M. Meindl, K. Niederhoff, H. Einwachter, P. Rehling *et al.*, 2003 Pex8p: an intraperoxisomal organizer of the peroxisomal import machinery. *Mol. Cell* 11: 635–646.
- Agrawal, G., S. Joshi, and S. Subramani, 2011 Cell-free sorting of peroxisomal membrane proteins from the endoplasmic reticulum. *Proc. Natl. Acad. Sci. USA* 108: 9113–9118.
- Aksam, E. B., A. Koek, J. A. Kiel, S. Jourdan, M. Veenhuis *et al.*, 2007 A peroxisomal lon protease and peroxisome degradation by autophagy play key roles in vitality of *Hansenula polymorpha* cells. *Autophagy* 3: 96–105.
- Aksam, E. B., B. de Vries, I. J. van der Klei, and J. A. Kiel, 2009 Preserving organelle vitality: peroxisomal quality control mechanisms in yeast. *FEMS Yeast Res.* 9: 808–820.
- Albertini, M., P. Rehling, R. Erdmann, W. Girzalsky, J. A. Kiel *et al.*, 1997 Pex14p, a peroxisomal membrane protein binding both receptors of the two PTS-dependent import pathways. *Cell* 89: 83–92.
- Albertyn, J., S. Hohmann, J. M. Thevelein, and B. A. Prior, 1994 GPD1, which encodes glycerol-3-phosphate dehydrogenase, is essential for growth under osmotic stress in *Saccharomyces cerevisiae*, and its expression is regulated by the high-osmolarity glycerol response pathway. *Mol. Cell. Biol.* 14: 4135–4144.
- Al-Feel, W., S. S. Chirala, and S. J. Wakil, 1992 Cloning of the yeast FAS3 gene and primary structure of yeast acetyl-CoA carboxylase. *Proc. Natl. Acad. Sci. USA* 89: 4534–4538.
- Anderson, R. M., K. J. Bitterman, J. G. Wood, O. Medvedik, and D. A. Sinclair, 2003 Nicotinamide and PNC1 govern lifespan extension by calorie restriction in *Saccharomyces cerevisiae*. *Nature* 423: 181–185.
- Antonio, E. A., C. Brees, E. Baumgart-Vogt, T. Hongu, S. J. Huybrechts *et al.*, 2009 Small G proteins in peroxisome biogenesis: the potential involvement of ADP-ribosylation factor 6. *BMC Cell Biol.* 10: 58.
- Antononkov, V. D., S. Mindthoff, S. Grunau, R. Erdmann, and J. K. Hiltunen, 2009 An involvement of yeast peroxisomal channels in transmembrane transfer of glyoxylate cycle intermediates. *Int. J. Biochem. Cell Biol.* 41: 2546–2554.
- Athenstaedt, K., and G. Daum, 1997 Biosynthesis of phosphatidic acid in lipid particles and endoplasmic reticulum of *Saccharomyces cerevisiae*. *J. Bacteriol.* 179: 7611–7616.
- Athenstaedt, K., and G. Daum, 1999 Phosphatidic acid, a key intermediate in lipid metabolism. *Eur. J. Biochem.* 266: 1–16.
- Athenstaedt, K., and G. Daum, 2000 1-Acyldihydroxyacetone-phosphate reductase (Ayr1p) of the yeast *Saccharomyces cerevisiae* encoded by the open reading frame YIL124w is a major component of lipid particles. *J. Biol. Chem.* 275: 235–240.
- Athenstaedt, K., and G. Daum, 2003 YMR313c/TGL3 encodes a novel triacylglycerol lipase located in lipid particles of *Saccharomyces cerevisiae*. *J. Biol. Chem.* 278: 23317–23323.
- Athenstaedt, K., and G. Daum, 2005 Tgl4p and Tgl5p, two triacylglycerol lipases of the yeast *Saccharomyces cerevisiae* are localized to lipid particles. *J. Biol. Chem.* 280: 37301–37309.
- Athenstaedt, K., and G. Daum, 2006 The life cycle of neutral lipids: synthesis, storage and degradation. *Cell. Mol. Life Sci.* 63: 1355–1369.
- Athenstaedt, K., and G. Daum, 2011 Lipid storage: Yeast we can! *Eur. J. Lipid Sci. Technol.* 113: 1188–1197.
- Athenstaedt, K., D. Zweytick, A. Jandrositz, S. D. Kohlwein, and G. Daum, 1999 Identification and characterization of major lipid particle proteins of the yeast *Saccharomyces cerevisiae*. *J. Bacteriol.* 181: 6441–6448.
- Avers, C. J., and M. Federman, 1968 The occurrence in yeast of cytoplasmic granules which resemble microbodies. *J. Cell Biol.* 37: 555–559.
- Azevedo, J. E., and W. Schliebs, 2006 Pex14p, more than just a docking protein. *Biochim. Biophys. Acta* 1763: 1574–1584.
- Barreto, L., A. Garcera, K. Jansson, P. Sunnerhagen, and E. Herrero, 2006 A peroxisomal glutathione transferase of *Saccharomyces cerevisiae* is functionally related to sulfur amino acid metabolism. *Eukaryot. Cell* 5: 1748–1759.
- Bartoszewska, M., L. Opalinski, M. Veenhuis, and I. J. van der Klei, 2011 The significance of peroxisomes in secondary metabolite biosynthesis in filamentous fungi. *Biotechnol. Lett.* 33(10): 1921–1931.
- Bartoszewska, M., C. Williams, A. Kikhney, L. Opalinski, C. W. van Roermund *et al.*, 2012 Peroxisomal proteostasis involves a Lon family protein that functions as protease and chaperone. *J. Biol. Chem.* 287: 27380–27395.
- Bellu, A. R., M. Komori, I. J. van der Klei, J. A. Kiel, and M. Veenhuis, 2001 Peroxisome biogenesis and selective degradation converge at Pex14p. *J. Biol. Chem.* 276: 44570–44574.
- Benghezal, M., C. Roubaty, V. Veepuri, J. Knudsen, and A. Conzelmann, 2007 SLC1 and SLC4 encode partially redundant acyl-coenzyme A 1-acylglycerol-3-phosphate O-acyltransferases of budding yeast. *J. Biol. Chem.* 282: 30845–30855.
- Binns, D., T. Januszewski, Y. Chen, J. Hill, V. S. Markin *et al.*, 2006 An intimate collaboration between peroxisomes and lipid bodies. *J. Cell Biol.* 173: 719–731.
- Binns, D., S. Lee, C. L. Hilton, Q. X. Jiang, and J. M. Goodman, 2010 Seipin is a discrete homooligomer. *Biochemistry* 49: 10747–10755.
- Black, P. N., and C. C. DiRusso, 2007 Yeast acyl-CoA synthetases at the crossroads of fatty acid metabolism and regulation. *Biochim. Biophys. Acta* 1771: 286–298.

- Globel, F., and R. Erdmann, 1996 Identification of a yeast peroxisomal member of the family of AMP-binding proteins. *Eur. J. Biochem.* 240: 468–476.
- Bonekamp, N. A., A. Volk, H. D. Fahimi, and M. Schrader, 2009 Reactive oxygen species and peroxisomes: struggling for balance. *Biofactors* 35: 346–355.
- Bottger, G., P. Barnett, A. T. Klein, A. Kragt, H. F. Tabak *et al.*, 2000 *Saccharomyces cerevisiae* PTS1 receptor Pex5p interacts with the SH3 domain of the peroxisomal membrane protein Pex13p in an unconventional, non-PXXP-related manner. *Mol. Biol. Cell* 11: 3963–3976.
- Boumann, H. A., M. J. Damen, C. Versluis, A. J. Heck, B. de Kruijff *et al.*, 2003 The two biosynthetic routes leading to phosphatidylcholine in yeast produce different sets of molecular species. Evidence for lipid remodeling. *Biochemistry* 42: 3054–3059.
- Bozaquel-Morais, B. L., J. B. Madeira, C. M. Maya-Monteiro, C. A. Masuda, and M. Montero-Lomeli, 2010 A new fluorescence-based method identifies protein phosphatases regulating lipid droplet metabolism. *PLoS ONE* 5(10): e13692.
- Brackmann, C., and J. Norbeck, M. Åkeson, D. Bosch, C. Larsson *et al.*, 2009 CARS microscopy of lipid stores in yeast: the impact of nutritional state and genetic background. *J. Raman Spectrosc.* 40: 748–756.
- Brasaemle, D. L., 2007 The perilipin family of structural lipid droplet proteins: stabilization of lipid droplets and control of lipolysis. *J. Lipid Res.* 48: 2547–2559.
- Braschi, E., V. Goyon, R. Zunino, A. Mohanty, L. Xu *et al.*, 2010 Vps35 mediates vesicle transport between the mitochondria and peroxisomes. *Curr. Biol.* 20: 1310–1315.
- Bratschi, M. W., D. P. Burrowes, A. Kulaga, J. F. Cheung, A. L. Alvarez *et al.*, 2009 Glycerol-3-phosphate acyltransferases *gat1p* and *gat2p* are microsomal phosphoproteins with differential contributions to polarized cell growth. *Eukaryot. Cell* 8: 1184–1196.
- Brocard, C., and A. Hartig, 2006 Peroxisome targeting signal 1: Is it really a simple tripeptide? *Biochim. Biophys. Acta* 1763: 1565–1573.
- Brocard, C., F. Kragler, M. M. Simon, T. Schuster, and A. Hartig, 1994 The tetratricopeptide repeat-domain of the PAS10 protein of *Saccharomyces cerevisiae* is essential for binding the peroxisomal targeting signal-SKL. *Biochem. Biophys. Res. Commun.* 204: 1016–1022.
- Brocard, C., G. Lametschwandtner, R. Koudelka, and A. Hartig, 1997 Pex14p is a member of the protein linkage map of Pex5p. *EMBO J.* 16: 5491–5500.
- Brown, T. W., V. I. Titorenko, and R. A. Rachubinski, 2000 Mutants of the *Yarrowia lipolytica* PEX23 gene encoding an integral peroxisomal membrane peroxin mislocalize matrix proteins and accumulate vesicles containing peroxisomal matrix and membrane proteins. *Mol. Biol. Cell* 11: 141–152.
- Burg, J. S., and P. J. Espenshade, 2011 Regulation of HMG-CoA reductase in mammals and yeast. *Prog. Lipid Res.* 50: 403–410.
- Carman, G. M., and G. S. Han, 2006 Roles of phosphatidate phosphatase enzymes in lipid metabolism. *Trends Biochem. Sci.* 31: 694–699.
- Carman, G. M., and G. S. Han, 2009 Phosphatidic acid phosphatase, a key enzyme in the regulation of lipid synthesis. *J. Biol. Chem.* 284: 2593–2597.
- Carman, G. M., and G. S. Han, 2011 Regulation of phospholipid synthesis in the yeast *Saccharomyces cerevisiae*. *Annu. Rev. Biochem.* 80: 859–883.
- Carman, G. M., and W. Wu, 2007 Lipid phosphate phosphatases from *Saccharomyces cerevisiae*, 305–315 in *Methods Enzymol.* edited by H. A. Brown. Academic Press, New York.
- Cartwright, J. L., L. Gasmi, D. G. Spiller, and A. G. McLennan, 2000 The *Saccharomyces cerevisiae* PCD1 gene encodes a peroxisomal nudix hydrolase active toward coenzyme A and its derivatives. *J. Biol. Chem.* 275: 32925–32930.
- Cepinska, M. N., M. Veenhuis, I. J. van der Klei, and S. Nagotu, 2011 Peroxisome fission is associated with reorganization of specific membrane proteins. *Traffic* 12: 925–937.
- Chang, J., F. D. Mast, A. Fagarasanu, D. A. Rachubinski, G. A. Eitzen *et al.*, 2009 Pex3 peroxisome biogenesis proteins function in peroxisome inheritance as class V myosin receptors. *J. Cell Biol.* 187: 233–246.
- Chapman, K. D., J. M. Dyer, and R. T. Mullen, 2012 Biogenesis and functions of lipid droplets in plants: Thematic Review Series: Lipid Droplet Synthesis and Metabolism: from Yeast to Man. *J. Lipid Res.* 53: 215–226.
- Chellappa, R., P. Kandasamy, C. S. Oh, Y. Jiang, M. Vemula *et al.*, 2001 The membrane proteins, Spt23p and Mga2p, play distinct roles in the activation of *Saccharomyces cerevisiae* OLE1 gene expression. Fatty acid-mediated regulation of Mga2p activity is independent of its proteolytic processing into a soluble transcription activator. *J. Biol. Chem.* 276: 43548–43556.
- Chen, Q., M. Kazachkov, Z. Zheng, and J. Zou, 2007 The yeast acylglycerol acyltransferase LCA1 is a key component of Lands cycle for phosphatidylcholine turnover. *FEBS Lett.* 581: 5511–5516.
- Choi, H. S., W. M. Su, J. M. Morgan, G. S. Han, Z. Xu *et al.*, 2011 Phosphorylation of phosphatidate phosphatase regulates its membrane association and physiological functions in *Saccharomyces cerevisiae*: Identification of SER602, THR723, and SER744 as the sites phosphorylated by CDC28 (CDK1)-encoded cyclin-dependent kinase. *J. Biol. Chem.* 286: 1486–1498.
- Choudhary, V., N. Jacquier, and R. Schneiter, 2011 The topology of the triacylglycerol synthesizing enzyme Lro1 indicates that neutral lipids can be produced within the luminal compartment of the endoplasmic reticulum: Implications for the biogenesis of lipid droplets. *Commun. Integr. Biol.* 4: 781–784.
- Connerth, M., K. Grillitsch, H. Kofeler, and G. Daum, 2009 Analysis of lipid particles from yeast. *Methods Mol. Biol.* 579: 359–374.
- Connerth, M., T. Czabany, A. Wagner, G. Zellnig, E. Leitner *et al.*, 2010 Oleate inhibits steryl ester synthesis and causes liposensitivity in yeast. *J. Biol. Chem.* 285: 26832–26841.
- Csaki, L. S., and K. Reue, 2010 *Lipins: multifunctional lipid metabolism proteins*. *Annu. Rev. Nutr.* 30: 257–272.
- Czabany, T., K. Athenstaedt, and G. Daum, 2007 Synthesis, storage and degradation of neutral lipids in yeast. *Biochim. Biophys. Acta* 1771: 299–309.
- Czabany, T., A. Wagner, D. Zwegtlick, K. Lohner, E. Leitner *et al.*, 2008 Structural and biochemical properties of lipid particles from the yeast *Saccharomyces cerevisiae*. *J. Biol. Chem.* 283: 17065–17074.
- Daum, G., A. Wagner, T. Czabany, and K. Athenstaedt, 2007a Dynamics of neutral lipid storage and mobilization in yeast. *Biochimie* 89: 243–248.
- Daum, G., A. Wagner, T. Czabany, K. Grillitsch, and K. Athenstaedt, 2007b Lipid storage and mobilization pathways in yeast. *Novartis Found. Symp.* 286: 142–151; discussion 151–144, 162–143, 196–203.
- Debelyy, M. O., H. W. Platta, D. Saffian, A. Hensel, S. Thoms *et al.*, 2011 Ubp15p, a ubiquitin hydrolase associated with the peroxisomal export machinery. *J. Biol. Chem.* 286: 28223–28234.
- De Duve, C., 1965 The separation and characterization of subcellular particles. *Harvey Lect.* 59: 49–87.
- Delille, H. K., and M. Schrader, 2008 Targeting of hFis1 to peroxisomes is mediated by Pex19p. *J. Biol. Chem.* 283: 31107–31115.
- De Smet, C. H., E. Vittone, M. Scherer, M. Houweling, G. Liebisch *et al.*, 2012 The yeast acyltransferase Sct1p regulates fatty acid

- desaturation by competing with the desaturase Ole1p. *Mol. Biol. Cell* 23: 1146–1156.
- De Virgilio, C., 2012 The essence of yeast quiescence. *FEMS Microbiol. Rev.* 36: 306–339.
- de Vries, B., J. A. Kiel, R. Scheek, M. Veenhuis, and I. J. van der Klei, 2007 A conserved alpha helical domain at the N-terminus of Pex14p is required for PTS1 and PTS2 protein import in *Hansenula polymorpha*. *FEBS Lett.* 581: 5627–5634.
- Dhavale, T., and G. Jedd, 2007 The fungal Woronin body, pp. 87–96 in *Biology of the Fungal Cell*, edited by R. J. Howard, and N. A. R. Gow. Springer, Berlin.
- Dickson, R. C., 2010 Roles for sphingolipids in *Saccharomyces cerevisiae*. *Adv. Exp. Med. Biol.* 688: 217–231.
- Distel, B., and A. Kragt, 2006 Purification of yeast peroxisomes. *Methods Mol. Biol.* 313: 21–26.
- Dixit, E., S. Boulant, Y. Zhang, A. S. Lee, C. Odendall *et al.*, 2010 Peroxisomes are signaling platforms for antiviral innate immunity. *Cell* 141: 668–681.
- Einwachter, H., S. Sowinski, W. H. Kunau, and W. Schliebs, 2001 *Yarrowia lipolytica* Pex20p, *Saccharomyces cerevisiae* Pex18p/Pex21p and mammalian Pex5pL fulfill a common function in the early steps of the peroxisomal PTS2 import pathway. *EMBO Rep.* 2: 1035–1039.
- Ejsing, C. S., J. L. Sampaio, V. Surendranath, E. Duchoslav, K. Ekroos *et al.*, 2009 Global analysis of the yeast lipidome by quantitative shotgun mass spectrometry. *Proc. Natl. Acad. Sci. USA* 106: 2136–2141.
- Emanuelsson, O., A. Elofsson, G. von Heijne, and S. Cristobal, 2003 In silico prediction of the peroxisomal proteome in fungi, plants and animals. *J. Mol. Biol.* 330: 443–456.
- Erdmann, R., and G. Blobel, 1995 Giant peroxisomes in oleic acid-induced *Saccharomyces cerevisiae* lacking the peroxisomal membrane protein Pmp27p. *J. Cell Biol.* 128: 509–523.
- Erdmann, R., M. Veenhuis, D. Mertens, and W. H. Kunau, 1989 Isolation of peroxisome-deficient mutants of *Saccharomyces cerevisiae*. *Proc. Natl. Acad. Sci. USA* 86: 5419–5423.
- Erdmann, R., F. F. Wiebel, A. Flessau, J. Rytka, A. Beyer *et al.*, 1991 PAS1, a yeast gene required for peroxisome biogenesis, encodes a member of a novel family of putative ATPases. *Cell* 64: 499–510.
- Espenshade, P. J., and A. L. Hughes, 2007 Regulation of sterol synthesis in eukaryotes. *Annu. Rev. Genet.* 41: 401–427.
- Fagarasanu, A., M. Fagarasanu, G. A. Eitzen, J. D. Aitchison, and R. A. Rachubinski, 2006 The peroxisomal membrane protein Inp2p is the peroxisome-specific receptor for the myosin V motor Myo2p of *Saccharomyces cerevisiae*. *Dev. Cell* 10: 587–600.
- Fagarasanu, A., F. D. Mast, B. Knoblach, Y. Jin, M. J. Brunner *et al.*, 2009 Myosin-driven peroxisome partitioning in *S. cerevisiae*. *J. Cell Biol.* 186: 541–554.
- Fakas, S., C. Konstantinou, and G. M. Carman, 2011a DGK1-encoded diacylglycerol kinase activity is required for phospholipid synthesis during growth resumption from stationary phase in *Saccharomyces cerevisiae*. *J. Biol. Chem.* 286: 1464–1474.
- Fakas, S., Y. Qiu, J. L. Dixon, G. S. Han, K. V. Ruggles *et al.*, 2011b Phosphatidate phosphatase activity plays key role in protection against fatty acid-induced toxicity in yeast. *J. Biol. Chem.* 286: 29074–29085.
- Farese, R. V. Jr, and T. C. Walther, 2009 Lipid droplets finally get a little R-E-S-P-E-C-T. *Cell* 139: 855–860.
- Fei, W., and H. Yang, 2012 Genome-wide screens for gene products regulating lipid droplet dynamics. *Methods Cell Biol.* 108: 303–316.
- Fei, W., G. Shui, B. Gaeta, X. Du, L. Kuerschner *et al.*, 2008 Fld1p, a functional homologue of human seipin, regulates the size of lipid droplets in yeast. *J. Cell Biol.* 180: 473–482.
- Fei, W., X. Du, and H. Yang, 2011a Seipin, adipogenesis and lipid droplets. *Trends Endocrinol. Metab.* 22: 204–210.
- Fei, W., H. Li, G. Shui, T. S. Kapterian, C. Bielby *et al.*, 2011b Molecular characterization of seipin and its mutants: implications for seipin in triacylglycerol synthesis. *J. Lipid Res.* 52: 2136–2147.
- Fei, W., G. Shui, Y. Zhang, N. Krahmer, C. Ferguson *et al.*, 2011c A role for phosphatidic acid in the formation of “supersized” lipid droplets. *PLoS Genet.* 7: e1002201.
- Fransen, M., I. Vastiau, C. Brees, V. Brys, G. P. Mannaerts *et al.*, 2004 Potential role for Pex19p in assembly of PTS-receptor docking complexes. *J. Biol. Chem.* 279: 12615–12624.
- Freitag, J., J. Alst, and M. Bölker, 2012 Cryptic peroxisomal targeting via alternative splicing and stop codon read-through in fungi. *Nature* 485: 522–525.
- Fujimoto, T., Y. Ohsaki, J. Cheng, M. Suzuki, and Y. Shinohara, 2008 Lipid droplets: a classic organelle with new outfits. *Histochem. Cell Biol.* 130: 263–279.
- Gallo, C. M., D. L. Smith, Jr, and J. S. Smith, 2004 Nicotinamide clearance by Pnc1 directly regulates Sir2-mediated silencing and longevity. *Mol. Cell Biol.* 24: 1301–1312.
- Garbarino, J., M. Padamsee, L. Wilcox, P. M. Oelkers, D. D’Ambrosio *et al.*, 2009 Sterol and diacylglycerol acyltransferase deficiency triggers fatty acid-mediated cell death. *J. Biol. Chem.* 284: 30994–31005.
- Garg, A., 2004 Acquired and inherited lipodystrophies. *N. Engl. J. Med.* 350: 1220–1234.
- Garg, A., and A. K. Agarwal, 2009 Lipodystrophies: disorders of adipose tissue biology. *Biochim. Biophys. Acta* 1791: 507–513.
- Gaspar, M. L., S. A. Jesch, R. Viswanatha, A. L. Antosh, W. J. Brown *et al.*, 2008 A block in endoplasmic reticulum-to-Golgi trafficking inhibits phospholipid synthesis and induces neutral lipid accumulation. *J. Biol. Chem.* 283: 25735–25751.
- Gaspar, M. L., H. F. Hofbauer, S. D. Kohlwein, and S. A. Henry, 2011 Coordination of storage lipid synthesis and membrane biogenesis: evidence for cross-talk between triacylglycerol metabolism and phosphatidylinositol synthesis. *J. Biol. Chem.* 286: 1696–1708.
- Gatto, G. J. Jr, B. V. Geisbrecht, S. J. Gould, and J. M. Berg, 2000 Peroxisomal targeting signal-1 recognition by the TPR domains of human PEX5. *Nat. Struct. Biol.* 7: 1091–1095.
- Geuze, H. J., J. L. Murk, A. K. Stroobants, J. M. Griffith, M. J. Kleijmeer *et al.*, 2003 Involvement of the endoplasmic reticulum in peroxisome formation. *Mol. Biol. Cell* 14: 2900–2907.
- Ghaemmaghami, S., W. K. Huh, K. Bower, R. W. Howson, A. Belle *et al.*, 2003 Global analysis of protein expression in yeast. *Nature* 425: 737–741.
- Girzalsky, W., P. Rehling, K. Stein, J. Kipper, L. Blank *et al.*, 1999 Involvement of Pex13p in Pex14p localization and peroxisomal targeting signal 2-dependent protein import into peroxisomes. *J. Cell Biol.* 144: 1151–1162.
- Glover, J. R., D. W. Andrews, and R. A. Rachubinski, 1994 *Saccharomyces cerevisiae* peroxisomal thiolase is imported as a dimer. *Proc. Natl. Acad. Sci. USA* 91: 10541–10545.
- Goffeau, A., B. G. Barrell, H. Bussey, R. W. Davis, B. Dujon *et al.*, 1996 Life with 6000 genes. *Science* 274(5287): 546; 563–567.
- Goldfischer, S., C. L. Moore, A. B. Johnson, A. J. Spiro, M. P. Valsamis *et al.*, 1973 Peroxisomal and mitochondrial defects in the cerebro-hepato-renal syndrome. *Science* 182: 62–64.
- Goodman, J. M., 2008 The gregarious lipid droplet. *J. Biol. Chem.* 283: 28005–28009.
- Goodman, J. M., 2009 Demonstrated and inferred metabolism associated with cytosolic lipid droplets. *J. Lipid Res.* 50: 2148–2156.
- Gould, S. J., and C. S. Collins, 2002 Opinion: peroxisomal-protein import: Is it really that complex? *Nat. Rev. Mol. Cell Biol.* 3: 382–389.

- Gould, S. G., G. A. Keller, and S. Subramani, 1987 Identification of a peroxisomal targeting signal at the carboxy terminus of firefly luciferase. *J. Cell Biol.* 105: 2923–2931.
- Gould, S. J., G. A. Keller, M. Schneider, S. H. Howell, L. J. Garrard *et al.*, 1990 Peroxisomal protein import is conserved between yeast, plants, insects and mammals. *EMBO J.* 9: 85–90.
- Grillitsch, K., M. Connerth, H. Kofeler, T. N. Arrey, B. Rietschel *et al.*, 2011 Lipid particles/droplets of the yeast *Saccharomyces cerevisiae* revisited: lipidome meets proteome. *Biochim. Biophys. Acta* 1811: 1165–1176.
- Gross, D. A., E. L. Snapp, and D. L. Silver, 2010 Structural insights into triglyceride storage mediated by fat storage-inducing transmembrane (FIT) protein 2. *PLoS ONE* 5: e10796.
- Gross, D. A., C. Zhan, and D. L. Silver, 2011 Direct binding of triglyceride to fat storage-inducing transmembrane proteins 1 and 2 is important for lipid droplet formation. *Proc. Natl. Acad. Sci. USA* 108: 19581–19586.
- Grunau, S., W. Schliebs, R. Linnepe, C. Neufeld, C. Cizmowski *et al.*, 2009 Peroxisomal targeting of PTS2 pre-import complexes in the yeast *Saccharomyces cerevisiae*. *Traffic* 10: 451–460.
- Gunkel, K., R. van Dijk, M. Veenhuis, and I. J. van der Klei, 2004 Routing of *Hansenula polymorpha* alcohol oxidase: an alternative peroxisomal protein-sorting machinery. *Mol. Biol. Cell* 15: 1347–1355.
- Guo, Y., K. R. Cordes, R. V. Farese Jr, and T. C. Walther, 2009 Lipid droplets at a glance. *J. Cell Sci.* 122: 749–752.
- Gurvitz, A., and H. Rottensteiner, 2006 The biochemistry of oleate induction: transcriptional upregulation and peroxisome proliferation. *Biochim. Biophys. Acta* 1763: 1392–1402.
- Gurvitz, A., A. M. Mursula, A. I. Yagi, A. Hartig, H. Ruis *et al.*, 1999 Alternatives to the isomerase-dependent pathway for the beta-oxidation of oleic acid are dispensable in *Saccharomyces cerevisiae*. Identification of YOR180c/DCI1 encoding peroxisomal delta(3,5)-delta(2,4)-dienoyl-CoA isomerase. *J. Biol. Chem.* 274: 24514–24521.
- Haan, G. J., R. J. Baerends, A. M. Krikken, M. Otzen, M. Veenhuis *et al.*, 2006 Reassembly of peroxisomes in *Hansenula polymorpha* pex3 cells on reintroduction of Pex3p involves the nuclear envelope. *FEMS Yeast Res.* 6: 186–194.
- Habeler, G., K. Natter, G. G. Thallinger, M. E. Crawford, S. D. Kohlwein *et al.*, 2002 YPLdb: the Yeast Protein Localization database. *Nucleic Acids Res.* 30: 80–83.
- Haemmerle, G., T. Moustafa, G. Woelkart, S. Buttner, A. Schmidt *et al.*, 2011 ATGL-mediated fat catabolism regulates cardiac mitochondrial function via PPAR-alpha and PGC-1. *Nat. Med.* 17: 1076–1085.
- Han, G. S., and G. M. Carman, 2010 Characterization of the human LPIN1-encoded phosphatidate phosphatase isoforms. *J. Biol. Chem.* 285: 14628–14638.
- Han, G. S., W. I. Wu, and G. M. Carman, 2006 The *Saccharomyces cerevisiae* lipin homolog is a Mg²⁺-dependent phosphatidate phosphatase enzyme. *J. Biol. Chem.* 281: 9210–9218.
- Han, G. S., S. Sinioglou, and G. M. Carman, 2007 The cellular functions of the yeast lipin homolog Pah1p are dependent on its phosphatidate phosphatase activity. *J. Biol. Chem.* 282: 37026–37035.
- Han, G. S., L. O'Hara, G. M. Carman, and S. Sinioglou, 2008a An unconventional diacylglycerol kinase that regulates phospholipid synthesis and nuclear membrane growth. *J. Biol. Chem.* 283: 20433–20442.
- Han, G. S., L. O'Hara, S. Sinioglou, and G. M. Carman, 2008b Characterization of the yeast DGK1-encoded CTP-dependent diacylglycerol kinase. *J. Biol. Chem.* 283: 20443–20453.
- Hapala, I., E. Marza, and T. Ferreira, 2011 Is fat so bad? Modulation of endoplasmic reticulum stress by lipid droplet formation. *Biol. Cell* 103: 271–285.
- Hasslacher, M., A. S. Ivessa, F. Paltauf, and S. D. Kohlwein, 1993 Acetyl-CoA carboxylase from yeast is an essential enzyme and is regulated by factors that control phospholipid metabolism. *J. Biol. Chem.* 268: 10946–10952.
- Hazra, P. P., I. Suriapranata, W. B. Snyder, and S. Subramani, 2002 Peroxisome remnants in pex3delta cells and the requirement of Pex3p for interactions between the peroxisomal docking and translocation subcomplexes. *Traffic* 3: 560–574.
- Heier, C., U. Taschler, S. Rengachari, M. Oberer, H. Wolinski *et al.*, 2010 Identification of Yju3p as functional orthologue of mammalian monoglyceride lipase in the yeast *Saccharomyces cerevisiae*. *Biochim. Biophys. Acta* 1801: 1063–1071.
- Henry, S. A., S. D. Kohlwein, and G. M. Carman, 2012 Metabolism and regulation of glycerolipids in the yeast *Saccharomyces cerevisiae*. *Genetics* 190: 317–349.
- Hettema, E. H., C. W. van Roermund, B. Distel, M. van den Berg, C. Vilela *et al.*, 1996 The ABC transporter proteins Pat1 and Pat2 are required for import of long-chain fatty acids into peroxisomes of *Saccharomyces cerevisiae*. *EMBO J.* 15: 3813–3822.
- Hettema, E. H., W. Girzalsky, M. van Den Berg, R. Erdmann, and B. Distel, 2000 *Saccharomyces cerevisiae* pex3p and pex19p are required for proper localization and stability of peroxisomal membrane proteins. *EMBO J.* 19: 223–233.
- Hodges, B. D. M., and C. C. Wu, 2010 Proteomic insights into an expanded cellular role for cytoplasmic lipid droplets. *J. Lipid Res.* 51: 262–273.
- Hoepfner, D., M. van den Berg, P. Philippsen, H. F. Tabak, and E. H. Hettema, 2001 A role for Vps1p, actin, and the Myo2p motor in peroxisome abundance and inheritance in *Saccharomyces cerevisiae*. *J. Cell Biol.* 155: 979–990.
- Hoepfner, D., D. Schildknecht, I. Braakman, P. Philippsen, and H. F. Tabak, 2005 Contribution of the endoplasmic reticulum to peroxisome formation. *Cell* 122: 85–95.
- Hofmann, K., 2000 A superfamily of membrane-bound O-acyltransferases with implications for wnt signaling. *Trends Biochem. Sci.* 25: 111–112.
- Hoppe, T., K. Matuschewski, M. Rape, S. Schlenker, H. D. Ulrich *et al.*, 2000 Activation of a membrane-bound transcription factor by regulated ubiquitin/proteasome-dependent processing. *Cell* 102: 577–586.
- Horvath, S. E., A. Wagner, E. Steyrer, and G. Daum, 2011 Metabolic link between phosphatidylethanolamine and triacylglycerol metabolism in the yeast *Saccharomyces cerevisiae*. *Biochim. Biophys. Acta* 1811: 1030–1037.
- Hronska, L., Z. Mrozova, M. Valachovic, and I. Hapala, 2004 Low concentrations of the non-ionic detergent Nonidet P-40 interfere with sterol biogenesis and viability of the yeast *Saccharomyces cerevisiae*. *FEMS Microbiol. Lett.* 238: 241–248.
- Huber, A., J. Koch, F. Kragler, C. Brocard, and A. Hartig, 2012 A subtle interplay between three Pex11 proteins shapes de novo formation and fission of peroxisomes. *Traffic* 13: 157–167.
- Huh, W. K., J. V. Falvo, L. C. Gerke, A. S. Carroll, R. W. Howson *et al.*, 2003 Global analysis of protein localization in budding yeast. *Nature* 425: 686–691.
- Islinger, M., K. W. Li, J. Seitz, A. Volk, and G. H. Luers, 2009 Hitchhiking of Cu/Zn superoxide dismutase to peroxisomes: evidence for a natural piggyback import mechanism in mammals. *Traffic* 10: 1711–1721.
- Ivniński-Steele, I., A. R. Holmes, E. Lamping, B. C. Monk, R. D. Cannon *et al.*, 2009 Identification of Nile red as a fluorescent substrate of the *Candida albicans* ATP-binding cassette transporters Cdr1p and Cdr2p and the major facilitator superfamily transporter Mdr1p. *Anal. Biochem.* 394: 87–91.
- Jacquier, N., V. Choudhary, M. Mari, A. Toulmay, F. Reggiori *et al.*, 2011 Lipid droplets are functionally connected to the endo-

- plasmic reticulum in *Saccharomyces cerevisiae*. *J. Cell Sci.* 124: 2424–2437.
- Jain, S., N. Stanford, N. Bhagwat, B. Seiler, M. Costanzo *et al.*, 2007 Identification of a novel lysophospholipid acyltransferase in *Saccharomyces cerevisiae*. *J. Biol. Chem.* 282: 30562–30569.
- Jandrositz, A., J. Petschnigg, R. Zimmermann, K. Natter, H. Scholze *et al.*, 2005 The lipid droplet enzyme Tgl1p hydrolyzes both steryl esters and triglycerides in the yeast, *Saccharomyces cerevisiae*. *Biochim. Biophys. Acta* 1735: 50–58.
- Jones, J. M., J. C. Morrell, and S. J. Gould, 2004 PEX19 is a predominantly cytosolic chaperone and import receptor for class 1 peroxisomal membrane proteins. *J. Cell Biol.* 164: 57–67.
- Jung, S., M. Marelli, R. A. Rachubinski, D. R. Goodlett, and J. D. Aitchison, 2010 Dynamic changes in the subcellular distribution of Gpd1p in response to cell stress. *J. Biol. Chem.* 285: 6739–6749.
- Kadereit, B., P. Kumar, W. J. Wang, D. Miranda, E. L. Snapp *et al.*, 2008 Evolutionarily conserved gene family important for fat storage. *Proc. Natl. Acad. Sci. USA* 105: 94–99.
- Kaeberlein, M., M. McVey, and L. Guarente, 1999 The SIR2/3/4 complex and SIR2 alone promote longevity in *Saccharomyces cerevisiae* by two different mechanisms. *Genes Dev.* 13: 2570–2580.
- Kal, A. J., A. J. van Zonneveld, V. Benes, M. van den Berg, M. G. Koerkamp *et al.*, 1999 Dynamics of gene expression revealed by comparison of serial analysis of gene expression transcript profiles from yeast grown on two different carbon sources. *Mol. Biol. Cell* 10: 1859–1872.
- Kal, A. J., E. H. Hettema, M. van den Berg, M. G. Koerkamp, L. van Ijlst *et al.*, 2000 In silico search for genes encoding peroxisomal proteins in *Saccharomyces cerevisiae*. *Cell Biochem. Biophys.* 32(Spring): 1–8.
- Kaliszewski, P., and T. Żołądek, 2008 The role of Rsp5 ubiquitin ligase in regulation of diverse processes in yeast cells. *Acta Biochim. Pol.* 55: 649–662.
- Kals, M., K. Natter, G. G. Thallinger, Z. Trajanoski, and S. D. Kohlwein, 2005 YPL.db2: the Yeast Protein Localization database, version 2.0. *Yeast* 22: 213–218.
- Karanasios, E., G. S. Han, Z. Xu, G. M. Carman, and S. Siniosoglou, 2010 A phosphorylation-regulated amphipathic helix controls the membrane translocation and function of the yeast phosphatidate phosphatase. *Proc. Natl. Acad. Sci. USA* 107: 17539–17544.
- Karnik, S. K., and R. N. Trelease, 2005 Arabidopsis peroxin 16 coexists at steady state in peroxisomes and endoplasmic reticulum. *Plant Physiol.* 138: 1967–1981.
- Kiel, J. A., K. Emmrich, H. E. Meyer, and W. H. Kunau, 2005a Ubiquitination of the peroxisomal targeting signal type 1 receptor, Pex5p, suggests the presence of a quality control mechanism during peroxisomal matrix protein import. *J. Biol. Chem.* 280: 1921–1930.
- Kiel, J. A., M. Otzen, M. Veenhuis, and I. J. van der Klei, 2005b Obstruction of polyubiquitination affects PTS1 peroxisomal matrix protein import. *Biochim. Biophys. Acta* 1745: 176–186.
- Kiel, J. A., M. Veenhuis, and I. J. van der Klei, 2006 PEX genes in fungal genomes: common, rare or redundant. *Traffic* 7: 1291–1303.
- Kienesberger, P. C., M. Oberer, A. Lass, and R. Zechner, 2009 Mammalian patatin domain containing proteins: a family with diverse lipolytic activities involved in multiple biological functions. *J. Lipid Res.* 50(Suppl.): S63–S68.
- Klein, A. T., M. van den Berg, G. Bottger, H. F. Tabak, and B. Distel, 2002 *Saccharomyces cerevisiae* acyl-CoA oxidase follows a novel, non-PTS1, import pathway into peroxisomes that is dependent on Pex5p. *J. Biol. Chem.* 277: 25011–25019.
- Klionsky, D. J., A. M. Cuervo, W. A. Dunn Jr., B. Levine, I. van der Klei *et al.*, 2007 How shall I eat thee? Autophagy 3: 413–416.
- Klose, C., M. A. Surma, M. J. Gerl, F. Meyenhofer, A. Shevchenko *et al.*, 2012 Flexibility of a eukaryotic lipidome: insights from yeast lipidomics. *PLoS ONE* 7: e35063.
- Knoblauch, B., and R. A. Rachubinski, 2010 Phosphorylation-dependent activation of peroxisome proliferator protein PEX11 controls peroxisome abundance. *J. Biol. Chem.* 285: 6670–6680.
- Kobayashi, S., A. Tanaka, and Y. Fujiki, 2007 Fis1, DLP1, and Pex11p coordinately regulate peroxisome morphogenesis. *Exp. Cell Res.* 313: 1675–1686.
- Koffel, R., and R. Schneiter, 2006 Yeh1 constitutes the major steryl ester hydrolase under heme-deficient conditions in *Saccharomyces cerevisiae*. *Eukaryot. Cell* 5: 1018–1025.
- Koffel, R., R. Tiwari, L. Falquet, and R. Schneiter, 2005 The *Saccharomyces cerevisiae* YLL012/YEH1, YLR020/YEH2, and TGL1 genes encode a novel family of membrane-anchored lipases that are required for steryl ester hydrolysis. *Mol. Cell. Biol.* 25: 1655–1668.
- Kohlwein, S. D., 2010a Obese and anorexic yeasts: experimental models to understand the metabolic syndrome and lipotoxicity. *Biochim. Biophys. Acta* 1801: 222–229.
- Kohlwein, S. D., 2010b Triacylglycerol homeostasis: insights from yeast. *J. Biol. Chem.* 285: 15663–15667.
- Kohlwein, S. D., and J. Petschnigg, 2007 Lipid-induced cell dysfunction and cell death: lessons from yeast. *Curr. Hypertens. Rep.* 9: 455–461.
- Komori, M., S. W. Rasmussen, J. A. Kiel, R. J. Baerends, J. M. Cregg *et al.*, 1997 The *Hansenula polymorpha* PEX14 gene encodes a novel peroxisomal membrane protein essential for peroxisome biogenesis. *EMBO J.* 16: 44–53.
- Kragler, F., A. Langeder, J. Raupachova, M. Binder, and A. Hartig, 1993 Two independent peroxisomal targeting signals in catalase A of *Saccharomyces cerevisiae*. *J. Cell Biol.* 120: 665–673.
- Kragt, A., T. Voorn-Brouwer, M. van den Berg, and B. Distel, 2005 Endoplasmic reticulum-directed Pex3p routes to peroxisomes and restores peroxisome formation in a *Saccharomyces cerevisiae* pex3Delta strain. *J. Biol. Chem.* 280: 34350–34357.
- Krahmer, N., Y. Guo, R. V. Farese Jr, and T. C. Walther, 2009 SnapShot: Lipid Droplets. *Cell* 139: 1024–1024.e102.
- Krikken, A. M., M. Veenhuis, and I. J. van der Klei, 2009 *Hansenula polymorpha* pex11 cells are affected in peroxisome retention. *FEBS J.* 276: 1429–1439.
- Kumar, A., S. Agarwal, J. A. Heyman, S. Matson, M. Heidtman *et al.*, 2002 Subcellular localization of the yeast proteome. *Genes Dev.* 16: 707–719.
- Kunau, W. H., V. Dommès, and H. Schulz, 1995 β -Oxidation of fatty acids in mitochondria, peroxisomes, and bacteria: a century of continued progress. *Prog. Lipid Res.* 34: 267–342.
- Kunze, M., I. Pracharoenwattana, S. M. Smith, and A. Hartig, 2006 A central role for the peroxisomal membrane in glyoxylate cycle function. *Biochim. Biophys. Acta* 1763: 1441–1452.
- Kurat, C. F., K. Natter, J. Petschnigg, H. Wolinski, K. Scheuringer *et al.*, 2006 Obese yeast: triglyceride lipolysis is functionally conserved from mammals to yeast. *J. Biol. Chem.* 281: 491–500.
- Kurat, C. F., H. Wolinski, J. Petschnigg, S. Kaluarachchi, B. Andrews *et al.*, 2009 Cdk1/Cdc28-dependent activation of the major triacylglycerol lipase Tgl4 in yeast links lipolysis to cell-cycle progression. *Mol. Cell* 33: 53–63.
- Kuravi, K., S. Nagotu, A. M. Krikken, K. Sjollem, M. Deckers *et al.*, 2006 Dynamin-related proteins Vps1p and Dnm1p control peroxisome abundance in *Saccharomyces cerevisiae*. *J. Cell Sci.* 119: 3994–4001.
- Kurbatova, E., M. Otzen, and I. J. van der Klei, 2009 p24 proteins play a role in peroxisome proliferation in yeast. *FEBS Lett.* 583: 3175–3180.

- Lam, S. K., N. Yoda, and R. Schekman, 2010 A vesicle carrier that mediates peroxisome protein traffic from the endoplasmic reticulum. *Proc. Natl. Acad. Sci. USA* 107: 21523–21528.
- Lambkin, G. R., and R. A. Rachubinski, 2001 *Yarrowia lipolytica* cells mutant for the peroxisomal peroxin Pex19p contain structures resembling wild-type peroxisomes. *Mol. Biol. Cell* 12: 3353–3364.
- Lametschwandtner, G., C. Brocard, M. Fransen, P. Van Veldhoven, J. Berger *et al.*, 1998 The difference in recognition of terminal tripeptides as peroxisomal targeting signal 1 between yeast and human is due to different affinities of their receptor Pex5p to the cognate signal and to residues adjacent to it. *J. Biol. Chem.* 273: 33635–33643.
- Lass, A., R. Zimmermann, M. Oberer, and R. Zechner, 2011 Lipolysis: a highly regulated multi-enzyme complex mediates the catabolism of cellular fat stores. *Prog. Lipid Res.* 50: 14–27.
- Lazarow, P. B., 2006 The import receptor Pex7p and the PTS2 targeting sequence. *Biochim. Biophys. Acta* 1763: 1599–1604.
- Lazarow, P. B., 2011 Viruses exploiting peroxisomes. *Curr. Opin. Microbiol.* 14: 458–469.
- Lazarow, P. B., and Y. Fujiki, 1985 Biogenesis of peroxisomes. *Annu. Rev. Cell Biol.* 1: 489–530.
- Leber, R., E. Zinser, G. Zellnig, F. Paltauf, and G. Daum, 1994 Characterization of lipid particles of the yeast, *Saccharomyces cerevisiae*. *Yeast* 10: 1421–1428.
- Leber, R., E. Zinser, C. Hrastnik, F. Paltauf, and G. Daum, 1995 Export of steryl esters from lipid particles and release of free sterols in the yeast, *Saccharomyces cerevisiae*. *Biochim. Biophys. Acta* 1234: 119–126.
- Leber, R., K. Landl, E. Zinser, H. Ahorn, A. Spok *et al.*, 1998 Dual localization of squalene epoxidase, Erg1p, in yeast reflects a relationship between the endoplasmic reticulum and lipid particles. *Mol. Biol. Cell* 9: 375–386.
- Leon, S., and S. Subramani, 2007 A conserved cysteine residue of *Pichia pastoris* Pex20p is essential for its recycling from the peroxisome to the cytosol. *J. Biol. Chem.* 282: 7424–7430.
- Li, D., and R. Roberts, 2001 WD-repeat proteins: structure characteristics, biological function, and their involvement in human diseases. *Cell. Mol. Life Sci.* 58: 2085–2097.
- Lingard, M. J., S. K. Gidda, S. Bingham, S. J. Rothstein, R. T. Mullen *et al.*, 2008 Arabidopsis PEROXIN11c-e, FISSION1b, and DYNAMIN-RELATED PROTEIN3A cooperate in cell cycle-associated replication of peroxisomes. *Plant Cell* 20: 1567–1585.
- Listenberger, L. L., X. Han, S. E. Lewis, S. Cases, R. V. Farese Jr. *et al.*, 2003 Triglyceride accumulation protects against fatty acid-induced lipotoxicity. *Proc. Natl. Acad. Sci. USA* 100: 3077–3082.
- Liu, H., X. Tan, M. Veenhuis, D. McCollum, and J. M. Cregg, 1992 An efficient screen for peroxisome-deficient mutants of *Pichia pastoris*. *J. Bacteriol.* 174: 4943–4951.
- Malanovic, N., I. Streith, H. Wolinski, G. Rechberger, S. D. Kohlwein *et al.*, 2008 S-adenosyl-L-homocysteine hydrolase, key enzyme of methylation metabolism, regulates phosphatidylcholine synthesis and triacylglycerol homeostasis in yeast: implications for homocysteine as a risk factor of atherosclerosis. *J. Biol. Chem.* 283: 23989–23999.
- Managadze, D., C. Wurtz, S. Wiese, M. Schneider, W. Girzalsky *et al.*, 2010 Identification of PEX33, a novel component of the peroxisomal docking complex in the filamentous fungus *Neurospora crassa*. *Eur. J. Cell Biol.* 89: 955–964.
- Manjithaya, R., T. Y. Nazarko, J. C. Farre, and S. Subramani, 2010 Molecular mechanism and physiological role of pexophagy. *FEBS Lett.* 584: 1367–1373.
- Marelli, M., J. J. Smith, S. Jung, E. Yi, A. I. Nesvizhskii *et al.*, 2004 Quantitative mass spectrometry reveals a role for the GTPase Rho1p in actin organization on the peroxisome membrane. *J. Cell Biol.* 167: 1099–1112.
- Marr, N., J. Foglia, M. Terebiznik, K. Athenstaedt, and V. Zaremb-berg, 2012 Controlling lipid fluxes at glycerol-3-phosphate acyltransferase step in yeast: unique contribution of Gat1p to oleic acid-induced lipid particle formation. *J. Biol. Chem.* 287: 10251–10264.
- Marshall, P. A., Y. I. Krimkevich, R. H. Lark, J. M. Dyer, M. Veenhuis *et al.*, 1995 Pmp27 promotes peroxisomal proliferation. *J. Cell Biol.* 129: 345–355.
- Martin, C. E., C. S. Oh, P. Kandasamy, R. Chellapa, and M. Vemula, 2002 Yeast desaturases. *Biochem. Soc. Trans.* 30: 1080–1082.
- Martin, C. E., C. S. Oh, and Y. Jiang, 2007 Regulation of long chain unsaturated fatty acid synthesis in yeast. *Biochim. Biophys. Acta* 1771: 271–285.
- Martin, W., and M. Muller, 1998 The hydrogen hypothesis for the first eukaryote. *Nature* 392: 37–41.
- Matsushita, M., and J. Nikawa, 1995 Isolation and characterization of a SCT1 gene which can suppress a choline-transport mutant of *Saccharomyces cerevisiae*. *J. Biochem.* 117: 447–451.
- Matsuzaki, T., and Y. Fujiki, 2008 The peroxisomal membrane protein import receptor Pex3p is directly transported to peroxisomes by a novel Pex19p- and Pex16p-dependent pathway. *J. Cell Biol.* 183: 1275–1286.
- McCammon, M. T., M. Veenhuis, S. B. Trapp, and J. M. Goodman, 1990 Association of glyoxylate and beta-oxidation enzymes with peroxisomes of *Saccharomyces cerevisiae*. *J. Bacteriol.* 172: 5816–5827.
- McNew, J. A., and J. M. Goodman, 1994 An oligomeric protein is imported into peroxisomes in vivo. *J. Cell Biol.* 127: 1245–1257.
- Meijer, W. H., I. J. van der Klei, M. Veenhuis, and J. A. Kiel, 2007 ATG genes involved in non-selective autophagy are conserved from yeast to man, but the selective Cvt and pexophagy pathways also require organism-specific genes. *Autophagy* 3: 106–116.
- Meinecke, M., C. Cizmowski, W. Schliebs, V. Kruger, S. Beck *et al.*, 2010 The peroxisomal importomer constitutes a large and highly dynamic pore. *Nat. Cell Biol.* 12: 273–277.
- Moir, R. D., D. A. Gross, D. L. Silver, and I. M. Willis, 2012 SCS3 and YFT2 Link Transcription of Phospholipid Biosynthetic Genes to ER Stress and the UPR. *PLoS Genet.* 8(8): e1002890.
- Moscicka, K. B., S. H. Klompmaker, D. Wang, I. J. van der Klei, and E. J. Boekema, 2007 The *Hansenula polymorpha* peroxisomal targeting signal 1 receptor, Pex5p, functions as a tetramer. *FEBS Lett.* 581: 1758–1762.
- Motley, A. M., and E. H. Hetteema, 2007 Yeast peroxisomes multiply by growth and division. *J. Cell Biol.* 178: 399–410.
- Motley, A. M., G. P. Ward, and E. H. Hetteema, 2008 Dnm1p-dependent peroxisome fission requires Caf4p, Mdv1p and Fis1p. *J. Cell Sci.* 121: 1633–1640.
- Mullner, H., and G. Daum, 2004 Dynamics of neutral lipid storage in yeast. *Acta Biochim. Pol.* 51: 323–347.
- Mullner, H., D. Zweytick, R. Leber, F. Turnowsky, and G. Daum, 2004 Targeting of proteins involved in sterol biosynthesis to lipid particles of the yeast *Saccharomyces cerevisiae*. *Biochim. Biophys. Acta* 1663: 9–13.
- Mullner, H., G. Deutsch, E. Leitner, E. Ingolic, and G. Daum, 2005 YEH2/YLR020c encodes a novel steryl ester hydrolase of the yeast *Saccharomyces cerevisiae*. *J. Biol. Chem.* 280: 13321–13328.
- Munck, J. M., A. M. Motley, J. M. Nuttall, and E. H. Hetteema, 2009 A dual function for Pex3p in peroxisome formation and inheritance. *J. Cell Biol.* 187: 463–471.
- Murphy, D. J., 2001 The biogenesis and functions of lipid bodies in animals, plants and microorganisms. *Prog. Lipid Res.* 40: 325–438.
- Murphy, S., S. Martin, and R. G. Parton, 2009 Lipid droplet-organelle interactions: sharing the fats. *Biochim. Biophys. Acta* 1791: 441–447.

- Nagotu, S., A. M. Krikken, M. Otzen, J. A. Kiel, M. Veenhuis *et al.*, 2008a Peroxisome fission in *Hansenula polymorpha* requires Mdv1 and Fis1, two proteins also involved in mitochondrial fission. *Traffic* 9: 1471–1484.
- Nagotu, S., R. Saraya, M. Otzen, M. Veenhuis, and I. J. van der Klei, 2008b Peroxisome proliferation in *Hansenula polymorpha* requires Dnm1p which mediates fission but not de novo formation. *Biochim. Biophys. Acta* 1783: 760–769.
- Natter, K., P. Leitner, A. Faschinger, H. Wolinski, S. McCraith *et al.*, 2005 The spatial organization of lipid synthesis in the yeast *Saccharomyces cerevisiae* derived from large scale green fluorescent protein tagging and high resolution microscopy. *Mol. Cell. Proteomics* 4: 662–672.
- Neuberger, G., S. Maurer-Stroh, B. Eisenhaber, A. Hartig, and F. Eisenhaber, 2003a Motif refinement of the peroxisomal targeting signal 1 and evaluation of taxon-specific differences. *J. Mol. Biol.* 328: 567–579.
- Neuberger, G., S. Maurer-Stroh, B. Eisenhaber, A. Hartig, and F. Eisenhaber, 2003b Prediction of peroxisomal targeting signal 1 containing proteins from amino acid sequence. *J. Mol. Biol.* 328: 581–592.
- Neufeld, C., F. V. Filipp, B. Simon, A. Neuhaus, N. Schuller *et al.*, 2009 Structural basis for competitive interactions of Pex14 with the import receptors Pex5 and Pex19. *EMBO J.* 28: 745–754.
- Niederhoff, K., N. M. Meindl-Beinker, D. Kerksen, U. Perband, A. Schafer *et al.*, 2005 Yeast Pex14p possesses two functionally distinct Pex5p and one Pex7p binding sites. *J. Biol. Chem.* 280: 35571–35578.
- Ntamack, A. G., I. V. Karpichev, S. J. Gould, G. M. Small, and H. Schulz, 2009 Oleate beta-oxidation in yeast involves thioesterase but not Yor180c protein that is not a dienoyl-CoA isomerase. *Biochim. Biophys. Acta* 1791: 371–378.
- Nuttley, W. M., A. M. Brade, G. A. Eitzen, M. Veenhuis, J. D. Aitchison *et al.*, 1994 PAY4, a gene required for peroxisome assembly in the yeast *Yarrowia lipolytica*, encodes a novel member of a family of putative ATPases. *J. Biol. Chem.* 269: 556–566.
- Nyathi, Y., and A. Baker, 2006 Plant peroxisomes as a source of signalling molecules. *Biochim. Biophys. Acta* 1763: 1478–1495.
- Oeljeklaus, S., B. S. Reinartz, J. Wolf, S. Wiese, J. Tonillo *et al.*, 2012 Identification of core components and transient interactors of the peroxisomal importomer by dual-track stable isotope labeling with amino acids in cell culture analysis. *J. Proteome Res.* 11: 2567–2580.
- Oelkers, P., A. Tinkelenberg, N. Erdeniz, D. Cromley, J. T. Billheimer *et al.*, 2000 A lecithin cholesterol acyltransferase-like gene mediates diacylglycerol esterification in yeast. *J. Biol. Chem.* 275: 15609–15612.
- Oelkers, P., D. Cromley, M. Padamsee, J. T. Billheimer, and S. L. Sturley, 2002 The DGA1 gene determines a second triglyceride synthetic pathway in yeast. *J. Biol. Chem.* 277: 8877–8881.
- O'Hara, L., G. S. Han, P. C. Sew, N. Grimsey, G. M. Carman *et al.*, 2006 Control of phospholipid synthesis by phosphorylation of the yeast lipin Pah1p/Smp2p Mg²⁺-dependent phosphatidate phosphatase. *J. Biol. Chem.* 281: 34537–34548.
- Olofsson, S. O., P. Boström, L. Andersson, M. Rutberg, J. Perman *et al.*, 2009 Lipid droplets as dynamic organelles connecting storage and efflux of lipids. *Biochim. Biophys. Acta* 1791: 448–458.
- Opalinski, L., J. A. Kiel, T. G. Homan, M. Veenhuis, and I. J. van der Klei, 2010 *Penicillium chrysogenum* Pex14/17p: a novel component of the peroxisomal membrane that is important for penicillin production. *FEBS J.* 277: 3203–3218.
- Opalinski, L., J. A. Kiel, C. Williams, M. Veenhuis, and I. J. van der Klei, 2011 Membrane curvature during peroxisome fission requires Pex11. *EMBO J.* 30: 5–16.
- Opperdoes, F. R., and P. Borst, 1977 Localization of nine glycolytic enzymes in a microbody-like organelle in *Trypanosoma brucei*: the glycosome. *FEBS Lett.* 80: 360–364.
- Otzen, M., U. Perband, D. Wang, R. J. Baerends, W. H. Kunau *et al.*, 2004 *Hansenula polymorpha* Pex19p is essential for the formation of functional peroxisomal membranes. *J. Biol. Chem.* 279: 19181–19190.
- Otzen, M., D. Wang, M. G. Lunenborg, and I. J. van der Klei, 2005 *Hansenula polymorpha* Pex20p is an oligomer that binds the peroxisomal targeting signal 2 (PTS2). *J. Cell Sci.* 118: 3409–3418.
- Otzen, M., A. M. Krikken, P. Z. Ozimek, E. Kurbatova, S. Nagotu *et al.*, 2006 In the yeast *Hansenula polymorpha*, peroxisome formation from the ER is independent of Pex19p, but involves the function of p24 proteins. *FEMS Yeast Res.* 6: 1157–1166.
- Otzen, M., R. Rucktäschel, S. Thoms, K. Emmrich, A. M. Krikken *et al.*, 2012 Pex19p contributes to peroxisome inheritance in the association of peroxisomes to Myo2p. *Traffic* 13(7): 947–959.
- Outeiro, T. F., and S. Lindquist, 2003 Yeast cells provide insight into alpha-synuclein biology and pathobiology. *Science* 302: 1772–1775.
- Ozimek, P., P. Kotter, M. Veenhuis, and I. J. van der Klei, 2006 *Hansenula polymorpha* and *Saccharomyces cerevisiae* Pex5p's recognize different, independent peroxisomal targeting signals in alcohol oxidase. *FEBS Lett.* 580: 46–50.
- Pagac, M., H. V. de la Mora, C. Duperrex, C. Roubaty, C. Vionnet *et al.*, 2011 Topology of 1-acyl-sn-glycerol-3-phosphate acyltransferases SLC1 and ALE1 and related membrane-bound O-acyltransferases (MBOATs) of *Saccharomyces cerevisiae*. *J. Biol. Chem.* 286: 36438–36447.
- Palmieri, L., H. Rottensteiner, W. Girzalsky, P. Scarcia, F. Palmieri *et al.*, 2001 Identification and functional reconstitution of the yeast peroxisomal adenine nucleotide transporter. *EMBO J.* 20: 5049–5059.
- Parish, R. W., 1975 The isolation and characterization of peroxisomes (microbodies) from baker's yeast, *Saccharomyces cerevisiae*. *Arch. Microbiol.* 105: 187–192.
- Pascual, F., and G. M. Carman, 2012 Phosphatidate phosphatase, a key regulator of lipid homeostasis. *Biochim. Biophys. Acta* (in press).
- Perkold, A., B. Zechmann, G. Daum, and G. Zellnig, 2007 Organelle association visualized by three-dimensional ultrastructural imaging of the yeast cell. *FEMS Yeast Res.* 7: 629–638.
- Perry, R. J., F. D. Mast, and R. A. Rachubinski, 2009 Endoplasmic reticulum-associated secretory proteins Sec20p, Sec39p, and Dsl1p are involved in peroxisome biogenesis. *Eukaryot. Cell* 8: 830–843.
- Peterfy, M., J. Phan, P. Xu, and K. Reue, 2001 Lipodystrophy in the fld mouse results from mutation of a new gene encoding a nuclear protein, lipin. *Nat. Genet.* 27: 121–124.
- Petschnigg, J., H. Wolinski, D. Kolb, G. Zelling, C. F. Kurat *et al.*, 2009 Good fat, essential cellular requirements for triacylglycerol synthesis to maintain membrane homeostasis in yeast. *J. Biol. Chem.* 284: 30981–30993.
- Pires, J. R., X. Hong, C. Brockmann, R. Volkmer-Engert, J. Schneider-Mergener *et al.*, 2003 The ScPex13p SH3 domain exposes two distinct binding sites for Pex5p and Pex14p. *J. Mol. Biol.* 326: 1427–1435.
- Platta, H. W., F. El Magraoui, B. E. Baumer, D. Schlee, W. Girzalsky *et al.*, 2009 Pex2 and pex12 function as protein-ubiquitin ligases in peroxisomal protein import. *Mol. Cell. Biol.* 29: 5505–5516.
- Poirier, Y., V. D. Antonenkov, T. Glumoff, and J. K. Hiltunen, 2006 Peroxisomal beta-oxidation: a metabolic pathway with multiple functions. *Biochim. Biophys. Acta* 1763: 1413–1426.

- Prein, B., K. Natter, and S. D. Kohlwein, 2000 A novel strategy for constructing N-terminal chromosomal fusions to green fluorescent protein in the yeast *Saccharomyces cerevisiae*. *FEBS Lett.* 485: 29–34.
- Ptacek, J., G. Devgan, G. Michaud, H. Zhu, X. Zhu *et al.*, 2005 Global analysis of protein phosphorylation in yeast. *Nature* 438: 679–684.
- Pu, J., C. W. Ha, S. Zhang, J. P. Jung, W. K. Huh *et al.*, 2011 Interactomic study on interaction between lipid droplets and mitochondria. *Protein Cell* 2: 487–496.
- Rajakumari, S., and G. Daum, 2010a Janus-faced enzymes yeast Tgl3p and Tgl5p catalyze lipase and acyltransferase reactions. *Mol. Biol. Cell* 21: 501–510.
- Rajakumari, S., and G. Daum, 2010b Multiple functions as lipase, steryl ester hydrolase, phospholipase, and acyltransferase of Tgl4p from the yeast *Saccharomyces cerevisiae*. *J. Biol. Chem.* 285: 15769–15776.
- Rajakumari, S., K. Grillitsch, and G. Daum, 2008 Synthesis and turnover of non-polar lipids in yeast. *Prog. Lipid Res.* 47: 157–171.
- Rajakumari, S., R. Rajasekharan, and G. Daum, 2010 Triacylglycerol lipolysis is linked to sphingolipid and phospholipid metabolism of the yeast *Saccharomyces cerevisiae*. *Biochim. Biophys. Acta* 1801: 1314–1322.
- Rape, M., T. Hoppe, I. Gorr, M. Kalocay, H. Richly *et al.*, 2001 Mobilization of processed, membrane-tethered SPT23 transcription factor by CDC48(UFD1/NPL4), a ubiquitin-selective chaperone. *Cell* 107: 667–677.
- Raychaudhuri, S., B. P. Young, P. J. Espenshade, and C. J. Loewen, 2012 Regulation of lipid metabolism: a tale of two yeasts. *Curr. Opin. Cell Biol.*
- Reue, K., and P. Zhang, 2008 The lipin protein family: dual roles in lipid biosynthesis and gene expression. *FEBS Lett.* 582: 90–96.
- Rhodin, J. A. G., 1954 *Correlation of Ultrastructural Organization and Function in Normal and Experimentally Changed Proximal Convolutated Tubule Cells of the Mouse Kidney*. Doctoral Thesis, Karolinska Institutet, Stockholm.
- Riekhof, W. R., J. Wu, J. L. Jones, and D. R. Voelker, 2007 Identification and characterization of the major lysophosphatidylethanolamine acyltransferase in *Saccharomyces cerevisiae*. *J. Biol. Chem.* 282: 28344–28352.
- Rockenfeller, P., J. Ring, V. Muschett, A. Beranek, S. Buettner *et al.*, 2010 Fatty acids trigger mitochondrion-dependent necrosis. *Cell Cycle* 9: 2836–2842.
- Roggenkamp, R., S. Numa, and E. Schweizer, 1980 Fatty acid-requiring mutant of *Saccharomyces cerevisiae* defective in acetyl-CoA carboxylase. *Proc. Natl. Acad. Sci. USA* 77: 1814–1817.
- Rottensteiner, H., K. Stein, E. Sonnenhol, and R. Erdmann, 2003 Conserved function of pex11p and the novel pex25p and pex27p in peroxisome biogenesis. *Mol. Biol. Cell* 14: 4316–4328.
- Sakai, Y., M. Oku, I. J. van der Klei, and J. A. Kiel, 2006 Pexophagy: autophagic degradation of peroxisomes. *Biochim. Biophys. Acta* 1763: 1767–1775.
- Saleem, R. A., B. Knoblach, F. D. Mast, J. J. Smith, J. Boyle *et al.*, 2008 Genome-wide analysis of signaling networks regulating fatty acid-induced gene expression and organelle biogenesis. *J. Cell Biol.* 181: 281–292.
- Saleem, R. A., R. Long-O'Donnell, D. J. Dilworth, A. M. Armstrong, A. P. Jamakhandi *et al.*, 2010 Genome-wide analysis of effectors of peroxisome biogenesis. *PLoS ONE* 5: e11953.
- Salomons, F. A., J. A. Kiel, K. N. Faber, M. Veenhuis, and I. J. van der Klei, 2000 Overproduction of Pex5p stimulates import of alcohol oxidase and dihydroxyacetone synthase in a *Hansenula polymorpha* Pex14 null mutant. *J. Biol. Chem.* 275: 12603–12611.
- Sandager, L., M. H. Gustavsson, U. Ståhl, A. Dahlqvist, E. Wiberg *et al.*, 2002 Storage lipid synthesis is non-essential in yeast. *J. Biol. Chem.* 277: 6478–6482.
- Santos-Rosa, H., J. Leung, N. Grimsey, S. Peak-Chew, and S. Siniosoglou, 2005 The yeast lipin Smp2 couples phospholipid biosynthesis to nuclear membrane growth. *EMBO J.* 24: 1931–1941.
- Saraya, R., M. N. Cepinska, J. A. Kiel, M. Veenhuis, and I. J. van der Klei, 2010 A conserved function for Inp2 in peroxisome inheritance. *Biochim. Biophys. Acta* 1803: 617–622.
- Saraya, R., A. M. Krikken, M. Veenhuis, and I. J. van der Klei, 2011 Peroxisome reintroduction in *Hansenula polymorpha* requires Pex25 and Rho1. *J. Cell Biol.* 193: 885–900.
- Schafer, A., D. Kerssen, M. Veenhuis, W. H. Kunau, and W. Schliebs, 2004 Functional similarity between the peroxisomal PTS2 receptor binding protein Pex18p and the N-terminal half of the PTS1 receptor Pex5p. *Mol. Cell. Biol.* 24: 8895–8906.
- Schafer, H., K. Nau, A. Sickmann, R. Erdmann, and H. E. Meyer, 2001 Identification of peroxisomal membrane proteins of *Saccharomyces cerevisiae* by mass spectrometry. *Electrophoresis* 22: 2955–2968.
- Schaffer, J. E., 2003 Lipotoxicity: when tissues overeat. *Curr. Opin. Lipidol.* 14: 281–287.
- Scharnewski, M., P. Pongdontri, G. Mora, M. Hoppert, and M. Fulda, 2008 Mutants of *Saccharomyces cerevisiae* deficient in acyl-CoA synthetases secrete fatty acids due to interrupted fatty acid recycling. *FEBS J.* 275: 2765–2778.
- Schell-Steven, A., K. Stein, M. Amoros, C. Landgraf, R. Volkmer-Engert *et al.*, 2005 Identification of a novel, intraperoxisomal pex14-binding site in pex13: association of pex13 with the docking complex is essential for peroxisomal matrix protein import. *Mol. Cell. Biol.* 25: 3007–3018.
- Schliebs, W., and W. H. Kunau, 2004 Peroxisome membrane biogenesis: the stage is set. *Curr. Biol.* 14: R397–R399.
- Schliebs, W., and W. H. Kunau, 2006 PTS2 co-receptors: diverse proteins with common features. *Biochim. Biophys. Acta* 1763: 1605–1612.
- Schliebs, W., W. Girzalsky, and R. Erdmann, 2010 Peroxisomal protein import and ERAD: variations on a common theme. *Nat. Rev. Mol. Cell Biol.* 11: 885–890.
- Schneider, R., B. Brugger, R. Sandhoff, G. Zellnig, A. Leber *et al.*, 1999 Electrospray ionization tandem mass spectrometry (ESI-MS/MS) analysis of the lipid molecular species composition of yeast subcellular membranes reveals acyl chain-based sorting/remodeling of distinct molecular species en route to the plasma membrane. *J. Cell Biol.* 146: 741–754.
- Schueller, N., S. J. Holton, K. Fodor, M. Milewski, P. Konarev *et al.*, 2010 The peroxisomal receptor Pex19p forms a helical mPTS recognition domain. *EMBO J.* 29: 2491–2500.
- Schweiger, M., A. Lass, R. Zimmermann, T. O. Eichmann, and R. Zechner, 2009 Neutral lipid storage disease: genetic disorders caused by mutations in adipose triglyceride lipase/PNPLA2 or CGI-58/ABHD5. *Am. J. Physiol. Endocrinol. Metab.* 297: E289–E296.
- Sere, Y. Y., M. Regnacq, J. Colas, and T. Berges, 2010 A *Saccharomyces cerevisiae* strain unable to store neutral lipids is tolerant to oxidative stress induced by alpha-synuclein. *Free Radic. Biol. Med.* 49: 1755–1764.
- Shimozawa, N., T. Tsukamoto, Y. Suzuki, T. Orii, Y. Shirayoshi *et al.*, 1992 A human gene responsible for Zellweger syndrome that affects peroxisome assembly. *Science* 255: 1132–1134.
- Shiozawa, K., P. V. Konarev, C. Neufeld, M. Wilmanns, and D. I. Svergun, 2009 Solution structure of human Pex5.Pex14.PTS1 protein complexes obtained by small angle X-ray scattering. *J. Biol. Chem.* 284: 25334–25342.
- Siniosoglou, S., H. Santos-Rosa, J. Rappsilber, M. Mann, and E. Hurt, 1998 A novel complex of membrane proteins required for formation of a spherical nucleus. *EMBO J.* 17: 6449–6464.

- Skoneczny, M., A. Chelstowska, and J. Rytka, 1988 Study of the coinduction by fatty acids of catalase A and acyl-CoA oxidase in standard and mutant *Saccharomyces cerevisiae* strains. *Eur. J. Biochem.* 174: 297–302.
- Smith, J. J., M. Marelli, R. H. Christmas, F. J. Vizeacoumar, D. J. Dilworth *et al.*, 2002 Transcriptome profiling to identify genes involved in peroxisome assembly and function. *J. Cell Biol.* 158: 259–271.
- Sorger, D., and G. Daum, 2002 Synthesis of triacylglycerols by the acyl-coenzyme A:diacyl-glycerol acyltransferase Dga1p in lipid particles of the yeast *Saccharomyces cerevisiae*. *J. Bacteriol.* 184: 519–524.
- Sorger, D., and G. Daum, 2003 Triacylglycerol biosynthesis in yeast. *Appl. Microbiol. Biotechnol.* 61: 289–299.
- Sorger, D., K. Athenstaedt, C. Hrstnik, and G. Daum, 2004 A yeast strain lacking lipid particles bears a defect in ergosterol formation. *J. Biol. Chem.* 279: 31190–31196.
- South, S. T., E. Baumgart, and S. J. Gould, 2001 Inactivation of the endoplasmic reticulum protein translocation factor, Sec61p, or its homolog, Ssh1p, does not affect peroxisome biogenesis. *Proc. Natl. Acad. Sci. USA* 98: 12027–12031.
- Spandl, J., D. J. White, J. Peychl, and C. Thiele, 2009 Live cell multicolor imaging of lipid droplets with a new dye, LD540. *Traffic* 10: 1579–1584.
- Spanova, M., T. Czabany, G. N. Zellnig, E. Leitner, I. Hapala *et al.*, 2010 Effect of lipid particle biogenesis on the subcellular distribution of squalene in the yeast *Saccharomyces cerevisiae*. *J. Biol. Chem.* 285: 6127–6133.
- Spanova, M., D. Zwegytick, K. Lohner, L. Klug, E. Leitner *et al.*, 2012 Influence of squalene on lipid particle/droplet and membrane organization in the yeast *Saccharomyces cerevisiae*. *Biochim. Biophys. Acta* 1821: 647–653.
- Stanley, W. A., F. V. Philipp, P. Kursula, N. Schuller, R. Erdmann *et al.*, 2006 Recognition of a functional peroxisome type 1 target by the dynamic import receptor pex5p. *Mol. Cell* 24: 653–663.
- Stanley, W. A., K. Fodor, M. A. Marti-Renom, W. Schliebs, and M. Wilmanns, 2007 Protein translocation into peroxisomes by ring-shaped import receptors. *FEBS Lett.* 581: 4795–4802.
- Stein, K., A. Schell-Steven, R. Erdmann, and H. Rottensteiner, 2002 Interactions of Pex7p and Pex18p/Pex21p with the peroxisomal docking machinery: implications for the first steps in PTS2 protein import. *Mol. Cell. Biol.* 22: 6056–6069.
- Steinberg, S. J., G. Dodt, G. V. Raymond, N. E. Braverman, A. B. Moser *et al.*, 2006 Peroxisome biogenesis disorders. *Biochim. Biophys. Acta* 1763: 1733–1748.
- Stevens, P., I. Monastyrska, A. N. Leao-Helder, I. J. van der Klei, M. Veenhuis *et al.*, 2005 *Hansenula polymorpha* Vam7p is required for macropexophagy. *FEMS Yeast Res.* 5: 985–997.
- Strating, J. R., T. G. Hafmans, and G. J. Martens, 2009 Functional diversity among p24 subfamily members. *Biol. Cell* 101: 207–219.
- Stukey, J. E., V. M. McDonough, and C. E. Martin, 1989 Isolation and characterization of OLE1, a gene affecting fatty acid desaturation from *Saccharomyces cerevisiae*. *J. Biol. Chem.* 264: 16537–16544.
- Stukey, J. E., V. M. McDonough, and C. E. Martin, 1990 The OLE1 gene of *Saccharomyces cerevisiae* encodes the delta 9 fatty acid desaturase and can be functionally replaced by the rat stearyl-CoA desaturase gene. *J. Biol. Chem.* 265: 20144–20149.
- Szymanski, K. M., D. Binns, R. Bartz, N. V. Grishin, W. P. Li *et al.*, 2007 The lipodystrophy protein seipin is found at endoplasmic reticulum lipid droplet junctions and is important for droplet morphology. *Proc. Natl. Acad. Sci. USA* 104: 20890–20895.
- Tabak, H. F., A. van der Zand, and I. Braakman, 2008 Peroxisomes: minted by the ER. *Curr. Opin. Cell Biol.* 20: 393–400.
- Tam, Y. Y., and R. A. Rachubinski, 2002 *Yarrowia lipolytica* cells mutant for the PEX24 gene encoding a peroxisomal membrane peroxin mislocalize peroxisomal proteins and accumulate membrane structures containing both peroxisomal matrix and membrane proteins. *Mol. Biol. Cell* 13: 2681–2691.
- Tam, Y. Y., J. C. Torres-Guzman, F. J. Vizeacoumar, J. J. Smith, M. Marelli *et al.*, 2003 Pex11-related proteins in peroxisome dynamics: a role for the novel peroxin Pex27p in controlling peroxisome size and number in *Saccharomyces cerevisiae*. *Mol. Biol. Cell* 14: 4089–4102.
- Tam, Y. Y., A. Fagarasanu, M. Fagarasanu, and R. A. Rachubinski, 2005 Pex3p initiates the formation of a preperoxisomal compartment from a subdomain of the endoplasmic reticulum in *Saccharomyces cerevisiae*. *J. Biol. Chem.* 280: 34933–34939.
- Taylor, K. M., C. P. Kaplan, X. Gao, and A. Baker, 1996 Localization and targeting of isocitrate lyases in *Saccharomyces cerevisiae*. *Biochem. J.* 319(Pt. 1): 255–262.
- Tehlivets, O., 2011 Homocysteine as a risk factor for atherosclerosis: Is its conversion to s-adenosyl-L-homocysteine the key to deregulated lipid metabolism? *J. Lipids* 2011: 702853.
- Tehlivets, O., M. Hasslacher, and S. D. Kohlwein, 2004 S-adenosyl-L-homocysteine hydrolase in yeast: key enzyme of methylation metabolism and coordinated regulation with phospholipid synthesis. *FEBS Lett.* 577: 501–506.
- Tehlivets, O., K. Scheuringer, and S. D. Kohlwein, 2007 Fatty acid synthesis and elongation in yeast. *Biochim. Biophys. Acta* 1771: 255–270.
- Thoms, S., M. O. Debelyy, K. Nau, H. E. Meyer, and R. Erdmann, 2008 Lpx1p is a peroxisomal lipase required for normal peroxisome morphology. *FEBS J.* 275: 504–514.
- Titorenko, V. I., and S. R. Terlecky, 2011 Peroxisome metabolism and cellular aging. *Traffic* 12: 252–259.
- Titorenko, V. I., J. J. Smith, R. K. Szilard, and R. A. Rachubinski, 1998 Pex20p of the yeast *Yarrowia lipolytica* is required for the oligomerization of thiolase in the cytosol and for its targeting to the peroxisome. *J. Cell Biol.* 142: 403–420.
- Titorenko, V. I., H. Chan, and R. A. Rachubinski, 2000 Fusion of small peroxisomal vesicles in vitro reconstructs an early step in the in vivo multistep peroxisome assembly pathway of *Yarrowia lipolytica*. *J. Cell Biol.* 148(1): 29–44.
- Tolbert, N. E., and E. Essner, 1981 Microbodies: peroxisomes and glyoxysomes. *J. Cell Biol.* 91: 271s–283s.
- Tower, R. J., A. Fagarasanu, J. D. Aitchison, and R. A. Rachubinski, 2011 The peroxin Pex34p functions with the Pex11 family of peroxisomal divisional proteins to regulate the peroxisome population in yeast. *Mol. Biol. Cell* 22: 1727–1738.
- Tsvetanova, N. G., D. M. Klass, J. Salzman, and P. O. Brown, 2010 Proteome-wide search reveals unexpected RNA-binding proteins in *Saccharomyces cerevisiae*. *PLoS ONE* 5: pii: e12671.
- Ubersax, J. A., E. L. Woodbury, P. N. Quang, M. Paraz, J. D. Blethrow *et al.*, 2003 Targets of the cyclin-dependent kinase Cdk1. *Nature* 425: 859–864.
- Urquhart, A. J., D. Kennedy, S. J. Gould, and D. I. Crane, 2000 Interaction of Pex5p, the type 1 peroxisome targeting signal receptor, with the peroxisomal membrane proteins Pex14p and Pex13p. *J. Biol. Chem.* 275: 4127–4136.
- van der Klei, I. J., and M. Veenhuis, 2006a PTS1-independent sorting of peroxisomal matrix proteins by Pex5p. *Biochim. Biophys. Acta* 1763: 1794–1800.
- van der Klei, I. J., and M. Veenhuis, 2006b Yeast and filamentous fungi as model organisms in microbody research. *Biochim. Biophys. Acta* 1763: 1364–1373.
- van der Klei, I. J., W. Harder, and M. Veenhuis, 1991 Methanol metabolism in a peroxisome-deficient mutant of *Hansenula polymorpha*: a physiological study. *Arch. Microbiol.* 156: 15–23.

- van der Klei, I. J., R. E. Hilbrands, G. J. Swaving, H. R. Waterham, E. G. Vrieling *et al.*, 1995 The *Hansenula polymorpha* PER3 gene is essential for the import of PTS1 proteins into the peroxisomal matrix. *J. Biol. Chem.* 270: 17229–17236.
- van der Klei, I. J., R. E. Hilbrands, J. A. Kiel, S. W. Rasmussen, J. M. Cregg *et al.*, 1998 The ubiquitin-conjugating enzyme Pex4p of *Hansenula polymorpha* is required for efficient functioning of the PTS1 import machinery. *EMBO J.* 17: 3608–3618.
- van der Zand, A., I. Braakman, and H. F. Tabak, 2010 Peroxisomal membrane proteins insert into the endoplasmic reticulum. *Mol. Biol. Cell* 21: 2057–2065.
- van der Zand, A., J. Gent, I. Braakman, and H. F. Tabak, 2012 Biochemically distinct vesicles from the endoplasmic reticulum fuse to form peroxisomes. *Cell* 13(149): 397–409.
- van Roermund, C. W., H. R. Waterham, L. Ijlst, and R. J. Wanders, 2003 Fatty acid metabolism in *Saccharomyces cerevisiae*. *Cell. Mol. Life Sci.* 60: 1838–1851.
- van Zutphen, T., R. J. Baerends, K. A. Susanna, A. de Jong, O. P. Kuipers *et al.*, 2010 Adaptation of *Hansenula polymorpha* to methanol: a transcriptome analysis. *BMC Genomics* 11: 1.
- Veenhuis, M., M. Mateblowski, W. H. Kunau, and W. Harder, 1987 Proliferation of microbodies in *Saccharomyces cerevisiae*. *Yeast* 3: 77–84.
- Veenhuis, M., G. Sulter, I. van der Klei, and W. Harder, 1989 Evidence for functional heterogeneity among microbodies in yeasts. *Arch. Microbiol.* 151: 105–110.
- Vizeacoumar, F. J., J. C. Torres-Guzman, Y. Y. Tam, J. D. Aitchison, and R. A. Rachubinski, 2003 YHR150w and YDR479c encode peroxisomal integral membrane proteins involved in the regulation of peroxisome number, size, and distribution in *Saccharomyces cerevisiae*. *J. Cell Biol.* 161: 321–332.
- Vizeacoumar, F. J., J. C. Torres-Guzman, D. Bouard, J. D. Aitchison, and R. A. Rachubinski, 2004 Pex30p, Pex31p, and Pex32p form a family of peroxisomal integral membrane proteins regulating peroxisome size and number in *Saccharomyces cerevisiae*. *Mol. Biol. Cell* 15: 665–677.
- Wagner, A., K. Grillitsch, E. Leitner, and G. Daum, 2009 Mobilization of steryl esters from lipid particles of the yeast *Saccharomyces cerevisiae*. *Biochim. Biophys. Acta* 1791: 118–124.
- Walther, T. C., and R. V. Farese, Jr., 2009 The life of lipid droplets. *Biochim. Biophys. Acta* 1791: 459–466.
- Walther, T. C., and R. V. Farese Jr., 2012 Lipid droplets and cellular lipid metabolism. *Annu. Rev. Biochem.* 81: 687–714.
- Walton, P. A., P. E. Hill, and S. Subramani, 1995 Import of stably folded proteins into peroxisomes. *Mol. Biol. Cell* 6: 675–683.
- Wanders, R. J., and H. R. Waterham, 2006 Biochemistry of mammalian peroxisomes revisited. *Annu. Rev. Biochem.* 75: 295–332.
- Wanders, R. J., S. Ferdinandusse, P. Brites, and S. Kemp, 2010 Peroxisomes, lipid metabolism and lipotoxicity. *Biochim. Biophys. Acta* 1801: 272–280.
- Wang, D., N. V. Visser, M. Veenhuis, and I. J. van der Klei, 2003 Physical interactions of the peroxisomal targeting signal 1 receptor pex5p, studied by fluorescence correlation spectroscopy. *J. Biol. Chem.* 278: 43340–43345.
- Watkins, P. A., J. F. Lu, S. J. Steinberg, S. J. Gould, K. D. Smith *et al.*, 1998 Disruption of the *Saccharomyces cerevisiae* FAT1 gene decreases very long-chain fatty acyl-CoA synthetase activity and elevates intracellular very long-chain fatty acid concentrations. *J. Biol. Chem.* 273: 18210–18219.
- Wibel, F. F., and W. H. Kunau, 1992 The Pas2 protein essential for peroxisome biogenesis is related to ubiquitin-conjugating enzymes. *Nature* 359: 73–76.
- Williams, C., and B. Distel, 2006 Pex13p: Docking or cargo handling protein? *Biochim. Biophys. Acta* 1763: 1585–1591.
- Williams, C., M. van den Berg, and B. Distel, 2005 *Saccharomyces cerevisiae* Pex14p contains two independent Pex5p binding sites, which are both essential for PTS1 protein import. *FEBS Lett.* 579: 3416–3420.
- Williams, C., M. van den Berg, E. Geers, and B. Distel, 2008 Pex10p functions as an E3 ligase for the Ubc4p-dependent ubiquitination of Pex5p. *Biochem. Biophys. Res. Commun.* 374: 620–624.
- Wolinski, H., and S. D. Kohlwein, 2008 Microscopic analysis of lipid droplet metabolism and dynamics in yeast. *Methods Mol. Biol.* 457: 151–163.
- Wolinski, H., K. Natter, and S. D. Kohlwein, 2009a The fidgety yeast: focus on high-resolution live yeast cell microscopy. *Methods Mol. Biol.* 548: 75–99.
- Wolinski, H., U. Petrovic, M. Mattiazzi, J. Petschnigg, B. Heise *et al.*, 2009b Imaging-based live cell yeast screen identifies novel factors involved in peroxisome assembly. *J. Proteome Res.* 8: 20–27.
- Wolinski, H., D. Kolb, S. Hermann, R. I. Koning, and S. D. Kohlwein, 2011 A role for seipin in lipid droplet dynamics and inheritance in yeast. *J. Cell Sci.* 124: 3894–3904.
- Wolinski, H., K. Bredies, and S. D. Kohlwein, 2012 Quantitative imaging of lipid metabolism in yeast: from 4D analysis to high content screens of mutant libraries. *Methods Cell Biol.* 108: 345–365.
- Woods, A., M. R. Munday, J. Scott, X. Yang, M. Carlson *et al.*, 1994 Yeast SNF1 is functionally related to mammalian AMP-activated protein kinase and regulates acetyl-CoA carboxylase in vivo. *J. Biol. Chem.* 269: 19509–19515.
- Yan, M., D. A. Rachubinski, S. Joshi, R. A. Rachubinski, and S. Subramani, 2008 Dysferlin domain-containing proteins, Pex30p and Pex31p, localized to two compartments, control the number and size of oleate-induced peroxisomes in *Pichia pastoris*. *Mol. Biol. Cell* 19: 885–898.
- Yang, H., M. Bard, D. A. Bruner, A. Gleeson, R. J. Deckelbaum *et al.*, 1996 Sterol esterification in yeast: a two-gene process. *Science* 272: 1353–1356.
- Yang, L., Y. Ding, Y. Chen, S. Zhang, C. Huo *et al.*, 2012 The proteomics of lipid droplets: structure, dynamics, and functions of the organelle conserved from bacteria to humans. *J. Lipid Res.*
- Yi, E. C., M. Marelli, H. Lee, S. O. Purvine, R. Aebersold *et al.*, 2002 Approaching complete peroxisome characterization by gas-phase fractionation. *Electrophoresis* 23: 3205–3216.
- Yu, C., N. J. Kennedy, C. C. Chang, and J. A. Rothblatt, 1996 Molecular cloning and characterization of two isoforms of *Saccharomyces cerevisiae* acyl-CoA:sterol acyltransferase. *J. Biol. Chem.* 271: 24157–24163.
- Yu, H., P. Braun, M. A. Yildirim, I. Lemmens, K. Venkatesan *et al.*, 2008 High-quality binary protein interaction map of the yeast interactome network. *Science* 322: 104–110.
- Zaman, S., S. I. Lippman, X. Zhao, and J. R. Broach, 2008 How *Saccharomyces* responds to nutrients. *Annu. Rev. Genet.* 42: 27–81.
- Zanghellini, J., K. Natter, C. Jungreuthmayer, A. Thalhammer, C. F. Kurat *et al.*, 2008 Quantitative modeling of triacylglycerol homeostasis in yeast: metabolic requirement for lipolysis to promote membrane lipid synthesis and cellular growth. *FEBS J.* 275: 5552–5563.
- Zanghellini, J., D. Ruckerbauer, F. Wodlei, H. H. von Grünberg, and C. Jungreuthmayer, 2010a Phospholipid demixing: molecular interpretation of lipid droplet biogenesis, pp. 1–28 in *Advances in Planar Lipid Bilayers and Liposomes* Elsevier: Amsterdam; New York.
- Zanghellini, J., F. Wodlei, and H. H. von Grünberg, 2010b Phospholipid demixing and the birth of a lipid droplet. *J. Theor. Biol.* 264: 952–961.
- Zaremborg, V., and C. R. McMaster, 2002 Differential partitioning of lipids metabolized by separate yeast glycerol-3-phosphate acyltransferases reveals that phospholipase D generation of

- phosphatidic acid mediates sensitivity to choline-containing lysolipids and drugs. *J. Biol. Chem.* 277: 39035–39044.
- Zechner, R., R. Zimmermann, T. O. Eichmann, S. D. Kohlwein, G. Haemmerle *et al.*, 2012 FAT SIGNALS: lipases and lipolysis in lipid metabolism and signaling. *Cell Metab.* 15: 279–291.
- Zehmer, J. K., R. Bartz, P. Liu, and R. G. Anderson, 2008 Identification of a novel N-terminal hydrophobic sequence that targets proteins to lipid droplets. *J. Cell Sci.* 121: 1852–1860.
- Zheng, Z., and J. Zou, 2001 The initial step of the glycerolipid pathway: identification of glycerol 3-phosphate/dihydroxyacetone phosphate dual substrate acyltransferases in *Saccharomyces cerevisiae*. *J. Biol. Chem.* 276: 41710–41716.
- Zimmermann, R., J. G. Strauss, G. Haemmerle, G. Schoiswohl, R. Birner-Gruenberger *et al.*, 2004 Fat mobilization in adipose tissue is promoted by adipose triglyceride lipase. *Science* 306: 1383–1386.
- Zinser, E., C. D. Sperka-Gottlieb, E. V. Fasch, S. D. Kohlwein, F. Paltauf *et al.*, 1991 Phospholipid synthesis and lipid composition of subcellular membranes in the unicellular eukaryote *Saccharomyces cerevisiae*. *J. Bacteriol.* 173: 2026–2034.
- Zinser, E., F. Paltauf, and G. Daum, 1993 Sterol composition of yeast organelle membranes and subcellular distribution of enzymes involved in sterol metabolism. *J. Bacteriol.* 175: 2853–2858.
- Zutphen, T., M. Veenhuis, and I. J. van der Klei, 2008 Pex14 is the sole component of the peroxisomal translocon that is required for pexophagy. *Autophagy* 4: 63–66.
- Zwart, K., M. Veenhuis, and W. Harder, 1979 Biogenesis and breakdown of peroxisomes in the yeast *Hansenula polymorpha* in relation to environmental changes. *Antonie van Leeuwenhoek* 45: 331–332.
- Zweytick, D., K. Athenstaedt, and G. Daum, 2000a Intracellular lipid particles of eukaryotic cells. *Biochim. Biophys. Acta* 1469: 101–120.
- Zweytick, D., E. Leitner, S. D. Kohlwein, C. Yu, J. Rothblatt *et al.*, 2000b Contribution of Are1p and Are2p to steryl ester synthesis in the yeast *Saccharomyces cerevisiae*. *Eur. J. Biochem.* 267: 1075–1082.

Communicating editor: T. N. Davis