

RESEARCH ARTICLE

Open Access

Pandemic influenza A/H1N1 virus infection and *TNF*, *LTA*, *IL1B*, *IL6*, *IL8*, and *CCL* polymorphisms in Mexican population: a case-control study

Guadalupe Morales-García^{1†}, Ramcés Falfán-Valencia², Román Alejandro García-Ramírez^{1†}, Ángel Camarena², Alejandra Ramirez-Venegas², Manuel Castillejos-López², Martha Pérez-Rodríguez¹, César González-Bonilla³, Concepción Grajales-Muñiz³, Víctor Borja-Aburto³ and Juan Manuel Mejía-Aranguré^{1*}

Abstract

Background: Some patients have a greater response to viral infection than do others having a similar level of viral replication. Hypercytokinemia is the principal immunopathological mechanism that contributes to a severer clinical course in cases of influenza A/H1N1. The benefit produced, or damage caused, by these cytokines in severe disease is not known. The genes that code for these molecules are polymorphic and certain alleles have been associated with susceptibility to various diseases. The objective of the present study was to determine whether there was an association between polymorphisms of *TNF*, *LTA*, *IL1B*, *IL6*, *IL8*, and *CCL1* and the infection and severity of the illness caused by the pandemic A/H1N1 in Mexico in 2009.

Methods: Case-control study. The cases were patients confirmed with real time PCR with infection by the A/H1N1 pandemic virus. The controls were patients with infection like to influenza and non-familial healthy contacts of the patients with influenza. Medical history and outcome of the disease was registered. The DNA samples were genotyped for polymorphisms *TNF* rs361525, rs1800629, and rs1800750; *LTA* rs909253; *IL1B* rs16944; *IL6* rs1818879; *IL8* rs4073; and *CCL1* rs2282691. Odds ratio (OR) and the 95% confidence interval (95% CI) were calculated. The logistic regression model was adjusted by age and severity of the illness in cases.

Results: Infection with the pandemic A/H1N1 virus was associated with the following genotypes: *TNF* rs361525 AA, OR = 27.00; 95% CI = 3.07–1248.77; *LTA* rs909253 AG (OR = 4.33, 95% CI = 1.82–10.32); *TNF* rs1800750 AA (OR = 4.33, 95% CI = 1.48–12.64); additionally, *LTA* rs909253 AG showed a limited statistically significant association with mortality ($p = 0.06$, OR = 3.13). Carriers of the *TNF* rs1800629 GA genotype were associated with high levels of blood urea nitrogen ($p = 0.05$); those of the *TNF* rs1800750 AA genotype, with high levels of creatine phosphokinase ($p = 0.05$). The *IL1B* rs16944 AA genotype was associated with an elevated number of leukocytes ($p < 0.001$) and the *IL8* rs4073 AA genotype, with a higher value for P_aO_2 mm Hg.

Conclusion: The polymorphisms of genes involved in the inflammatory process contributed to the severity of the clinical behavior of infection by the pandemic influenza A/H1N1 virus.

Keywords: *TNF*, *IL1B*, *IL8*, *IL6*, *LTA*, *CCL1*, Influenza AH1N1

* Correspondence: juan.mejiaa@imss.gob.mx

†Equal contributors

¹Coordinación de Investigación en Salud, Instituto Mexicano del Seguro Social (IMSS), Torre Academia Nacional de Medicina 4to piso, Av. Cuauhtémoc 330, 06720 México, DF, México

Full list of author information is available at the end of the article

Background

The influenza A/H1N1 virus pandemic of 2009 started in Mexico and then spread worldwide, with an alert level of pandemic phase 6 declared by the World Health Organization (WHO) in June of that year [1]. Although the majority of those infected by the influenza A/H1N1 virus presented mild symptoms that were self-limiting, a subgroup of patients followed an adverse clinical course, thus requiring a greater level of medical attention and more aggressive management [2]. In Mexico the lethal rate was estimated to be 1.2% for cases of influenza-like illness (ILI) and 5% for confirmed cases of influenza A/H1N1 [3]. Some co-morbidities (e.g., immunosuppression, pre-existing pulmonary disease, cardiac disease, diabetes, asthma that requires regular medical attention, smoking, and obesity) have been demonstrated to increase the risk of hospitalization for infection with influenza A/H1N1 [4,5]. The risk is also augmented in the second and third trimesters of pregnancy and when treatment with oseltamivir is prescribed five days after the onset of the illness [6]. In addition, the results of certain laboratory tests (e.g., lactate dehydrogenase (LDH), >600 IU/L; hypoxemia (P_aO_2 , <60 mm Hg); C-reactive protein (CRP), 10 mg/dl; and leukopenia <5000/mm³) have been associated with greater mortality from infection by influenza A/H1N1 virus [7,8].

Through experimental and clinical studies, it has been determined that the most important pathological mechanism in this infection is systemic dysregulation of the inflammatory response, which is correlated with the severity and progression of the illness [9,10]. The secretion of cytokines by infected cells appears to be necessary for the initiation of the immunological response that controls the replication of the virus [11]; in addition, the presence of immunopathological mechanisms, such as hypercytokinemia ("cytokine storm"), generally is considered to contribute to the severest evolution of the infection [11-13]. Elevated levels of pro-inflammatory cytokines and chemokines (e.g., TNF α , IFN γ , IL-1, IL-6, IL-8, IL-9, IL-12 IL-15, and IL-17) have been found, up to ten days after the onset of symptoms, in the plasma of patients with acute respiratory distress syndrome (ARDS) caused by influenza A/H1N1 [9,10,14]. The genes that code for these molecules are polymorphic and certain alleles have been associated with susceptibility to various diseases that cover a wide range of pathologies, from infectious to oncological, including pulmonary and systemic diseases [15-30]. The role that the polymorphisms of the genes encoding these cytokines play in the severity of the disease is not clear.

The extensive polymorphism of these molecules may be associated with the high mortality rate during the 2009 influenza A/H1N1 pandemic in Mexico. Because we think that genetic factors of the host may influence

the nature and intensity of the inflammatory immune response, the objective of this study was to determine whether the polymorphisms of genes that are associated with inflammation may be associated with the development of the infection and with the clinical severity in Mexican mestizo patients with influenza A/H1N1.

Methods

The rapid QuickVue Influenza A+B test (Quidel, San Diego, CA, USA) was used to analyze the nasopharyngeal swab samples obtained from the patients in 94 cases of suspected infection by influenza A/H1N1, following the recommendations for collection and testing of the U.S. Centers for Disease Control and Prevention (CDC) and of the WHO [31,32]. The patients were then separated into two groups: those positive for influenza A/H1N1 (A/H1N1 group) and those negative, as having influenza-like illness (ILI group). A total of 44 patients were positive for the influenza A/H1N1 virus (A/H1N1 group); the remaining 50 patients who had tested negative for the influenza A/H1N1 virus, were diagnosed as having influenza-like illness (ILI).

In addition, a group of 176 asymptomatic healthy contacts (AHC group) voluntarily participated in the study. Those in the AHC group, although not biologically related to any patient, were in personal contact with at least one of the patients during the period of the illness. To ensure that those in the AHC group had been exposed to the virus, their titers of anti-influenza A/H1N1 antibodies were determined. To evaluate the presence of antibody, we use haemagglutination inhibition technique (HAI); contacts exhibited significant titers of specific anti-A/H1N1 antibodies, supporting the fact that they were in contact with the A/H1N1 virus. By serially diluted aliquots of serum samples; those individuals with titers greater than 1:16 were considered positive for A/H1N1 infection/exposure. The two patient groups and the AHC group were Mexican mestizos (age range: 18–85 years). All patients suspected of influenza A/H1N1 infection were treated with anti-viral therapy (oseltamivir) upon admittance at the hospital.

The information collected in this study included demographic, clinical history, laboratory test data, pharmacological treatment, and follow-up. This study was approved by the Institutional Committee of Science and Bioethics of the National Institute of Respiratory Diseases (code B05-10). The study protocol was explained to all participants and signed informed consent was duly obtained from each participant. This information was obtained by means of a clinical form in accord with Official Mexican Standards Mexicana NOM-168-SSA1-1998; topics covered included age; gender; tobacco smoking; body-mass index (BMI; patients with BMI >30 k/m²); and disease morbidity (pulmonary,

hepatic, renal, cardiac, neurological diseases, diabetes mellitus, hypertension, and cancer). The symptoms evaluated were fever, cough, rhinorrhea, dyspnea, nasal congestion, thoracic pain, headache, diarrhea, and vomiting. The begin of anti-viral therapy was evaluated in relation to the days with previous symptomatology. The laboratory parameters included were leukocyte titer; lactate dehydrogenase (LDH); creatine phosphokinase (CPK); blood urea nitrogen (BUN); and arterial gases (with P_aO_2 <60 mm Hg defined as severe disease). Pneumonia was verified by radiological findings. All patients admitted to the intensive care unit (ICU) and those put on assisted mechanical ventilation (AMV) were identified.

Genotyping of allelic variants (single nucleotide polymorphism, SNPs)

The DNA samples were genotyped for polymorphisms *TNF* rs361525, rs1800629, and rs1800750; *LTA* rs909253; *IL1B* rs16944; *IL6* rs1818879; *IL8* rs4073; and *CCL1* rs2282691 using Taqman commercial probes (Applied Biosystems, USA) the primers listed in Table 1. In brief, the procedure for the real time PCR was the following: 15 ng DNA; 15 µL of Taqman universal PCR master mix (Roche NJ, USA) and 6.5 µL of each probes. The conditions for amplification were the following: 94°C (3 min), 61°C (1 min), and 72°C (1 min); followed by 35 cycles of 94°C (1 min), 61°C (1 min), and 72°C (1 min); and a final cycle of 94°C (1 min), 61°C (1 min), and 72°C (5 min). The genetic data for the SNPs were analyzed and are listed in Table 2.

Analysis

When the polymorphisms were evaluated, the ancestral genotype was used for comparison. The odds ratio (OR) and the 95% confidence interval (95% CI) were calculated. In the logistic regression model, the OR was adjusted by age and severity of the illness. The χ^2 test was used to evaluate the differences between the proportions of the groups. The differences among the clinical parameters, continuous variables, and polymorphisms

Table 2 Genetic data of the single nucleotide polymorphisms (SNPs) analyzed

SNP	Gene			
	Symbol	Location	Position	Alleles
rs1800750	<i>TNF</i>	-376	Promoter	G/A
rs1800629	"	-308	Promoter	G/A
rs361525	"	-238	Promoter	G/A
rs909253	<i>LTA</i>	+252	Intronic	C/T
rs16944	<i>IL1B</i>	-511	Promoter	G/A
rs1818879	<i>IL6</i>	5845	3'UTR	A/G
rs4073	<i>IL8</i>	-251	Promoter	A/T
rs2282691	<i>CCL1</i>	29712422	Intronic	A/T

were evaluated by using the Mann-Whitney *U* test. Software packages SPSS 19 (IBM, Chicago, IL) and Epi-Info 6.04b were used (Atlanta, CDC).

Results

The patients were separated into two groups depending whether they were positive or negative for influenza A/H1N1, thereby forming the groups, influenza A/H1N1 and ILI, respectively. In both the A/H1N1 and ILI groups, the majority was male (68.0% and 58.0%, respectively), whereas in the AHC group, 61.93% were female. The mean age of the A/H1N1 and ILI groups was <45 years (68.18% and 60.0%, respectively), compared to that of the AHC group, 56.82% of who were 45-64 years of age.

The data concerning the demographics, co-morbidities, and symptomatology for both groups of patients are presented in Table 3. Interestingly, the mortality was higher for patients with infection by influenza A/H1N1 than for those with ILI (34.09% vs. 4.0%, respectively; $P < 0.001$); similarly, hospitalization in the ICU was more frequent for the influenza A/H1N1 group (54.55% vs. 30.0%, respectively; $p = 0.01$). No statistically significant differences were found for any of the other variables analyzed. Of the influenza A/H1N1 patients, 93.3% had radiographic signs of pulmonary compromise, with a statistically significant difference

Table 1 DNA samples were genotyped by using Taqman commercial probes

Gene (reference SNP)	Probe
<i>TNF</i> (rs1800750)	GAGGCAATAAGACCCCTCGGAATC[A/G]GAGCAGCTGTCAATTGCAGGAGCT
<i>TNF</i> (rs1800629)	GAGGCAATAGGTTTTGAGGGGCATG[A/G]GGACGGGTTCCAGCTCCAGGGTCC
<i>TNF</i> (rs361525)	GGCCAGAAAGACCCCTCGGAATC[A/G]GAGCAGGGAGGATGGGGAGTGTGA
<i>LTA</i> (rs909253)	AAGCCTTAAACCTAGGGCATAACA[C/T]TTGATAATTACCCCTCCAGGGTCCGTT
<i>IL1B</i> (rs16944)	TACCTTGGGTGCTGTCTCTGCCTC[G/A]GGAGCTCTCTGTCAATTGCAGGAGC
<i>IL6</i> (rs1818879)	AGACGAGCTGGCGCAGTGGCTCAC[A/G]CCTATAATCCAGCACTTTGGGAGG
<i>IL8</i> (rs4073)	TTATCTAGAAAATAAAAAGCATAACA[A/T]TTGATAATTACCAAATTGTGGAGC
<i>CCL1</i> (rs2282691)	AAAAAGCCTTAAATACTGACTGGT[A/T]TTGTGAAAGCTACTCCAATTAAGTTT

SNP Single nucleotide polymorphisms.

Table 3 Demographical and clinical characteristics of influenza A/H1N1 patients, influenza-like illness (ILI) patients, and healthy control subjects

Characteristic	A/H1N1 patients	ILI patients	Healthy controls	p
	(Total: 44) n (%)	(Total: 50) n (%)	(Total: 176) n (%)	
Sex				
Male	30 (68.18)	29 (58.0)	67 (38.07)	
Female	14 (31.82)	21 (42.0)	109 (61.93)	
Age				
<45	30 (68.18)	30 (60.0)	54 (30.68)	
45-64	11 (25.00)	13 (26.0)	100 (56.82)	
≥65	3 (6.82)	7 (14.0)	22 (12.50)	
BMI ≥30	18 (40.91)	21 (42.0)	46 (26.42)	
Mortality	15 (34.09)	2 (4.0)		<0.001*
P_aO₂ <60 mm Hg (severe)	19 (43.18)	24 (48.0)		
ICU	24 (54.55)	15 (30.0)		0.01*
Co-morbidities				
Neurological disease	25 (56.82)	22 (44.0)		
Asthma	2 (4.55)	6 (12.0)		
Cancer	1 (2.27)	2 (4.0)		
Hypertension	8 (18.18)	7 (14.0)		
Smoking	21 (47.73)	29 (58.0)		
Symptomatology				
Fever (>38°C)	33 (75.00)	42 (84.0)		
Cough	23 (52.27)	31 (62.0)		
Nasal Congestion	5 (11.36)	2 (4.0)		
Rhinorrhea	12 (27.27)	20 (40.0)		
Dyspnea	33 (75.00)	35 (70.0)		

*When comparing A/H1N1 patients vs. ILI patients. BMI: Body mass index; P_aO₂: partial pressure of oxygen in arterial blood; ICU: Intensive Care Unit. Results were considered statistically significant when P was <0.05.

between the two groups of patients (p = 0.043; data not shown).

Analysis of genetic association

Genotyping was carried out for eight SNPs from six genes, the protein products of which have been associated with inflammatory processes. The genetic information of the polymorphisms that were evaluated is presented in Table 4. An appreciation of the genetic contribution to the risk of infection by influenza A/H1N1 was obtained by evaluating genotype and alleles of both patient groups and comparing the frequencies of the genotypes and alleles with those of the AHC group.

For the A/H1N1 and AHC groups, of the eight SNPs used, 24 genotype products were generated; in the ILI group, only 23 genotypes were determined, as the genotype AA does not exist for the rs361525 of the gene *TNF*. For the A/H1N1 group, five genotypes associated with risk were identified (p <0.05; OR >2.0). Of particular interest was the finding of homozygous A genotype

in SNPs rs361525 and rs1800750 in *TNF*, both with values of OR >5.0. In addition, the genotypes rs2282691 AA, rs4073 AT, and rs909253 CT (*CCL1*, *IL8*, and *LTA*, respectively) demonstrated statistically significant association with risk. For the ILI group, three of the five associations previously reported for the A/H1N1 group were found. It is of note that these associations, which coincided in both patient groups, showed statistically significant data (p values and OR) that were very similar. On the other hand, our findings demonstrated the existence of three genotypes with association to protection (p <0.05; OR ≤1.0): *IL1B* rs16944 AA, *LTA* rs909253, and *TNF* rs1800750 GG, in both patient groups, as compared to the AHC group (Table 4).

In the analysis of alleles, two signs that showed statistically significant association with risk were found: allele A of rs2282691 of *CCL1* was shown to be increased in both patient groups (A/H1N1, p <0.05; ILI, p <0.01), when the allelic frequency (AF) of each patient group was compared with that of the AHC group (OR = 2.15

Table 4 Genetic frequency of the genotypes of eight SNPs in six genes and their association with infection by influenza A/H1N1 virus

Gene and genotype	Genetic frequency								
	A/H1N1 group	AHC group	p	OR	95% CI	ILI group	p	OR	95% CI
<i>CCL1</i> rs2282691									
AA	0.643	0.402	0.0085	2.67	1.26-5.81	0.630	0.0096	2.53	1.23-5.30
TA	0.286	0.456				0.304			
TT	0.071	0.142				0.065			
<i>IL1B</i> rs16944									
GA	0.535	0.449				0.583			
GG	0.442	0.364				0.375			
AA	0.023	0.188	0.015	0.10	0.00-0.66	0.042	0.024	0.19	0.02-0.79
<i>IL8</i> rs4073									
AT	0.659	0.445	0.0231	2.40	1.12-5.32	0.500			
AA	0.244	0.396				0.386			
TT	0.098	0.159				0.114			
<i>IL6</i> rs1818879									
GG	0.439	0.268				0.381			
AG	0.390	0.530				0.452			
AA	0.171	0.202				0.167			
<i>LTA</i> rs909253									
CT	0.535	0.244	0.0004	3.56	1.68-7.54	0.540	0.00014	3.63	1.79-7.38
TT	0.302	0.488	0.0432	0.45	0.20-0.97	0.320	0.0517*	0.49	0.24-1.00
CC	0.163	0.267				0.140			
<i>TNF</i>									
rs361525									
GG	0.818	0.884				0.894			
AA	0.136	0.006	0.00031	27.0	3.07-1248.77	0.000			
GA	0.045	0.110				0.106			
rs1800629									
GG	0.932	0.946				0.911			
GA	0.068	0.054				0.089			
AA	0.000	0.000							
rs1800750									
GG	0.789	0.949	0.0035	0.20	0.06-0.66	0.830	0.0116	0.26	0.08-0.84
AA	0.158	0.028	0.005	6.41	1.51-27.92	0.128	0.0128	5.00	1.20-21.61
GA	0.053	0.023				0.043			

A/H1N1 group: patient infected with influenza A/H1N1 virus; AHC group: asymptomatic, healthy contacts; ILI group: patients with influenza-like illness; OR: odds ratio; CI: confidence interval. Results were considered statistically significant when p was <0.05.

and 2.11, respectively). Also, rs1800750 allele A, one of the three polymorphisms evaluated for *TNF*, showed a similar behavior, that is, an association with risk in both patient groups (A/H1N1, p <0.01; ILI, p <0.01). No other association was found for the remaining six SNPs analyzed (Table 5).

Association of genotypes with clinical variables

Mortality

The genotype AG of rs909253 in *LTA* showed a tendency to be associated with mortality (OR = 3.13), with p = 0.06

just above the limit of statistical significance. Table 6 presents the data for the different genotypes evaluated with respect to mortality. The data corresponding to the allelic OR were carried out for both alleles; however, only those carried out for the ancestral allele are shown. No data for the remaining alleles or genotypes studied were statistically significant. The logistic regression analysis, after adjusting for levels of P_aO₂ and for admission to the ICU, showed no statistically significant association between the genotypes and mortality.

Table 5 Risk of influenza A/H1N1 infection in relation to the different polymorphisms studied

Gen/allele	Risk of influenza A/H1N1 infection		
	p	OR (95% CI)	OR allelic (95% CI)
<i>CCL1</i>			2.31 (1.25-4.31)
rs2282691			
AT	0.85	1.14 (0.29-4.38)	
AA	0.12	2.75 (0.78-9.72)	
<i>IL1B</i>			1.62 (0.92-2.88)
rs16944			
GG	0.04	8.57 (1.11-66.46)	
AG	0.06	7.33 (0.95-56.41)	
<i>IL8</i>			1.45 (0.80-2.63)
rs4073			
TT	0.96	0.97 (0.29-3.28)	
AT	0.17	1.73 (0.79-3.74)	
<i>IL6</i>			1.01 (0.59-1.71)
r1818879			
AG	0.2	0.61 (0.29-1.30)	
AA	0.45	0.69 (0.26-1.82)	
<i>LTA</i>			0.72 (0.42-1.26)
rs909253			
GG	0.66	1.25 (0.46-3.41)	
AG	<0.001	3.3 (1.51-7.20)	
<i>TNF</i>			1.25 (0.79-1.99)
rs1800750			
AG	0.27	2.54 (0.49-13.24)	
AA	0.02	3.52 (1.24-10.03)	
rs1800629			1.10 (0.30-3.91)
AG	0.88	1.1 (0.30-4.02)	
rs361525			3.34 (1.56-7.08)
AG	0.37	0.5 (0.11-2.24)	
AA	<0.001	34.8 (4.06-297.87)	

OR: odds ratio; CI: confidence interval; allelic OR: comparison with ancestral gene. Results were considered statistically significant when P was <0.05.

Polymorphisms and laboratory tests

No changes in laboratory parameters were found with respect to the genotypes of the SNPs in *TNF* rs361525, *LTA* rs909253, *CCL1* rs2282691, or *IL6* rs1818879 (Additional file 1: Table S1). For the homozygous genotype AA of rs16944 in *IL1B*, an elevated number of leukocytes was found (mean: $44.9 \times 10^3/\text{mm}^3$), whereas for the heterozygous genotype AG and the homozygous genotype GG, the mean titer values were $8.62 \times 10^3/\text{mm}^3$ and $6.96 \times 10^3/\text{mm}^3$, respectively ($p < 0.001$). The genotype AG in rs1800629 in *TNF* was associated with elevated levels of BUN (33mg/dl), whereas for the genotype GG, the BUN levels were lower ($p = 0.05$) (Additional file 1: Table S1). The genotype AA of rs1800750 in *TNF* showed high levels of

Table 6 Risk of death from influenza A/H1N1 in relation to the different polymorphisms studied

Gen/allele	p	OR (CI 95%)	OR allelic (CI 95%)
<i>CCL1</i> rs2282691			1.63 (0.63-4.40)
AT	0.40	0.4 (0.07-2.79)	
AA	0.90	1.1 (0.22-5.31)	
<i>IL1B</i> rs16944			1.55 (0.56-4.50)
GG	0.70	1.6 (0.17-14.40)	
AG	0.30	2.8 (0.35-23.02)	
<i>IL8</i> rs4073			1.41 (0.53-3.78)
TT	0.70	0.7 (0.07-6.11)	
AT	0.40	1.7 (0.51-5.76)	
<i>IL6</i> rs1818879			0.96 (0.40-2.28)
AG	0.30	0.5 (0.15-1.70)	
AA	0.50	0.5 (0.10-2.76)	
<i>LTA</i> rs909253			0.46 (0.16-1.27)
GG	0.50	0.5 (0.05-4.27)	
AG	0.06	3.1 (0.93-10.53)	
<i>TNF</i>			
rs1800750			1.21 (0.55-2.61)
AG	0.30	3.5 (0.39-31.76)	
AA	0.20	2.9 (0.58-14.51)	
rs1800629			2.39 (0.51-11.03)
AG	0.30	2.5 (0.51-12.26)	
rs361525			0.95 (0.21-4.16)
AG	1.00		
AA	0.40	2.8 (0.31-24.74)	

OR: odds ratio; CI: confidence intervals; allelic OR: comparison with ancestral gene. Results were considered statistically significant when P was <0.05.

CPK (mean: 2246.25 IU/L), whereas the genotypes GG and AG had mean values of 366.32 IU/L and 450.00 IU/L, respectively ($p=0.05$).

Discussion

Although it has been reported that some of the genes involved in inflammation are associated with pulmonary and infectious diseases [16,17,21,25-28], to date, no direct association between these polymorphisms and infection by influenza A/H1N1 virus has been reported. In the present study, the logistic regression analysis of cases and controls showed that *TNF* rs361525 (AA), *rs1800750* (AA), and *LTA* rs909253 (AG) were associated with high risk of infection by pandemic influenza A/H1N1. Although mortality of the A/H1N1 patients was greater than that of the ILI patients ($p < 0.001$), only genotype AG of *LTA* rs909253 demonstrated an association with mortality ($p = 0.06$) just missing being statistically significant. This finding may indicate that being a carrier of genotype AG *LTA* at rs909253 entails a poorer prognosis for this illness.

Some biomarkers, such as CRP, LDH, leukopenia, and hypoxemia [7,8] have been considered as predictors for

severe illness by influenza A/H1N1 virus. In this study, we found that elevated values of laboratory test parameters (BUN, CPK, and leukocyte titer) were associated with *TNF* (rs1800629 AG, rs1800750 AA), and *IL1B* (genotype AA). These results may indicate that being a carrier of these polymorphisms in particular may be associated with severer organic damage by infection by influenza A/H1N1 virus. However, in the case of *IL8*, the homozygous AA appeared to offer a certain degree of protection against severe illness; this may be explained, in part, by the association of a higher concentration of oxygen in the blood ($P_aO_2 >60$ mm Hg) with this genotype.

Although susceptibility for presenting an adverse clinical course (i.e., sepsis, septic shock, multiples organ failure, or death) varies due to different degrees of inflammatory response [30], genetic factors of the host may influence the nature and intensity of this response. The present study is the first to demonstrate that the polymorphisms in genes related to the inflammatory response may, in some manner, be influencing the risk of infection by influenza A/H1N1 virus and of death from this illness. One possible mechanism may be the formation of disequilibria in the bond between alleles, creating haplotypes that differentially affect the expression and activity of cytokines and chemokines, thereby resulting in a severer clinical course for the infection.

To avoid variability in our results, we studied a homogeneous population of Mexican mestizos. Although the sampling size was a limitation in our study, we were able to demonstrate statistically significant differences in the distribution of genotypes in terms of infection, mortality, and biomarkers between cases and controls, thus suggesting a strong association with the illness. However, infection by influenza A/H1N1 virus is a very complex illness that involves not only the association of environmental factors and the genetic make-up and biology of the individual per se, but also the presence of comorbidities that may contribute to a greater severity of the illness [4,5].

We had limitations. For example, contacts exhibited significant titers of specific anti-A/H1N1 antibodies, supporting the fact that they were in contact with the A/H1N1 virus. Additionally is necessary clear that is not a patients group, we just include unrelated contacts in this study (e.g. family in law persons, home workers, etc.). They were in close contact with patients when the latter exhibited acute respiratory illness. None of these household contacts developed respiratory illness. However, the presence of antibody would not confirm infection with A/H1N1 virus because there is the likelihood of cross-reactivity [33]. But, when a person with positivity of antibodies that had contact with an A/H1N1 virus infected patient, the infection cannot be excluded

[34]. The gold standard for identifying the infection to A/H1N1 virus is real time PCR test; however to identify the presence of the virus in symptomatic patients is necessary that the patients be assessed during the first days from the beginning of the disease, the probability of identifying the virus by molecular test in asymptomatic patients is very low and not practical [35].

With the high mortality rate from the 2009 influenza A/H1N1 pandemic in Mexico [31], the approach used in this work could acquire great importance, as the study of polymorphisms may be useful in predicting the conduct that the infection would follow during future outbreaks of influenza A/H1N1 in Mexico.

Conclusions

The *TNF* polymorphisms studied were associated with risk of infection by influenza A/H1N1 virus during the pandemic in Mexico in 2009. These genetic variants may contribute to the severest clinical manifestations in Mexican mestizos.

Additional file

Additional file 1: Table S1. Comparison of laboratory findings in relation to the polymorphisms studied in A/H1N1 patients and controls.

Abbreviations

AHC: Asymptomatic healthy contacts; ARDS: Acute respiratory distress syndrome; BMI: Body mass index; BUN: Blood urea nitrogen; CCL1: Chemokine (C-C motif) ligand 1; CDC, U.S: Centers for Disease Control and Prevention; CI: 95% Confidence interval; CPK: Creatine phosphokinase; CRP: C-reactive protein; GLR: Global lethal rate; ICU: Intensive Care Unit; IFN: Interferon; IL: Interleukin; ILI: Influenza-like illness; LDH: Lactate dehydrogenase; LTA: Lymphotoxin α ; OR: Odds ratio; P_aO_2 : Partial pressure of oxygen in arterial blood; SNP: Single nucleotide polymorphism; TNF: Tumor necrosis factor; WHO: World Health Organization.

Competing interests

The authors declare that they have no conflict of interests.

Authors' contributions

GMG, RFV, RAGR, AC, and MCL carried out the study and participated in its design and co-ordination, in the molecular assays, and in the preparation of the manuscript. ARV participated in the design and co-ordination of the study, compiled data, and helped prepare the manuscript. MPR, CGB, CGM, VBA, and JMMA participated in the design and co-ordination of the study, carried out the study and the statistical analysis, and helped prepare the manuscript. All authors read and approved the final version of the manuscript.

Acknowledgments

This study was funded by the Consejo Nacional de Ciencia y Tecnología (CONACyT), México and by CONACyT-Básica, SALUD2009-C02-127089. and CONACyT-FOSIS SALUD2009-C02-126699. The authors thank Veronica Yakoleff for translating the original Spanish manuscript and for helpful comments. The authors thank the Coordinación de Investigación en Salud of the IMSS for covering the cost of the translation and publication.

Author details

¹Coordinación de Investigación en Salud, Instituto Mexicano del Seguro Social (IMSS), Torre Academia Nacional de Medicina 4to piso, Av. Cuauhtémoc 330, 06720 México, DF, México. ²Instituto Nacional de Enfermedades Respiratorias Ismael Cosío Villegas, Tlalpan 4502, 14080,

México, DF, México. ³Coordinación de Vigilancia Epidemiológica y Apoyo en Contingencias, IMSS, Mier y Pesado 120, 03100 México, DF, México.

Received: 24 February 2012 Accepted: 6 November 2012
Published: 13 November 2012

References

- Bautista E, Chotpitayasonndh T, Gao Z, Harper SA, Shaw M, Uyeki TM, Zaki SR, Hayden FG, Hui DS, Kettner JD, Kumar A, Lim M, Shindo N, Penn C, Nicholson KG: **Clinical aspects of pandemic 2009 influenza A (H1N1) virus infection.** *N Engl J Med* 2010, **362**:1708–1719.
- Singanayagam A, Singanayagam A, Wood V, Chalmers JD: **Factors associated with severe illness in pandemic 2009 influenza A (H1N1) infection: implications for triage in primary and secondary care.** *J Infect* 2011, **63**:243–251.
- Chowell G, Echeverría-Zuno S, Viboud C, Simonsen L, Tamerius J, Miller MA, Borja-Aburto VH: **Characterizing the epidemiology of the 2009 influenza A/H1N1 pandemic in Mexico.** *PLoS Med* 2011, **8**:e1000436.
- Ward KA, Spokes PJ, McNulty JM: **Case-control study of risk factors for hospitalization caused by pandemic (H1N1) 2009.** *Emerg Infect Dis* 2011, **17**:1409–1416.
- Fezeu L, Julia C, Henegar A, Bitu J, Hu FB, Grobbee DE, Kengne AP, Hercberg S, Czernichow S: **Obesity is associated with higher risk of intensive care unit admission and death in influenza A (H1N1) patients: a systematic review and meta-analysis.** *Obes Rev* 2011, **12**:653–659.
- Yu H, Feng Z, Uyeki TM, Liao Q, Zhou L, Feng L, Ye M, Xiang N, Huai Y, Yuan Y, Jiang H, Zheng Y, Gargiullo P, Peng Z, Feng Y, Zheng J, Xu C, Zhang Y, Shu Y, Gao Z, Yang W, Wang Y: **Risk factors for severe illness with 2009 pandemic influenza A (H1N1) virus infection in China.** *Clin Infect Dis* 2011, **52**:457–465.
- Reyes S, Montulla B, Martínez R, Córdoba J, Molina JM, Martí V, Martínez A, Ramírez P, Menéndez R: **Risk factors of H1N1 etiology in pneumonia and impact on mortality.** *Respir Med* 2011, **105**:1404–1411.
- Wen Y, Deng BC, Zhou Y, Wang Y, Cui W, Wang W, Liu P: **Immunological features in patients with pneumonitis due to influenza A H1N1 infection.** *J Investig Allergol Clin Immunol* 2011, **21**:44–50.
- To KK, Hung IF, Li IW, Lee KL, Koo CK, Yan WW, Liu R, Ho KY, Chu KH, Watt CL, Luk WK, Lai KY, Chow FL, Mok T, Buckley T, Chan JF, Wong SS, Zheng B, Chen H, Lau CC, Tse H, Cheng VC, Chan KH, Yuen KY: **Delayed clearance of viral load and marked cytokine activation in severe cases of pandemic H1N1 2009 influenza virus infection.** *Clin Infect Dis* 2010, **50**:850–859.
- Bermejo-Martin JF, de Lejarazu RO, Pumarola T, Rello J, Almansa R, Ramirez P, Martin-Loeches I, Varillas D, Gallegos MC, Serón C, Micheloud D, Gomez JM, Tenorio-Abreu A, Ramos MJ, Molina ML, Huidobro S, Sanchez E, Gordón M, Fernández V, Del Castillo A, Marcos MA, Villanueva B, López CJ, Rodríguez-Domínguez M, Galan JC, Cantón R, Lieter A, Rojo S, Eiros JM, Hinojosa C: **Th1 and Th17 hypercytokinemia as early host response signature in severe pandemic influenza.** *Crit Care* 2009, **13**:R201. doi:10.1186/cc8208.
- Michaelis M, Doerr HW, Cinatl J Jr: **Of chickens and men: avian influenza in humans.** *Curr Mol Med* 2009, **9**:131–151.
- de Jong MD, Simmons CP, Thanh TT, Hien VM, Smith GJ, Chau TN, Hoang DM, Chau NV, Khanh TH, Dong VC, Qui PT, Cam BV, Ha DQ, Guan Y, Peiris JS, Chinh NT, Hien TT, Farrar J: **Fatal outcome of human influenza A (H5N1) is associated with high viral load of human influenza A.** *Nat Med* 2006, **12**:1203–1207.
- Mainers TR, Szretter KJ, Perrone L, Belsler JA, Bright RA, Zeng H, Tumpey TM, Katz JM: **Pathogenesis of emerging avian influenza viruses in mammals and the host innate immune response.** *Immunol Rev* 2008, **225**:68–84.
- Hagau N, Slavcovici A, Gonganau DN, Oltean S, Dirzu DS, Brezozski ES, Maxim M, Ciuce C, Mlesnita M, Gavrus L, Laslo C, Hagau R, Petrescu M, Studnicska DM: **Clinical aspects and cytokine response in severe H1N1 influenza A virus infection.** *Crit Care* 2010, **14**:R203. doi:10.1186/cc9324.
- Wang B, Wang J, Zheng Y, Zhou S, Zhen J, Wang F, Ma X, Zeng Z, HBV Study Consortium: **TNF-alpha-238 and -308 polymorphisms with different outcomes of persistent hepatitis B virus infection in china.** *Pathology* 2010, **42**:674–680.
- Alassani B, Ogwaro KM, Shrestha S, Tang J, Breen EC, Wong HL, Jacobson LP, Rabkin CS, Ambinder RF, Martinez-Maza O, Kaslow RA: **The major histocompatibility complex conserved extended haplotype 8.1 in AIDS-related non-Hodgkin lymphoma.** *J Acquir Immune Defic Syndr* 2009, **52**:170–179.
- Gaudet MM, Egan KM, Lissowska J, Newcomb PA, Brinton LA, Titus-Ernstoff L, Yeager M, Chanock S, Welch R, Peplonska B, Trentham-Dietz A, Garcia-Closas M: **Genetic variation in tumor factor and lymphotoxin-alpha (TNA-LTA) and breast cancer risk.** *Human Genet* 2007, **121**:483–490.
- Partida-Rodríguez O, Torres J, Flores-Luna L, Camorlinga M, Nieves-Ramirez M, Lazcano E, Perez-Rodríguez M: **Polymorphism in TNF and HSP-70 show a significant association with gastric cancer and duodenal ulcer.** *Int J Cancer* 2010, **126**:1861–1868.
- Ned RM, Yesupriya A, Imperatore G, Smelser D, Moonesinghe R, Chang MH, Dowling NF: **nflammation gene variants and susceptibility to albuminuria in U.S. population: analysis in the Third National Health and Nutrition Examination Survey (NHANES III), 1991–94.** *BMC Med Genet* 2010, **11**:155.
- Martinez-Carrillo DN, Garza-Gonzalez E, Betancout-Linares R, Mónico-Manzano T, Antúnez-Rivera C, Roman-Roman A, Flores-Alfaro E, Illades-Aguar B, Fernández-Tilapa G: **Association of ILB -511C/-31T haplotypes and *Helicobacter Pylori* VacA genotypes with gastric ulcer and chronic gastritis.** *BMC Gastroenterol* 2010, **10**:126.
- Vazarova B, Fernández-Real JM, Knowler WC, Gallart L, Hanson RL, Guber JD, Ricart W, Vendrell J, Richart C, Tataranni PA, Wolford JK: **The interleukin-6 (-174) G/C promoter polymorphism is associated with type-2 diabetes mellitus in Native Americans and Caucasians.** *Hum Genet* 2003, **112**:409–413.
- Qi L, Zhang C, van Dam RM, Hu FB: **Interleukin-6 genetic variability and adiposity: association in two prospective cohorts and systematic review in 26,944 individuals.** *J Clin Endocrinol Metab* 2007, **92**:3618–3625.
- Cheng CH, Lee YS, Tsau YK, Lin TY: **Genetic polymorphisms and susceptibility to parenchymal renal infection among pediatric patients.** *Pediatr Infect Dis J* 2011, **30**:309–314.
- Heinzmann A, Ahlert I, Kurtz T, Berner R, Deichmann KA: **Association study suggests opposite effects of polymorphism within IL8 on bronchial asthma and respiratory syncytial virus bronchiolitis.** *J Allergy Clin Immunol* 2004, **114**:671–676.
- Hacking D, Knight JC, Rockett K, Brown H, Frampton J, Kwiatkowski DP, Hull J, Udalova IA: **Increased *in vivo* transcription of an IL-8 haplotype associated with respiratory syncytial virus disease-susceptibility.** *Genes Immun* 2004, **5**:274–282.
- Hull J, Thomson A, Kwiatkowski D: **Association of respiratory syncytial virus bronchiolitis with the interleukin 8 gene region in UK families.** *Thorax* 2000, **55**:1023–1027.
- Takabatake N, Shibata Y, Abe S, Wada T, Machiva J, Igarashi A, Tokairin Y, Ji G, Sato H, Sata M, Takeishi Y, Emi M, Muramatsu M, Kubota I: **A single nucleotide polymorphism in the CCL1 gene predicts acute exacerbations in chronic obstructive pulmonary disease.** *Am J Respir Crit Care Med* 2006, **174**:875–885.
- Camarena A, Juárez AM, Estrada A, Carrillo G, Falfán R, Zuñiga J, Navarro C, Granados J, Selman M: **Major histocompatibility complex and tumor necrosis factor-α polymorphisms in pigeon breeder's disease.** *Am J Respir Crit Care Med* 2001, **163**:1528–1533.
- Nieves ME, Partida O, Alegre P, Tapia MC, Pérez M: **Characterization of single-nucleotide polymorphisms in the tumor necrosis factor α promoter region and lymphotoxin α in squamous intraepithelial lesion, precursors of cervical cancer.** *Transl Oncol* 2011, **4**:336–344.
- Paskullin DD, Fallavena P, Paludo F, Borges T, Picanco J, Dias F, Alho C: **TNF -308G > a promoter polymorphism (rs1800626) and outcome from critical illness.** *Braz J Infect Dis* 2011, **15**:231–238.
- Echeverría-Zuno S, Mejía-Aranguré JM, Mar-Obeso AJ, Grajales-Muñiz C, Robles-Pérez E, González-León M, Ortega-Alvarez MC, Gonzalez-Bonilla C, Rascón-Pacheco RA, Borja-Aburto VH: **Infection and death from influenza A H1N1 virus in Mexico: a retrospective analysis.** *Lancet* 2009, **374**:2072–2079.
- WHO information for laboratory diagnosis of pandemic (H1N1) 2009 virus in humans—update; 2009. http://www.who.int/csr/resources/publications/swineflu/WHO_Diagnostic_RecommendationsH1N1_20090521.pdf (accessed May 26, 2009).
- Hancock K, Veguilla V, Lu X, Zhong W, Butler EN, Sun H, Liu F, Dong L, DeVos JR, Gargiullo PM, Brammer TL, Cox NJ, Tumpey TM, Katz JM: **Cross-**

reactive antibody responses to the 2009 pandemic H1N1 influenza virus. *N Engl J Med* 2009, **361**:1945–1952.

34. Baguelin M, Hoschler K, Stanford E, Waight P, Hardelid P, Andrews N, Miller E: **Age-specific incidence of A/H1N1 2009 influenza infection in England from sequential antibody prevalence data using likelihood-based estimation.** *PLoS One* 2011, **6**:e17074.
35. Papenburg J, Baz M, Hamelin MÈ, Rhéaume C, Carbonneau J, Ouakki M, Rouleau I, Hardy I, Skowronski D, Roger M, Charest H, De Serres G, Boivin G: **Household transmission of the 2009 pandemic A/H1N1 influenza virus: elevated laboratory-confirmed secondary attack rates and evidence of asymptomatic infections.** *Clin Infect Dis* 2010, **51**:1033–1041.

doi:10.1186/1471-2334-12-299

Cite this article as: Morales-García *et al.*: Pandemic influenza A/H1N1 virus infection and *TNF*, *LTA*, *IL1B*, *IL6*, *IL8*, and *CCL* polymorphisms in Mexican population: a case-control study. *BMC Infectious Diseases* 2012 **12**:299.

**Submit your next manuscript to BioMed Central
and take full advantage of:**

- Convenient online submission
- Thorough peer review
- No space constraints or color figure charges
- Immediate publication on acceptance
- Inclusion in PubMed, CAS, Scopus and Google Scholar
- Research which is freely available for redistribution

Submit your manuscript at
www.biomedcentral.com/submit

