

The optimal treatment of type II and III odontoid fractures in the elderly: a systematic review

Jeroen G. J. Huybregts · Wilco C. H. Jacobs ·
Carmen L. A. M. Vleggeert-Lankamp

Received: 22 May 2012 / Accepted: 16 July 2012 / Published online: 2 September 2012
© Springer-Verlag 2012

Abstract

Purpose Odontoid fractures are the most common cervical spine fractures in the elderly. As the population ages, their incidence is expected to increase progressively. The optimal treatment of this condition is still the subject of controversy. The objective of this review is to summarize and compare the outcome of surgical and conservative interventions in the elderly (≥ 65 years).

Methods A comprehensive search was conducted in nine databases of medical literature, supplemented by reference and citation tracking. Clinical status was considered the primary outcome. Fracture union and stability rates were considered secondary outcomes.

Results A total of nineteen studies met the inclusion criteria. All studies were performed retrospectively and were of limited quality. There was insufficient data, especially from direct comparisons, to determine the difference in clinical outcome between surgical and conservative interventions. Osseous union was achieved in 66–85 % of surgically treated patients and in 28–44 % of conservatively treated patients. Fracture stability was achieved in 82–97 % of surgically patients and in 53–79 % of conservatively treated patients.

Conclusions There was insufficient data to determine a potential difference in clinical outcome between different treatment groups. Surgically treated patients showed higher osseous union rates compared to conservatively treated

patients, possibly because of different selection mechanisms. The majority of patients appears to achieve fracture stability regardless of the applied treatment. A prospective trial with appropriate sample size is needed to identify the optimal treatment of odontoid fractures in the elderly and predictors for the success of either one of the available treatments.

Keywords Odontoid fractures · Elderly · Surgical treatment · Conservative treatment · Systematic review

Introduction

Odontoid fractures account for 9–18 % of all cervical spine fractures and are most frequently caused by either hyperextension or hyperflexion [5, 14, 20, 22, 34, 37]. In the elderly, odontoid fractures are the most common cervical spine fractures [17, 18, 20, 22, 29, 30, 35]. Moreover, as the population ages, these fractures will become increasingly relevant to clinical practice [37]. The optimal treatment of odontoid fractures in the elderly is, however, still subject to controversy. This age group typically suffers from an increased risk of operation complications when treated surgically as well as from an increased risk of non-union and prolonged treatment duration when treated conservatively.

The treatment for patients with fractures of the odontoid process should be based on fracture pattern (such as defined by Anderson and D'Alonzo [1]), patient age, neurological deficits and the patient's medical condition [20]. Many factors have to be taken into account to find the right balance between fracture healing and treatment complications. Based on these factors, the decision for either surgical or conservative treatment is made.

J. G. J. Huybregts (✉) · W. C. H. Jacobs ·
C. L. A. M. Vleggeert-Lankamp
Department of Neurosurgery, Leiden University Medical Center,
Albinusdreef 2, PO Box 9600, 2300 RC Leiden,
The Netherlands
e-mail: j.g.j.huybregts@lumc.nl

Surgical intervention is in most cases performed by anterior odontoid screw fixation or posterior atlantoaxial arthrodesis. The general presumption is that a surgical intervention leads to a stable cervical spine. However, the condition of the patient may deteriorate by undergoing (major) cervical spine surgery. Especially in the very old patient (≥ 80 years of age), a surgical intervention leads to significant risks for the patient.

An alternative to surgical stabilization is conservative treatment, involving rigid or non-rigid immobilization. Such treatment is often proposed to avoid the complications that may accompany spine surgery. However, this may result in non-union and prolonged fracture instability, requiring secondary surgery. This unnecessarily lengthens treatment duration and, worse, can cause significant deterioration of the cervical spine anatomy.

The objective of this review is to summarize and compare the outcome of surgical and conservative treatments for type II and III odontoid fractures in the elderly (≥ 65 years), focusing primarily on clinical outcome and secondarily on fracture union and stability rates.

Methods

Search methods for identification of studies

A comprehensive and systematic search was conducted in nine databases of medical literature: MEDLINE, EMBASE, Cochrane Central Register of Controlled Trials, Web of Science, CINAHL, Academic Search Premier, ScienceDirect, PEDro and RCT-registers up to April 2012. The MEDLINE search strategy is given in Table 1. The search strategy was adapted for the other databases. No restriction was made with regard to language or date. ‘Os odontoideum’ was included in the search, as this term is sometimes incorrectly used to describe odontoid fractures. References from the included studies were also screened in order to identify additional primary studies not previously identified. Two review authors (JH, WJ) working independently from one another examined titles and abstracts

Table 1 The MEDLINE search strategy

(“Odontoid fractures” OR “Odontoid fracture” OR “Os odontoideum” OR “c2 fracture” OR “c2 fractures” OR “c-2 fracture” OR “c-2 fractures” OR “dens fracture” OR “dens fractures” OR ((Fractures OR fracture OR fracture* OR injury OR injuries OR “Spinal Injuries”[mesh]) AND (odontoid OR odontoid* OR dens)) OR “Odontoid Process/injuries”[mesh]) AND (“Aged”[mesh] OR Aged OR “Aged, 80 and over”[mesh] OR Elderly OR Old* OR Elder* OR Geriatric*) NOT (“child”[mesh] NOT “aged”[mesh])

from the electronic search. Full articles were obtained if necessary. The third review author (CV) was consulted, if consensus was not reached.

Criteria for considering studies for this review

References were selected if they met all of the following inclusion criteria:

- The study design was either a randomised controlled trial, a non-randomised study with (concurrent or historical) control group or a case series
- A minimum of ten subjects was included
- The patients included suffered from acute, isolated type II or III odontoid fractures with or without associated luxation
- The patients included were at least 65 years old or their data could be extracted separately from studies that also involved younger subjects
- The criteria for inclusion were explicit (e.g. age range given, co-morbidity)
- Number of patients deceased or lost to follow-up reported or included in appropriate statistical analysis
- The study evaluated the results of any surgical and/or conservative treatment
- Results were given for each distinct treatment in the study
- The outcome(s) evaluated included at least one of the main clinically relevant outcome measures (e.g. using NDI, Smiley-Webster scale, CSOQ) and/or radiologically assessed union or stability data
- Patients were not treated for odontoid fractures in the past that were unrelated to the reporting study
- Patients did not suffer from systemic co-morbidity (e.g. rheumatoid arthritis)
- The follow-up period was at least 2 weeks
- The paper was published in a peer-reviewed journal

Clinical status was considered the primary outcome. The Neck Disability Index (NDI) was expected to be the most commonly used tool to assess clinical outcome. Therefore, the minimal clinically important difference (MCID/MCIC) for the NDI was predetermined to be 7.5 [6, 8, 38, 39].

Fracture union and stability rates were considered the secondary outcomes.

Assessment of risk of bias for the included studies

Two review authors (JH, CV) working independently of each other conducted the risk-of-bias assessment and data extraction. Risk of bias of the individual studies was assessed with methodology scores based on the type of study concerned: Cochrane form II for RCTs [9], Cochrane

form III for cohort studies [10], Cochrane form IV for patient control studies [11] and a self-designed appraisal form for uncontrolled case series based on three other studies [7, 12, 36]. The criteria for risk-of-bias assessments of case series are given in Table 2. Items were scored as positive if they fulfilled the criterium, negative when bias was likely or marked as inconclusive if there was insufficient information. Differences in the scoring of the risk-of-bias assessment and data extraction were discussed during a consensus meeting. If an item was scored positive, one point was awarded. The number of positively scored items was summated per study.

Data collection and analysis

Data was extracted onto separate, pre-developed forms depending on the type of study concerned. From each study, both demographic/descriptive data (e.g. study population, types of treatment, types of outcomes assessed, sample size, age, gender) and quantitative data regarding primary and secondary outcomes were extracted. With sufficient clinically and statistically homogeneous and comparable reported outcomes, data was planned to be pooled with the aid of Revman 5 for studies with control group and Excel for case series. To identify publication

Table 2 Criteria for risk-of-bias assessment (scored as ‘yes’, ‘no’, ‘inconclusive’ or ‘not applicable’)

	Criteria for ‘yes’
Key criteria	
Clear study objective (B)	Goal of the study mentioned and motivated
Criteria for inclusion explicit (B)	Inclusion criteria mentioned
Fractures appropriately described (B)	Classification system and radiological tools used mentioned
Distinction type II/III appropriate (B)	Classification system and radiological tools used mentioned
Mean age (range) (B)	Mean age and age range reported or computable
Selection bias ruled out	Methods for patient selection and inclusion mentioned
Mean follow-up (range) (R)	Follow-up data reported or computable
Surgical treatment(s) specified (B)	Types of performed surgical interventions described
Conservative treatment(s) specified (B)	Types of performed conservative interventions described
Clear criteria for measuring outcomes (B)	Outcome measures mentioned
Clinically relevant outcomes (e.g. NDI, Smiley-Webster Scale, CSOQ) (B)	Clinical outcome systematically evaluated
When clinical outcome reported: Is pre-treatment neurol status stated (B)	Pre-treatment status reported for comparison to post-treatment status
Results for surgical/conservative treatment separately given (R)	If applicable; results for treatments separately reported
Selective loss-to-follow-up ruled out (R), scored as:	Number of patients lost to follow-up reported including its causes
‘yes’ (2 pt)	
‘no, l.t.f.u. <20 %, but may not be a selective’ (1 pt)	
‘no, l.t.f.u. >20 %’ (0 pt) → exclusion criterium	
‘too little information/not described’ (0 pt)	
Other criteria	
Valid statistical analysis undertaken (R)	Statistical analyses carried out; if impossible: ‘NA’
Number of men and women given (B)	Gender distribution of included patients reported or extractable
Clinical evaluation independent of treating physician (R)	Evaluation carried out by independent party
Radiological evaluation independent and blinded to clinical results (R)	Evaluation carried out by independent party
Independence of investigators stated (R)	Independence specifically stated (no vested interest)
Quantification of outcomes (R), scored as:	Categorized according to the scale used for outcome measures
‘yes, >5 scale-classification (3 pt)	
‘yes, <5 scale-classification (2 pt)	
‘yes, descriptive’ (1 pt)	
‘no’ (0 pt)	
‘too little information’ (0 pt)	

B Concerning baseline data

R Concerning results

bias, funnel plots were planned to be examined. Secondary analysis was to be carried out based on fracture union and stability rates. Subgroup analysis was planned to be carried out based on fracture type.

Results

Search and selection results

The initial search yielded 1,487 unique references; 159 studies were either discussed during the consensus meeting or a full-text version of the article was obtained. Additionally, reference and citation tracking were carried out that yielded no further references. A total of 22 studies was initially identified. Three of these studies were subsequently excluded [28, 31, 32]. The study by Platzer et al. [31] was excluded because the patient cohort in this study was believed to overlap with the patient cohort described in the other study by Platzer that was used for this review [30]. The study by Reinhold et al. [32] was excluded because loss-to-follow-up was 49 %, which was considered to induce too much bias. The study by Müller et al. [28]

was excluded as the outcome was not adequately described and relevant data could not be extracted.

A total of 19 studies (all were case series) was eventually identified for this review, 5 with control groups (Fig. 1). Of these 19 studies, 11 systematically reported clinical outcome and hence were primarily included. The other eight studies that only reported union and stability rates were secondarily included. Only five studies compared outcomes for surgical treatment to conservative treatment in the elderly; in two of these studies, clinical outcome was assessed, four studies reported union data and all five reported data regarding fracture stability. Seventeen studies were published in English, one in French and one in German. The main characteristics of the included studies are given in Table 3. The results of the data extraction are given in Tables 4 and 5.

Risk-of-bias assessment

The results of the risk-of-bias assessment are given in Table 6. All studies were retrospective case series, with their associated limitations (such as missing data and variability in outcome assessment). Most baseline demographic data

Fig. 1 Flow chart for inclusion of studies




Table 3 Characteristics of included studies

	Inclusion criteria	Interventions	Sample size	Number of pt. followed-up	μ Age (year) (range) $\mu = 79.7$	Sex		CT/MRI for diagnosis
						Male	Female	
Primary included								
Seybold [33]	Type II–III	S & C	22	18	77.9 (65–93)	10	12	?
Hénaux [18]	Type IIB, >80 year	S	11	9	85.4 (82–93)	4	7	Yes
Koech [22]	Type II, >65 year	C	57	42	80 (67–91)	9	33	Yes
Platzer [30]	Type I–III, >65 year, “complete sets of clinical data”, surgical treatment	S	62	56	71.4 (66–83)	25	31	Yes
Osti [29]	Type II, >65 year	S	35	25	79.6 (65.1–94)	18	15	Yes
Hou [19]	Type II, >65 year, anterior screw fixation	S	43	42	80.6 (65–92)	24	19	?
Kaminski [German] [21]	OF/pseudoarthrosis, >70 year	S	36	28	80.1 (70–93)	14	22	Yes
Butler [5]	Type II, external immobilization	C	14	14	? (?–80)	Cannot retrieve		Yes
Lefranc [French] [25]	Type II–III, >70 year	S & C	27	22	80.7 (70–97)	13	14	?
Molinari JS DT [26]	Type II, >65 year, $\geq 50\%$ odontoid displacement, posterior fusion	S	26	21	79 (65–90)	14	12	?
Molinari ESJ [27]	Type II, >65 year, $\leq 50\%$ odontoid displacement, external immobilization	C	34	32	84 (71–99)	17	17	?
Secondary included								
Dailey [13]	Type II–III, >70 year	S	57	42	81.2 (70–96)	27	30	?
Kuntz [24]	Type II, >65 year	S & C	20	20	78.9 (66–92)	12	8	Yes
Koivikko [23]	Type II	C	25	25	? (65–94)	Cannot retrieve		?
Börm [34]	Type II, >70 year	S	15	15	81 (70–?)	Cannot retrieve		Yes
Andersson [2]	Type II–III, >65 year	S & C	29	27	78 (66–99)	11	18	?
Berlemann [3]	Type II, >65 year	S	19	19	75 (65–87)	10	9	?
Hanigan [15]	Type II–III, >80 year	S & C	19	19	86.2 (80–99)	9	10	?
Hanssen [16]	Type II–III	C	18	11	76 (65–87)	10	8	?

S surgical, C conservative, ? cannot tell or not retrievable

Table 4 Results of data extraction

	μ $\mu = 29$	Adverse effects		Baseline clinical status		Clinical outcome	
		Surgical	Conservative	Surgical	Conservative	Surgical	Conservative
Primary included							
Seybold [33]	21 (0–109)			–		1.25 (SW)	1.92 (SW)
Hénaux [18]	34 (2–116)			All ASIA E		2.18 (SW)	
Koeh [22]	Median = 24 (9–24)		1 × cervical collar-related decubitus ulceration, 2 × halo-vest-related pressure ulceration, 1 × decreased halothoracic pin tension	–			2.05 (SW)
Platzler [30]	24 (24–24)	4 × mortality, 3 × reoperations 8.6 % mortality		ASA average: 2.2		1.68 (SW)	
Osti [29]	74 (9.6–128)			All ASIA E		1.65 (SW)	
Hou [19]	21.3 (18–24)			Frankel: 1 × C, 3 × D		1.67 (Robinson's criteria)	
Kaminski [German] [21]	Clinical = 38, radiological = 23 (12–72)	2 × screw misplaced, 1 × wound infection		1 × ASIA D		Pain: 27/28 good-very good and 1 pt required pain killers, Activity: 20/28 very good, 6/20 good, 1/20 satisfying, 1/20 inadequate	
Butler [5]	66* (cannot retrieve)			–			CSOQ: neck pain (33.88 ± 27.08), shoulder/arm pain (35.10 ± 28.20), phys. symp. (38.57 ± 26.75), funct. disability (33.93 ± 18.04), psych. distress (45.24 ± 24.64) Rankin score average: 0.7
Lefranc [French] [25]	12 (cannot retrieve)			Neck pain, retropharyng. hematoma, neurol. signs		Rankin score average: 0.1	

Table 4 continued

	μ Follow-up (mo)—(range)	Adverse effects		Baseline clinical status		Clinical outcome	
		Surgical	Conservative	Surgical	Conservative	Surgical	Conservative
Molinari JSJT [26]	14 (1–48)	19.2 % mortality	–	–	18.1 (NDI)	–	–
Molinari ESJ [27]	15 (2–48)	–	11.8 % mortality, 5.9 % collar-related complications	–	–	–	15.7 (NDI)
Secondary included							
Dailey [13]	15 (3–62)	1 × myocardial infarction	–	NA	NA	–	NA
Kuntz [24]	Surg = 17, cons = 13 (0–36)	1 × revision surgery, 1 × death after surgery	1 × death	NA	NA	–	NA
Koivikko [23]	12.2* (2–82)	1 × surgery related death	–	NA	NA	–	NA
Börm [4]	17* (cannot retrieve)	10 × died from unrelated causes during the follow-up period [S or C?]	–	NA	NA	–	NA
Andersson [2]	51 (24–89)	–	–	NA	NA	–	NA
Berlemann [3]	Clinical = 54, radiological = 30 (3.6–132)	1 × hematoma requiring reoperation	–	NA	NA	–	NA
Hanigan [15]	Surg = 44.8, cons = 19.9 (5–72)	1 × mortality	5 × mortality	NA	NA	–	NA
Hanssen [16]	31 (6–60)	–	–	NA	NA	–	NA

* For entire pt group studied (i.e. including pt <65 years), ASA American Spinal Injury Association Impairment Scale, ASA American Society of Anesthesiologists Physical Status Classification System, SW Smiley-Webster score, CSOQ Cervical Spine Outcomes Questionnaire, NA not applicable/available

Table 5 Results of data extraction—fracture healing

	Osseous union						Fracture stability					
	Surgical			Conservative			Surgical			Conservative		
	Union	Non-union	Cannot tell	Union	Non-union	Cannot tell	Stable	Unstable	Cannot tell	Stable	Unstable	Cannot tell
Primary included												
Seybold [33]	4	5	2				2		2	11		7
Hénaux [18]				17	25	15	9		2	41	1	15
Koech [22]	52	4	6				52		10			
Platzer [30]	18	7	10				18		17			
Osti [29]	36	6	1				41	1	1			
Hou [19]	28	0	8				28	0	8			
Kaminski [German] [21]				9	5					13	1	
Butler [5]				6	4	7	10	0		10	0	7
Lefranc [French] [25]	9	1					10	0				
Molinari JSDT [26]	7	14	5				21	0	5			
Molinari ESJ [27]				2	30	2				11	21	2
Secondary included												
Dalley [13]	24	18	15				34	8	15			
Kuntz [24]	4		2	3	6	5	4		2	6	3	5
Koivikko [23]				6	19					6		19
Börm [4]	11	4					13	2				
Andersson [2]	14	1	3	2	6	3	14	1	3	2	6	3
Berlemann [3]	16	2	1				17	1	1			
Hangan [15]	2	3		5	4	5	5	0		8	1	5
Hanssen [16]				8	3	7				9	1	8
SUM	225	65	53	58	102	44	268	13	44	117	34	71
% Including 'cannot tell' (sensitivity)			66 %			28 %			82 %			53 %
% Excluding 'cannot tell'			78 %			36 %			95 %			77 %
% In studies comparing S to C treatment			85 %			44 %			97 %			79 %

Table 6 Results of risk-of-bias assessment of the individual studies

	Primary included										
	Seybold [33]	Hénaux [18]	Koech [22]	Platzer [30]	Osti [29]	Hou [19]	Kaminski [21]	Butler [5]	Lefranc [25]	Molinari JSDT [26]	Molinari ESJ [27]
Key criteria											
Clear study objective	+	+	+	+	+	+	+	+	-	+	+
Criteria for inclusion explicit	+	+	+	-	+	+	+	+	+	+	+
Fractures appropriately described	+	+	+	+	+	+	-	+	+	+	+
Distinction Type II/III appropriate	+	+	+	+	+	+	-	+	+	?	?
Mean age (range)	+	+	+	+	+	+	+	+	+	+	+
Selection bias ruled out	+	+	+	-	+	+	+	+	+	+	+
Mean follow-up (range)	+	+	+	+	+	+	+	+	+	+	+
Surgical treatm. specified	+	+	NA	+	+	+	+	NA	+	+	NA
Conservative treatm. specified	-	NA	+	NA	NA	NA	NA	+	+	NA	+
Clear criteria for measuring outcomes	+	+	+	+	+	+	+	+	+	+	+
Clinically relevant outcomes	+	+	+	+	+	+	+	+	+	+	+
When clinical outcome reported: Is pre-treatment neurol status stated	-	+	-	+	+	+	+	-	+	-	-
Results for surg/cons treatm. separately given	+	NA	NA	NA	NA	NA	NA	NA	+	NA	NA
Selective l.t.f.u. ruled out	No, l.t.f.u. >20 %	Yes	Yes	Yes	No, l.t.f.u. >20 %	Yes	No, l.t.f.u. <20 %	Yes	Yes	Yes	Yes
Other criteria											
Valid statistical analysis	+	NA	NA	+	+	NA	NA	+	+	+	+
Number of men and women given or extractable	-	+	+	+	+	+	+	+	+	+	+
Clinical evaluation independent of treating physician	?	?	?	?	?	?	?	?	?	?	?
Radiological evaluation independent and blinded	?	?	?	?	?	?	?	?	?	?	?
Independence of investigators stated	-	+	+	+	+	+	+	-	-	+	+
Quantification of outcomes	Yes, <5 scale	Yes, <5 scale	Yes, <5 scale	Yes, <5 scale	Yes, <5 scale	Yes, <5 scale	Yes, <5 scale	Yes, >5 scale	Yes, >5 scale	Yes, <5 scale	Yes, <5 scale
Results											
Selection bias ruled out	Yes	Yes	Yes	No	Yes	Yes	Yes	Yes	Yes	Yes	Yes
Baseline (max 11 points)	8	10	9	9	10	10	8	9	10	8	8
Results (max 11 points)	5	6	6	7	5	6	5	7	8	7	7

Table 6 continued

	Secondary included							
	Dailey [13]	Kuntz [24]	Koivikko [23]	Börm [4]	Andersson [2]	Berlemann [3]	Hanigan [15]	Hanssen [16]
Key criteria								
Clear study objective	+	+	+	+	+	+	+	+
Criteria for inclusion explicit	+	+	+	+	+	+	+	+
Fractures appropriately described	+	+	-	+	+	+	+	+
Distinction Type II/III appropriate	+	+	+	+	+	+	+	+
Mean age (range)	+	+	+	+	+	+	+	+
Selection bias ruled out	+	+	-	+	+	+	-	+
Mean follow-up (range)	+	+	+	-	+	+	+	+
Surgical treatm. specified	+	+	-	+	+	+	+	NA
Conservative treatm. specified	NA	+	+	NA	+	NA	+	+
Clear criteria for measuring outcomes	+	+	+	+	+	+	+	+
Clinically relevant outcomes	-	-	-	+NE	+NE	+NE	+NE	-
When clinical outcome reported: Is pre-treatment neuro status stated	NA	NA	NA	+NE	+NE	+NE	-	NA
Results for surg/cons treatm. separately given	NA	+	NA	NA	+	NA	+	NA
Selective I.t.f.u. ruled out	No, I.t.f.u. <20 %	Yes	Yes	Yes	Yes	Yes	Yes	No, I.t.f.u. <20 %
Other criteria								
Valid statistical analysis	NA	+	+	+	-	-	-	-
Number of men and women given or extractable	+	+	+	+	+	+	+	+
Clinical evaluation independent of treating physician	?	?	?	?	?	?	?	?
Radiological evaluation independent and blinded	?	?	?	?	?	?	?	?
Independence of investigators stated	-	-	+	-	-	-	-	-
Quantification of outcomes	Yes, <5 scale	Yes, descr.	Yes, descr.	Yes, <5 scale	Yes, <5 scale	Yes, descr.	Yes, descr.	Yes, descr.
Results								
Selection bias ruled out	Yes	Yes	No	Yes	Yes	Yes	No	Yes
Baseline (max 11 points)	8	9	7	10	11	10	10	8
Results (max 11 points)	4	5	6	5	6	4	6	3

+ Yes, - no, ? cannot tell, NA not applicable, NE not extractable for review purposes

were adequately described in all studies, with scores ranging from 7 to 10 on a 11-point scale. However, baseline data regarding clinical status were poorly reported. Only six of the primary included studies systematically reported baseline clinical status and they all applied different tools for the pre- and post-treatment measurements. Data concerning the results of the studies were less extensively reported than baseline data. Scores ranged between 3 and 8 on a 11-point scale.

Clinical outcome

A variety of tools was used to assess clinical outcome. The NDI, which was expected to be widely applied, was used in only two of the included studies. The Smiley-Webster scale was the most commonly applied tool to assess clinical outcome, used in five studies. The Cervical Spine Outcomes Questionnaire was used in one study, as were the Robinson's criteria, Rankin score and a self-designed pain-and-activity scoring scale. These variations made it impossible for statistical analyses to be carried out and limited the comparisons that could be made.

Surgery versus conservative care in individual studies

Of the eleven studies reporting clinical outcome, only two studies compared surgical to conservative treatment [25, 33]. However, these two studies applied different tools to assess clinical outcome, using the Rankin score [25] (0 = without symptoms; 5 = major handicap) and Smiley-Webster (SW) scale [33] (1 = excellent; 4 = poor), respectively, and hence could not adequately be compared. One of these studies [25] showed statistically less morbidity in the surgery group ($p = 0.037$), but no significant difference in non-union at fracture site ($p = 0.64$) except for type II fractures ($p = 0.028$) [25]. However, this was a very small study, which makes these estimates unstable. The other study showed a slightly better clinical outcome in surgically compared to conservatively treated patients (average Smiley-Webster score 1.25 and 1.92, respectively). Statistical analysis of these results could not be performed due to the limited number of patients involved ($n = 17$) [33].

Outcome compared between studies

In the two studies applying the NDI, groups were evidently different; patients in one group had $\geq 50\%$ odontoid displacement and were treated surgically [26], those in the other group had $\leq 50\%$ odontoid displacement and were treated by external immobilization [27], still showing a slightly better outcome for surgically treated patients (18.1 and 15.7, respectively). This difference was, however, not

clinically relevant. In the five studies evaluating the clinical outcome using the Smiley-Webster scale, both surgically and conservatively treated patients had an intermediate outcome [18, 22, 29, 30, 33]. The SW score for surgically treated patients averaged at 1.71, the SW score for conservatively treated patients averaged at 2.02. The remaining studies all used different instruments to assess outcome and hence could not be compared.

Osseous union

In eighteen studies reporting extractable union rates, four compared surgical to conservative treatment. In these four studies, union was achieved in 85 % (29/34) of the surgically treated patients and in 44 % (16/36) of the conservatively treated patients [2, 15, 24, 25]. The results were mainly based on X-ray data. In the individual studies no statistical analysis could be performed due to the small number of patients. In a comparison of all patients included in the eighteen studies, union was achieved in 66 % (225/343) of surgically treated patients and 28 % (58/204) of conservatively treated patients. When patients whose outcome was unclear are left out of the calculation, union was achieved in 78 % (225/290) of surgically treated and in 36 % (58/160) of conservatively treated patients. Again, in the individual studies no statistical analysis could be carried out because of the limited number of patients included.

Fracture stability

All nineteen studies reported extractable stability rates, showing fracture stability in the majority of patients regardless of their treatment. Stability was in most studies assessed with dynamic X-ray. In five studies comparing surgically to conservatively treated patients, stability was achieved in 97 % (35/36) and 79 % (37/47) of cases, respectively [2, 15, 24, 25, 33]. In a comparison of all patients included in the nineteen studies, stability was achieved in 82 % (268/325) of surgically treated patients and in 53 % (117/222) of conservatively treated patients. When patients whose outcome was unclear were left out of the calculation, stability was achieved in 95 % (268/281) of surgically treated patients and in 77 % (117/151) of conservatively treated patients.

Discussion

The studies identified for this review were all performed retrospectively, mostly without the use of a control group. Due to the small numbers of patients and the rather poor quality of the acquired data, no strong recommendations can be made with regard to the optimal treatment. A variety of tools was used to assess clinical outcome. The Smiley-

Webster scale, the only instrument used in multiple studies, showed a slightly better outcome for surgically treated patients than for conservatively treated patients. However, pre-treatment status was poorly reported and the grounds for starting specific treatments are unknown. Osseous union in surgically treated patients was twice as high as in conservatively treated patients, but if non-union leads to stability (e.g. fibrous union), fracture stability is a more important and relevant parameter to evaluate. The results of this review indicate that fracture stability can be achieved in the majority of patients, regardless of the applied treatment or presence of osseous union. Adverse effects were common in both treatment groups, with mortality rates as high as 20 %. Complications related to the operation, such as screw misplacement, were the most common complications in surgically treated patients. Device-related complications, such as ulcerations, were the most common complication in patients treated conservatively.

Patient age in the included studies was more or less comparable. Mean age for surgically treated patients was 78.9 years. Mean age for conservatively treated patients was 80.0 years. In all but two studies, patients over 65 or 70 years old were included. Two studies included only patients aged over 80 years of age (in one of these studies, patients were treated surgically, in the other, patients were treated both surgically and conservatively). Surgically and conservatively treated groups described in the included studies may, however, not be comparable due to potential differences in a variety of other patient characteristics (i.e. co-morbidity, osteoporosis, severity of comminution). Outcome diversification per age group amongst the elderly was absent and needs further study. It is often postulated that treatment outcome depends on patient age [20]. Other factors must, however, play a role as well, as different studies have shown different outcomes for the same treatment, which cannot be explained by patient age alone (e.g. in the studies by Koech and Molinari [22, 27]).

No consensus exists as to the exact goal of treatment (debate remains as to whether it should be osseous union, fracture stability, or clinical outcome), nor as to how outcome should be measured. Many studies used osseous union rates as primary outcome, fibrous union was under-exposed and the correlation with clinical outcome or fracture stability was not properly studied. The results of this review show no evidence that clinical outcome correlates better to fracture union than to fracture stability (i.e. that the quality of union, whether it be osseous or fibrous, is influential on clinical outcome).

Strengths and limitations

The limited number and poor quality of the included studies limits the strength of the results found in this

review. Furthermore, type II and III fractures were analysed as one group. However, an evident type II fracture is more often treated surgically, whereas an evident type III fracture is most often treated conservatively. This may have flattened the findings. This choice was made as differentiation of these types is often difficult to make and would even further reduce the amount of data that could be used for this review, as this differentiation was also not made in all of the included studies. As a result, the subgroup analysis that was planned to be carried out based on fracture type has also lapsed.

Conclusions

Implications for clinical practice

Different treatment options for this condition have been extensively reported and their efficacy debated. Based on this review, the following conclusions can be drawn based on the three main outcome parameters.

1. Clinical outcome: There is insufficient data available to determine a potential difference in clinical outcome between surgical and conservative interventions in the elderly with isolated odontoid fractures
2. Osseous union: Surgically treated patients appear to show higher osseous union rates compared to conservatively treated patients, although selection mechanisms, especially regarding fracture type, might explain this difference
3. Stability: The majority of patients appears to achieve fracture stability regardless of the applied treatment

Implications for research

A prospective study with appropriate sample size is necessary to identify the optimal treatment of odontoid fractures in the elderly and predictors for the success of either one of the available treatments.

Acknowledgments The authors are thankful to Mr. Jan Schoones of the Walaeus Library for carrying out the database search.

Conflict of interest None.

References

1. Anderson LD, D'Alonzo RT (1974) Fractures of the odontoid process of the axis. *J Bone Joint Surg Am* 56(8):1663–1674
2. Andersson S, Rodrigues M, Olerud C (2000) Odontoid fractures: high complication rate associated with anterior screw fixation in the elderly. *Eur Spine J* 9(1):56–59

3. Berlemann U, Schwarzenbach O (1997) Dens fractures in the elderly: results of anterior screw fixation in 19 elderly patients. *Acta Orthop Scand* 68(4):319–324
4. Borm W, Kast E, Richter HP, Mohr K (2003) Anterior screw fixation in type II odontoid fractures: is there a difference in outcome between age groups? *Neurosurgery* 52(5):1089–1092
5. Butler JS, Dolan RT, Burbridge M et al (2010) The long-term functional outcome of type II odontoid fractures managed non-operatively. *Eur Spine J* 19(10):1635–1642
6. Carreon LY, Glassman SD, Campbell MJ, Anderson PA (2010) Neck Disability Index, short form-36 physical component summary, and pain scales for neck and arm pain: the minimum clinically important difference and substantial clinical benefit after cervical spine fusion. *Spine J* 10(6):469–474
7. Chan K, Bhandari M (2011) Three-minute critical appraisal of a case series article. *Indian J Orthop* 45(2):103–104
8. Cleland JA, Fritz JM, Whitman JM, Palmer JA (2006) The reliability and construct validity of the Neck Disability Index and patient specific functional scale in patients with cervical radiculopathy. *Spine (Phila Pa 1976)* 31(5):598–602
9. Cochrane (2012) Cochrane form II for RCTs. <http://dcc.cochrane.org/sites/dcc.cochrane.org/files/uploads/RCT.pdf>. Accessed 1 March 2012
10. Cochrane (2012) Cochrane form III for cohort studies. <http://dcc.cochrane.org/sites/dcc.cochrane.org/files/uploads/cohort.pdf>. Accessed 1 March 2012
11. Cochrane (2012) Cochrane form IV for patient control studies. <http://dcc.cochrane.org/sites/dcc.cochrane.org/files/uploads/patient-controlonderzoek.pdf>. Accessed 1 March 2012
12. Cowley DE (1995) Prostheses for primary total hip replacement: a critical appraisal of the literature. *Int J Technol Assess Health Care* 11(4):770–778
13. Dailey AT, Hart D, Finn MA, Schmidt MH, Apfelbaum RI (2010) Anterior fixation of odontoid fractures in an elderly population. *J Neurosurg Spine* 12(1):1–8
14. Grauer JN, Shafi B, Hilibrand AS et al (2005) Proposal of a modified, treatment-oriented classification of odontoid fractures. *Spine J* 5(2):123–129
15. Hanigan WC, Powell FC, Elwood PW, Henderson JP (1993) Odontoid fractures in the elderly patient. *Clin Res* 41(3):A710
16. Hanssen AD, Cabanela ME (1987) Fractures of the dens in adult patients. *J Trauma* 27(8):928–934
17. Harrop JS, Hart R, Anderson PA (2010) Optimal treatment for odontoid fractures in the elderly. *Spine* 35(21):S219–S227
18. Henaux PL, Cueff F, Diabira S et al (2011) Anterior screw fixation of type IIB odontoid fractures in octogenarians. *Eur Spine J* 21(2):335–339
19. Hou Y, Yuan W, Wang X (2011) Clinical evaluation of anterior screw fixation for elderly patients with type II odontoid fractures. *J Spinal Disord Tech* 24(8):75–81
20. Hsu WK, Anderson PA (2010) Odontoid fractures: update on management. *J Am Acad Orthopaedic Surg* 18(7):383–394
21. Kaminski A, Gstrein A, Muhr G, Muller EJ (2008) Transarticular C1–C2 screw fixation: results of unstable odontoid fractures and pseudarthrosis in the elderly. *Unfallchirurg* 111(3):167–172
22. Koech F, Ackland HM, Varma DK, Williamson OD, Malham GM (2008) Nonoperative management of type II odontoid fractures in the elderly. *Spine* 33(26):2881–2886
23. Koivikko MP, Kiuru MJ, Koskinen SK, Myllynen P, Santavirta S, Kivisaari L (2004) Factors associated with nonunion in conservatively treated type-II fractures of the odontoid process. *J Bone Joint Surg Br* 86(8):1146–1151
24. Kuntz C, Mirza SK, Jarell AD, Chapman JR, Shaffrey CI, Newell DW (2000) Type II odontoid fractures in the elderly: early failure of nonsurgical treatment. *Neurosurg Focus* 8(6):e7
25. Lefranc M, Peltier J, Fichten A, Desenclos C, Toussaint P, Le GD (2009) Odontoid process fracture in elderly patients over 70 years: morbidity, handicap, and role of surgical treatment in a retrospective series of 27 cases. *Neurochirurgie* 55(6):543–550
26. Molinari RW, Dahl J, Gruhn WL, Molinari WJ (2011) Functional outcomes, morbidity, mortality, and fracture healing in 26 consecutive geriatric odontoid fracture patients treated with posterior fusion. *J Spinal Disord Tech* 19(5):1229–1234
27. Molinari RW, Khera OA, Gruhn WL, McAssey RW (2011) Rigid cervical collar treatment for geriatric type II odontoid fractures. *Eur Spine J* 21(5):855–862
28. Muller EJ, Wick M, Russe O, Muhr G (1999) Management of odontoid fractures in the elderly. *Eur Spine J* 8(5):360–365
29. Osti M, Philipp H, Meusbürger B, Benedetto KP (2011) Analysis of failure following anterior screw fixation of Type II odontoid fractures in geriatric patients. *Eur Spine J* 20(11):1915–1920
30. Platzer P, Thalhammer G, Oberleitner G, Schuster R, Vecsei V, Gaebler C (2007) Surgical treatment of dens fractures in elderly patients. *J Bone Joint Surg Am* 89(8):1716–1722
31. Platzer P, Thalhammer G, Ostermann R, Wieland T, Vecsei V, Gaebler C (2007) Anterior screw fixation of odontoid fractures comparing younger and elderly patients. *Spine (Phila Pa 1976)* 32(16):1714–1720
32. Reinhold M, Bellabarba C, Bransford R et al (2011) Radiographic analysis of type II odontoid fractures in a geriatric patient population: description and pathomechanism of the “Geier”-deformity. *Eur Spine J* 20(11):1928–1939
33. Seybold EA, Bayley JC (1998) Functional outcome of surgically and conservatively managed dens fractures. *Spine (Phila Pa 1976)* 23(17):1837–1845
34. Shears E, Armitstead CP (2008) Surgical versus conservative management for odontoid fractures. *Cochrane Database Syst Rev* 4:CD005078
35. Tashjian RZ, Majercik S, Biffi WL, Palumbo MA, Cioffi WG (2006) Halo-vest immobilization increases early morbidity and mortality in elderly odontoid fractures. *J Trauma* 60(1):199–203
36. University of Wales, College of Medicine (2011). Critical appraisal form. http://www.core-info.cardiff.ac.uk/thermal/FULL_COTS_critical%20appraisal%20FORM%20B%202007.pdf. Accessed 2 Sept 2011
37. White AP, Hashimoto R, Norvell DC, Vaccaro AR (2010) Morbidity and mortality related to odontoid fracture surgery in the elderly population. *Spine* 35(9):S146–S157
38. Young BA, Walker MJ, Strunce JB, Boyles RE, Whitman JM, Childs JD (2009) Responsiveness of the Neck Disability Index in patients with mechanical neck disorders. *Spine J* 9(10):802–808
39. Young IA, Cleland JA, Michener LA, Brown C (2010) Reliability, construct validity, and responsiveness of the neck disability index, patient-specific functional scale, and numeric pain rating scale in patients with cervical radiculopathy. *Am J Phys Med Rehabil* 89(10):831–839