

Laminin 332 processing impacts cellular behavior

Patricia Rousselle^{1,*} and Konrad Beck²

¹SFR BioSciences Gerland-Lyon Sud; Institut de Biologie et Chimie des Protéines; UMR 5305, CNRS; Université Lyon 1; Lyon, France; ²Cardiff University School of Dentistry; Cardiff, UK

Keywords: laminin 332, LG45 domain, LE domain, syndecan, basement membrane, keratinocyte

Abbreviations: ECM, extracellular matrix; LG, laminin globular domain; pre-laminin, precursor-laminin; MMP, matrix metalloproteinase; mTLD, mammalian tolloid; BMP-1, bone morphogenic protein 1; JEB, junctional epidermolysis bullosa; pdb, protein data bank

Laminin 332, composed of the $\alpha 3$, $\beta 3$ and $\gamma 2$ chains, is an epithelial-basement membrane specific laminin variant. Its main role in normal tissues is the maintenance of epithelial-mesenchymal cohesion in tissues exposed to external forces, including skin and stratified squamous mucosa. After being secreted and deposited in the extracellular matrix, laminin 332 undergoes physiological maturation processes consisting in the proteolytic processing of domains located within the $\alpha 3$ and the $\gamma 2$ chains. These maturation events are essential for laminin 332 integration into the basement membrane where it plays an important function in the nucleation and maintenance of anchoring structures. Studies in normal and pathological situations have revealed that laminin 332 can trigger distinct cellular events depending on the level of its proteolytic cleavages. In this review, the biological and structural characteristics of laminin 332 domains are presented and we discuss whether they trigger specific functions.

Laminins are large extracellular glycoproteins that are important components of all basement membranes. They are involved in several biological processes, including self polymerization, binding to the extracellular matrix (ECM) and cellular interactions.^{1,2} All laminins are composed of three different gene products, termed α , β and γ chains assembled into a cross-shaped heterotrimer $\alpha\beta\gamma$. The three chains assemble within the endoplasmic reticulum through their C-terminal domains to form a triple stranded α -helical coiled coil rod.^{3,4} Sixteen laminin isoforms of different subunit composition selected from five individual α chains ($\alpha 1$ to $\alpha 5$), three β chains ($\beta 1$ to $\beta 3$) and three γ chains ($\gamma 1$ to $\gamma 3$), are known with variable cell and tissue specific expression, and they are differentially recognized by cellular receptors.⁵ All laminin α chains possess a large globule at the carboxyl-terminal end that consists of five similar domains LG1 to LG5 each containing about 200 residues.^{6,7}

Laminin 332 and Its Maturation Events

Laminin 332, composed of the $\alpha 3$, $\beta 3$ and $\gamma 2$ chains, is an epithelial-basement membrane specific variant. The $\alpha 3$ chain is encoded by the LAMA3 gene, which has two transcript variants $\alpha 3A$ and $\alpha 3B$. As the truncated LAMA3A variant is expressed and incorporated into laminin 332 heterotrimers,⁸ we will refer to the $\alpha 3A\beta 3\gamma 2$ trimer as representing laminin 332 in this review. Its main role in normal tissues is the maintenance of epithelial-mesenchymal cohesion in tissues exposed to external forces including skin and stratified squamous mucosa.^{9,10} To fulfill this function, laminin 332 undergoes several physiological post-translational processing events allowing its supramolecular integration into the basement membrane. In other situations such as wound healing or carcinoma, additional cleavages may take place delivering pro-migratory signals to cells therefore compromising the adhesive function of the protein.

Laminin 332 processing and integration into the skin basement membrane. In skin, laminin 332 is synthesized by keratinocytes as a high molecular weight precursor protein of 460 kDa. After secretion and deposition into the ECM, the $\alpha 3$ and $\gamma 2$ chains undergo maturation events consisting in specific processing to smaller forms.^{4,11} The 190–200 kDa $\alpha 3$ chain ($\alpha 3_{200}$) can be successively processed at both C- and N-terminal extremities producing 165 ($\alpha 3_{165}$) and 145 kDa ($\alpha 3_{145}$) maturation products. The 155 kDa $\gamma 2$ chain ($\gamma 2_{155}$) is processed in the N-terminal region leading to a 105 kDa product ($\gamma 2_{105}$).

Processing of the $\alpha 3$ chain consists of cleavage of the C-terminal globular domains 4 and 5 (LG45) within the spacer between LG3 and LG4 (Fig. 1A and B).^{12,13} In vitro studies using human laminin 332 have revealed that enzymes involved in the processing include plasmin,¹² MMP-2, MT1-MMP and the C-proteinase family of enzymes, especially mammalian tolloid (mTLD) and the bone morphogenic protein 1 (BMP-1).^{14,15} In addition, thrombin was shown to have the potency to cleave LG45 specifically.¹⁶ Alternatively, there may be other mechanisms that control the rate of laminin $\alpha 3$ LG45 processing, such as the tissue plasminogen proteolytic cascade.¹² N-terminal amino acid sequencing of human LG45 purified from the conditioned medium of either primary or immortalized human keratinocytes and of the human gastric adenocarcinoma cells STKM-1 revealed that the $\alpha 3$ chain is cleaved between Q₁₃₃₇ and D₁₃₃₈ within the

*Correspondence to: Patricia Rousselle; Email: p.rousselle@ibcp.fr
Submitted: 07/16/12; Revised: 11/12/12; Accepted: 12/05/12
<http://dx.doi.org/10.4161/cam.23132>

hinge region between LG3 and LG4 suggesting that a proteolytic cleavage site of the LG45 domain matches the minimal consensus sequence LLQD (Fig. 1B).^{17,18} It is not known what endopeptidase catalyzes the hydrolysis of this cleavage sequence, and the existence of additional proteolytic cleavage sites has been suspected both in the hinge and the adjacent regions of the LG3 and LG4 domains.^{7,13,14,19} This hypothesis is compatible with the spacer length of the laminin $\alpha 3$ chain, which is longer than those of the $\alpha 1$, $\alpha 2$ and $\alpha 5$ chains.⁷ Punctual mutations within the spacer region and/or deletion of the sequence LLQD did not protect the $\alpha 3$ LG3-LG4 linker from cleavage.^{14,19} As proteolytic processing of LG domains occurs in all α chains except $\alpha 1$, the substitution of the $\alpha 3$ LG3-LG4 hinge with that of the $\alpha 1$ chain was an elegant way to render the laminin $\alpha 3$ chain uncleavable.¹⁹ This was also the case when a 46 amino-acid portion within the $\alpha 3$ LG3-LG4 hinge was deleted.¹⁶ These LG45-uncleavable $\alpha 3$ chain constructs were expressed in human¹⁹ or mouse¹⁶ skin keratinocytes deficient for expression of the $\alpha 3$ chain, and in both cases the resulting LG45-uncleavable heterotrimeric laminin 332 was deposited in the ECM. Information gained from the structure of the mouse $\alpha 2$ chain LG45 domain pair revealed an unusual path of the LG3-LG4 linker as it forms an inter-domain disulfide bridge to LG5.²⁰ Being integrated into the LG45 tandem the linker is therefore an integral part of the structure, a feature that is conserved in all laminin α chains. It is tempting to speculate that proteinases may have easy access to the LG3-LG4 cleavage sites located within a well-exposed linker.⁷

Further cleavage occurs in the N-terminal LE region of the $\alpha 3$ chain releasing the full short arm which might be important for laminin 332 function (Fig. 1A).^{14,21} A mutation causing an N-terminal deletion of 226 amino acids in the human $\alpha 3A$ isoform was identified in the laryngo-oncho-cutaneous syndrome, a rare autosomal recessive disorder characterized by chronic production of vascularized granulation tissue.²² Cultured keratinocytes obtained from patients revealed that heterotrimeric laminin 332 carrying the mutant $\alpha 3$ chain is assembled and secreted. The mutated laminin was regularly expressed in the skin basement membrane and only very subtle ultrastructural changes were seen such as focal widening of the lamina lucida in places where hemidesmosome plaques were smaller. These studies suggest that the laminin $\alpha 3$ N-terminal domain may be a key regulator of the granulation tissue response during wound healing.

In human skin laminin 332, the processing of the $\gamma 2_{155}$ chain leads to the $\gamma 2_{105}$ subunit and was shown to be achieved by BMP-1 and mTLD metalloproteinases.^{11,14,15,23,24} The N-terminal processing is complex as it occurs within a disulfide-linked loop of the L4 domain followed by reshuffling of disulfide bonds for release of the cleaved fragments.²⁴ The immunohistochemical analysis of laminin 332 in mTLD/BMP-1-deficient mouse skin revealed a strong expression of the precursor $\gamma 2$ chain confirming that these enzymes are involved in the $\gamma 2$ chain maturation in vivo.²⁵ In contrast, the processing of the LG45 domain in the $\alpha 3$ chain seemed, at least partially, unaffected reinforcing the hypothesis that several enzymes are involved in this maturation process.

Laminin 332 was shown to be the major component of anchoring filaments in skin⁹ where it mediates cell adhesion via interaction of the $\alpha 3$ carboxyl-terminal LG1–3 triplet domain with both $\alpha 3\beta 1$ and $\alpha 6\beta 4$ integrins,^{26–29} while the N-terminal short arms connect to basement membrane components. Laminin 332 can be incorporated into the basement membrane through at least two mechanisms. The first involves cross-linking of laminin 332 with laminin 311 ($\alpha 3\beta 1\gamma 1$) in the skin basement membrane.²¹ The complex of laminin 311 with laminin 332 is most likely to derive from an interaction of domain LN in the $\beta 3$ chain of laminin 332 with domain LE of the $\alpha 3$ chain short arm in laminin 311 (see below). As seen on rotary shadowing electron micrographs of the complex, the LN domain of the short arm interacts with a laminin 311 domain near the intersection of the laminin 311 short arms.²¹ These complexes are most likely stabilized by a disulfide bridge between an unpaired cysteine in domain LN of the $\beta 3$ -chain and domain LE of processed $\alpha 3$ in laminin 311. A similar complex between laminin 332 and laminin 321 ($\alpha 3\beta 2\gamma 1$) is found in the basement membrane of the amnions. According to the 3-arm interaction hypothesis of laminin polymerization,³⁰ the dimers could self-associate. The second mechanism reports a direct interaction between anchoring filaments and anchoring fibrils. Anchoring fibrils are disulfide bond stabilized dimers of type VII collagen.³¹ Monomeric laminin 332 as well as the laminin 332/311 dimer directly bind the N-terminal globular domain NC1 of type VII collagen.^{10,32} The interaction is likely to occur within the short arm of the $\beta 3$ and/or $\gamma 2$ subunit.³²

Other processing events in laminin 332. In addition to the laminin 332 maturation aimed at regulating the skin basement membrane structural integrity, other studies have reported two specific migration-inducing MMP-2 cleavages of the laminin $\gamma 2$ N-termini producing $\gamma 2$ chain fragments of 100 and 80 kDa.³³ These $\gamma 2$ processing could also be carried by MT1-MMP in colon and breast carcinoma cells.³⁴ Both enzymes, that were first shown to cleave rat laminin 332, were proposed to also cut down human laminin 332 in both physiological and pathological situations such as tumorigenesis.^{35,36} MMP-3, -12, -13, -19 and -20 were also shown to process the $\gamma 2$ chain inducing epithelial cell migration.^{37,38} Migration events might therefore result from the interaction of newly exposed laminin domains with cell surface signaling receptors. Rat recombinant $\gamma 2LEb1-LEb4'$ was shown to bind and activate the EGF receptor to stimulate cell migration.³⁹ Studies with cancer cells have reported that MMP-7, MT1-MMP and hepsin cleave laminin $\beta 3$ leading to increased migration.^{40–42} The molecular mechanisms of laminin 332 in squamous carcinoma have been reviewed in detail by Marinkovich.⁴³

Laminin 332 Sequence-Structure Relationship

The structure of the heterotrimeric laminin 332 molecule consisting of disulfide-linked $\alpha 3$, $\beta 3$ and $\gamma 2$ chains was first revealed by rotary shadowing electron microscopy, which showed a rod 107 nm in length terminated by one large and two small globules at opposite ends⁹ which correspond to the C-terminal




Figure 1. Structure of human laminin 332 and its physiological maturation process. (A) Laminin 332 is composed of three subunits α 3A, β 3 and γ 2. Each chain is composed of different domains that are indicated. Domain L4 in the γ 2 chain corresponds to an LE domain with a ~180 residues insert between the third and fourth cysteine of the canonical 8-Cys pattern. LE4' and LEb4' of the α 3A and γ 2 chain are truncated LE repeats containing only the first four and six cysteines, respectively, of the pattern. Based on the odd number of cysteines, β 3LN, α 3ALE4', β 3LE6 and γ 2LEb1 (after maturation), and the coiled coil of α 3 have free SH groups as indicated. The coiled coil is stabilized by disulfide bonds at the N-terminus (dotted lines) though the connectivity is yet unknown. The large LG structure located at the C-terminal end of the α 3 chain contains five repeating LG domains. The first three repeats (LG1–3) interact with α 3 β 1, α 6 β 1 and α 6 β 4 integrins while the last two (LG45) contain binding sites for syndecan-1 and -4. Laminin 332 is synthesized as a precursor molecule that undergoes maturation by proteolytic processings at the α 3A chain N- and C-terminus as well as at the γ 2 chain N-terminal extremity. The cleavage sites are indicated by arrows as well as enzymes involved identified so far. (B) Schematic structures of LG. The five LG domains contain numerous cysteines that are numbered. Disulfide bridged and free cysteines (SH) within the human sequence are indicated. At the bottom, an alignment of the human and mouse LG3-LG4 linker is shown. Each sequence displays the same cleavage site (underlined) and a cysteine (printed in inverse font) that forms an inter-domain disulfide bond to LG5.²⁰ The sequence P₁₃₁₂PPFLLMLLKGSTR supposed to be crucial for integrin α 3 β 1 binding⁵⁸ is double underlined.

LG and N-terminal LN and LE/L4 domains, respectively (Fig. 1A). This length is consistent with the later determined sequences which indicate that ca. 570 residues per chain contribute to the three stranded α -helical coiled coil structure giving a length of ~80 nm, and ~7 tandem LE domains accounting for a further 20 nm. Circular dichroism spectroscopy revealed an α -helical content of ~30%⁴⁴ similar to that determined for laminin 111⁴⁵ suggesting that most of this secondary structure contribution relates to the coiled coil. Assembly of the laminin 332 trimer occurs in the endoplasmic reticulum where first a disulfide linked β 3- γ 2 dimer is formed to which the α 3 chain aligns via the coiled coil. Trimer formation is a prerequisite for transport to the Golgi complex and secretion but does not require N-linked glycosylation.⁴

Laminin β 3 LN domain. So far no high resolution structures of laminin β 3 domains have been solved. However, based on sequence similarity, some insight into the molecular organization of various domains can be anticipated from homology modeling. Recently, the crystal structure of the N-terminal LN domain together with four adjacent LE modules of the mouse laminin β 1 chain has been solved (pdb code: 4aqs).⁴⁶ The corresponding regions of human laminin β 3 can be fitted to this structure (Fig. 2) suggesting that LN (residues 18–248; accession number NP_000219) forms a β -sandwich consisting of eight β -strands folded into a jelly roll motif. The first LE domain (res. 249–314) is tightly integrated into LN, which starts with a disulfide-bonded reverse turn connected by C₂₁–C₂₆, and the N-terminal region (res. 18–28) associates to LE1. Pro₂₈, which is conserved in all laminin LN sequences, fits into a pocket of LE1. Further disulfide bonds can be expected between C₄₇–C₆₉, C₅₆–C₆₆ and C₁₅₃–C₁₇₃ (Fig. 2A). This leaves C₆₇, which is unique for β 3 LN domains and surface accessible (Fig. 2B), in a reduced state making it available for covalently binding to other proteins like laminin 331. The N-linked glycosylation sites found within β 1 LN at N₁₂₀ and in γ 1 LN at N₅₈/N₁₃₂ are absent in the β 3 chain. The critical residues D₁₀₆ and T₁₁₄ of γ 1 LN involved in calcium binding,⁴⁶ which might be important for laminin polymerization, are replaced by S₇₀ and R₇₆ in the β 3 chain indicating that this site will not be functional.

Laminin 332 LE and L4 domains. A common feature of LE modules is their connectivity of the eight cysteine residues as C1–C3, C2–C4, C5–C6 and C7–C8. The loop formed between C5–C6 usually consists of eight residues with a glycine in position six. In human laminin 332, the following LE domains deviate from the 8-Cys pattern (Fig. 1A): β 3-LE6 preceding the coiled coil contains one extra cysteine (C₅₇₂) between C7 and C8 whereas the corresponding region within α 3A-LE4' is truncated after the fourth cysteine (C₁₈₅; accession number AAA59483) followed by an irregular cysteine (C₁₉₉) just before C₂₀₂/C₂₀₅, which are presumably forming intrachain disulfides at the N-terminus of the coiled coil. The α 3A chain is processed just after LE4' between K₁₉₁ and D₁₉₂.¹⁴ LEB1 following domain L4 of the human γ 2 chain is missing the fifth cysteine of the 8-Cys pattern and does not contain the glycine residue N-terminal to the sixth cysteine (C₄₄₂; accession number NP_005553). This probably results in an increased mobility of the loop between C₄₃₁ and C₄₄₂ allowing

for protease susceptibility. Indeed, major processing occurs by cleavage of the G₄₃₄-D₄₃₅ site (Fig. 1A).²³

The L4 domain separating LEa3 and LEB1 can be regarded as a specialized LE domain with a large, ~180 residues long insert between the third (C₁₉₆) and fourth (C₃₈₂) cysteine of the regular pattern. In case of the γ 2 chains, the L4 insert is highly conserved from human down to *Xenopus* and zebrafish, and contains one unique cysteine absent in all other L4 domains. For L4-LEB1 it was shown that this C₃₀₉ is covalently linked with the last cysteine C₄₅₉ of LEB1, which contains an unusual S-S bridge between C₄₄₂–C₄₄₅ corresponding to C6–C7. After cleavage by BMP-1 between G₄₃₄ and D₄₃₅, protein disulfide isomerase was required to release the fragments.²⁴ It remains unclear whether after processing C₄₅₉ stays in a reduced state, or whether disulfide reshuffling might form the usual C7–C8 connectivity and leaves C6, i.e., C₄₄₂, with a free SH group.²⁴ The mouse L4-LEB1 domain pair was modified probably via one or both of the two putative NxT acceptor sites for N-linked glycosylation, which are also found in the human sequence (N₃₄₂ and N₃₆₂). The N₃₄₂xT site is conserved down to zebrafish. Mouse L4-LEB1 binds heparin, nidogen-1 (but not -2), and fibulin-1 and -2; it remains unclear whether these interactions occur between L4 or LEB1, or whether epitopes from both are required.²⁴

In conclusion, besides the β 3 LN domain, each of the three irregular LE modules, α 3A-LE4', β 3-LE6 and γ 2-LEB1 after processing, have one free cysteine which could participate either in interchain disulfide bonding or covalent binding to other proteins like laminins 311 and/or 321. Based on the canonical cysteine pattern, the most likely residues are α 3A-C₁₉₉, β 3-C₅₇₂ and γ 2-C₄₄₂ or γ 2-C₄₅₉.

Laminin 332 coiled coil and L β domain. Carboxyl terminal to the LE modules, the three laminin chains have a C-x-x-C motif (α 3A: C₂₀₂-C₂₀₅; β 3: C₅₈₁-C₅₈₂; γ 2: C₆₀₉-C₆₁₂) which cysteines are conserved in all laminins and most likely serve for interchain covalent linkage of the heterotrimer. The actual disulfide pattern has not yet been established. The following ~570 residues associate via formation of an α -helical coiled coil represented on the sequence level by a 3–4 heptad pattern of hydrophobic residues.³ As it is the case for all β chains, the β 3 coiled coil is interrupted by domain L β ~200 residues away from the C-x-x-C motif containing six cysteines (human β 3: C₇₈₈-C₈₁₈) probably forming intrachain disulfide bonds. The α 3 chain coiled coil contains one cysteine (human α 3: C₅₅₁) about 50 nm from the C-x-x-C motif, which has no adjacent counterpart in the other two chains. The coiled coil region is terminated by cysteines close to the C-terminus (human β 3: C₁₁₇₁; γ 2: C₁₁₈₄), which based on similarity to the β 1 and γ 1 chains most probably form an interchain disulfide bond.⁴⁸

Besides its structural importance for assembly, it has recently been found that in the absence of the LG domains the laminin 111 coiled coil has anti-adhesive properties. When cultured on such truncated laminin 111, cell adhesion and spreading were inhibited; genes compatible with a pro-migratory and pro-invasive function like MMPs and various matricellular proteins were upregulated.⁴⁷ No comparable studies have yet been performed for laminin 332, but an antagonistic interplay between coiled-coil




Figure 2. Structural model of the human laminin $\beta 3$ LN-LE1–4 domains. (A) The structure shows residues Q₁₈ to Q₄₆₄ as a cartoon with red cylinders and blue arrows indicating α -helices and β -strands, respectively. Disulfide bonded cysteines within the LE domains are depicted in ball-and-stick presentation using different colors. Domain boundaries are indicated by dashed lines. P₂₈ fits into a pocket shared by the LN and LE1 domain, and the N-terminal residues are interacting with LE1. Disulfides and the free cysteine residue of the LN domain are indicated. (B) The electrostatic surface potential is represented in the same orientation and on the same scale as in (A). Negative and positive potentials are shown in red and blue, respectively. Homology based modeling was performed using Swiss-Model¹⁰⁶ in manual alignment mode with the mouse $\beta 1$ structure (pdb code 4aqs⁴⁶) as a template.

and LG domains (see below) could be of importance in cancer metastasis.

Laminin $\alpha 3$ LG domains. The large globule seen in electron micrographs at the tip of the long arm consists of five LG domains formed by the α chains. Domains LG4–LG5 are separated from LG1 to LG3 by a flexible linker (Fig. 1B). As these domains are involved in many interactions (see below), particular interest has been focused on their structural characterization (for a review, see ref. 7). X-ray structures have been solved for the mouse $\alpha 5$ LG1–LG3 triplet (pdb code: 2wjs),⁴⁹ mouse $\alpha 1$ and $\alpha 2$ LG45 pair (2jd4,⁵⁰ 1dyk,²⁰ 1okq⁵¹), and mouse $\alpha 2$ LG5 (1qu0⁵²) which can

be used to model the corresponding $\alpha 3$ LG1–LG5 domains (Fig. 3). LG domains fold to a β -sandwich consisting of 14 strands, where the two sheets form a concave and convex interface. Close to the C-terminus, each LG sequence contains two cysteines ~ 30 residues apart, which form intradomain disulfide bonds (Fig. 3A and C). In contrast to electron micrographs suggesting close proximity of LG1 to LG3,^{53,54} the $\alpha 2$ LG1–3 structure shows LG1 dissociated from the LG2–LG3 pair.⁴⁹ Previous work suggested that trimerization of the α with β and γ chains is required for the LG domain to exert its integrin binding activity, and that specifically a glutamic acid two residues




Figure 3. For figure lenged, see next page.

Figure 3 (See previous page). Structural models of human laminin $\alpha 3$ LG domains. (A) The structure of LG1 to LG3 encompassing residues M₇₇₁ to S₁₃₀₈ is shown as a cartoon as described for Figure 2. All cysteine residues are depicted in space filling (CPK) style. The disulfide bridge between C₉₄₁ and C₉₇₁ (red dotted line) could not be correctly modeled as the template sequence has a shorter loop between the corresponding residues. All other cysteines form disulfide bridges except for C₈₄₀ in LG1 which side chain points into the inner space of the β -sandwich. (B) The electrostatic surface potential is represented in the same orientation and on the same scale as in (A). Negative and positive potentials are shown in red and blue, respectively. (C) A model of LG45 corresponding to residues C₁₃₅₄ to Q₁₇₁₃ is shown with all cysteines in CPK style. C₁₃₅₄ from the loop region connecting LG3 and LG4 forms an interdomain disulfide bond with C₁₆₁₇. C₁₄₄₃ in LG4 and C₁₅₈₇ in LG5 are pointed to the space between the β sheets of the sandwich structures and are not surface accessible, whereas C₁₅₀₇-C₁₅₃₀ and C₁₆₈₂-C₁₇₁₀ form disulfide bonds. (D) A model of LG4 interacting with a heparin-like carbohydrate chain consisting of four GlcNS(6S)-IdoA(2S) disaccharides (ball-and-stick representation with a surface envelope) is shown. The side chains of K₁₄₃₄, R₁₄₃₆, and K₁₄₃₈, which were shown to be critically involved in syndecan-1 interaction are depicted in red, whereas those of K₁₄₄₀ and K₁₄₄₂ involved in syndecan-4 interaction are shown in green.⁵⁹ The ϵ -amino and guanidinium groups of the lysine and arginine residues, respectively, point to the sulfate groups of the carbohydrate chains. (E) The surface view of the model shown in (D) indicates the tight packing of the heparin fragment into the groove formed by the concave surface of the LG4 β -sandwich with positive surface potential spaced to fit the distance of sulfate groups. The template structures used for modeling were mouse laminin $\alpha 2$ LG1-LG3 (pdb code: 2wjs)⁴⁹ and mouse laminin $\alpha 1$ LG4-LG5 (2jd4, chain B).⁵⁰ Docking of a heparin fragment (1hpn)⁶⁰ to LG4 was performed with PatchDock.¹⁰⁷

apart from the C-terminus of the γ chain is essential.⁵⁵ This residue is indeed highly conserved in the $\gamma 1$ and $\gamma 2$, but not $\gamma 3$ subunits. A lack of integrin $\alpha 3\beta 1$ binding activity in the absence of at least short $\beta 3$ and $\gamma 2$ segments has been shown for $\alpha 3$ LG1-LG3.^{54,56} The absence of such crucial parts of the β and γ chains could result in the open LG1-LG2 conformation (Fig. 3A and B) and might not reflect the conformation of the native molecule.⁴⁹

The LG3 and LG4 domains are connected by a flexible linker, which was realized very early by the proteolytic susceptibility of mouse laminin 111.⁴⁵ Based on the C- and N-terminal residues, which can be resolved in the mouse $\alpha 5$ LG1-3 (2wjs)⁴⁹ and $\alpha 1$ LG4-5 (2jd4)⁵⁰ structures, respectively, for human $\alpha 3$ the termini of the corresponding loop can be defined as N₁₃₁₀ and D₁₃₅₂ (Fig. 1B). Close to the middle, $\alpha 3$ LG is processed between Q₁₃₃₇ and D₁₃₃₈.^{17,57} This leaves a segment P₁₃₁₂PFLMLLKSTR attached to LG3 which, based on a recombinantly overexpressed LG and synthetic peptide approach, was reported as crucial for integrin $\alpha 3\beta 1$ binding.⁵⁸ C-terminally truncated peptides, however, abolished this activity. The difference of this part in mouse $\alpha 3$ (PPFLMLFKSPKG, Fig. 1B), however, would suggest that this interaction is rather species specific.

Based on the structures of the mouse $\alpha 1$ and $\alpha 2$ LG4-5 pairs, modeling predicts that human $\alpha 3$ C₁₃₅₄ within the loop region forms a disulfide bond with C₁₆₁₇ in LG5 (Fig. 3C). Corresponding cysteines within the loop and LG5 are found in all laminin α chains down to *Drosophila* and *C. elegans*. Interestingly, human $\alpha 3$ LG4 (C₁₄₄₃) and $\alpha 1$ LG3 contain further cysteines in a similar position, which are not conserved in mouse and rat. Within the model (Fig. 3C), the C₁₄₄₃ and C₁₅₈₇ SH groups point into the hydrophobic core of the β sandwich.

The crystal structures of mouse $\alpha 5$ LG1, LG2, and $\alpha 1$, $\alpha 2$ LG4, LG5 contain tightly bound calcium or magnesium ions interacting with pairs of aspartate residues.^{20,49-51} Our human $\alpha 3$ LG models (Fig. 3) suggest that these cation binding sites are not conserved. Although within LG1 and LG2 D₈₂₇ and D₁₀₉₀, respectively, are present, their partner residues are changed to N₉₁₀ and R₁₀₂₃. Within LG4 and LG5, the corresponding residues are replaced by S₁₄₁₃/S₁₄₈₁ and K₁₅₈₄/S₁₆₅₄. This does not exclude that $\alpha 3$ LG domains might contain other divalent cation binding sites, but so far no conclusive experimental data exist.

Due to their physiological relevance, many studies have been performed to elucidate the interaction of laminins with proteoglycans. As a model for the complex glycosaminoglycan chains, most experiments are performed with heparin. We have recently reported on the specific interaction of human $\alpha 3$ LG45 with syndecans and found that K₁₄₃₄, R₁₄₃₆ and K₁₄₃₈ are crucial for syndecan-1 binding, whereas K₁₄₄₀ and K₁₄₄₂ show a preference for syndecan-4 (see below).⁵⁹ Using the modeled $\alpha 3$ LG4 domain and a truncated heparin-like structure (pdb code 1hpn),⁶⁰ docking simulations predict that the carbohydrate chain aligns to the concave surface of LG4 with the positively charged residues filling the grooves formed by the repeating sulfates (Fig. 3D and E).

Functions of LG45 in Precursor $\alpha 3$

A potential function for the tandem LG45 domains was initially suspected based on the ability of laminin 332 to trigger distinct cellular events depending on the level of processing of its $\alpha 3$ chain. A form of laminin 332 that lacked LG45 was found in mature basement membranes, where it was shown to play an important function in the nucleation and maintenance of anchoring structures through $\alpha 3\beta 1$ and $\alpha 6\beta 4$ integrin interactions.⁶¹⁻⁶³ In contrast, laminin 332 with intact LG45 ($\alpha 3_{200}$) was found in migratory/remodelling situations such as epidermal repair.⁶⁴⁻⁶⁶ Indeed, laminin 332 with an $\alpha 3_{200}$ chain was found in the ECM of keratinocytes migrating on collagen I or after stimulation by TGF- $\beta 1$.^{67,68} In vivo, epidermal injury activates the transcription and deposition of laminin 332 into the provisional matrix by the leading keratinocytes in the process of epidermal outgrowth and migration at the wound edge.^{8,61,69} Noteworthy, $\alpha 3_{200}$ laminin 332 is found in this provisional matrix but is absent from mature basement membranes.^{16,61,62} Recently, laminin 332 comprising an $\alpha 3_{200}$ chain was proposed to be involved in the invasion of squamous cell carcinomas in vivo.⁷⁰ Polyclonal antibodies targeting the LG45 domains induced squamous cell carcinoma apoptosis in vivo and thus inhibited tumor proliferation.⁷⁰

Function in laminin 332/311 deposition into the ECM. A function for LG45 in the deposition of laminin 332 in the ECM has been first hypothesized by Carter and coworkers,¹⁶ who

showed that exogenous human $\alpha 3_{200}$ laminin 332, but not $\alpha 3_{165}$ laminin 332, was trapped by cultured LAMA3 deficient mouse keratinocytes and deposited within their ECM. Moreover, when these keratinocytes were transfected with constructs encoding either an LG45-uncleavable or a precleaved $\alpha 3$ chain, they deposited a larger amount of the LG45-uncleavable laminin 332 into their ECM as compared with the LG45 pre-cleaved laminin 332 that was preferentially secreted into the culture medium. The importance of LG45 in laminin 332 deposition was confirmed by a study reporting that a mutant laminin 332 lacking the LG45 domain, stably expressed in laminin 332 null keratinocytes derived from a patient with junctional epidermolysis bullosa (JEB) with underlying LAMA3 gene mutations, was less retained in the ECM but rather found in the culture medium as compared with the WT.⁷⁰ Co-expression of LG45 in these $\alpha 3_{165}$ laminin expressing keratinocytes enhanced laminin 332 deposition into the ECM, reinforced keratinocyte adhesion to the ECM and decreased migration. In these cells, LG45 co-localized with the $\alpha 6$ integrin subunit in stable adhesion contacts in a manner comparable to the LG45 domain of WT laminin 332 keratinocytes.⁷⁰

Another study, however, found no difference in laminin 332 deposited by keratinocytes expressing either LG45 uncleavable or pre-cleaved laminin 332.¹⁹ Skin equivalents epithelialized with keratinocytes expressing the $\alpha 3_{165}$ laminin 332 displayed a dermal-epidermal junction identical to that obtained with wild type keratinocytes suggesting that $\alpha 3_{165}$ laminin 332 was properly deposited into the basement membrane or at least in an amount sufficient to allow dermal-epidermal cohesion.¹⁹ Besides, skin equivalents designed with keratinocytes expressing the LG45 uncleavable laminin 332 revealed absence of hemidesmosomes accompanied by an appreciable thickening of the lamina lucida likely secondary to the reduced cohesion of the tissue.¹⁹ These experiments show that LG45 cleavage appears to be a prerequisite for hemidesmosome formation in skin equivalent models.

These studies show that LG45 has the ability to integrate into the keratinocyte ECM either on its own or when present in the $\alpha 3_{200}$ chain. It targets $\alpha 6\beta 4$ integrin containing stable adhesion contacts suggesting that it may play a role in $\alpha 6\beta 4$ integrin clustering in vitro. Although LG45 removal decreased laminin 332 deposition in the ECM, it did not fully prevent it, suggesting that other mechanisms may contribute in controlling laminin 332 deposition.

Defective $\alpha 3$ and $\gamma 2$ processing was seen in cylindromatosis, a rare genetic human disorder characterized by the occurrence of multiple irregular benign epithelial tumors in the upper dermis.⁶³ These nodules display a dramatically enlarged basement membrane (up to 4.3 μm thickness) as well as ultrastructural abnormalities as no clear lamina densa could be detected. The different integrin receptors are found in improper ratios as $\beta 1$ integrins are upregulated while $\alpha 6\beta 4$ integrin expression and hemidesmosome numbers are decreased.⁶³ Despite a massive accumulation of laminin 332 throughout the entire basal lamina, a thin labeling of the LG45 domain detected at the interface between cylindroma cells and the basal lamina suggests that the LG45 processing may have been delayed. It is not clear whether

laminin 332 accumulation in the basement membrane is related to this processing defect. Only one study reported a missense mutation in the LAMA3 gene affecting LG4 in a patient with a mild non-Herlitz JEB phenotype.⁷¹ This mutation resulting in G1506E triggers an imperfect local protein folding that, without impairing trimerization of the coiled coil, causes laminin 332 intracellular accumulation within the endoplasmic reticulum. Only a small amount of the laminin 332 harboring the mutated $\alpha 3$ chain is secreted and physiologically processed thus providing partial adhesion functions and explaining the mild phenotype. Therefore structural changes caused by mutations of this highly conserved residue throughout laminin LG4 domains highlights potential important functions of LG45 in laminin secretion.

When a laminin $\alpha 3$ cDNA was transfected into HT1080 cells, the exogenous $\alpha 3$ chain was assembled with the endogenous $\beta 3/\gamma 2$ and $\beta 1/\gamma 1$ to produce laminin 332 and laminin 311 heterotrimers.⁷² Out of the two laminin isoforms found in the culture medium, laminin 311 was found primarily with a precursor $\alpha 3_{200}$ chain while the laminin 332 was found with an $\alpha 3_{165}$ chain.⁷² Sigle et al.¹⁶ showed that exogenous human $\alpha 3_{200}$ laminin 311 was not trapped by cultured LAMA3 deficient mouse keratinocytes nor deposited within their ECM like $\alpha 3_{200}$ laminin 332 (see above). When these keratinocytes were transfected with constructs encoding an LG45-uncleavable $\alpha 3$ chain, they deposited large amount of LG45-uncleavable laminin 332 within their ECM, and the LG45-uncleavable laminin 311 was found exclusively in the culture medium.¹⁶ These results suggest that the two laminin isoforms integrate within the basement membrane through different mechanisms. That $\alpha 3_{200}$ laminin 311 was not deposited within the ECM first shows that LG45 could not fulfil this event on its own. It also reinforces the hypothesis that a protein domain present in laminin 332 but absent in laminin 311, possibly in the $\gamma 2$ chain,⁷³ may contribute to the laminin deposition.

Since laminin 311 forms a complex with laminin 332 in vivo, it is tempting to speculate that laminin 311 deposition within the ECM might depend on its covalent association with laminin 332. Most interestingly, co-polymers of perlecan with mature laminin 311 were identified in ECM formed by alveolar epithelial cells.^{74,75} Through nidogen linkage to the laminin 311 $\gamma 1$ chain, perlecan was proposed to nucleate formation of laminin 311 fibrils.

Function in cell adhesion and migration. It has been proposed that precursor laminin 332, together with integrin $\alpha 3\beta 1$, plays a central role in the polarization and migration of cells.^{12,66} Processing of the laminin 332 α chain was suggested to alter its binding ability to various integrins by differential exposure of binding sites/sequences to integrin receptors. Cell adhesion studies have shown that immobilized purified $\alpha 3_{200}$ laminin 332 promotes adhesion of keratinocytes, but to a lesser extent than that obtained with mature laminin 332.⁷⁶ Further analysis revealed that the $\alpha 3\beta 1$ integrin was the major receptor to interact with $\alpha 3_{200}$ laminin 332, while both $\alpha 3\beta 1$ and $\alpha 6\beta 4$ integrins were involved in cell adhesion to $\alpha 3_{165}$ laminin 332.⁷⁶ These findings correlate with previous results showing that cells plated on an $\alpha 3_{200}$ laminin 332 rich matrix failed to form $\alpha 6\beta 4$

Table 1. Heparin binding sites identified within human and mouse LG45 domains

Laminin $\alpha 3$ LG45 heparin binding sites	Method used for identification	Binding partner	References
Human			
N ₁₄₁₂ SFMALYLSKGR	Peptide	Syndecan-2 Syndecan-4	82
L ₁₄₂₈ GTDGKKLRIKSKEKNDG	LG45/heparin cross-link	Heparin	85
K ₁₄₃₃ KLRIK	Directed mutagenesis	Syndecan-1	59
R ₁₄₃₆ IKSKEK	Directed mutagenesis	Syndecan-4	59
L ₁₄₈₈ GSPPSGKPKSL	LG45/heparin cross-link	Heparin	85
V ₁₆₁₀ TPKQSLCD	LG45/heparin cross-link	Heparin	85
Mouse			
K ₁₃₃₆ ARFNVNQLLQD	Peptide	Unknown receptor	81
K ₁₃₉₈ PRLQFSLDIQT	Peptide	Unknown receptor	81
D ₁₆₂₉ GQWHSVTSIK	Peptide	Unknown receptor	81

integrin-dependent hemidesmosomes, whereas plasmin cleavage of the LG45 domain inhibited migration and promoted hemidesmosome formation.¹² These data were corroborated by studies showing that $\alpha 3_{200}$ laminin 332 is a preferential ligand for integrin $\alpha 3\beta 1$ and/or $\alpha 6\beta 1$, whereas processed $\alpha 3$ LG domains are the preferred ligand for integrin $\alpha 6\beta 4$.⁶⁵ Besides, the analysis of the binding capacity of soluble recombinant $\alpha 3\beta 1$ integrin to a mini-laminin 332 comprising the five LG domains has revealed an increased affinity for integrin binding when LG45 was cleaved off.⁵⁶ Information gained from the structural model of the LG predicts that LG1–3 have the shape of a cloverleaf from which LG45 extends.⁷ The specific length of the LG3-LG4 linker results in LG5 being closer to the LG1–3 cloverleaf than LG4 suggesting that the integrin-binding domain may be differently exposed in $\alpha 3_{200}$ and $\alpha 3_{165}$ laminin 332. Other studies have contradicted this hypothesis as both $\alpha 6\beta 4$ and $\alpha 3\beta 1$ integrins were found to interact with an LG45-uncleavable laminin 332¹⁶ and were colocalized in precursor laminin 332 trails left behind migrating cells.⁵⁷ Moreover, $\alpha 6\beta 4$ integrin containing stable anchoring complexes⁷⁷ were detected in cultured keratinocytes expressing precursor- or LG45-uncleavable laminin 332^{16,19} in which LG45 and $\alpha 6\beta 4$ integrins colocalized.⁷⁰ Although the $\beta 4$ integrin was correctly expressed and localized at the apical side of the epidermal basal layer, LG45-uncleavable laminin 332 expressing keratinocytes failed to form hemidesmosomes when seeded over a dermal equivalent model.¹⁹

Other studies have reported that soluble LG45 may have a function when released from laminin 332. Experiments have shown that adding either recombinant $\alpha 3$ LG45 or LG4 to the culture medium could induce migration of keratinocytes in a MMP-9-dependent manner.⁷⁸ In similar experiments, LG4 induced activation of pro-migratory MMP-1.⁷⁹ Moreover, Ras/I κ B α transformed keratinocytes that lacked the LG45 domain showed a deficiency in MMP-9 and MMP-1 expression. This deficiency was reversed by replacing LG45 through retroviral transduction.⁷⁰

Binding sites within $\alpha 3$ LG45. The mechanism underlying the function of the LG45 domain in laminin 332 remains poorly understood. Several heparin-binding sites have been identified in

the $\alpha 3$ LG45 sequence (Table 1). Some of these conferred heparin-dependent cell adhesion properties, which suggested that this region in laminin 332 could interact with a heparan sulfate proteoglycan receptor.^{29,59} An early report described the characterization of a heparin-binding synthetic peptide corresponding to residues K₁₃₉₈PRLQFSLDIQT derived from the murine $\alpha 1$ LG4 sequence.⁸⁰ This peptide induces adhesion of the melanoma cells B16F10 through an heparan sulfate proteoglycan type receptor that remains to be identified. A peptidic screening of the murine $\alpha 3$ LG45 domains allowed identification of three heparin binding sites with cell adhesion properties.⁸¹ Peptidic screening of the human sequence lead to the identification of a motif that included residues N₁₄₁₂SFMALYLSKGR, which was shown to induce syndecan-2 and -4 mediated cell adhesion, neurite outgrowth and MMP-1 and -9 secretion.^{78,79,82,83} Further work suggested that this motif also induced keratinocyte migration by triggering syndecan-4 clustering and subsequent $\beta 1$ integrin activation.⁸⁴ Later, three novel heparin-binding sites were identified based on cross-linking the native protein to heparin beads.⁸⁵ Recently, through a site-directed mutagenesis approach to alter the most critical basic residues in a recombinant LG45 protein, we identified a unique heparin-binding site surrounded by a track of converging low affinity, positively charged residues (Fig. 3D and E).⁴⁹ We further showed that this K₁₄₃₃KLRIKSKEK sequence region, which matches one of the sites identified by Vives et al.⁸⁵ harbours distinctive syndecan-1 and -4 interaction sites (Table 1). Besides, our group has reported that syndecan-1 is the cellular receptor involved in cell adhesion to the $\alpha 3$ LG45 domain^{57,86} and this interaction may participate in keratinocyte migration by supporting the formation of actin-based cellular protrusions (Fig. 4A and B).^{76,87} The development of these membrane protrusions remarkably requires dephosphorylation of tyrosine residues in the cytoplasmic tail of syndecan-1, a condition essential for syntenin-1 recruitment.^{87,88} In the epidermis, syndecan-1 is located in the pericellular region of keratinocytes and displays a modest expression in the basal cell layer, which becomes increasingly intense in the suprabasal layers. Remarkably, syndecan-1 is strongly induced in wound edge keratinocytes during wound healing.⁸⁹⁻⁹¹ This elevated expression

in keratinocytes appears to be specific to syndecan-1 as no changes have been detected for other syndecans.⁹² Moreover, syndecan-1 deficient mice display a defect in keratinocyte proliferation and migration during wound healing.⁹³ In light of these data, syndecan family members stimulate interest as potential laminin 332 co-receptors. Recent data has provided evidence that when human squamous carcinoma A431 cells are plated on laminin 332, syndecan-1 forms a complex with the $\alpha 6 \beta 4$ integrin. This triggers Fyn-mediated phosphorylation of the $\beta 4$ integrin cytoplasmic tail activating PI3K- and Akt-mediated signaling, protecting the cells against apoptosis.⁹⁴ This is of particular interest as LG45 in $\alpha 3_{200}$ laminin 332 was proposed as an essential PI3K pathway activation promoter.⁷⁰

Functions of Precursor $\gamma 2$ N-Terminal Domain

It has been shown that in human skin basement membrane, the N-terminal domain of the $\gamma 2$ chain is absent suggesting that the $\gamma 2$ processing could have an important physiological role.^{11,63} In contrast to these findings, antibodies specific for the released L4 domain were shown to label the basal epidermal cells in mouse skin.²⁴ A positive signal was also found in the basement membranes of early stages of skin development suggesting that the precursor $\gamma 2$ chain or its released N-terminal domain expression might persist *in vivo*.²⁴ The analysis of mTLD/BMP-1 deficient mice revealed that although defaults in skin cohesion were not clinically apparent, ultrastructural examination of the basement membrane showed places where anchoring structures were disconnected at the level of the lamina densa with presence of rudimentary hemidesmosomes.^{15,25} A strong expression of the precursor $\gamma 2$ chain suggested that $\gamma 2$ chain maturation takes part in the basement membrane cohesion and stability. Besides,

defective $\gamma 2$ processing was found in the rare inherited cylindromatosis disease (see above), in which abnormal basement membranes and excessive ectopic expression of collagen VII were found.^{63,96} The defective processing of the $\gamma 2$ chain may have resulted in a deficient interaction with its basement membrane partners collagen VII causing defaults in anchoring fibrils linkage.^{32,95}

Numerous studies conducted with normal breast epithelial and carcinoma cells have shown that cell migration-inducing functions of the $\gamma 2$ chain rely on its proteolysis.³⁵ Recently, it has been reported that a syndecan-1 interaction with the N-terminal domain in precursor $\gamma 2$ could negatively regulate $\beta 4$ integrin phosphorylation leading to enhanced carcinoma cell adhesion and decreased motility⁹⁷ reinforcing the idea that the precursor $\gamma 2$ chain does not support cell migration. In contrast, other reports have suggested that precursor $\gamma 2$ may play a role in keratinocyte migration during the wound healing process. An interaction of its N-terminal portion with $\alpha 2 \beta 1$ integrin was shown to occur during TGF- $\beta 1$ induced keratinocyte migration *in vitro* and was proposed to take place at the wound margin of the skin healing process *in vivo*.⁶⁸ Keratinocytes at the edge of wounds made in early passage cultures shown to co-express the laminin 332 $\gamma 2$ chain and the cell cycle inhibitor p16^{INK4a}, displayed increased directional motility.⁹⁸ Additional studies revealed that this keratinocyte hypermotility/growth arrest response relied on both the precursor state of the laminin 332 $\gamma 2$ chain and the participation of a serum co-factor through a TGF β receptor I dependent mechanism.⁹⁹ This apparent contradiction in experimental findings is likely due to tissue specific differences. A study indicated that the L4 module mediates the integration of laminin 332 into the extracellular matrix through intermolecular interactions.⁷³

The question arises whether the heparin-binding domains located within the $\alpha 3$ LG45 domains and the N-terminal portion of the $\gamma 2$ chain are involved in laminin 332 deposition. Heparin was shown to inhibit laminin 332 deposition within the ECM suggesting that a heparan sulfate proteoglycan might be involved in binding to both precursor $\alpha 3$ and $\gamma 2$.¹⁵ Null mutation in syndecan-1¹⁰⁰ and -4¹⁰¹ has not reported defects in laminin deposition, which suggests that any of these two receptors is solely responsible for this mechanism. Generation of double syndecan-1 and -4 knockout mice would reveal whether both these receptors are implicated. Future studies should examine whether a heparan sulfate proteoglycan cell surface receptor or a basement membrane components are implicated in precursor laminin 332 deposition. Interestingly, a recent study has reported that the perlecan colocalizes with laminin 332 at the wound margin of healing full thickness wounds generated in mice suggesting proximity of these two components during basement membrane remodelling.¹⁰² However, as perlecan was shown not to be essential for matrix assembly in perlecan-null mouse embryos,¹⁰³ its presence in the ECM might not be an absolute requirement for laminin 332 deposition. Considering the important function of laminin 332 in skin homeostasis, several extracellular ligands might be involved in its deposition in the basement membrane.




Figure 4. Syndecan-1 mediated cell adhesion to laminin 332 LG45 domain. (A) Fibrosarcoma HT1080 cells were plated on surfaces coated with purified LG45 domain for 1 h, fixed and stained with phalloidin-FITC. The analysis of the actin staining revealed an organization of the cytoskeleton in radial arrays of microspikes, protrusive adhesion structures known to be involved in transient cell-substratum interactions. (B) Confocal image of HT1080 cells expressing GFP-syndecan-1 plated on the LG45 domain. The phase contrast image shows the typical induced cell shape characterized by the formation of spike-like actin-based protrusive structures. The receptor syndecan-1 (green) promotes cell adhesion to the laminin fragment and aligns along the actin filaments, which are organized into parallel bundles. Bars, 10 μ m.

Conclusions

Comprehension of the physiological significance of laminin 332 maturation events remains an open issue. Numerous studies have focused on the LG45 and γ 2 N-terminal domains suggesting their involvement in diverse important functions such as laminin secretion, deposition and/or retention within the ECM; others have suggested a participation in epithelial cell adhesion and migration processes. However, their exact function remains unknown. While in vitro data sometimes disagree due to the cellular system's diversity, in vivo models firmly demonstrate that defects in laminin 332 maturation impede correct basement assembly as well as hemidesmosome formation. Molecular interaction abnormalities occur at the level of cellular receptors and basement membrane components causing both functional and organisational anomalies. Molecular characterization and structural definition of laminin domains open interesting and

every so often unexpected questions. From its identification in the 1990s^{9,26,104,105} to nowadays, laminin 332 has stimulated incessantly increasing interest in the scientific community due to its multifunctional properties and its involvement in human physiological processes and pathologies.

Disclosure of Potential Conflicts of Interest

No potential conflicts of interest were disclosed.

Acknowledgments

We thank Dr. François Letourneur (CGMC, Lyon) for his help in designing the syndecan-1 GFP constructs. Original work by the authors was financially supported by the Agence Nationale pour la Recherche grants Chemispiké and adhesionMab, the Association pour la Recherche sur le Cancer and the Ligue Nationale contre le Cancer. We acknowledge the contribution of the PLATIM platform of SFR Biosciences Gerland-Lyon Sud (UMS344/US8).

References

- Domogatskaya A, Rodin S, Tryggvason K. Functional diversity of laminins. *Annu Rev Cell Dev Biol* 2012; 28:523-53; PMID:23057746; <http://dx.doi.org/10.1146/annurev-cellbio-101011-155750>
- Miner JH, Yurchenco PD. Laminin functions in tissue morphogenesis. *Annu Rev Cell Dev Biol* 2004; 20:255-84; PMID:15473841; <http://dx.doi.org/10.1146/annurev.cellbio.20.010403.094555>
- Beck K, Dixon TW, Engel J, Parry DA. Ionic interactions in the coiled-coil domain of laminin determine the specificity of chain assembly. *J Mol Biol* 1993; 231:311-23; PMID:8510149; <http://dx.doi.org/10.1006/jmbi.1993.1284>
- Matsui C, Wang CK, Nelson CF, Bauer EA, Hoeffler WK. The assembly of laminin-5 subunits. *J Biol Chem* 1995; 270:23496-503; PMID:7559513; <http://dx.doi.org/10.1074/jbc.270.40.23496>
- Durbeq J. Laminins. *Cell Tissue Res* 2010; 339:259-68; PMID:19693542; <http://dx.doi.org/10.1007/s00441-009-0838-2>
- Aumailley M, Bruckner-Tuderman L, Carter WG, Deutzmann R, Edgar D, Ekblom P, et al. A simplified laminin nomenclature. *Matrix Biol* 2005; 24:326-32; PMID:15979864; <http://dx.doi.org/10.1016/j.matbio.2005.05.006>
- Timpl R, Tisi D, Talts JF, Andac Z, Sasaki T, Hohenester E. Structure and function of laminin LG modules. *Matrix Biol* 2000; 19:309-17; PMID:10963991; [http://dx.doi.org/10.1016/S0945-053X\(00\)00072-X](http://dx.doi.org/10.1016/S0945-053X(00)00072-X)
- Ryan MC, Tizard R, VanDevanter DR, Carter WG. Cloning of the LamA3 gene encoding the α 3 chain of the adhesive ligand epiligrin. Expression in wound repair. *J Biol Chem* 1994; 269:22779-87; PMID:8077230
- Rousselle P, Lunstrum GP, Keene DR, Burgeson RE. Kalinin: an epithelium-specific basement membrane adhesion molecule that is a component of anchoring filaments. *J Cell Biol* 1991; 114:567-76; PMID:1860885; <http://dx.doi.org/10.1083/jcb.114.3.567>
- Rousselle P, Keene DR, Ruggiero F, Champlaud MF, Rest M, Burgeson RE. Laminin 5 binds the NC-1 domain of type VII collagen. *J Cell Biol* 1997; 138:719-28; PMID:9245798; <http://dx.doi.org/10.1083/jcb.138.3.719>
- Marinkovich MP, Lunstrum GP, Burgeson RE. The anchoring filament protein kalinin is synthesized and secreted as a high molecular weight precursor. *J Biol Chem* 1992; 267:17900-6; PMID:1517226
- Goldfinger LE, Stack MS, Jones JC. Processing of laminin-5 and its functional consequences: role of plasmin and tissue-type plasminogen activator. *J Cell Biol* 1998; 141:255-65; PMID:9531563; <http://dx.doi.org/10.1083/jcb.141.1.255>
- Tsubota Y, Yasuda C, Kariya Y, Ogawa T, Hirotsuki T, Mizushima H, et al. Regulation of biological activity and matrix assembly of laminin-5 by COOH-terminal, LG4-5 domain of α 3 chain. *J Biol Chem* 2005; 280:14370-7; PMID:15695818; <http://dx.doi.org/10.1074/jbc.M413051200>
- Amano S, Scott IC, Takahara K, Koch M, Champlaud MF, Gerecke DR, et al. Bone morphogenetic protein 1 is an extracellular processing enzyme of the laminin 5 γ 2 chain. *J Biol Chem* 2000; 275:22728-35; PMID:10806203; <http://dx.doi.org/10.1074/jbc.M002345200>
- Veitch DP, Nokelainen P, McGowan KA, Nguyen TT, Nguyen NE, Stephenson R, et al. Mammalian tolloid metalloproteinase, and not matrix metalloproteinase 2 or membrane type 1 metalloproteinase, processes laminin-5 in keratinocytes and skin. *J Biol Chem* 2003; 278:15661-8; PMID:12473650; <http://dx.doi.org/10.1074/jbc.M210588200>
- Sigle RO, Gil SG, Bhattacharya M, Ryan MC, Yang TM, Brown TA, et al. Globular domains 4/5 of the laminin α 3 chain mediate deposition of precursor laminin 5. *J Cell Sci* 2004; 117:4481-94; PMID:15316072; <http://dx.doi.org/10.1242/jcs.01310>
- Tsubota Y, Mizushima H, Hirotsuki T, Higashi S, Yasumitsu H, Miyazaki K. Isolation and activity of proteolytic fragment of laminin-5 α 3 chain. *Biochem Biophys Res Commun* 2000; 278:614-20; PMID:11095958; <http://dx.doi.org/10.1006/bbrc.2000.3851>
- Décline F, Okamoto O, Mallein-Gerin F, Helbert B, Bernaud J, Rigal D, et al. Keratinocyte motility induced by TGF-beta1 is accompanied by dramatic changes in cellular interactions with laminin 5. *Cell Motil Cytoskeleton* 2003; 54:64-80; PMID:12451596; <http://dx.doi.org/10.1002/cm.10086>
- Baudoin C, Fantin L, Meneguzzi G. Proteolytic processing of the laminin α 3 G domain mediates assembly of hemidesmosomes but has no role on keratinocyte migration. *J Invest Dermatol* 2005; 125:883-8; PMID:16297184; <http://dx.doi.org/10.1111/j.0022-202X.2005.23881.x>
- Tisi D, Talts JF, Timpl R, Hohenester E. Structure of the C-terminal laminin G-like domain pair of the laminin α 2 chain harbouring binding sites for α -dystroglycan and heparin. *EMBO J* 2000; 19:1432-40; PMID:10747011; <http://dx.doi.org/10.1093/emboj/19.7.1432>
- Champlaud MF, Lunstrum GP, Rousselle P, Nishiyama T, Keene DR, Burgeson RE. Human amnion contains a novel laminin variant, laminin 7, which like laminin 6, covalently associates with laminin 5 to promote stable epithelial-stromal attachment. *J Cell Biol* 1996; 132:1189-98; PMID:8601594; <http://dx.doi.org/10.1083/jcb.132.6.1189>
- McLean WH, Irvine AD, Hamill KJ, Whittock NV, Coleman-Campbell CM, Mellerio JE, et al. An unusual N-terminal deletion of the laminin α 3a isoform leads to the chronic granulation tissue disorder laryngo-onycho-cutaneous syndrome. *Hum Mol Genet* 2003; 12:2395-409; PMID:12915477; <http://dx.doi.org/10.1093/hmg/ddg234>
- Vailly J, Verrando P, Champlaud MF, Gerecke D, Wagman DW, Baudoin C, et al. The 100-kDa chain of nicein/kalinin is a laminin B2 chain variant. *Eur J Biochem* 1994; 219:209-18; PMID:8306988; <http://dx.doi.org/10.1111/j.1432-1033.1994.tb19932.x>
- Sasaki T, Göhring W, Mann K, Brakebusch C, Yamada Y, Fässler R, et al. Short arm region of laminin-5 γ 2 chain: structure, mechanism of processing and binding to heparin and proteins. *J Mol Biol* 2001; 314:751-63; PMID:11733994; <http://dx.doi.org/10.1006/jmbi.2001.5176>
- Suzuki N, Labosky PA, Furuta Y, Hargett L, Dunn R, Fogo AB, et al. Failure of ventral body wall closure in mouse embryos lacking a procollagen C-proteinase encoded by Bmp1, a mammalian gene related to Drosophila tolloid. *Development* 1996; 122:3587-95; PMID:8951074
- Carter WG, Ryan MC, Gahr PJ. Epiligrin, a new cell adhesion ligand for integrin α 3 β 1 in epithelial basement membranes. *Cell* 1991; 65:599-610; PMID:2032285; [http://dx.doi.org/10.1016/0092-8674\(91\)90092-D](http://dx.doi.org/10.1016/0092-8674(91)90092-D)
- Sonnenberg A, de Melker AA, Martínez de Velasco AM, Janssen H, Calafat J, Niessen CM. Formation of hemidesmosomes in cells of a transformed murine mammary tumor cell line and mechanisms involved in adherence of these cells to laminin and kalinin. *J Cell Sci* 1993; 106:1083-102; PMID:8126093
- Rousselle P, Aumailley M. Kalinin is more efficient than laminin in promoting adhesion of primary keratinocytes and some other epithelial cells and has a different requirement for integrin receptors. *J Cell Biol* 1994; 125:205-14; PMID:8138572; <http://dx.doi.org/10.1083/jcb.125.1.205>

29. Mizushima H, Takamura H, Miyagi Y, Kikkawa Y, Yamanaka N, Yasumitsu H, et al. Identification of integrin-dependent and -independent cell adhesion domains in COOH-terminal globular region of laminin-5 α 3 chain. *Cell Growth Differ* 1997; 8: 979-87; PMID:9300180
30. Cheng YS, Champlaud MF, Burgeson RE, Marinkovich MP, Yurchenco PD. Self-assembly of laminin isoforms. *J Biol Chem* 1997; 272:31525-32; PMID:9395489; <http://dx.doi.org/10.1074/jbc.272.50.31525>
31. Burgeson RE. Type VII collagen, anchoring fibrils, and epidermolysis bullosa. *J Invest Dermatol* 1993; 101: 252-5; PMID:8370960; <http://dx.doi.org/10.1111/1523-1747.ep12365129>
32. Chen M, Marinkovich MP, Veis A, Cai X, Rao CN, O'Toole EA, et al. Interactions of the amino-terminal noncollagenous (NC1) domain of type VII collagen with extracellular matrix components. A potential role in epidermal-dermal adherence in human skin. *J Biol Chem* 1997; 272:14516-22; PMID:9169408; <http://dx.doi.org/10.1074/jbc.272.23.14516>
33. Giannelli G, Falk-Marzillier J, Schiraldi O, Stetler-Stevenson WG, Quaranta V. Induction of cell migration by matrix metalloproteinase-2 cleavage of laminin-5. *Science* 1997; 277:225-8; PMID:9211848; <http://dx.doi.org/10.1126/science.277.5323.225>
34. Koshikawa N, Giannelli G, Cirulli V, Miyazaki K, Quaranta V. Role of cell surface metalloproteinase MT1-MMP in epithelial cell migration over laminin-5. *J Cell Biol* 2000; 148:615-24; PMID:10662785; <http://dx.doi.org/10.1083/jcb.148.3.615>
35. Guess CM, Quaranta V. Defining the role of laminin-332 in carcinoma. *Matrix Biol* 2009; 28:445-55; PMID:19686849; <http://dx.doi.org/10.1016/j.matbio.2009.07.008>
36. Tsuruta D, Kobayashi H, Imanishi H, Sugawara K, Ishii M, Jones JC. Laminin-332-integrin interaction: a target for cancer therapy? *Curr Med Chem* 2008; 15: 1968-75; PMID:18691052; <http://dx.doi.org/10.2174/092986708785132834>
37. Pirtilä E, Sharabi A, Salo T, Quaranta V, Tu H, Heljasvaara R, et al. Matrix metalloproteinases process the laminin-5 γ 2-chain and regulate epithelial cell migration. *Biochem Biophys Res Commun* 2003; 303: 1012-7; PMID:12684035; [http://dx.doi.org/10.1016/S0006-291X\(03\)00452-2](http://dx.doi.org/10.1016/S0006-291X(03)00452-2)
38. Sadowski T, Dietrich S, Koschinsky F, Ludwig A, Proksch E, Titz B, et al. Matrix metalloproteinase 19 processes the laminin 5 γ 2 chain and induces epithelial cell migration. *Cell Mol Life Sci* 2005; 62: 870-80; PMID:15868410; <http://dx.doi.org/10.1007/s00018-005-4478-8>
39. Schenk S, Hintermann E, Bilban M, Koshikawa N, Hojilla C, Khokha R, et al. Binding to EGF receptor of a laminin-5 EGF-like fragment liberated during MMP-dependent mammary gland involution. *J Cell Biol* 2003; 161:197-209; PMID:12695504; <http://dx.doi.org/10.1083/jcb.200208145>
40. Udayakumar TS, Chen ML, Bair EL, Von Bredow DC, Cress AE, Nagle RB, et al. Membrane type-1-matrix metalloproteinase expressed by prostate carcinoma cells cleaves human laminin-5 β 3 chain and induces cell migration. *Cancer Res* 2003; 63:2292-9; PMID:12727852
41. Remy L, Trespeuch C, Bachy S, Scoazec JY, Rousselle P. Matrilysin 1 influences colon carcinoma cell migration by cleavage of the laminin-5 β 3 chain. *Cancer Res* 2006; 66:11228-37; PMID:17145868; <http://dx.doi.org/10.1158/0008-5472.CAN-06-1187>
42. Tripathi M, Potdar AA, Yamashita H, Weidow B, Cummings PT, Kirchhofer D, et al. Laminin-332 cleavage by matriptase alters motility parameters of prostate cancer cells. *Prostate* 2011; 71:184-96; PMID:20672321; <http://dx.doi.org/10.1002/pros.21233>
43. Marinkovich MP. Tumour microenvironment: laminin 332 in squamous-cell carcinoma. *Nat Rev Cancer* 2007; 7:370-80; PMID:17457303; <http://dx.doi.org/10.1038/nrc2089>
44. Rousselle P, Golbik R, van der Rest M, Aumailley M. Structural requirement for cell adhesion to kalinin (laminin-5). *J Biol Chem* 1995; 270:13766-70; PMID:7775432; <http://dx.doi.org/10.1074/jbc.270.23.13766>
45. Ott U, Odermatt E, Engel J, Furthmayr H, Timpl R. Protease resistance and conformation of laminin. *Eur J Biochem* 1982; 123:63-72; PMID:7040076; <http://dx.doi.org/10.1111/j.1432-1033.1982.tb06499.x>
46. Carafoli F, Hussain SA, Hohenester E. Crystal structures of the network-forming short-arm tips of the laminin β 1 and γ 1 chains. *PLoS One* 2012; 7: e42473; PMID:22860131; <http://dx.doi.org/10.1371/journal.pone.0042473>
47. Santos-Valle P, Guijarro-Muñoz I, Cuesta AM, Alonso-Camino V, Villate M, Alvarez-Cienfuegos A, et al. The heterotrimeric laminin coiled-coil domain exerts anti-adhesive effects and induces a pro-invasive phenotype. *PLoS One* 2012; 7:e39097; PMID:22723936; <http://dx.doi.org/10.1371/journal.pone.0039097>
48. Paulsson M, Deutzmann R, Timpl R, Dalzoppo D, Odermatt E, Engel J. Evidence for coiled-coil α -helical regions in the long arm of laminin. *EMBO J* 1985; 4: 309-16; PMID:3848400
49. Carafoli F, Clout NJ, Hohenester E. Crystal structure of the LG1-3 region of the laminin α 2 chain. *J Biol Chem* 2009; 284:22786-92; PMID:19553699; <http://dx.doi.org/10.1074/jbc.M109.026658>
50. Harrison D, Hussain SA, Combs AC, Ervasti JM, Yurchenco PD, Hohenester E. Crystal structure and cell surface anchorage sites of laminin α 1LG4-5. *J Biol Chem* 2007; 282:11573-81; PMID:17307732; <http://dx.doi.org/10.1074/jbc.M610657200>
51. Wizemann H, Garbe JH, Friedrich MV, Timpl R, Sasaki T, Hohenester E. Distinct requirements for heparin and α -dystroglycan binding revealed by structure-based mutagenesis of the laminin α 2 LG4-LG5 domain pair. *J Mol Biol* 2003; 332:635-42; PMID:12963372; [http://dx.doi.org/10.1016/S0022-2836\(03\)00848-9](http://dx.doi.org/10.1016/S0022-2836(03)00848-9)
52. Hohenester E, Tisi D, Talts JF, Timpl R. The crystal structure of a laminin G-like module reveals the molecular basis of α -dystroglycan binding to laminins, perlecan, and agrin. *Mol Cell* 1999; 4:783-92; PMID:10619025; [http://dx.doi.org/10.1016/S1097-2765\(00\)80388-3](http://dx.doi.org/10.1016/S1097-2765(00)80388-3)
53. Beck K, Hunter I, Engel J. Structure and function of laminin: anatomy of a multidomain glycoprotein. *FASEB J* 1990; 4:148-60; PMID:2404817
54. Navdaev A, Heitmann V, Desantana Evangelista K, Mörgelin M, Wegener J, Eble JA. The C-terminus of the γ 2 chain but not of the β 3 chain of laminin-332 is indirectly but indispensably necessary for integrin-mediated cell reactions. *Exp Cell Res* 2008; 314:489-97; PMID:18045589; <http://dx.doi.org/10.1016/j.yexcr.2007.10.027>
55. Ido H, Nakamura A, Kobayashi R, Ito S, Li S, Futaki S, et al. The requirement of the glutamic acid residue at the third position from the carboxyl termini of the laminin γ chains in integrin binding by laminins. *J Biol Chem* 2007; 282:11144-54; PMID:17307733; <http://dx.doi.org/10.1074/jbc.M609402200>
56. Künneken K, Pohlentz G, Schmidt-Hederich A, Odenthal U, Smyth N, Peter-Katalinic J, et al. Recombinant human laminin-5 domains. Effects of heterotrimerization, proteolytic processing, and N-glycosylation on α 3 β 1 integrin binding. *J Biol Chem* 2004; 279:5184-93; PMID:14612440; <http://dx.doi.org/10.1074/jbc.M310424200>
57. Décline F, Okamoto O, Mallein-Gerin F, Helbert B, Bernaud J, Rigal D, et al. Keratinocyte motility induced by TGF- β 1 is accompanied by dramatic changes in cellular interactions with laminin 5. *Cell Motil Cytoskeleton* 2003; 54:64-80; PMID:12451596; <http://dx.doi.org/10.1002/cm.10086>
58. Kim JM, Park WH, Min BM. The PPFLLMLLKGSTR motif in globular domain 3 of the human laminin-5 α 3 chain is crucial for integrin α 3 β 1 binding and cell adhesion. *Exp Cell Res* 2005; 304:317-27; PMID:15707596; <http://dx.doi.org/10.1016/j.yexcr.2004.11.009>
59. Carulli S, Beck K, Dayan G, Boulesteix S, Lortat-Jacob H, Rousselle P. Cell surface proteoglycans syndecan-1 and -4 bind overlapping but distinct sites in laminin α 3 LG45 protein domain. *J Biol Chem* 2012; 287: 12204-16; PMID:22351752; <http://dx.doi.org/10.1074/jbc.M111.300061>
60. Mulloy B, Forster MJ, Jones C, Davies DB. N.m.r. and molecular-modelling studies of the solution conformation of heparin. *Biochem J* 1993; 293:849-58; PMID:8352752
61. Lampe PD, Nguyen BP, Gil S, Usui M, Olerud J, Takada Y, et al. Cellular interaction of integrin α 3 β 1 with laminin 5 promotes gap junctional communication. *J Cell Biol* 1998; 143:1735-47; PMID:9852164; <http://dx.doi.org/10.1083/jcb.143.6.1735>
62. Goldfinger LE, Hopkinson SB, deHart GW, Collawn S, Couchman JR, Jones JC. The α 3 laminin subunit, α 6 β 4 and α 3 β 1 integrin coordinately regulate wound healing in cultured epithelial cells and in the skin. *J Cell Sci* 1999; 112:2615-29; PMID:10413670
63. Tunggul L, Ravaux J, Pesch M, Smola H, Krieg T, Gaill F, et al. Defective laminin 5 processing in cylindroma cells. *Am J Pathol* 2002; 160:459-68; PMID:11839566; [http://dx.doi.org/10.1016/S0002-9440\(10\)64865-1](http://dx.doi.org/10.1016/S0002-9440(10)64865-1)
64. Ryan MC, Lee K, Miyashita Y, Carter WG. Targeted disruption of the LAMA3 gene in mice reveals abnormalities in survival and late stage differentiation of epithelial cells. *J Cell Biol* 1999; 145:1309-23; PMID:10366601; <http://dx.doi.org/10.1083/jcb.145.6.1309>
65. Hintermann E, Quaranta V. Epithelial cell motility on laminin-5: regulation by matrix assembly, proteolysis, integrins and erbB receptors. *Matrix Biol* 2004; 23:75-85; PMID:15246107; <http://dx.doi.org/10.1016/j.matbio.2004.03.001>
66. Frank DE, Carter WG. Laminin 5 deposition regulates keratinocyte polarization and persistent migration. *J Cell Sci* 2004; 117:1351-63; PMID:14996912; <http://dx.doi.org/10.1242/jcs.01003>
67. Nguyen BP, Ryan MC, Gil SG, Carter WG. Deposition of laminin 5 in epidermal wounds regulates integrin signaling and adhesion. *Curr Opin Cell Biol* 2000; 12:554-62; PMID:10978889; [http://dx.doi.org/10.1016/S0955-0674\(00\)00131-9](http://dx.doi.org/10.1016/S0955-0674(00)00131-9)
68. Decline F, Rousselle P. Keratinocyte migration requires α 2 β 1 integrin-mediated interaction with the laminin 5 γ 2 chain. *J Cell Sci* 2001; 114:811-23; PMID:11171386
69. Kainulainen T, Hakkinen L, Hamidi S, Larjava K, Kallioinen M, Peltonen J, et al. Laminin-5 expression is independent of the injury and the microenvironment during reepithelialization of wounds. *J Histochem Cytochem* 1998; 46:353-60; PMID:9487117; <http://dx.doi.org/10.1177/002215549804600309>
70. Tran M, Rousselle P, Nokelainen P, Tallapragada S, Nguyen NT, Fincher EF, et al. Targeting a tumor-specific laminin domain critical for human carcinogenesis. *Cancer Res* 2008; 68:2885-94; PMID:18413757; <http://dx.doi.org/10.1158/0008-5472.CAN-07-6160>

71. Scaturro M, Posteraro P, Mastrogiacomo A, Zaccaria ML, De Luca N, Mazzanti C, et al. A missense mutation (G1506E) in the adhesion G domain of laminin-5 causes mild junctional epidermolysis bullosa. *Biochem Biophys Res Commun* 2003; 309:96-103; PMID:12943669; [http://dx.doi.org/10.1016/S0006-291X\(03\)01533-X](http://dx.doi.org/10.1016/S0006-291X(03)01533-X)
72. Hirotsaki T, Tsubota Y, Kariya Y, Moriyama K, Mizushima H, Miyazaki K. Laminin-6 is activated by proteolytic processing and regulates cellular adhesion and migration differently from laminin-5. *J Biol Chem* 2002; 277:49287-95; PMID:12379663; <http://dx.doi.org/10.1074/jbc.M111096200>
73. Gagnoux-Palacios L, Allegra M, Spirito F, Pommeret O, Romero C, Ortonne JP, et al. The short arm of the laminin $\gamma 2$ chain plays a pivotal role in the incorporation of laminin 5 into the extracellular matrix and in cell adhesion. *J Cell Biol* 2001; 153:835-50; PMID:11352943; <http://dx.doi.org/10.1083/jcb.153.4.835>
74. Jones JC, Lane K, Hopkinson SB, Lecuona E, Geiger RC, Dean DA, et al. Laminin-6 assembles into multimolecular fibrillar complexes with perlecan and participates in mechanical-signal transduction via a dystroglycan-dependent, integrin-independent mechanism. *J Cell Sci* 2005; 118:2557-66; PMID:15928048; <http://dx.doi.org/10.1242/jcs.02395>
75. Hamill KJ, Kligys K, Hopkinson SB, Jones JC. Laminin deposition in the extracellular matrix: a complex picture emerges. *J Cell Sci* 2009; 122:4409-17; PMID:19955338; <http://dx.doi.org/10.1242/jcs.041095>
76. Bachy S, Letourneur F, Rousselle P. Syndecan-1 interaction with the LG4/5 domain in laminin-332 is essential for keratinocyte migration. *J Cell Physiol* 2008; 214:238-49; PMID:17579341; <http://dx.doi.org/10.1002/jcp.21184>
77. Carter WG, Kaur P, Gil SG, Gahr PJ, Wayner EA. Distinct functions for integrins $\alpha 3 \beta 1$ in focal adhesions and $\alpha 6 \beta 4$ /bullous pemphigoid antigen in a new stable anchoring contact (SAC) of keratinocytes: relation to hemidesmosomes. *J Cell Biol* 1990; 111:3141-54; PMID:2269668; <http://dx.doi.org/10.1083/jcb.111.6.3141>
78. Momota Y, Suzuki N, Kasuya Y, Kobayashi T, Mizoguchi M, Yokoyama F, et al. Laminin $\alpha 3$ LG4 module induces keratinocyte migration: involvement of matrix metalloproteinase-9. *J Recept Signal Transduct Res* 2005; 25:1-17; PMID:15960391; <http://dx.doi.org/10.1081/RRS-200047870>
79. Utani A, Momota Y, Endo H, Kasuya Y, Beck K, Suzuki N, et al. Laminin $\alpha 3$ LG4 module induces matrix metalloproteinase-1 through mitogen-activated protein kinase signaling. *J Biol Chem* 2003; 278:34483-90; PMID:12826666; <http://dx.doi.org/10.1074/jbc.M304827200>
80. Hoffman MP, Engbring JA, Nielsen PK, Vargas J, Steinberg Z, Karmand AJ, et al. Cell type-specific differences in glycosaminoglycans modulate the biological activity of a heparin-binding peptide (RKRLQVQLSIRT) from the G domain of the laminin $\alpha 1$ chain. *J Biol Chem* 2001; 276:22077-85; PMID:11304538; <http://dx.doi.org/10.1074/jbc.M100774200>
81. Urushibata S, Katagiri F, Takaki S, Yamada Y, Fujimori C, Hozumi K, et al. Biologically active sequences in the mouse laminin $\alpha 3$ chain G domain. *Biochemistry* 2009; 48:10522-32; PMID:19807150; <http://dx.doi.org/10.1021/bi901421t>
82. Utani A, Nomizu M, Matsuura H, Kato K, Kobayashi T, Takeda U, et al. A unique sequence of the laminin $\alpha 3$ G domain binds to heparin and promotes cell adhesion through syndecan-2 and -4. *J Biol Chem* 2001; 276:28779-88; PMID:11373281; <http://dx.doi.org/10.1074/jbc.M101420200>
83. Kato K, Utani A, Suzuki N, Mochizuki M, Yamada M, Nishi N, et al. Identification of neurite outgrowth promoting sites on the laminin $\alpha 3$ chain G domain. *Biochemistry* 2002; 41:10747-53; PMID:12196012; <http://dx.doi.org/10.1021/bi020180k>
84. Araki E, Momota Y, Togo T, Tanioka M, Hozumi K, Nomizu M, et al. Clustering of syndecan-4 and integrin $\beta 1$ by laminin $\alpha 3$ chain-derived peptide promotes keratinocyte migration. *Mol Biol Cell* 2009; 20:3012-24; PMID:19403697; <http://dx.doi.org/10.1091/mbc.E08-09-0977>
85. Vivès RR, Crubler E, Andrieu JP, Gagnon J, Rousselle P, Lortat-Jacob H. A novel strategy for defining critical amino acid residues involved in protein/glycosaminoglycan interactions. *J Biol Chem* 2004; 279:54327-33; PMID:15485868; <http://dx.doi.org/10.1074/jbc.M409760200>
86. Okamoto O, Bachy S, Odenthal U, Bernard J, Rigal D, Lortat-Jacob H, et al. Normal human keratinocytes bind to the $\alpha 3$ LG4/5 domain of unprocessed laminin-5 through the receptor syndecan-1. *J Biol Chem* 2003; 278:44168-77; PMID:12947106; <http://dx.doi.org/10.1074/jbc.M300726200>
87. Sulka B, Lortat-Jacob H, Terreux R, Letourneur F, Rousselle P. Tyrosine phosphorylation of the syndecan-1 PDZ binding domain regulates syntenin-1 recruitment. *J Biol Chem* 2009; 284:10659-71; PMID:19228696; <http://dx.doi.org/10.1074/jbc.M807643200>
88. Rousselle P, Letourneur F. Mysterious tasks of tyrosines in syndecan-1 cytoplasmic tail. *ScientificWorldJournal* 2009; 9:629-32; PMID:19618090; <http://dx.doi.org/10.1100/tsw.2009.87>
89. Elenius K, Vainio S, Laato M, Salmivirta M, Thesleff I, Jalkanen M. Induced expression of syndecan in healing wounds. *J Cell Biol* 1991; 114:585-95; PMID:1860887; <http://dx.doi.org/10.1083/jcb.114.3.585>
90. Oksala O, Salo T, Tammi R, Häkkinen L, Jalkanen M, Inki P, et al. Expression of proteoglycans and hyaluronan during wound healing. *J Histochem Cytochem* 1995; 43:125-35; PMID:7529785; <http://dx.doi.org/10.1177/43.2.7529785>
91. Jaakkola P, Kontusaari S, Kauppi T, Määttä A, Jalkanen M. Wound reepithelialization activates a growth factor-responsive enhancer in migrating keratinocytes. *FASEB J* 1998; 12:959-69; PMID:9707168
92. Gallo R, Kim C, Kokenyesi R, Adzick NS, Bernfield M. Syndecans-1 and -4 are induced during wound repair of neonatal but not fetal skin. *J Invest Dermatol* 1996; 107:676-83; PMID:8875948; <http://dx.doi.org/10.1111/1523-1747.ep12365571>
93. Stepp MA, Gibson HE, Gala PH, Iglesia DD, Pajoohesh-Ganji A, Pal-Ghosh S, et al. Defects in keratinocyte activation during wound healing in the syndecan-1-deficient mouse. *J Cell Sci* 2002; 115:4517-31; PMID:12414997; <http://dx.doi.org/10.1242/jcs.00128>
94. Wang H, Leavitt L, Ramaswamy R, Rapraeger AC. Interaction of syndecan and $\alpha 6 \beta 4$ integrin cytoplasmic domains: regulation of ErbB2-mediated integrin activation. *J Biol Chem* 2010; 285:13569-79; PMID:20181947; <http://dx.doi.org/10.1074/jbc.M110.102137>
95. Rousselle P, Keene DR, Ruggiero F, Champlaud MF, Rest M, Burgeson RE. Laminin 5 binds the NC-1 domain of type VII collagen. *J Cell Biol* 1997; 138:719-28; PMID:9245798; <http://dx.doi.org/10.1083/jcb.138.3.719>
96. Bruckner-Tuderman L, Pfaltz M, Schnyder UW. Cylindroma overexpresses collagen VII, the major anchoring fibril protein. *J Invest Dermatol* 1991; 96:729-34; PMID:1850773; <http://dx.doi.org/10.1111/1523-1747.ep12470965>
97. Ogawa T, Tsubota Y, Hashimoto J, Kariya Y, Miyazaki K. The short arm of laminin gamma2 chain of laminin-5 (laminin-332) binds syndecan-1 and regulates cellular adhesion and migration by suppressing phosphorylation of integrin $\beta 4$ chain. *Mol Biol Cell* 2007; 18:1621-33; PMID:17314405; <http://dx.doi.org/10.1091/mbc.E06-09-0806>
98. Natarajan E, Saeb M, Crum CP, Woo SB, McKee PH, Rheinwald JG. Co-expression of p16(INK4A) and laminin 5 gamma2 by microinvasive and superficial squamous cell carcinomas in vivo and by migrating wound and senescent keratinocytes in culture. *Am J Pathol* 2003; 163:477-91; PMID:12875969; [http://dx.doi.org/10.1016/S0002-9440\(10\)63677-2](http://dx.doi.org/10.1016/S0002-9440(10)63677-2)
99. Natarajan E, Omobono JD, 2nd, Guo Z, Hopkinson S, Lazar AJ, Brenn T, et al. A keratinocyte hypermotility/growth-arrest response involving laminin 5 and p16INK4A activated in wound healing and senescence. *Am J Pathol* 2006; 168:1821-37; PMID:16723698; <http://dx.doi.org/10.2353/ajpath.2006.051027>
100. Stepp MA, Liu Y, Pal-Ghosh S, Jurjus RA, Tadvalkar G, Sekaran A, et al. Reduced migration, altered matrix and enhanced TGFbeta1 signaling are signatures of mouse keratinocytes lacking Sdc1. *J Cell Sci* 2007; 120:2851-63; PMID:17666434; <http://dx.doi.org/10.1242/jcs.03480>
101. Echtermeyer F, Streit M, Wilcox-Adelman S, Saoncella S, Denhez F, Detmar M, et al. Delayed wound repair and impaired angiogenesis in mice lacking syndecan-4. *J Clin Invest* 2001; 107:R9-14; PMID:11160142; <http://dx.doi.org/10.1172/JCI10559>
102. Botta A, Deltel F, Mettouchi A, Vieira A, Estrach S, Négroni L, et al. Confluence switch signaling regulates ECM composition and the plasmin proteolytic cascade in keratinocytes. *J Cell Sci* 2012; 125:4241-52; PMID:22641690; <http://dx.doi.org/10.1242/jcs.096289>
103. Costell M, Gustafsson E, Aszódi A, Mörögelin M, Bloch W, Hunziker E, et al. Perlecan maintains the integrity of cartilage and some basement membranes. *J Cell Biol* 1999; 147:1109-22; PMID:10579729; <http://dx.doi.org/10.1083/jcb.147.5.1109>
104. Verrando P, Blanchet-Bardon C, Pisani A, Thomas L, Cambazard F, Eady RA, et al. Monoclonal antibody GB3 defines a widespread defect of several basement membranes and a keratinocyte dysfunction in patients with lethal junctional epidermolysis bullosa. *Lab Invest* 1991; 64:85-92; PMID:1990210
105. Miyazaki K, Kikkawa Y, Nakamura A, Yasumitsu H, Umeda M. A large cell-adhesive scatter factor secreted by human gastric carcinoma cells. *Proc Natl Acad Sci U S A* 1993; 90:11767-71; PMID:8265624; <http://dx.doi.org/10.1073/pnas.90.24.11767>
106. Arnold K, Bordoli L, Kopp J, Schwede T. The SWISS-MODEL workspace: a web-based environment for protein structure homology modelling. *Bioinformatics* 2006; 22:195-201; PMID:16301204; <http://dx.doi.org/10.1093/bioinformatics/bti770>
107. Schneidman-Duhovny D, Inbar Y, Nussinov R, Wolfson HJ. PatchDock and SymmDock: servers for rigid and symmetric docking. *Nucleic Acids Res* 2005; 33(Web Server issue):W363-7; PMID:15980490; <http://dx.doi.org/10.1093/nar/gki481>