

RESEARCH ARTICLE

Open Access

Quantitative PCR of ear discharge from Indigenous Australian children with acute otitis media with perforation supports a role for *Alloiococcus otitidis* as a secondary pathogen

Robyn L Marsh^{1,2*}, Michael J Binks^{1,2}, Jemima Beissbarth^{1,2}, Peter Christensen^{1,2}, Peter S Morris^{1,2,3}, Amanda J Leach^{1,2} and Heidi C Smith-Vaughan^{1,2}

Abstract

Background: Otitis media is endemic in remote Indigenous communities of Australia's Northern Territory. *Alloiococcus otitidis* is an outer ear commensal and putative middle ear pathogen that has not previously been described in acute otitis media (AOM) in this population. The aims of this study were to determine the presence, antibiotic susceptibility and bacterial load of *A. otitidis* in nasopharyngeal and ear discharge swabs collected from Indigenous Australian children with AOM with perforation.

Methods: Paired nasopharyngeal and ear discharge swabs from 27 children with AOM with perforation were tested by *A. otitidis* quantitative PCR (qPCR). Positive swabs were cultured for 21 days. Total and respiratory pathogen bacterial loads in *A. otitidis*-positive swabs were determined by qPCR.

Results: *A. otitidis* was detected by qPCR in 11 ear discharge swabs from 10 of 27 (37%) children, but was not detected in paired nasopharyngeal swabs. *A. otitidis* was cultured from 5 of 11 qPCR-positive swabs from four children. All *A. otitidis* isolates had minimum inhibitory concentrations consistent with macrolide resistance. All *A. otitidis* qPCR-positive swabs were culture-positive for other bacteria. *A. otitidis* bacterial load ranged from 2.2×10^4 - 1.1×10^8 cells/swab (median 1.8×10^5 cells/swab). The relative abundance of *A. otitidis* ranged from 0.01% to 34% of the total bacterial load (median 0.7%). In 6 of 11 qPCR-positive swabs the *A. otitidis* relative abundance was <1% and in 5 of 11 it was between 2% and 34%. The *A. otitidis* bacterial load and relative abundance measures were comparable to that of *Haemophilus influenzae*.

Conclusions: *A. otitidis* can be a dominant species in the bacterial communities present in the ear discharge of Indigenous children with AOM with perforation. The absence of *A. otitidis* in nasopharyngeal swabs suggests the ear canal as the likely primary reservoir. The significance of *A. otitidis* at low relative abundance is unclear; however, at higher relative abundance it may be contributing to the associated inflammation. Further studies to better understand *A. otitidis* as a secondary otopathogen are warranted, particularly in populations at high-risk of progression to chronic suppurative otitis media and where macrolide therapies are being used.

Keywords: *Alloiococcus otitidis*, Otitis media, Acute otitis media with perforation, Indigenous Australian children, Bacterial load

* Correspondence: robyn.marsh@menzies.edu.au

¹Child Health Division, Menzies School of Health Research, Darwin, Australia

²Institute of Advanced Studies, Charles Darwin University, Darwin, Australia

Full list of author information is available at the end of the article

Background

Otitis media is endemic in remote Indigenous communities in Australia's Northern Territory. In this population, middle ear infection begins in the first weeks and months of life [1] and often progresses to perforation and chronic suppurative otitis media (CSOM) [2]. Cross-sectional studies have reported perforation in up to 24% of Indigenous children in the Northern Territory [3], exceeding the rate of 4% recommended by the World Health Organization as requiring urgent public health attention [4]. Standard antibiotic therapies for acute otitis media (AOM) have limited effect in this population. A randomised controlled trial (RCT) of antibiotic treatments for AOM in Indigenous children reported clinical treatment failure in 50% and 54% of children treated with single dose azithromycin or 7 days amoxicillin, respectively (RD, -4%, [95% CI, -15% to 7%]; $P = 0.504$) [5]. High clinical failure rates following antibiotic treatment are likely due to the dense, polymicrobial nature of otitis media in this population [6].

Alloiococcus otitidis is a slow-growing, strictly aerobic, Gram-positive coccus [7]. Culture-based detection of this species requires prolonged incubation for 5-14 days in an aerobic environment [8-10], and thus, *A. otitidis* will not be detected by the standard culture conditions used to identify *Streptococcus pneumoniae* or *Haemophilus influenzae*.

A. otitidis has not previously been considered in AOM affecting Indigenous Australian children; although one Australian study reported 45% Indigenous children with otitis media with effusion (OME) had *A. otitidis*-culture positive middle ear specimens [8]. Qualitative PCR-based studies from other populations have reported *A. otitidis* in MEF collected by tympanocentesis from 20-61% of children with chronic OME [11-14] and 25-50% of children with AOM [13,15-17].

Despite its detection in middle ear fluid (MEF), a primary pathogenic role for *A. otitidis* remains controversial. Data supporting a pathogenic role for *A. otitidis* include its detection in MEF from children with OME [14,16,18] and AOM [17] (in the absence of recognised otopathogens), and its capacity to provoke an inflammatory response [18-21]. One study of children with AOM reported IL-8, IL-1- β and IL-6 levels in *A. otitidis*-positive MEF to be similar to that of *S. pneumoniae*-positive specimens [22].

The reservoir of *A. otitidis* associated with otitis media is unclear. *A. otitidis* has been detected by culture or PCR in 7-12% of nasopharyngeal swabs from paediatric and adult patients with upper respiratory infection or otitis media [9,16,23]. It has also been detected by culture or PCR in 14-83% of ear canal swabs from healthy volunteers [9,24-26], suggesting *A. otitidis* detection in MEF may reflect specimen contamination by normal

canal flora. However, both PCR and culture-based studies have reported *A. otitidis* in MEF collected by tympanocentesis after careful cleaning and disinfection of the canal and tympanic membrane [8,14,22,27].

Although *A. otitidis* antibiotic minimum inhibitory concentration (MIC) breakpoints have not been defined [28], MIC levels consistent with macrolide resistance in *S. pneumoniae* have been reported for 11 of 20 (55%) [8] and 11 of 12 (92%) [29] *A. otitidis* isolates from two studies.

Azithromycin is an emerging AOM therapy for Indigenous Australian children [5]. Whether *A. otitidis* is associated with AOM in this population (and the significance of any association) is currently unclear. The aims of this study were: i) to determine the prevalence of *A. otitidis* in nasopharyngeal and ear discharge swabs from Indigenous Australian children with AOM with perforation; ii) to determine the antibiotic susceptibility of *A. otitidis* isolates to penicillin and macrolides; and iii) to determine the relative abundance of *A. otitidis* in polymicrobial specimens.

Methods

Ethical approval

Ethical approval for this study (HREC 07/85) was provided by the Human Research Ethics Committee of the Northern Territory Department of Health and Menzies School of Health Research, which includes an Aboriginal Sub-committee.

Otitis media definitions

Otitis media definitions were as previously described [5]. AOM with perforation was defined as the presence of middle ear discharge for less than six weeks, and perforation covering less than 2% of the pars tensa of the tympanic membrane. Acute otitis media without perforation was defined as a bulging tympanic membrane and a type B tympanogram. OME was defined as an intact and non-bulging tympanic membrane and type B tympanogram.

Specimens

This study retrospectively tested paired nasopharyngeal and ear discharge swabs collected at enrolment into a randomised controlled trial (RCT) of amoxicillin or azithromycin treatment for acute otitis media, conducted between March 2003 and July 2005 [5]. Eligible children were Indigenous, living in a remote community, between 6 months and 6 years of age, and had not received antibiotics in the 7 days preceding specimen collection [5]. Children were only selected for inclusion in the *A. otitidis* study if parental consent was given for their future use in related studies.

The RCT enrolled 320 children and included 49 children with a diagnosis of AOM with perforation. Of these 49 children, paired baseline nasopharyngeal and ear discharge swabs from 27 Indigenous children (median age 1.2 years, range 0.5-4.0 years) with AOM with perforation were available for *A. otitidis* testing. Four children had bilateral perforation. Thus, the sample set included 27 nasopharyngeal and 31 ear discharge swabs (58 swabs in total). As part of the original RCT, 12 children were treated with amoxicillin (50mg/kg/day in two divided doses for a minimum of 7 days) and 15 received azithromycin (30mg/kg as a single dose). Tympanic membrane perforation had not resolved in 21 of 27 children when reviewed 6-11 days after antibiotic treatment commenced. In the six remaining children, 3 had AOM without perforation (2 azithromycin treated) and 3 had OME (2 azithromycin treated) when reviewed.

Nasopharyngeal swabs were collected as previously described [5,30]. Ear discharge swabs were collected after the ear canal was cleaned to remove any visible debris and pus. The ear discharge was then sampled under direct vision (Welch Allyn Lumiview) by positioning the swab as close to the perforation as possible. Swabs were placed in 1mL skim milk-tryptone-glucose-glycerol broth [31] and frozen immediately at -20°C before being transported to the laboratory in a liquid nitrogen dry shipper [30] for long-term at storage -70°C.

DNA extraction

Total DNA was extracted from the swabs as previously described [6] with some modification. Briefly, cellular material in a 200µL aliquot of each swab was pelleted by centrifugation. Pellets were resuspended with 200µL of enzymatic lysis buffer (containing 36.25mM phosphate buffer, 1mg/mL lysozyme (Sigma®), 0.75mg/mL mutanolysin (Sigma®), and 2mg/mL Proteinase K(QIAGEN)) then incubated at 56°C for 45 min. 10µL of 20% w/v sodium dodecyl sulfate (Amresco) was added, followed by gentle mixing at room temperature for 2 min. If the solution did not become clear, 20µL of 20mg/mL Proteinase K was added and the samples were incubated at 56°C for 10 min. 4µL of 100µg/mL RNase A (QIAGEN) was added and the samples were incubated at 70°C for 10 min. 200µL of 100% ethanol (Merck) was added and the samples were pulse vortexed for 15 s. DNA was then extracted using QIAamp DNA Mini Kit (QIAGEN) as per the manufacturer's tissue protocol. DNA was eluted with 200µL of Buffer AE (QIAGEN).

For bacterial isolates, up to 100 colonies were harvested from purity plates and suspended in sterile water. Cells were pelleted by centrifugation and DNA was extracted as described above. The DNA concentration was determined using PicoGreen® reagent (Molecular Probes) as per the manufacturer's instructions with

fluorescence measured using a Victor3™ 1420 Multi-label counter (Perkin Elmer) and WorkOut software (Perkin Elmer; version 2.5).

General parameters for bacterial load qPCR assays

All qPCR assays were performed using Rotor-Gene 6000 real-time thermocyclers (Corbett Research). Quantitative standards were prepared by 1:10 serial dilution of reference isolate genomic DNA. qPCR raw data and standard curves were prepared using the Rotor-Gene 6000 software (Corbett Research; version 1.7). All standard curves were linear within the ranges described below. The qPCR efficiency and R² values were calculated using the "Auto-find threshold" function with default settings. All standards and samples were tested in duplicate. Results were only accepted where duplicates differed by ≤0.5 cycles. Each assay's limit of detection was defined as the lowest standard concentration at which specific amplification was detected in at least 95% of replicates [32].

A. otitidis qPCR

The *A. otitidis* qPCR used primers previously described by Hendolin *et al.* [33] (Table 1) which amplify a 265-bp *A. otitidis*-specific region of the 16S rRNA gene [GenBank: NR_026088] between positions 437-702, based on *E. coli* numbering. The qPCR was done using a SensiMix™ NoRef Kit (Bioline). Each 10µL qPCR included 1µM of each primer, 1X SensiMix™ NoRef reagent, 0.5X SYBR® Green I solution (Bioline), and 1µL template DNA. The reaction conditions were 95°C for 10 min followed by 35 cycles of 95°C for 15 s, 66°C for 30 s and 72°C for 15 s; then a final extension at 72°C for 1 min. This was followed by 50°C for 2 min before melt-curve analysis from 80°C-90°C in 1°C steps. Agarose gel electrophoresis was done to confirm amplification of the expected 265-bp product.

The standard curve (100pg-100fg) was prepared using genomic DNA from the *A. otitidis* reference isolate ATCC51267. The limit of detection was 100fg of genomic DNA, which is equivalent to ~50 genome copies based on an *A. otitidis* genome size of ~1.7-Mb [34]. The *A. otitidis* qPCR efficiency ranged from 0.87-0.90 and all standard curves had R² values >0.99. The no template control was negative in all except two instances in which non-specific amplification was detected below the assay's limit of detection – quantification cycle (Cq) 32-33 which was at least 5 cycles beyond the limit of detection, at a concentration less than one cell, suggesting detection of primer-dimers.

For swabs, positive results were defined as Cq-value less than or equal to that of the 100fg standard with a dissociation temperature (as determined by melt-curve analysis) within 0.5°C of the mean dissociation temperature of the assay's standards. Negative results were

Table 1 Primers and probes used in qPCR assays

Assay	Gene	Primer and Probe sequences	Position	Amplicon size (bp)	Reference
Ao	16S rRNA	Forward primer 5'- CTACGCATTTACCGCTACAC -3'	437-457	265	[33]
		Reverse primer 5'- GGGGAAGAACACGGATAGGA -3'	702-483		
TBL	16S rRNA	Forward primer 5'- TCCTACGGGAGGCAGCAGT -3'	331-349	466	[6,36]
		Reverse primer 5'- GGACTACCAGGTATCTAATCCTGTT -3'	797-772		
Hi	<i>hpd</i>	Forward primer 5'- GGTTAAATATGCCGATGGTGTG -3'	822-844	151	[37,38]
		Reverse primer 5'- TGCATCTTACGCACGGTGTGA -3'	972-953		
		Probe 5'Hex- TTGTGTACTCTCCGT "T" GGTAAGAAGAACTGCAC -SpC6 -3'*	928-896		
Spn	<i>lytA</i>	Forward primer 5'- TCTTACGCAATCTAGCAGATGAAGC -3'	306-326	101	[6]
		Reverse primer 5'- GTTGTGGTGGTTATTCTGTC -3'	406-386		
		Probe 5'- [6-FAM]-TTTGCCGAAAACGCTTGATACAGGG -[TAMRA] -3'	354-330		
Mc	<i>copB</i>	Forward primer 5'- GTGAGTGCCGCTTTTACAACC -3'	50-70	72	[6]
		Reverse primer 5'- TGATATCGCCTGCCAAGACAA -3'	121-102		

Ao = *A. otitidis*. TBL = Total bacterial load. Spn = *S. pneumoniae*. Hi = *H. influenzae*. Mc = *M. catarrhalis*. 16S rRNA gene position numbers are based on *E. coli* numbering. * Inverted commas in the primer sequence of the *hpd* probe indicate the position of the Black Hole Quencher.

recorded for samples with i) no amplification; or ii) amplification but dissociation temperature >0.5°C different to the mean dissociation temperature of the standards, consistent with primer-dimer or other non-specific amplicons. Results were considered equivocal when i) replicates gave a positive and a negative result; or ii) Cq-values were greater than that of the 100fg control but melt profiles were consistent with *A. otitidis*, suggesting detection of <50 cells. Equivocal PCRs were repeated with 2µL template DNA. Samples that remained equivocal upon repeat testing were recorded as *A. otitidis* negative.

Other bacterial load qPCR assays

H. influenzae, *S. pneumoniae*, *M. catarrhalis* and total bacterial loads were estimated as previously described [6,35,38] with modification as described in Additional file 1. Primers and probes for all assays are shown in Table 1.

A. otitidis culture and identification

Swabs qPCR-positive for *A. otitidis* were thawed on ice and vortexed. 10µL was then inoculated onto horse blood agar (Oxoid), and brain-heart-infusion agar (BHI; Oxoid) supplemented with 6.5% NaCl (Crown Scientific). Horse blood agar was selected as it had previously been used to culture *A. otitidis* from middle ear specimens [8]. As *A. otitidis* is salt tolerant [7], BHI agar with 6.5% NaCl was included as a potentially selective medium, as previously proposed by Tano *et al.* [9].

A. otitidis reference strain ATCC51267 was used as a positive control. Plates were incubated at 37°C in an ambient atmosphere and were read after 2, 5, 7, 9, 14 and 21 days of incubation. Small, off-white colonies [8] of similar appearance to the *A. otitidis* reference isolate were Gram stained and subcultured on horse blood

agar with incubation at 37°C for 48 hrs in an ambient atmosphere. Alpha-haemolytic, catalase-positive, oxidase-negative, Gram-positive cocci were considered presumptive *A. otitidis* [7]. Genomic DNA from presumptive *A. otitidis* colonies was extracted and identification was confirmed using the qPCR described above.

A. otitidis antibiotic susceptibility testing

Penicillin, erythromycin and azithromycin MICs were determined using E-Tests[®] as previously described [8]. Briefly, isolates were suspended in sterile saline to no. 3 MacFarland standard then inoculated onto horse blood agar plates. After drying, E-test[®] strips were applied and the plates incubated at 37°C in an ambient atmosphere for 48 hrs. These conditions are required to support sufficient *A. otitidis* growth to enable MIC determination [8]. As CLSI MIC breakpoints have not been determined for *A. otitidis*, the breakpoints of *S. pneumoniae* [28] were applied (consistent with previous studies [8,29]).

Culture for other bacterial species

Culture data for other bacterial species in *A. otitidis* qPCR-positive swabs were derived from the original RCT database. Culture had been performed as previously described [5,30] for 18-24 hrs and did not include testing for *A. otitidis*.

Results

qPCR detection of *A. otitidis*

A. otitidis was detected by qPCR in ear discharge swabs from 10 of 27 children (37%), including one child with bilateral detection (11 positive ear discharge swabs in total). In the remaining three children with bilateral perforation, two were qPCR-positive for *A. otitidis* in one ear only; and one was *A. otitidis* negative. As part of the

original RCT, five of the *A. otitidis*-positive children were treated with azithromycin and five treated with amoxycillin. When reviewed 6-11 days after treatment, 7 of the 10 *A. otitidis*-positive children continued to have a diagnosis of acute otitis media with perforation (3 azithromycin treated, 4 amoxycillin treated). *A. otitidis* was not detected by qPCR in nasopharyngeal swabs (0/27). No further testing of nasopharyngeal swabs was performed.

Culture of *A. otitidis* qPCR-positive swabs

A. otitidis was isolated from 5/11 qPCR-positive ear discharge swabs from 4 children after 2-14 days' incubation (Table 2). As *A. otitidis* is salt tolerant [7], BHI agar with 6.5% NaCl was used as a selective medium [9]; however, all cultures grew other bacterial species after two days' incubation, resulting in substantial overgrowth before day 21.

Susceptibility of *A. otitidis* isolates

Five *A. otitidis* isolates were available for antibiotic susceptibility testing (Table 2). Erythromycin MICs ranged from 3 - >256µg/mL. Azithromycin MICs ranged from 1.5 - >256µg/mL. Penicillin MICs ranged from 0.006-0.19µg/mL. If *S. pneumoniae* Clinical Laboratory Standards Institute (CLSI) MIC breakpoints [28] are applied, as previously described [8,29], all isolates were resistant to erythromycin and 4/5 were resistant to azithromycin. Using *S. pneumoniae* CLSI breakpoints for oral penicillin V, 3/5 isolates had intermediate penicillin susceptibility.

Other bacteria cultured from *A. otitidis*-positive ear discharge swabs

Previously recorded culture data for the 11 *A. otitidis* qPCR-positive swabs were reviewed to determine if other species were present. All *A. otitidis*-qPCR positive swabs were polymicrobial with at least 2-5 other species cultured (Table 3). In one child (Child G), *A. otitidis*, *S. pneumoniae*, *H. influenzae*, *M. catarrhalis*, *Staphylococcus* sp. and other unidentified bacteria were all

detected. Overall, *Staphylococcus* sp. were cultured from 10 of the 11 *A. otitidis* qPCR-positive swabs.

A. otitidis bacterial load and relative abundance in polymicrobial ear discharge specimens

As all *A. otitidis*-qPCR positive swabs were culture-positive for at least two other species, bacterial load measures were used to determine if *A. otitidis* was a minor or dominant constituent of the bacterial communities (Table 4). *A. otitidis* bacterial load in the 11 qPCR-positive swabs ranged from 2.2×10^4 - 1.1×10^8 cells/swab (median 1.8×10^5 cells/swab), while the total bacterial load estimates ranged from 1.5×10^7 - 8.1×10^8 cells/swab (median 9.4×10^7 cells/swab; Figure 1).

A. otitidis relative abundance was then determined by expressing its bacterial load as a percentage of the total bacterial load (Table 4). The relative abundance of *A. otitidis* ranged from 0.01-34% of the total bacterial load (median 0.7%). In 6 of 11 qPCR-positive swabs the *A. otitidis* relative abundance was <1% (median 0.2%). The *A. otitidis* bacterial load in these swabs ranged from 2.2×10^4 - 1.8×10^5 cells/swab and only one swab was positive by culture. In the remaining five swabs the relative abundance ranged from 2-34% of the total bacterial load (median 9%). These samples had the highest *A. otitidis* bacterial loads (2.9×10^6 - 1.1×10^8 cells/swab) and 4 of 5 were *A. otitidis*-positive by culture.

Bacterial load and relative abundance of *A. otitidis* compared to recognised otopathogens

H. influenzae, *S. pneumoniae* and *M. catarrhalis* bacterial loads were determined for the 11 *A. otitidis*-positive ear discharge swabs (Table 4 and Figure 1). Ten *A. otitidis*-positive swabs were also qPCR-positive for *H. influenzae*. Overall, *H. influenzae* and *A. otitidis* bacterial loads and relative abundances were similar (Table 4): the bacterial load ranged from 4.3×10^4 - 1.2×10^7 cells/swab for *H. influenzae* (median 7.4×10^5 cells/swab) and 2.2×10^4 - 1.1×10^8 cells/swab for *A. otitidis* (median 1.8×10^5 cells/swab). Maximum relative abundance was

Table 2 *A. otitidis* isolates cultured from ear discharge swabs

Sample	Days of culture until <i>A. otitidis</i> was isolated		Penicillin MIC	Erythromycin MIC	Azithromycin MIC
	HBA agar	BHI agar with 6.5% NaCl			
Child A (Left ear)	5	7	0.19	4	1.5
Child A (Right ear)	2	5	0.125	3	2
Child B	14	5	0.006	>256	128
Child C	Not detected*	9	0.19	>256	>256
Child D	7	Not detected	0.032	96	64

Culture and antibiotic susceptibility data for five isolates from four children (designated child A-D). All cultures were positive for other bacteria after two days' incubation. MIC values are in µg/mL. In the absence of *A. otitidis*-specific CLSI breakpoints, the MIC values used to define pneumococcal susceptibility were applied [28]. For oral penicillin V: sensitive ≤0.06µg/mL, intermediate susceptibility 0.12-1µg/mL, resistant ≥2µg/mL. For erythromycin: sensitive ≤0.25µg/mL, intermediate susceptibility 0.5µg/mL, resistant ≥1µg/mL. For azithromycin: sensitive ≤0.5µg/mL, intermediate susceptibility 1µg/mL, resistant ≥2µg/mL. HBA = Horse blood agar. *This plate was overgrown by a *Proteus* sp.

Table 3 Other bacteria previously cultured from *A. otitidis* qPCR-positive ear discharge specimens

Swab	<i>S. pneumoniae</i>	<i>H. influenzae</i>	<i>M. catarrhalis</i>	β -haemolytic streptococci	<i>Staphylococcus</i> sp.	<i>P. aeruginosa</i>	<i>Proteus</i> sp.	Other
Child A (Left ear)	-	+	-	-	-	-	-	+
Child A (Right ear)	-	+	-	-	+	-	-	-
Child B	-	-	n/a	-	+	n/a	+	+
Child C	-	-	n/a	-	+	-	+	+
Child D	-	-	-	+	+	-	-	+
Child E	+	+	-	-	+	-	-	-
Child F	-	-	n/a	-	+	n/a	+	+
Child G	+	+	+	-	+	-	-	+
Child H	-	-	-	-	+	-	-	+
Child I	-	-	-	-	+	+	-	+
Child J	-	-	-	-	+	+	-	+

These data were derived from the original RCT database. Culture data for *M. catarrhalis* and *P. aeruginosa* were not available for some swabs due to overgrowth by *Proteus* species (n/a = not available).

27% for *H. influenzae* versus 34% for *A. otitidis*. The relative abundance of *H. influenzae* was <1% in 5/10 *H. influenzae* qPCR-positive swabs. In 5/10 *H. influenzae* qPCR-positive swabs the *A. otitidis* bacterial load was higher than that of *H. influenzae*.

S. pneumoniae and *M. catarrhalis* were detected by qPCR in fewer *A. otitidis*-positive swabs (3/11 and 4/11, respectively) and were only present at <1% relative abundance (Table 4). *S. pneumoniae* bacterial load ranged from 3.5×10^4 - 1.8×10^5 cells/swab. *M. catarrhalis* bacterial load ranged from 4.3×10^4 - 5.9×10^5 cells/swab.

Discussion

A. otitidis is present in the ear discharge of Indigenous children with AOM with perforation

In this study, *A. otitidis* was detected by qPCR in ear discharge swabs from 10 of 27 children (37%). This rate is consistent with PCR-based AOM studies from Finland, the USA and Japan (25%, 32% and 50%, respectively) [13,15,16], and a culture-based study of chronic OME in Indigenous and non-Indigenous Australian

children (45% and 36%, respectively) [8] which tested MEF from children referred for myringotomy.

A. otitidis isolation from ear discharge swabs


Published incubation times for *A. otitidis* culture vary from 5-14 days [9,22]. Data from this study support culture for 14 days. It is important to note that unless extended incubation times are used, culture-based otitis media studies may not detect *A. otitidis*.

Using an extended culture procedure, we isolated *A. otitidis* from 5 of 11 qPCR-positive ear discharge swabs. Several factors may have contributed to the low *A. otitidis* isolation rate. Firstly, PCR-positive but culture-negative results may indicate detection of non-viable bacteria; however, it is generally accepted that DNA is quickly cleared from the middle ear [39-41] suggesting that a positive PCR is indicative of viable cells. Secondly, *A. otitidis* viability may have been affected by storage at -70°C for up to five years; however, viability of *S. pneumoniae* and *H. influenzae* has been demonstrated in swabs stored for up to 12 years at -70°C [42]

Table 4 *H. influenzae*, *S. pneumoniae*, and *M. catarrhalis* bacterial loads in *A. otitidis*-positive ear discharge swabs

	<i>A. otitidis</i>	<i>H. influenzae</i>	<i>S. pneumoniae</i>	<i>M. catarrhalis</i>
In <i>A. otitidis</i> -positive ED swabs (n = 11 [#])				
qPCR-positive swabs	11	10	3	4
Culture-positive swabs	5	4	2	1
Bacterial load range (cells/swab)	2.2×10^4 - 1.1×10^8	4.3×10^4 - 1.2×10^7	3.5×10^4 - 1.8×10^5	4.3×10^4 - 5.9×10^5
Number of swabs with bacterial load > 1×10^6 cells/swab	5	5	0	0
Number of swabs culture-positive and bacterial load > 1×10^6 cells/swab	4	3	0	0
Number of swabs with relative abundance <1%	6	5	3	4
Relative abundance range (%)	0.01-0.70	0.02-0.79	0.01-0.68	0.01-0.89
Number of swabs with relative abundance >1%	5	5	0	0
Relative abundance range (%)	2-34	4-27	0	0

[#]One child with bilateral. **A. otitidis* culture was only performed for qPCR-positive swabs.



suggesting good overall bacterial survival despite the prolonged storage.

Overgrowth by other bacterial species likely affected *A. otitidis* isolation. All cultures grew other species after two days' incubation, which potentially obscured the small *A. otitidis* colonies. Use of BHI agar with 6.5% NaCl as a selective medium was unsuccessful due to heavy overgrowth by other species after two days' incubation, despite the high salt concentration. Interestingly, successful *A. otitidis* culture was associated with higher bacterial loads (4 of 5 culture-positive swabs with bacterial load $\geq 1 \times 10^6$ cells/swab versus 1/6 culture-positive with bacterial load $< 1 \times 10^6$ cells/swab). The small study size limits further interpretation of this finding. Overall, our findings are consistent with the lower sensitivity of culture compared to PCR that has been documented for several bacteria in middle ear samples [43] and highlights the difficulties associated with *A. otitidis* culture, even when prolonged culture conditions are used.

Macrolide resistance in *A. otitidis* isolates

The five *A. otitidis* isolates had MIC values suggestive of macrolide resistance (erythromycin MICs 3 to $>256 \mu\text{g}/\text{mL}$; azithromycin MICs 1.5 to $>256 \mu\text{g}/\text{mL}$). If *S. pneumoniae* CLSI breakpoints are applied [8,29], 3 of 5 *A. otitidis* isolates also had intermediate level penicillin resistance. The validity of applying *S. pneumoniae* breakpoints to interpret *A. otitidis* MIC data remains to be determined, especially as atypical culture conditions are required to achieve sufficient growth [8]. Specific CLSI breakpoints are required to confirm macrolide resistance and reduced-penicillin susceptibility in *A. otitidis* isolates. Overall, our findings are consistent with previous studies describing macrolide resistance in *A. otitidis* [8,29].

Is *A. otitidis* contributing to middle ear pathology?

The clinical significance of *A. otitidis* in the ear discharge swabs is unclear. Several studies have found *A. otitidis* in the ear canal of healthy volunteers indicating that it is part of the normal canal flora [9,24-26]. Other studies using culture or species-specific PCRs have detected *A. otitidis* (in the absence of recognised pathogens) in MEF from children with otitis media collected after careful disinfection of the canal and tympanic membrane suggesting infection of the middle ear [8,27].

In this study, *A. otitidis* was not detected by PCR in any nasopharyngeal swabs. Previous PCR and culture-based studies from other populations have reported *A. otitidis* in 7-12% of nasopharyngeal swabs [9,16,23]. As our study was limited to 27 children, it is possible that *A. otitidis* nasopharyngeal carriage may be detected if a larger cohort were tested. However, failure to detect *A. otitidis* in nasopharyngeal swabs from any of the children with positive ear discharge swabs suggests it is unlikely to be a primary otopathogen in this population.

This prompts further consideration of *A. otitidis* in the canal flora. Staphylococcal sp. were cultured from 10 of the 11 *A. otitidis* qPCR-positive ear discharge swabs. This may indicate detection of canal flora; however, it was not possible to confirm this as paired canal swabs were not available. Furthermore, as the study children may have had a perforation for up to 6 weeks, secondary middle ear infection by these species cannot be excluded.

Bacterial load and relative abundance measures were used to better understand the significance of *A. otitidis* in the polymicrobial ear discharge swabs. *A. otitidis* bacterial load and relative abundance measures were similar to that of the otopathogen *H. influenzae* (in *A. otitidis*-positive swabs). For both species, relative abundance $< 1\%$ was associated with bacterial load $< 10^6$ cells/swab and most of these samples were qPCR-positive and culture-negative. This suggests a minor role for these species in these specimens, and may be indicative of canal flora (*H. influenzae* has also been reported in normal canal flora from healthy volunteers [26]). For the remaining specimens, relative abundance ranged from 2-34% and 4-27% for *A. otitidis* and *H. influenzae*, respectively. The significance of this finding is unclear as there are no data describing relative abundance thresholds, and the small size of this study limits further interpretation. However, a relative abundance of 34% is clearly suggestive of a dominant species. Furthermore, *A. otitidis* was present at a relative abundance greater than that of *H. influenzae* in 5/10 swabs positive for both species. These data suggest that, in these children, *A. otitidis* may have secondarily infected the middle ear following tympanic membrane perforation. This interpretation is consistent with the

hypothesis previously proposed by De Baere *et al.* [26] who reported substantially higher rates of *A. otitidis* in ear canal rather than nasopharyngeal swabs (83% versus 8%) and concluded that *A. otitidis* in MEF from children with OME most likely reflected either secondary infection following perforation or specimen contamination by canal flora. As Indigenous Australian children are at high-risk of progressing to CSOM [44], further research is warranted to better understand *A. otitidis* in secondary middle ear infection, particularly where macrolide antibiotics, such as azithromycin, are used to treat AOM with perforation [5].

Study limitations

This study retrospectively considered *A. otitidis* in polymicrobial ear discharge swabs collected from Indigenous Australian children with AOM with perforation. The study specimens had been stored for up to five years. While it is unclear if prolonged storage may have adversely affected qPCR detection of *A. otitidis*, culture-based studies have reported continued bacterial viability in upper respiratory swabs after up to 12 years storage at -70°C [42].

The study focused on Indigenous Australian children who are at high-risk of progression to CSOM, and thus the findings may not be generalisable to other populations. The small study size (27 children) also limits generalisation of the findings. Despite these limitations, it appears that *A. otitidis* is often present in the ear discharge of young Indigenous children with AOM with perforation (37% of our cohort).

The study was also potentially limited by the ear discharge sampling technique. Ear discharge swabs in this study were all collected from children with AOM with perforation affecting <2% of the pars tensa of the tympanic membrane for up to six weeks. Paired canal swabs were not collected. Canals were cleaned prior to sampling, but disinfection was not done. As the small size of the perforations prevented middle ear sampling, it is likely that some degree of contamination from canal flora will be present in the swabs. This limitation prevents definitive differentiation of canal flora from middle ear pathogens.

The qPCR described in this study is a modification of a previously published qualitative *A. otitidis* PCR [33]. The qPCR had efficiency of 0.87-0.90, slightly below the ideal range of 0.9-1.10 [45]. This may reflect the amplicon size which is approximately twice that recommended for qPCR [45]. This limitation was accepted as there are few data describing alternative *A. otitidis*-specific PCR targets and a qPCR efficiency of 0.8-0.9 is common to other bacterial load assays [46].

Conclusions

This study used qPCR to show that *A. otitidis* is present in ear discharge swabs from Indigenous Australian children with AOM with perforation. In a subset of ear discharge swabs we found high *A. otitidis* load and relative abundance. The absence of *A. otitidis* in nasopharyngeal swabs suggests *A. otitidis* may only have a role in secondary middle ear infection following tympanic membrane perforation. Larger longitudinal studies and treatment trials should include paired ear discharge, canal and nasopharyngeal swabs to further test this hypothesis. Such studies should be PCR-based and include culture to assess antibiotic susceptibility. Bacterial load and relative abundance estimates should be considered when investigating the significance of *A. otitidis* in polymicrobial contexts, particularly in children with existing perforations. In light of the co-detection of *Staphylococcus* sp. in 10 of the 11 *A. otitidis* qPCR-positive specimens, future studies should also consider the potential for polymicrobial secondary middle ear infection by these and other species. Where collection of MEF by tympanocentesis is not feasible, paired ear discharge and canal swabs may be helpful in distinguishing bacteria from the outer and middle ear. Additional studies investigating a role for *A. otitidis* as a secondary pathogen in CSOM are also warranted in populations with a high prevalence of spontaneous tympanic membrane perforation.

Additional file

Additional file 1: Other bacterial load qPCR assays. qPCR methods used to quantify *H. influenzae*, *S. pneumoniae*, *M. catarrhalis* and total bacterial loads.

Abbreviations

AOM: Acute otitis media; BHI: Brain heart infusion agar; CLSI: Clinical Laboratory Standards Institute; Cq: Quantification cycle; CSOM: Chronic suppurative otitis media; RCT: Randomised controlled trial; MEF: Middle ear fluid; MIC: Minimum inhibitory concentration; OME: Otitis media with effusion; qPCR: Quantitative PCR.

Competing interests

The authors declare that they have no competing interests.

Authors' contributions

RLM conceived the study, coordinated laboratory testing and prepared the manuscript. MJB participated in the study design, qPCR development, data analyses and manuscript preparation. JB and PC carried out the qPCR and assisted with data analyses. PSM and AJL participated in the study design and provided critical review of the manuscript. HSV participated in the study design, co-ordination and manuscript preparation. All authors read and approved the final manuscript.

Acknowledgements

We wish to acknowledge Kim Hare who assisted with MIC measurements and Allen Cheng for commenting on the manuscript draft. We also wish to acknowledge the families, health centres and schools from communities across the Northern Territory and South Australia that have participated in our studies. Financial support for this study was provided by the Channel 7

Children's Research Foundation; The Trust Foundation; and the National Health and Medical Research Council (Australia).

Author details

¹Child Health Division, Menzies School of Health Research, Darwin, Australia.
²Institute of Advanced Studies, Charles Darwin University, Darwin, Australia.
³Northern Territory Clinical School, Flinders University, Adelaide, Australia.

Received: 16 April 2012 Accepted: 21 August 2012

Published: 3 October 2012

References

1. Leach AJ, Boswell JB, Asche V, Nienhuys TG, Mathews JD: **Bacterial colonization of the nasopharynx predicts very early onset and persistence of otitis media in Australian aboriginal infants.** *Pediatr Infect Dis J* 1994, **13**:983–989.
2. Leach A, Wood Y, Gadil E, Stubbs E, Morris P: **Topical ciprofloxacin versus topical framycetin-gramicidin-dexamethasone in Australian aboriginal children with recently treated chronic suppurative otitis media: a randomized controlled trial.** *Pediatr Infect Dis J* 2008, **27**:692–698.
3. Morris PS, Leach AJ, Silberberg P, Mellon G, Wilson C, Hamilton E, Beissbarth J: **Otitis media in young Aboriginal children from remote communities in Northern and Central Australia: a cross-sectional survey.** *BMC Pediatr* 2005, **5**:27.
4. World Health Organization and the CIBA foundation: *Prevention of hearing impairment from chronic otitis media.* London, UK: Report of a WHO/CIBA Foundation workshop; 1996.
5. Morris PS, Gadil G, McCallum GB, Wilson CA, Smith-Vaughan HC, Torzillo P, Leach AJ: **Single-dose azithromycin versus seven days of amoxicillin in the treatment of acute otitis media in Aboriginal children (AATAAC): a double blind, randomised controlled trial.** *Med J Aust* 2010, **192**:24–29.
6. Smith-Vaughan H, Byun R, Nadkarni M, Jacques NA, Hunter N, Halpin S, Morris PS, Leach AJ: **Measuring nasal bacterial load and its association with otitis media.** *BMC Ear Nose and Throat Disord* 2006, **10**:10.
7. Aguirre M, Collins MD: **Phylogenetic analysis of *Alloioococcus otitis* gen. nov., sp. nov., an organism from human middle ear fluid.** *Int J Syst Evol Microbiol* 1992, **42**:79–83.
8. Ashhurst-Smith C, Hall ST, Walker P, Stuart J, Hansbro PM, Blackwell CC: **Isolation of *Alloioococcus otitidis* from Indigenous and non-Indigenous Australian children with chronic otitis media with effusion.** *FEMS Immunol Med Microbiol* 2007, **51**:163–170.
9. Tano K, von ER, Eriksson PO, Sjostedt A: ***Alloioococcus otitidis*-otitis media pathogen or normal bacterial flora?** *APMIS* 2008, **116**:785–790.
10. Harimaya A, Takada R, Somekawa Y, Fujii N, Himi T: **High frequency of *Alloioococcus otitidis* in the nasopharynx and in the middle ear cavity of otitis-prone children.** *International Journal of Pediatric Otorhinolaryngology* 2006, **70**:1009–1014.
11. Hendolin PH, Karkkainen U, Himi T, Markkanen A, Ylikoski J: **High incidence of *Alloioococcus otitis* in otitis media with effusion.** *Pediatr Infect Dis J* 1999, **18**:860–865.
12. Leskinen K, Hendolin P, Virolainen-Julkunen A, Ylikoski J, Jero J: **The clinical role of *Alloioococcus otitidis* in otitis media with effusion.** *Int J Pediatr Otorhi* 2002, **66**:41–48.
13. Harimaya A, Takada R, Hendolin PH, Fujii N, Ylikoski J, Himi T: **High incidence of *Alloioococcus otitidis* in children with otitis media, despite treatment with antibiotics.** *J Clin Microbiol* 2006, **44**:946–949.
14. Guvenc MG, Midilli K, Inci E, Kuskucu M, Tahamiler R, Ozergil E, Ergin S, Ada M, Altas K: **Lack of *Chlamydia pneumoniae* and predominance of *Alloioococcus otitidis* in middle ear fluids of children with otitis media with effusion.** *Auris Nasus Larynx* 2010, **37**:269–273.
15. Leskinen K, Hendolin P, Virolainen-Julkunen A, Ylikoski J, Jero J: ***Alloioococcus otitidis* in acute otitis media.** *Int J Pediatr Otorhi* 2004, **68**:51–56.
16. Kaur R, Adlowitz DG, Casey JR, Zeng M, Pichichero ME: **Simultaneous assay for four bacterial species including *Alloioococcus otitidis* using multiplex-PCR in children with culture negative acute otitis media.** *Pediatr Infect Dis J* 2010, **29**:741–745.
17. Neumark T, Ekblom M, Brudin L, Groth A, Eliasson I, Molstad S, Petersson AC, Tomngren A: **Spontaneously draining acute otitis media in children: An observational study of clinical findings, microbiology and clinical course.** *Scand J Infect Dis* 2011, **43**:891–898.
18. Faden H, Dryja D: **Recovery of a unique bacterial organism in human middle ear fluid and its possible role in chronic otitis media.** *J Clin Microbiol* 1989, **27**:2488–2491.
19. Kita H, Himi T, Fujii N, Ylikoski J: **Interleukin-8 secretion of human epithelial and monocytic cell lines induced by middle ear pathogens.** *Microbiol Immunol* 2000, **44**:511–517.
20. Himi T, Kita H, Mitsuzawa H, Harimaya A, Tarkkanen J, Hendolin P, Ylikoski J, Fujii N: **Effect of *Alloioococcus otitidis* and three pathogens of otitis media in production of interleukin-12 by human monocyte cell line.** *FEMS Immunol Med Microbiol* 2000, **29**:101–106.
21. Harimaya A, Takada R, Himi T, Yokota S, Fujii N: **Evidence of local antibody response against *Alloioococcus otitidis* in the middle ear cavity of children with otitis media.** *FEMS Immunol Med Microbiol* 2007, **49**:41–45.
22. Harimaya A, Fujii N, Himi T: **Preliminary study of proinflammatory cytokines and chemokines in the middle ear of acute otitis media due to *Alloioococcus otitidis*.** *Int J Pediatr Otorhi* 2009, **73**:677–680.
23. Takada R, Harimaya A, Yamazaki N, Himi T: **Detection of *Alloioococcus otitidis* and three middle ear pathogens in the nasopharynx and the middle ear effusion of otitis-prone children.** *International Congress Series* 2003, **1257**:213–215.
24. Stroman DW, Roland PS, Dohar J, Burt W: **Microbiology of normal external auditory canal.** *Laryngoscope* 2001, **111**:2054–2059.
25. Frank DN, Spiegelman GB, Davis W, Wagner E, Lyons E, Pace NR: **Culture-independent molecular analysis of microbial constituents of the healthy human outer ear.** *J Clin Microbiol* 2003, **41**:295–303.
26. De Baere T, Vanechoutte M, Deschaght P, Huyghe J, Dhooge I: **The prevalence of middle ear pathogens in the outer ear canal and the nasopharyngeal cavity of healthy young adults.** *Clin Microbiol Infect* 2010, **16**:1031–1035.
27. Khoramrooz SS, Mirsalehian A, Emaneini M, Jabalameli F, Aligholi M, Saedi B, Bazargani A, Taherikalani M, Borghaei P, Razmpa E: **Frequency of *Alloioococcus otitidis*, *Streptococcus pneumoniae*, *Moraxella catarrhalis* and *Haemophilus influenzae* in children with otitis media with effusion (OME) in Iranian patients.** *Auris Nasus Larynx* 2012, **39**:369–373.
28. Clinical and Laboratory Standards Institute: *Performance standards for antimicrobial susceptibility testing; eighteenth informational supplement, M100-S19 edn.* Wayne, PA, USA: Clinical and Laboratory Standards Institute; 2009.
29. Bosley GS, Whitney AM, Pruckler JM, Moss CW, Daneshvar M, Sih T, Talkington DF: **Characterization of ear fluid isolates of *Alloioococcus otitidis* from patients with recurrent otitis media.** *J Clin Microbiol* 1995, **33**:2876–2880.
30. Stubbs E, Hare K, Wilson C, Morris P, Leach AJ: ***Streptococcus pneumoniae* and noncapsular *Haemophilus influenzae* nasal carriage and hand contamination in children: a comparison of two populations at risk of otitis media.** *Pediatr Infect Dis J* 2005, **24**:423–428.
31. Gibson L, Khoury J: **Storage and survival of bacteria by ultra-freeze.** *Lett Appl Microbiol* 1986, **3**:127–129.
32. Bustin SA, Benes V, Garson JA, Hellemans J, Huggett J, Kubista M, Mueller R, Nolan T, Pfaffl MW, Shipley GL, Vandesompele J, Wittwer CT: **The MIQE guidelines: minimum information for publication of quantitative real-time PCR experiments.** *Clin Chem* 2009, **55**:611–622.
33. Hendolin PH, Markkanen A, Ylikoski J, Wahlfors JJ: **Use of multiplex PCR for simultaneous detection of four bacterial species in middle ear effusions.** *J Clin Microbiol* 1997, **35**:2854–2858.
34. McMichael JC, Zagursky RJ, Fletcher LD: *Alloioococcus otitidis* open reading frames (orfs) encoding polypeptide antigens, immunogenic compositions and uses thereof. [20050203280]. USA: Ref Type: Patent; 2005.
35. Binks M, Cheng A, Smith-Vaughan H, Sloots T, Nissen M, Whiley D, McDonnell J, Leach A: **Viral-bacterial co-infection in Australian Indigenous children with acute otitis media.** *BMC Infectious Diseases* 2011, **11**:161.
36. Nadkarni MA, Martin FE, Jacques NA, Hunter N: **Determination of bacterial load by real-time PCR using a broad-range (universal) probe and primers set.** *Microbiology* 2002, **148**:257–266.
37. Wang X, Mair R, Hatcher C, Theodore MJ, Edmond K, Wu HM, Harcourt BH, Carvalho M, Pimenta F, Nymadawa P, Altantssetseg D, Kirsch M, Satola SW, Cohn A, Messonnier NE, Mayer LW: **Detection of bacterial pathogens in Mongolia meningitis surveillance with a new real-time PCR assay to detect *Haemophilus influenzae*.** *Int J Med Microbiol* 2011, **301**:303–309.

38. Binks MJ, Temple B, Kirkham LA, Wiertsema SP, Dunne EM, Richmond PC, Leach AJ, Smith-Vaughan H: **Molecular surveillance of true nontypeable *Haemophilus influenzae*: an evaluation of PCR screening assays.** *PLoS ONE* 2011, **7**:e34083.
39. Aul JJ, Anderson KW, Wadowsky RM, Doyle WJ, Kingsley LA, Post JC, Ehrlich GD: **Comparative evaluation of culture and PCR for the detection and determination of persistence of bacterial strains and DNAs in the *Chinchilla laniger* model of otitis media.** *Ann Otol Rhinol Laryngol* 1998, **107**:508–513.
40. Post JC, Aul JJ, White GJ, Wadowsky RM, Zavoral T, Tabari R, Kerber B, Doyle WJ, Ehrlich GD: **PCR-based detection of bacterial DNA after antimicrobial treatment is indicative of persistent, viable bacteria in the chinchilla model of otitis media.** *Am J Otolaryngol* 1996, **17**:106–111.
41. Bakaletz LO: **Chinchilla as a robust, reproducible and polymicrobial model of otitis media and its prevention.** *Expert Rev Vaccines* 2009, **8**:1063–1082.
42. Hare KM, Smith-Vaughan HC, Leach AJ: **Viability of respiratory pathogens cultured from nasopharyngeal swabs stored for up to 12 years at –70 degrees in skim milk tryptone glucose glycerol broth.** *J Microbiol Meth* 2011, **86**:364–367.
43. Hall-Stoodley L, Hu FZ, Gieseke A, Nistico L, Nguyen D, Hayes J, Forbes M, Greenberg DP, Dice B, Burrows A, Wackym PA, Stoodley P, Post JC, Ehrlich GD, Kerschner JE: **Direct detection of bacterial biofilms on the middle-ear mucosa of children with chronic otitis media.** *JAMA* 2006, **296**:202–211.
44. World Health Organization: *Chronic suppurative otitis media. Burden of Illness and Management Options.* Geneva, Switzerland: World Health Organization; 2004.
45. D'haene B, Vandesompele J, Hellemans J: **Accurate and objective copy number profiling using real-time quantitative PCR.** *Methods* 2010, **50**:262–270.
46. Kais M, Spindler C, Kalin M, Ortqvist A, Giske CG: **Quantitative detection of *Streptococcus pneumoniae*, *Haemophilus influenzae*, and *Moraxella catarrhalis* in lower respiratory tract samples by real-time PCR.** *Diagn Microb Infect Dis* 2006, **55**:169–178.

doi:10.1186/1472-6815-12-11

Cite this article as: Marsh et al.: Quantitative PCR of ear discharge from Indigenous Australian children with acute otitis media with perforation supports a role for *Alloicoccus otitidis* as a secondary pathogen. *BMC Ear, Nose and Throat Disorders* 2012, **12**:11.

Submit your next manuscript to BioMed Central and take full advantage of:

- Convenient online submission
- Thorough peer review
- No space constraints or color figure charges
- Immediate publication on acceptance
- Inclusion in PubMed, CAS, Scopus and Google Scholar
- Research which is freely available for redistribution

Submit your manuscript at
www.biomedcentral.com/submit

