

ARTICLE

16p11.2–p12.2 duplication syndrome; a genomic condition differentiated from euchromatic variation of 16p11.2

John CK Barber^{*1}, Victoria Hall², Viv K Maloney³, Shuwen Huang², Angharad M Roberts⁴, Angela F Brady⁴, Nicki Foulds⁵, Beverley Bewes⁴, Marianne Volleth⁶, Thomas Liehr⁷, Karl Mehnert⁸, Mark Bateman³ and Helen White²

Chromosome 16 contains multiple copy number variations (CNVs) that predispose to genomic disorders. Here, we differentiate pathogenic duplications of 16p11.2–p12.2 from microscopically similar euchromatic variants of 16p11.2. Patient 1 was a girl of 18 with autism, moderate intellectual disability, behavioural difficulties, dysmorphic features and a 7.71-Mb (megabase pair) duplication (16:21 521 005–29 233 146). Patient 2 had a 7.81-Mb duplication (16:21 382 561–29 191 527), speech delay and obsessional behaviour as a boy and, as an adult, short stature, macrocephaly and mild dysmorphism. The duplications contain 65 coding genes of which *Polo-like kinase 1* (*PLK1*) has the highest likelihood of being haploinsufficient and, by implication, a triplosensitive gene. An additional 1.11-Mb CNV of 10q11.21 in Patient 1 was a possible modifier containing the *G-protein-regulated inducer of neurite growth 2* (*GPRIN2*) gene. In contrast, the euchromatic variants in Patients 3 and 4 were amplifications from a 945-kb region containing non-functional *immunoglobulin heavy chain* (*IGHV*), *hect domain pseudogene* (*HERC2P4*) and *TP53-inducible target gene 3* (*TP53TG3*) loci in proximal 16p11.2 (16:31 953 353–32 898 635). Paralogous pyrosequencing gave a total copy number of 3–8 in controls and 8 to > 10 in Patients 3 and 4. The 16p11.2–p12.2 duplication syndrome is a recurrent genomic disorder with a variable phenotype including developmental delay, dysmorphic features, mild to severe intellectual disability, autism, obsessive or stereotyped behaviour, short stature and anomalies of the hands and fingers. It is important to differentiate pathogenic 16p11.2–p12.2 duplications from harmless, microscopically similar euchromatic variants of proximal 16p11.2, especially at prenatal diagnosis.

European Journal of Human Genetics (2013) 21, 182–189; doi:10.1038/ejhg.2012.144; published online 25 July 2012

Keywords: chromosomes, human, pair 16; chromosome duplication; 16p11.2–p12.2; DNA array; euchromatic variant; copy number variation

INTRODUCTION

The proximal short arm of chromosome 16 contains a number of copy number variation (CNV) hotspots¹ that predispose to at least four reciprocal deletions and duplications within bands 16p11.2–16p12.2 (Figure 1). From centromere to telomere, these include first, the common microdeletions/duplications of ~600 kb in 16p11.2 associated with neurocognitive difficulties and obesity;^{2–4} second, the microscopically visible duplications of 8–9 Mb from 16p11.2 to 16p12.1/2 associated with developmental delay and autism^{5–11} and the reciprocal microdeletions associated with developmental delay, intellectual disability and subtle dysmorphic features;^{12,13} within these are, third, the distal (formerly atypical) microdeletions of ~220 kb in 16p11.2 associated with a phenotype that includes developmental delay or obesity but extends into the normal range^{14–16} and, fourth, the microdeletions/duplications of ~520 kb in 16p12.1 associated with developmental delay.^{1,17} Here, we report two new patients with cytogenetically visible duplications of 16p11.2–16p12.2 analysed using oligonucleotide array comparative

genomic hybridisation (oaCGH) and compared with 10 previous postnatal patients.^{5–11}

CNV is common with over 66 000 examples recorded in the Database of Genomic Variants (DGV) and an estimated 1% of the human population having a CNV > 1 Mb.¹ When copy number is high enough, rare CNVs of 8p23.1, 9p12, 9qh/q12, 9q13, 15q11.2 and 16p11.2 become visible in the light microscope and have been described as euchromatic variants.¹⁸ The euchromatic variants of 16p11.2 were originally defined using semi-quantitative cosmid fluorescence *in situ* hybridisation (FISH) and believed to result from constitutional cytogenetic amplification of paralogous sequences including non-functional *IGHV*, creatine transporter (*SLC6A10P*) and *TP53TG3* genes.^{18–20} Euchromatic variants of 16p11.2 can be mistaken for pathogenic duplications resulting in a termination of pregnancy in at least one documented case.²⁰ Here, we report two new cases in which bacterial artificial chromosome (BAC) FISH was used to assign the amplicons to an ~1-Mb region of proximal 16p11.2. Pyrosequencing²¹ was used to quantify amplicon

¹Department of Human Genetics and Genomic Medicine, Faculty of Medicine, University of Southampton, Southampton General Hospital, Southampton, Hampshire, UK; ²National Genetics Reference Laboratory (Wessex), Salisbury NHS Foundation Trust, Salisbury, Wiltshire, UK; ³Wessex Regional Genetics Laboratory, Salisbury NHS Foundation Trust, Salisbury, Wiltshire, UK; ⁴NW Thames Regional Genetics Service, Kennedy Galton Centre, Northwick Park and St Mark's Hospitals, Harrow, UK; ⁵Wessex Clinical Genetics Service, University Hospital Southampton NHS Foundation Trust, Princess Anne Hospital, Southampton, Hampshire, UK; ⁶Institut für Humangenetik, Otto-von-Guericke University, Magdeburg, Germany; ⁷Institute of Human Genetics, Jena University Hospital, Jena, Germany; ⁸Genetikum Neu-Ulm, Genetische Beratung und Diagnostik, Neu-Ulm, Germany

*Correspondence: Dr JCK Barber, Department of Human Genetics and Genomic Medicine, Faculty of Medicine, Southampton University School of Medicine, Southampton General Hospital, Tremona Road, Southampton, Hampshire SO16 6YD, UK. Tel +44 (0)2380 796586; Fax: +44 (0)2380 794760; E-mail: john.barber@soton.ac.uk
Received 21 December 2011; revised 2 May 2012; accepted 4 May 2012; published online 25 July 2012




Figure 1 UCSC browser screenshot annotated with extent of the duplications, triplication, deletions and euchromatic variants of proximal 16p in the context of the UCSC browser bands and data on genomic variants and segmental duplications (NCBI 36/hg18): included are the 16p11.2–p21.2 duplications in Patients 1 and 2 (light grey bars), the 16p11.2–p21.2 duplications of Finelli *et al*⁸ (light grey bar), the 16p11.2–16p12.1 triplication (intermediate grey bar) and duplication (light grey bar) of Ballif *et al*,¹⁰ subject 5, the duplications of Tabet *et al*,¹¹ the reciprocal 16p11.2–p21.2 ‘micro’deletions (dark grey bar with narrow bars at either end indicating variable extent), the 16p12.1 microdeletions and duplications (diagonally hatched light and dark grey bars), the distal (formerly atypical) 16p11.2 microdeletions (dark grey bar), the common 16p11.2 microdeletions/duplications (diagonally hatched light and dark grey bars) and the euchromatic amplification variants in Patients 3 and 4 (dotted bars on a light grey background). The approximate start and end points of bands, imbalances, variants and candidate genes (*CDR2*, *PLK1* and *SH2B1*) are given in Mb.

copy number and corroborate estimates made using semi-quantitative FISH in euchromatic variant carriers and controls.

MATERIALS AND METHODS

Conventional G-banded chromosome analysis was carried out in lymphocytes (Patients 1 and 2) and amniotic fluid cells (Patients 3 and 4) using standard techniques. FISH was carried out with non-functional *IGVH* (cos11, 13, 33, 97 and 98) and creatine transporter (c329B6) cosmids as before.¹⁹ Dual colour BAC FISH and oaCGH with a customised 4 × 44K Agilent oligonucleotide array (Santa Clara, CA, USA) were performed as described previously.²² Base pair coordinates correspond to NCBI release 36.1 (hg18). Copy number in euchromatic variant carriers and normal controls was determined by co-amplifying paralogous sequences at 16p11.2 and Xq28²³ and quantifying the paralogous sequence mismatches (PSM) between them using pyrosequencing according to the manufacturer’s instructions. The resultant PSM allele frequencies were calculated using the Pyrosequencing AQ software (Biotage AB, Uppsala, Sweden) and reflect the relative frequency of the paralogous sequence on each chromosome.

Clinical reports

Patient 1. This 18-year-old girl (Figures 2a–e) was the first child of healthy unrelated Caucasian parents. She had a younger brother aged 15 with normal development. The pregnancy was complicated by hyperemesis. She was born at 36 weeks gestation by spontaneous vaginal delivery weighing 2920 g (90th centile). Discharge was delayed by 5 days because of neonatal jaundice. She was a poor feeder owing to poor suck and also needed waking for feeds. She had poor weight gain and lactose intolerance. Weaning to solid food was slow and achieved by 3 years of age. Developmentally she sat at 8 months and walked at

13 months. By 18 months she had four words with meaning. During infancy, she was asthmatic and prone to respiratory tract infections but had outgrown this by the age of 9. She had otitis media and grommets inserted at the age of 3. In childhood she was a poor sleeper but this improved after the age of 12. She had vacant staring episodes but no seizures and her EEG was normal. She was toilet trained aged 7 years. She was small for her age until the age of 12 but has since caught up. She had onset of menarche aged 14 years. She has had orthodontic treatment since the age of 15 years. A Ruth Griffiths assessment at chronological age 51 months showed mental age results ranging from 28 to 40 months (sub quotient 55–78). From the age of 5 years she attended a special school for moderate learning difficulties. Since the age of 16 years she has been attending a transition college and doing a life skills course. She can dress herself, but needs help with small buttons.

From early on her behaviour was described as fixed and repetitive. She did not like plans to be changed. She would prepare well in advance for any event out of her normal routine and become fixated on it. She fixated on certain ideas or phrases and would repeat a phrase over and over to herself. She generally enjoyed meeting new people and talked very quickly to them without accounting for their lack of prior knowledge. She could be inappropriately familiar and affectionate with strangers. At other times she would happily entertain herself and not interact with others.

Clinical evaluation at the age of 15 revealed a weight of 48 kg (<25th centile), height of 158.2 cm (<25th centile) and an OFC of 55 cm (<50th centile). She had a prominent glabella, heavy eyebrows, synophrys, slight ptosis, left convergent squint, broad nasal bridge, an upturned nose with a slightly bulbous tip, bow shaped upper lip, full everted lower lip, prominent upper incisors, dental crowding and overfolded helices (Figures 2a–c). She had tapering, hypermobile fingers with brachydactyly, squared off finger tips (Figure 2d) and prominent foetal finger pads. She has had problems with painful flat feet and had bilateral medial bony

protuberances near the ankles for which she has worn orthotics (Figure 2e). Her skin was soft and velvety with wrinkled skin on the palms and soles. Her hands were clammy. She had a postural kyphosis. Neurological examination revealed poor balance and coordination.

She had an autism diagnostic interview-revised (ADI-R) assessment and observations by an occupational therapist, an art therapist and a consultant psychiatrist at the age of 16 years and 9 months and the results confirmed the diagnosis of autism. The assessment showed difficulties with non-verbal regulation of social interaction, difficulty in developing peer relationships and a lack of shared enjoyment and socio-emotional reciprocity. She had difficulty initiating and sustaining conversation in a reciprocal way, some stereotyped and repetitive speech and did not have the social awareness to avoid asking inappropriate things at inappropriate times. The assessment showed that she had more difficulties with minor changes in routine or environment when she was younger compared with when she was assessed. Her parents had made a great deal of effort to avoid any changes of her routine to prevent distress. She has become more aware of her difficulties with time




Figure 2 (a–e) The phenotype of Patient 1 aged 15 years: (a) Facial features showing heavy eyebrows, synophrys, slight ptosis, left convergent squint, broad nasal bridge, bow shaped upper lip and full lower lip; (b) lateral facial features showing overfolded helix, prominent glabella, upturned nose with a slightly bulbous tip and full everted lower lip; (c) oral features showing prominent upper incisors and dental crowding; (d) hands of the proband showing brachydactyly and squared off finger tips; and (e) feet of the proband showing bilateral medial protuberances near her ankles.




Figure 3 (a–d) G-banded partial karyotypes from Patients 1, 2, 3 and 4 with the normal chromosome (n) on the left and the duplicated (dup) or euchromatic variant (var) chromosome 16 on the right: (a) Patient 1 and (b) Patient 2 with the duplicated region of chromosome 16 indicated by the large black arrows; (c) Patient 3 and (d) Patient 4 with the euchromatic variant region of chromosome 16 indicated by the large black arrows. Note the similarity between the G-banded duplications in the Patients 1 and 2 (a, b) and the euchromatic variants in Patients 3 and 4 (c, d).

and she gets frustrated at not being able to do things alone. As a result, she has been verbally and physically aggressive towards her brother and peers at times. At college, one to one support is being put in place to try and help her behaviour. She enjoys reading and writing.

Patient 2 (*Database of Chromosomal Imbalance and Phenotype in Humans Using Ensembl Resources (DECIPHER) 2131*). This male patient was referred at the age of 45 so that his elderly parents could plan for his future. He had been a full-term normal delivery with a good birth weight and no very early health concerns. He was a little late to walk but his speech was markedly delayed and a diagnosis of glue ear was made. He attended special school from the age of 6 after an extra year at nursery school. He was a generally good natured boy with some obsessional behaviour. He had an abnormal fixation on buses and trucks, which involved large amounts of time engaged in spinning the wheels, but is not regarded as having had autism. He can now read well but cannot write more than a poor rendition of his name and is not good with numbers. He was unable to pass any exams but does have a number of skills such as excellent map-reading ability. He was generally healthy as a child but did have one episode of unexplained hypercalcemia. Since leaving school he has had jobs in sheltered employment. As an adult he has been well other than recurrent rectal prolapse. On examination, he is short at 158 cm (0.4th centile), has an OFC of 59 cm (98th centile) and is of average build. He has a wide mouth, a large furrowed tongue and broad alveolar margins. His hands are broad with a generalised brachydactyly and his digits are untapered. He has one large lipoma on his back and freckling on the palms and soles. A diagnosis of the Bannayan–Riley–Ruvalcaba syndrome was considered but molecular analysis of the *phosphatase and tensin homologue (PTEN)* gene was negative.

Patient 3. This healthy 43-year-old woman was in her third pregnancy and delivered a boy with normal measurements at the fortieth week of pregnancy (weight 3160 g, length 54 cm, OFC 34 cm and APGAR scores 10/10/10). The healthy boy is now 6 years old and will start school in the autumn of this year.

Patient 4. This patient was referred for prenatal diagnosis after two miscarriages. No further clinical details were available.

RESULTS

Patient 1

An extra G-dark band was present in the proximal short arm of chromosome 16 (Figure 3a). FISH with non-functional *IGVH* (cos33 and 98) and creatine transporter cosmids (c329B6) from proximal 16p11.2¹⁹ was normal but signals with BAC CTB2515A14 from 16p12.1 were duplicated (data not shown) (Table 1). OaCGH confirmed a duplication of 16p12.2–16p11.2 with a minimum size of 7.71 Mb and a maximum of 8.07 Mb (Figure 4e) (Table 1). An additional duplication CNV of band 10q11.21 had a minimum size of 1.11 Mb and a maximum of 2.47 Mb. Parental chromosomes were normal. The karyotype of the proband was: 46,XX, dup (16) (p11.2–p12.2)dn.ish dup (16) (p11.2–12.2)(CTB2515 A14 + +,CTB74E23 + RP11-165M2 +).arr 16p12.2p11.2(21 507 188

$\times 221\,521\,005\text{--}29\,233\,146 \times 3\,29\,581\,455 \times 2$), 10q11.21 (45 480 521 $\times 2$ 46 371 243-47 485 737 $\times 3$ 47 946 216 $\times 2$).

Patient 2 (DECIPHER 2131)

An extra G-dark band was present in the proximal short arm of chromosome 16 (Figure 3b) and duplicated signals were found using FISH with BAC RP11-142A12 from distal 16p12.1 (Figure 4a) (Table 1). OaCGH confirmed a duplication of 16p12.2-16p11.2 (Figure 4a) with a minimum of 7.81 Mb and a maximum of 8.37 Mb (Table 1). An additional duplication CNV of 1.52 Mb from Xp22.31 was found with a minimum size of 1.52 Mb. Parental chromosomes were normal and the karyotype of the proband was: 46,XY,dup (16) (p11.2-p12.2) dn.ish dup (16) (p11.2-12.2) (RP11-142A12,RP11-165M2 +).arr 16p12.1p11.2 (21 209 438 $\times 2$ 21 382 561-29 191 527 $\times 3$ 29 581 455 $\times 2$), Xp22.31(6 467 403-7 992 061) $\times 3$.

Patient 3

An extra GTG-positive band in proximal 16p was found during chromosome analysis of amniotic fluid cultures and lymphocytes from the mother (Figure 3c). The extra band was CBG negative but enhanced (enh) signals were seen using FISH with *IGVH* cosmids (cos11, 13, 33, 97 and 98) (data not shown)¹⁹ Enh signals were also seen with BACs RP11-408D2 (Figure 4b) and RP11-378C4. The results with BAC RP5-1044J9 were less convincing than those in Patient 4 implying an amplicon of 692 kb (Table 1). The duplicated signals with BAC RP11-410P5 are likely to represent independent variation as they are non-contiguous. The karyotype was 46,XX, var (16) (p11.2p11.2) mat.ish var (16) (RP11-410P5 + +,RP11-196G11 + ,RP5-1044J9?enh,RP11-408D2enh,RP11-378C4enh,RP11-488I20 + ,RP11-80F22 + ,RP11-65B9 +). Pyrosequencing gave a total diploid copy number of 8 (Supplementary Figure 1c) and, assuming that the normal chromosome had the median copy number of 2, the euchromatic variant chromosome would have at least six copies of the amplicon spanning 4.2 Mb.

Patient 4

An extra GTG-positive band was found during chromosome analysis of amniotic fluid cultures (Figure 3d). No information on the karyotypes of the parents was available. Enh signals were seen using FISH with *IGVH* cosmids (cos11, 33 and 98; data not shown)¹⁹ and an increased interval between the 16p BAC RP11-67I10 and a 16q11.2-microdissection probe (Figure 4c). Enh signals were also found with BACs RP5-1044J9, RP11-408D2 and RP11-378C4 (Figure 4d) (Table 1) implying an amplicon of 945 kb. Enh signals from non-contiguous BACs RP11-410P5 and RP11-80F22 (Table 1) were thought to be independent variation especially as RP11-80F22 is one of the commonest CNVs found using diagnostic oacGH.²⁴ The formal karyotype was: 46,XY,var(16) (p11.2p11.2). ish var (16) (RP11-410P5enh,RP11-196G11 + ,RP5-1044J9enh,RP11-408D2enh,RP11-378C4enh,RP11-488I20 + ,RP11-80F22enh,RP11-65B9 +). Pyrosequencing gave a total diploid copy number of > 10 (Supplementary Figure 1d) and, if the normal chromosome had the median copy number of 2, the euchromatic variant chromosome would have at least eight copies of the amplicon spanning 7.6 Mb.

Copy number in controls

Paralogous pyrosequencing gave total copy numbers of between three and eight in 46 controls (20 males and 26 females) with a median value of four (Supplementary Figures 1a and b).

DISCUSSION

Our results indicate how similar large single-copy duplications of 16p11.2-p12.2 and amplification variants of 16p11.2 appear under the light microscope (Figures 3a-d) and how distinct they are at the molecular level (Figure 1). The importance of differentiating duplications from variants has been illustrated by the termination of a pregnancy when a euchromatic variant was misinterpreted as a duplication.²⁰ The presence or absence of the pathogenic duplication can be tested using FISH, qPCR or microarrays (Table 1). The same methods

Table 1 FISH and oacGH results

Band	BAC/oligonucleotide	Position (Mb from telomere, hg18)	Patient 1	Patient 2	Patient 3	Patient 4
16p12.2	A_14_P201422	21 209 438		Normal		
16p12.2	A_14_P125957	21 382 561		Dup		
16p12.2	A_14_P202187	21 507 188	Normal	Dup	NT	NT
16p12.2	A_14_P110982	21 521 005	Dup	Dup	NT	NT
16p12.1	CTD-2515A14	24 637 484-24 856 494	Dup	NT	NT	NT
16p11.2	RP11-410P5	28 082 641-28 266 523	NT	NT	Dup	Enh
16p11.2	A_14_P114769	29 191 527	Dup	Dup	NT	NT
16p11.2	A_14_P107156	29 233 146	Dup	?Dup	NT	NT
16p11.2	A_14_P200817	29 581 455	Normal	Normal	NT	NT
16p11.2	RP11-74E23	29 550 782-29 724 963	Normal	NT	NT	NT
16p11.2	RP11-196G11	30 862 944-31 045 174	NT	NT	Normal	Normal
16p11.2	RP5-1044J9	31 953 353-32 079 506	NT	NT	?Enh	Enh
16p11.2	RP11-408D2	32 206 388-32 345 224	NT	NT	Enh	Enh
16p11.2	RP11-378C4	32 722 382-32 898 635	NT	NT	Enh	Enh
16p11.1/2	RP11-488I20	34 289 152-34 490 212	NT	NT	Normal	Normal
16p11.1	RP11-80F22	34 476 095-34 627 143	NT	NT	Normal	Enh^a
16p11.1	RP11-65B9	34 982 278-35 143 302	NT	NT	Normal	Normal
16cen	D16Z2		NT	NT	Normal	Normal
16qh	D16Z3		NT	NT	Normal	Normal
16q12.2	RP11-165M2	54 425 480-54 559 961	Normal	NT	NT	NT

Abbreviations: CNV, copy number variation; dup, duplication; enh, enhanced signal; FISH, fluorescence in situ hybridisation; NT, not tested; oacGH, oligonucleotide array comparative genomic hybridisation. The Ensembl Genome browser G-dark band assignments are in bold.

^aCommon benign CNV;²⁴ grey shading highlights the regions involved in pathogenic duplication or euchromatic variation.




Figure 4 (a–f) Dual colour BAC FISH results and oaCGH results: (a) Distinct pairs of signals from RP11-142A12 (red arrows) indicating the duplication in Patient 2; (b) enh (large red arrow) and normal signals (small red arrow) with RP11-408D2 (red) consistent with amplification in a metaphase cell from Patient 3; (c) increased interval (blue arrows) between BAC RP11-67110 (red) and the 16q11.2 microdissection probe (green) in a pair of chromosomes 16 from Patient 4; (d) enh (large red arrows) and normal signals (small red arrows) with RP11-408D2 (red) consistent with amplification in interphase and metaphase cells from Patient 4 (the centromeres of chromosomes 1, 5 and 19 have also been labelled in green in these images); (e) and (f) oaCGH results from Patients 1 (e) and 2 (f) analysed with the Agilent Analytics software. The vertical coloured bars and background represent the extent of the duplications adjacent to ideograms of chromosome 16 in the left hand panels, magnified in the right hand panels, with black dots representing oligonucleotides with normal copy number and red dots oligonucleotides with increased copy number. The colour reproduction of this figure is available at the *European Journal of Human Genetics* journal online.

can be used to obtain results consistent with a euchromatic variant (Figures 4a–d), but care is needed in the interpretation of results from this copy number variable region, qPCR can be sensitive to template DNA quality and targeted arrays may lack adequate coverage and require careful choice of controls. Accurate determination of copy numbers higher than 10 remains problematic even with the pyrosequencing assay used here. The less likely possibility of an unbalanced insertion would require a combination of techniques. Both pathogenic duplications of 16p11.2–p12.1 and euchromatic variants of 16p11.2 have been directly transmitted from parents to children but one of the parents has been affected, to some extent, in all the duplications.⁷

16p11.2–p21.2 duplication phenotype

Table 2 summarises the differing phenotypic consequences in Patients 1 and 2 together with seven individuals with duplications of 16p11–16p13,^{5–9} monozygotic twins with a slightly larger duplication¹¹ and a patient with a triplication and duplication (Ballif *et al*,¹⁰ subject 5). A prenatal case was excluded as no sonographic anomalies were found before termination of the pregnancy.²⁵ Only developmental and/or psychomotor delay was present in all patients. In one patient and the twins, normal

development was followed by a marked decline after the age of 2. Intellectual disability in nine patients varied from severe in four to mild or moderate in others. Autism or autistic behaviour was recorded in eight patients but only Patient 1, the twins and one other patient had a formal ADI-R or DSM IV diagnosis. Patient 2 was not thought to have autism as a child. Obsessive and/or stereotyped behaviour was found in seven patients, social problems in six, ADHD or hyperactivity in four and echolalia in three. MRI/CT scan anomalies were noted in four patients and epilepsy, seizures and/or EEG anomalies in three.

Dysmorphism was seen in 10 patients but absent in one mother and daughter diad.⁷ Common dysmorphisms included a depressed, broad or large nasal bridge, upslanting or narrow palpebral features, hypertelorism and a long or tented philtrum. Short stature was found in five patients and microcephaly in four but normal OFC was found in Patient 1 and macrocephaly in Patient 2. Tapering, long, short and/or hypermobile fingers were found in eight patients. Patient 1 also had the prominent fingertip pads noted before (Ballif *et al*,¹⁰ subject 5) as well as brachydactyly and medial bony protuberances on both feet. Skin syndactyly of the feet was seen in three patients. Recurrent infections and nystagmus or strabismus were found in four

Table 2 Phenotypic features found in patients with postnatal duplications of proximal 16p

Features	Case 1 ⁵	Case 2 ⁶	Reference 6	Mother ⁷	Daughter ^{a,7}	P1 ⁸	Reference 9	Subject 5 ¹⁰	P2, 3 ^{b,11}	Present P1	Present P2
Trisomic region	16p11-p13	16p11-p13	16p11.2-p12	16p11.2-p12.1	16p11.2-p12.1	16p11.2-p12.2	16p11.2-p13.1	16p11.2-p12.2	16p11.2-p12.2	16p11.2-p12.2	16p11.2-p12.2
~Extent (Mb)	35	2630	Full term	8	8	Full term	Full term	6 tpr;1 dup	8.95	7.71-8.07	7.81-8.37
Birth weight (g)	2200	2630	3700	35, 39 and 40	4050 (> p97)	2550 (p3)	2250	F	2450 (p4)	2920 (p90)	Full term
Sex (M/F)				F	F	M	F	F	M	F	M
Age at examination (years)			2	158 (< p3)	5	3, 25	1.5, 4	11	17	15	45
Height (cm)				52.5 (< p3)	103 (p3)	151 (< p3)	NR	-3 to -4 SDS	173 (mean)	158 (< p25)	45
Weight (kg)				LD +	16.3 (p50)	42 (< p3)	NR	-2 SDS	49 (-2 SDS)	48 (< p25)	p0.4
OFC	Micro			LD +	49 (p3)	54 (p25)	Micro	-1 SDS	55 (-1 SDS)	55 (< p50)	Macro
Dev/psychomotor delay	+	+	Behaviour	DSM-IV	After 24/12		+	++	After 26/12	ADI-R	+
Autism				DSM-IV	Partial		+	-	DSM-IV		+
Obsessive behaviours				+				+			+
Stereotyped behaviours				+				+			+
Echolalia				+				+			+
Social problems				+				+			+
ADHD											
Intellectual disability				Mild	Mild	Severe	+	Severe	Severe	Moderate	+
IQ				83	80	25 - > 39	+	42	+		
MRI/CT anomalies											
EEG anomalies			+								
Epilepsy/seizures			+								
Ataxia											
Hypotonia	+										
Dysmorphisms	+			None							
Depressed/large/broad nasal bridge	+										
Palpebral fissures	+		Narrow		Upslant						
Hypertelorism	+										
Long philtrum	+										
Round face	+										
Prominent glabella											
Anteverted nares											
Upper lip anomaly	+										
Low set ears	+										
Short neck	+										
Micro/retrognathia	+										
Nystagmus/strabismus	+										
Finger/hand anomalies	+										
Skin syndactyly of feet	+										
Recurrent infections	-										
Other single features				AD; EF; CD		CF	CPC; HT	HA	Early on	Early on	FT; L; MST

Abbreviations: AD, anxiety disorder; Brachyd; Brachydactyly; CD, cervical dystonia; CF, club feet; CPC, cleft palate; DSM-IV, diagnostic and statistical manual of mental disorders, fourth edition; EF, emotional flatness; EV, everted lower lip; F, female; FT, furrowed tongue; H, hyperactivity; HA, hyperacusis; HT, hypoplastic thumb; L, lipomata; LD, learning difficulties; M, male; macro, macrocephaly; MH, muscular hypotrophy; micro, microcephaly; MST, macrostomia; NV, non-verbal; P, Patient; p3, third centile etc; SD, sternum deformity; SDS, SD; VSD, ventral septal defect (in twin P2 that spontaneously closed); +, present; -, absent.

^aData added from Fieni *et al*,⁸ Patient 2.
^bP2,3 = monozygotic twins of Taber *et al*,¹¹ Patients 2 and 3.

Table 3 UCSC genes within the 945-kb euchromatic CNV region of 16p11.2 (16:31 953 353–32 898 635) (hg18)

No	Start (bp)	End (bp)	Size (bp)	UCSC gene ID	UCSC description
1	31 957 501	31 985 174	27 673	<i>IGHV</i> (uc010cat.1)	Immunoglobulin A heavy chain mRNA IgA5741-27
2	31 970 984	31 971 143	159	<i>IGHV3-30</i> (uc002ecu.1)	Immunoglobulin gamma heavy chain variable region mRNA for <i>IGHV3-30</i> gene, clone MGgis13b
3	31 978 049	31 978 136	87	<i>IGH</i> (uc002ecv.1)	IGHV VHDJ region mRNA, clone:H35
4	31 984 750	31 984 975	5	<i>IGH</i> (uc002ecw.1)	IGHV VHDJ region mRNA, clone:H155
5	32 070 110	32 071 375	1265	<i>HERC2P4</i> (uc002ecx.1)	hect domain and RLD 2 pseudogene 4 mRNA (cDNA clone IMAGE:5416019)
6	32 075 067	32 085 501	10 434	<i>X69637</i> (uc010cau.1)	mRNA sequence
7	32 172 145	32 174 741	2596	<i>TP53TG3b</i> (uc002ecy.2)	<i>TP53TG3b</i> mRNA
8	32 172 145	32 174 741	2596	<i>TP53TG3b</i> (uc010cav.1)	<i>TP53</i> -inducible gene 3b mRNA
9	32 181 732	32 184 170	2438	<i>BC042588</i> (uc002eda.2)	cDNA clone IMAGE:4826738
10	32 208 370	32 229 286	20 916	<i>BC041879</i> (uc002edb.2)	Similar to protein phosphatase 2A 48 kDa regulatory subunit isoform 1; serine/threonine protein phosphatase 2A, 48 kDa regulatory subunit; PP2A, subunit B, PR48 isoform; PP2A B subunit PR48; NY-REN-8 antigen, mRNA (cDNA clone IMAGE:5272051).
11	32 236 530	32 236 557	27	<i>DQ571479</i> (uc002edc.1)	DQ571479
12	32 592 342	32 596 379	4 037	<i>TP53TG3</i> (uc002edd.2)	<i>TP53TG3</i>
13	32 592 342	32 596 379	4 037	<i>TP53TG3b</i> (uc002ede.2)	<i>TP53TG3b</i> mRNA
14	32 688 520	32 688 546	26	<i>DQ574674</i> (uc002edg.1)	DQ574674
15	32 693 751	32 695 017	1 266	<i>BC038215</i> (uc010caw.1)	Hect domain and RLD 2 pseudogene 4 mRNA (cDNA clone IMAGE:5416019).
16	32 679 767	32 690 418	10 651	<i>LOC440366</i> (uc002edf.1)	Hect domain and RLD 2 pseudogene (LOC440366) non-coding RNA.
17	32 796 298	32 799 423	3 125	<i>SLC6A10P</i> (uc002edh.1)	Solute carrier family 6 (neurotransmitter transporter, creatine), member 10
18	32 796 298	32 803 964	7 666	<i>SLC6A10P</i> (uc002edi.1)	Pseudogene; cDNA FL43855 fis, clone TEST14007163, highly similar to Sodium- and chloride-dependent creatine transporter 2.
19	32 822 426	32 822 513	87	<i>IGHV</i> (uc002edj.1)	<i>IGHV</i> variable region mRNA, clone N85.

Abbreviations: CNV, copy number variant; *IGHV*, immunoglobulin heavy chain; *TP53TG3*, *TP53*-inducible target gene 3. Note: the graded background shading highlights loci that imply an inverted repeat structure; HGNC genes in bold.

patients each. Only one of the twins had a congenital heart defect (VSD) that resolved spontaneously.

Candidate genes

A minimum of 65 known coding genes are common to the duplication intervals in Patients 1 and 2 of which 11 are OMIM Morbid genes. Of these 11, many are autosomal recessive and only 3 have an estimated haploinsufficiency likelihood score (HLS), with implied dosage sensitivity, of <10%.²⁶ These include the *sarcoplasmic/endoplasmic reticulum calcium ATPase 1* (*ATP2A1*) gene (8.8%), associated with autosomal recessive Brody myopathy, and the *Amiloride-sensitive sodium channel subunits β* and γ genes *SCNN1B* (9.2%) and *SCNN1G* (8.8%) associated with the autosomal dominant hypertensive disorder Liddle syndrome. Of the other 54 genes, the *Polo-like kinase 1* (*PLK1*) gene has the lowest HLS of 0.4% and no other gene has an HLS of <10%. *PLK1* is a protein kinase superfamily member and the human form of *Drosophila polo*, which is an important cell cycle regulator implicated in mammals as both oncogene and tumour suppressor. *Polo* mutants can switch development from neurons to additional neuroblasts in *Drosophila*,²⁷ which suggests that dosage of *PLK1* might conceivably have an effect on neuronal development and microcephaly in humans.

Despite higher HLS scores, the *cerebellar degeneration-related protein 2* (*CDR2*) gene (46.9%) is a candidate for the 16p12.1 microdeletion syndrome²⁸ and might contribute to the developmental or behavioural problems when duplicated. Similarly, duplication of the *Ataxin-2-like protein* (*ATXNL2*) gene (31.5%) might be related to the ataxia seen in Patient 1 and Case 1 of Cohen *et al.*⁵ The *SH2B adapter protein 1* (*SH2B1*) gene (19.2%) is a candidate for the developmental delay and obesity associated with the 200-kb microdeletions of 16p11.2. This gene lies within the minimum extent of the present and published 16p11.2–p12.2 duplications (Figure 1)¹⁵ but not within the triplication that has a more severe but similar phenotype to the duplications reported here (Ballif *et al.*,¹⁰ subject 5).

Modifiers or second hits

The phenotype of patients with 520-kb microdeletions of 16p12.1 was more severe when a second large CNV (gain or loss >500 kb) was present and these microdeletions are thought to predispose to neuropsychiatric phenotypes in the presence of other large deletions or duplications.^{17,28} The genes within the remaining 7.2–7.3 Mb

minimum interval in Patients 1 and 2 may therefore be sufficient to explain the cognitive phenotype in 16p11.2–p12.2 duplication syndrome but both patients had additional CNVs >500 kb in size. In the female Patient 1 with autism, the additional 1.11-Mb duplication CNV of 10q11.2 overlaps multiple CNVs in the DGV but also contains a minimum of 9 genes. Of these, duplication of the *G-protein-regulated inducer of neurite growth 2* (*GPRIN2*) gene is a candidate modifier of severity in Rett syndrome²⁹ and might conceivably interact with other candidate autism genes in 16p11.2–p12.2 duplications. In the male Patient 2 without autism, an additional duplication CNV of 1.52 Mb in Xp22.31 contains a minimum of 5 genes, including the *steroid sulfatase* (*STS*) gene, but there is growing evidence that duplication CNVs containing *STS* are benign population variants.^{30,31} The inheritance pattern of these two CNVs was not determined.

Breakpoints

The breakpoints mapped in the present patients with oaCGH and the previously reported patients with BAC- or M-FISH^{8–11} correspond broadly to clusters of segmental duplications in proximal 16p (Figure 1). In both their families, Finelli *et al.*⁸ mapped the telomeric breakpoints to BAC RP11-98D10 in 16p12.2 (16:21 354 957–21 532 536) and the centromeric breakpoints to BAC RP11-368N21 in 16p11.2 (16:29 408 797–29 609 842). These authors also suggested that the duplications are mediated by non-allelic homologous recombination (NAHR) between a low copy repeat (LCR16v) in 16p11.2 and 16p12.2 that contains a truncated paralogue of the *SLCL7A5* gene from 16q24.3.⁸ Both the centromeric breakpoints in the present the Patients 1 and 2 lie within an interval of 390 kb in 16p11.2 (Table 1) that overlaps with BAC 368N21 and contains the *SLCL7A5P1* gene. The telomeric breakpoints lie within different 173 kb and 13.8 kb intervals in 16p12.2 (Table 1) that overlap with BAC 98D10 but do not contain a copy of the *SLCL7A5P1* gene or, in the case of Patient 1, correspond to any known segmental duplication. Similar variation has been recorded in the reciprocal deletion syndrome^{11–13} (Figure 1) but further high-resolution analysis of the breakpoints will be needed to establish whether all recurrent deletions and duplications of 16p11.2–16p12.2 arise by NAHR between a variety of repeats and/or via the more complex mechanisms being found in other genomic disorders.³²

Constitutional euchromatic amplification variants of 16p11.2

Including the present patients, euchromatic variants of 16p11.2 have been described in 14 families^{18,19} and have no known phenotypic consequences. The amplicons in Patients 3 and 4 have been approximately mapped with FISH (Table 1) and come from a 945-kb region of proximal 16p11.2 (16:31 953 353–32 898 635) that contains 19 UCSC (University of California Santa Cruz) sequences of which only two are HUGO Gene Nomenclature Committee (HGNC) genes (Table 2). The region is flanked by non-functional *immunoglobulin heavy chain (IGHV)* genes, with *hect domain pseudogenes (HERC2P4)* and *TP53-inducible target gene 3 (TP53TG3)* protein loci that imply an asymmetrical palindromic structure (Table 3) that might mediate amplification.³³

A diploid copy number of 4 in normal individuals and 10–12 in 16p11.2 euchromatic variant carriers was previously estimated using semi-quantitative cosmid FISH.¹⁹ The results of paralogous pyrosequencing (previously reported in abstract³⁴) showed a wider range of copy number from 3 to 8 in controls and 8 to >10 in 16p11.2 euchromatic variant carriers. We could not quantify copy numbers greater than 10 but accurate, cost effective and high-throughput methods are needed to determine the possible effect of large CNVs on quantitative phenotypic traits in large cohorts.³⁵

CONCLUSIONS

Our results indicate that the 16p11.2–p12.2 duplication syndrome is a recurrent genomic disorder characterised by developmental delay, autistic and/or repetitive behaviour, dysmorphic features, microcephaly, short stature and tapering fingers. Duplications of 16p11.2–p12.1 can be mistaken for harmless euchromatic variants of proximal 16p11.2 in the light microscope but are distinct at the molecular level and can be distinguished using FISH or aCGH.

CONFLICT OF INTEREST

The authors declare no conflict of interest.

ACKNOWLEDGEMENTS

We thank all four families concerned as well as Dr Christine Garrett and Dr Kay MacDermot for their clinical input on Patient 1. We are grateful to Dr Petra Muschke (Magdeburg) for the information on Patient 3 and for contacting her for the second blood sampling. We thank Dr Ilona Dietze-Armana for sending information on Patient 4. We are also grateful to Professor Hunt Willard for the D16Z2 centromeric probe. VH, SH and HW were supported by the UK Department of Health as part of the National Genetics Reference Laboratory (Wessex).

WEB RESOURCES

DGV: <http://projects.tcag.ca/variation/>

DECIPHER v5.0: <https://decipher.sanger.ac.uk/>

Ensembl Genome Browser: http://www.ensembl.org/Homo_sapiens/Info/Index

UCSC Genome Browser: <http://genome.ucsc.edu/>

- 1 Itsara A, Cooper GM, Baker C *et al*: Population analysis of large copy number variants and hotspots of human genetic disease. *Am J Hum Genet* 2009; **84**: 148–161.
- 2 Ghebranious N, Giampietro PF, Wesbrook FP *et al*: A novel microdeletion at 16p11.2 harbors candidate genes for aortic valve development, seizure disorder, and mild mental retardation. *Am J Med Genet Part A* 2007; **143A**: 1462–1471.
- 3 Weiss LA, Shen Y, Korn JM *et al*: Association between microdeletion and microduplication at 16p11.2 and autism. *N Engl J Med* 2008; **358**: 667–675.

- 4 Walters RG, Jacquemont S, Valsesia A *et al*: A new highly penetrant form of obesity due to deletions on chromosome 16p11.2. *Nature* 2010; **463**: 671–675.
- 5 Cohen MM, Lerner C, Balkin NE: Duplication of 16p from insertion of 16p into 16q with subsequent duplication due to crossing over within the inserted segment. *Am J Med Genet* 1983; **14**: 89–96.
- 6 Carrasco Juan JL, Cigudosa JC, Otero Gomez A *et al*: De novo trisomy 16p. *Am J Med Genet* 1997; **68**:219–221.
- 7 Engelen JJ, de Die-Smulders CE, Dirckx R *et al*: Duplication of chromosome region (16) (p11.2->p12.1) in a mother and daughter with mild mental retardation. *Am J Med Genet* 2002; **109**: 149–153.
- 8 Finelli P, Natacci F, Bonati MT *et al*: FISH characterisation of an identical (16) (p11.2p12.2) tandem duplication in two unrelated patients with autistic behaviour. *J Med Genet* 2004; **41**: e90.
- 9 Behjati F, Shafaghati Y, Firouzabadi SG *et al*: M-banding characterization of a 16p11.2p13.1 tandem duplication in a child with autism, neurodevelopmental delay and dysmorphism. *Eur J Med Genet* 2008; **51**: 608–614.
- 10 Ballif BC, Hornor SA, Jenkins E *et al*: Discovery of a previously unrecognized microdeletion syndrome of 16p11.2-p12.2. *Nat Genet* 2007; **39**: 1071–1073.
- 11 Tabet AC, Pilorge M, Delorme R *et al*: Autism multiplex family with 16p11.2p12.2 microduplication syndrome in monozygotic twins and distal 16p11.2 deletion in their brother. *Eur J Hum Genet* 2012; **20**: 540–546.
- 12 Battaglia A, Novelli A, Bernardini L *et al*: Further characterization of the new microdeletion syndrome of 16p11.2-p12.2. *Am J Med Genet Part A* 2009; **149A**: 1200–1204.
- 13 Hempel M, Rivera Brugués N, Wagenstaller J *et al*: Microdeletion syndrome 16p11.2-p12.2: clinical and molecular characterization. *Am J Med Genet Part A* 2009; **149A**: 2106–2112.
- 14 Bijlsma EK, Gijsbers AC, Schuurs-Hoeijmakers JH *et al*: Extending the phenotype of recurrent rearrangements of 16p11.2: deletions in mentally retarded patients without autism and in normal individuals. *Eur J Med Genet* 2009; **52**: 77–87.
- 15 Bachmann-Gagescu R, Mefford HC, Cowan C *et al*: Recurrent 200-kb deletions of 16p11.2 that include the SH2B1 gene are associated with developmental delay and obesity. *Genet Med* 2010; **12**: 641–647.
- 16 Barge-Schaapveld DQCM, Maas SM, Polstra A *et al*: The atypical 16p11.2 deletion: A not so atypical microdeletion syndrome? *Am J Med Genet Part A* 2011; **155**: 1066–1072.
- 17 Girirajan S, Rosenfeld JA, Cooper GM *et al*: A recurrent 16p12.1 microdeletion supports a two-hit model for severe developmental delay. *Nat Genet* 2010; **42**: 203–209.
- 18 Barber JC: Directly transmitted unbalanced chromosome abnormalities and euchromatic variants. *J Med Genet* 2005; **42**: 609–629.
- 19 Barber JC, Reed CJ, Dahoun SP *et al*: Amplification of a pseudogene cassette underlies euchromatic variation of 16p at the cytogenetic level. *Hum Genet* 1999; **104**: 211–218.
- 20 Lopez Pajares I, Villa O, Salido M *et al*: Euchromatic variant 16p+ implications in prenatal diagnosis. *Prenat Diagn* 2006; **26**: 535–538.
- 21 Ronaghi M, Uhlén M, Nyrén P: A sequencing method based on real-time pyrophosphate. *Science* 1998; **281**: 363–365.
- 22 Barber JC, Maloney VK, Huang S *et al*: 8p23.1 duplication syndrome; a novel genomic condition with unexpected complexity revealed by array CGH. *Eur J Hum Genet* 2008; **16**: 18–27.
- 23 Eichler EE, Lu F, Shen Y *et al*: Duplication of a gene rich cluster between 16p11.1 and Xq28: a novel pericentromeric directed mechanism for paralogous genome evolution. *Hum Mol Genet* 1996; **5**: 899–913.
- 24 Whitby H, Tsalenko A, Aston E *et al*: Benign copy number changes in clinical cytogenetic diagnostics by array CGH. *Cytogenet Genome Res* 2008; **123**: 94–101.
- 25 Bourthoumieu S, Esclaire F, Terro F *et al*: First prenatally diagnosed case of 16p11.2p12.1 duplication. *Prenat Diagn* 2008; **28**: 254–256.
- 26 Huang N, Lee I, Marcotte EM *et al*: Characterising and predicting haploinsufficiency in the human genome. *PLoS Genet* 2010; **6**: e1001154.
- 27 Wang H, Ouyang Y, Somers WG *et al*: Polo inhibits progenitor self-renewal and regulates Numb asymmetry by phosphorylating Pon. *Nature* 2007; **449**: 96–100.
- 28 Girirajan S, Eichler EE: A recurrent 16p12.1 microdeletion suggests a two-hit model for severe developmental delay. *Hum Mol Genet* 2010; **19**: R176–R187.
- 29 Artuso R, Papa FT, Grillo E *et al*: Investigation of modifier genes within copy number variations in Rett syndrome. *J Hum Genet* 2011; **56**: 508–515.
- 30 Li F, Shen Y, Köhler U *et al*: Interstitial microduplication of Xp22.31: causative of intellectual disability or benign copy number variant? *Eur J Med Genet* 2010; **53**: 93–99.
- 31 Furrow A, Theisen A, Velsher L *et al*: Duplication of the STS region in males is a benign copy-number variant. *Am J Med Genet Part A* 2011; **155A**: 1972–1975.
- 32 Hastings PJ, Lupski JR, Rosenberg SM *et al*: Mechanisms of change in gene copy number. *Nat Rev Genet* 2009; **10**: 551–564.
- 33 Guenthoer J, Diede SJ, Tanaka H *et al*: Assessment of palindromes as platforms for DNA amplification in breast cancer. *Genome Res* 2012; **22**: 232–245.
- 34 Hall V, Maloney VK, White H *et al*: The use of pyrosequencing to identify copy number variation of 16p11.2 in euchromatic variant carriers and the normal population. *J Med Genet* 2006; **43**(Suppl1): 3.26.
- 35 Fode P, Jespersgaard C, Hardwick RJ *et al*: Determination of beta-defensin genomic copy number in different populations: a comparison of three methods. *PLoS One* 2011; **6**: e16768.

Supplementary Information accompanies the paper on European Journal of Human Genetics website (<http://www.nature.com/ejhg>)