

RESEARCH

Open Access

Changes in the contents of four active diterpenoids at different growth stages in *Andrographis paniculata* (Burm.f.) Nees (*Chuanxinlian*)

Nanphanit Pholphana¹, Nuchanart Rangkadilok^{1,2}, Jitra Saehun¹, Supachai Ritruuechai¹ and Jutamaad Satayavivad^{1,2,3*}

Abstract

Background: The therapeutic activities of *Andrographis paniculata* are attributed to four major active diterpenoids: andrographolide (AP₁), 14-deoxy-11,12-didehydroandrographolide (AP₃), neoandrographolide (AP₄), and 14-deoxyandrographolide (AP₆). This study aims to quantify the four active diterpenoids in various plant organs of *A. paniculata* at different growth stages in greenhouse and field experiments, with a developed HPLC-diode array detector (HPLC-DAD) method for simultaneous determination of these diterpenoids.

Methods: Plants were grown in greenhouse and in field conditions, harvested at different growth stages, and separated into different organs for determination of the four active diterpenoids by an HPLC-DAD method.

Results: The most abundant diterpenoid was AP₆ between seedling and vegetative stages in the greenhouse experiment (13.38 to 23.71 mg/g in 2006 and 10.67 to 24.54 mg/g in 2007). High levels of AP₆ were also detected in leaves at the transfer stage in the greenhouse experiment (36.05 ± 0.69 mg/g) and field experiment (30.59 ± 1.39 mg/g). The levels of AP₆ then decreased as plants matured. The highest content of AP₄ was in cotyledons (16.65 ± 4.48 mg/g) at the transfer stage. The highest contents of AP₁ were detected in leaves at seed-forming stage in greenhouse experiment (24.72 ± 1.89 mg/g) and vegetative stage in field experiment (43.16 ± 0.92 mg/g). Flowers of *A. paniculata* contained high levels of AP₁ (21.42 ± 3.74 mg/g). AP₃ and AP₄ were at low levels in leaves at all growth stages.

Conclusion: In *A. paniculata*, AP₆ was at the highest level in leaves at transfer stage in both greenhouse and field experiments. AP₁ was at the highest level in leaves at vegetative stage and seed-forming stage in field and greenhouse experiments, respectively. The contents of AP₃ and AP₄ in leaves were low at all growth stages.

Background

Andrographis paniculata (Burm.f.) Nees, (Acanthaceae) (known as *Chuanxinlian* in China) is used in Scandinavian and Asian countries to treat the common cold, fever, and diarrhea [1]. The aerial part of *A. paniculata* is used in traditional Chinese medicine [2]. *A. paniculata* is beneficial to the liver, immune system, respiratory

system, and cardiovascular system, and shows anti-inflammatory, antimarial, antidiarrheal, hypoglycemic, anti-fertility, anticancer, and anti-HIV activities [1-4]. These therapeutic activities of *A. paniculata* are attributed to four active diterpenoids; andrographolide (AP₁), 14-deoxy-11,12-didehydroandrographolide (AP₃), neoandrographolide (AP₄), and 14-deoxyandrographolide (AP₆) [1-4], which exhibit different pharmacological activities [2,4]. AP₁ showed anti-inflammatory [5] and anticancer activities with stronger effects on diverse cancer cells than AP₃ and AP₆ [6]. It also showed cardioprotective properties [7,8]. Our previous studies indicated

* Correspondence: jutamaad@cri.or.th

¹Laboratory of Pharmacology, Chulabhorn Research Institute (CRI), Kamphaeng Phet 6 Road, Laksi, Bangkok 10210, Thailand

²Chulabhorn Graduate Institute (CGI), Kamphaeng Phet 6 Road, Laksi, Bangkok 10210, Thailand

Full list of author information is available at the end of the article

that AP₃ had a potent hypotensive effect [9] and higher anti-platelet activity than AP₁ [10]. AP₄ scavenged free radicals [11], and showed stronger anti-inflammatory activity than AP₁ by inhibiting nitric oxide production both *in vitro* and *ex vivo* [12]. AP₄ also had antimalarial activities [13] and hepatoprotective effects against carbon tetrachloride [14]. AP₆ is an effective antagonist of platelet activating factor-mediated processes in bovine neutrophils via its effects on calcium channels [15]. It also caused vasorelaxation of rat thoracic aorta [16] and relaxation of uterine smooth muscle via selectively blocking voltage-operated calcium channels [17].

There is a large market requirement for *A. paniculata* materials, with an estimated consumption in India (aerial part) of 250 tonnes per year [18]. *A. paniculata* is now commercially cultivated to meet the high demand [19]. There have been several studies on cultivation of *A. paniculata*. Kumar *et al.* [20] reported that 25°C was the optimal germination temperature (94.6% germination), while there was no germination at 40°C. The growth regulators abscisic acid and gibberellic acid also increased the AP₁ content [21]. There are few reports on the contents of active diterpenoids in *A. paniculata* during plant development. Bhan *et al.* [22] analyzed the contents of three diterpenoids in leaves at three harvesting dates. Prathan-turarug *et al.* [23] investigated the contents of AP₁ and AP₃ in field-grown *A. paniculata* harvested at the 60% flowering stage. Parasher *et al.* [24] reported that the maximum AP₁ content in leaves was at 120 days of maturity. There were large variations in the contents of three active diterpenoids (AP₁, AP₃, and AP₄) among different *A. paniculata* products in a Thai market [25]. Both the growing region and season strongly affect production of the diterpene lactones in this plant [1].

In this study, we developed a simple and rapid extraction method followed by simultaneous determination of the four active diterpenoids (AP₁, AP₃, AP₄, and AP₆) in different parts of *A. paniculata*. This simple reversed-phase HPLC-diode array detector (DAD) method was successfully applied to quantify the major components in *A. paniculata* at different growth stages. The method is sufficiently rapid, simple, sensitive, and cost-effective to be used for quality control of raw materials and herbal preparations [25]. DAD can be used to identify the active compounds based on their absorption spectra.

To our knowledge, there are no reports on the contents of the four active diterpenoids at different growth stages from seedling to mature stages of *A. paniculata*. In addition, there are few reports on the contents of these four active compounds in various plant organs at different growth stages. The aim of this study was to determine the changes in contents of the four active diterpenoids in various plant organs of *A. paniculata* at different growth

stages in greenhouse and field experiments at two different cultivation times: September 2006 and January 2007.

Methods

Chemicals

Four standard diterpenoids, andrographolide (AP₁, purity 99%), 14-deoxy-11,12-didehydroandrographolide (AP₃, purity 96%), neoandrographolide (AP₄, purity 99%), and 14-deoxyandrographolide (AP₆, purity 99%) were purified at our Institute by the Laboratory of Pharmacology and Laboratory of Natural Products, Chulabhorn Research Institute (CRI), Thailand, following a method published previously [26]. They were extracted from *A. paniculata* plant materials and identified by thin-layer chromatography (TLC), ultraviolet (UV) spectra, melting point (MP) analysis, infrared (IR) analysis, mass spectrometry (MS), and nuclear magnetic resonance (NMR) spectroscopy. The structures of these four active diterpenoids are shown in Figure 1. HPLC grade methanol and acetonitrile were obtained from Merck (Darmstadt, FR, Germany). High purity water obtained from a Milli-Q water purification system (Millipore, Bedford, MA, USA) was used in these experiments.

Plant materials

Greenhouse experiment

A. paniculata plants were identified by Dr. Wongsatit Chuakul and a voucher specimen was deposited at the Pharmaceutical Botany Mahidol Herbarium, Department




Figure 1 Structures of four main active diterpenoids in *Andrographis paniculata*. A. andrographolide (AP₁); B. 14-deoxy-11,12-didehydroandrographolide (AP₃); C. neoandrographolide (AP₄); and D. 14-deoxyandrographolide (AP₆).

of Pharmaceutical Botany, Faculty of Pharmacy, Mahidol University, Bangkok, Thailand (PBM 3760). In the greenhouse experiment, we studied two cultivation times: September 2006 and January 2007. The temperature ranged from 24.2 to 28.5°C and the relative humidity was from 60 to 70% in the greenhouse environment. Seeds were sown in a tray and grown in the greenhouse at CRI. Peat moss soil (Chia Tai, Bangkok, Thailand) was used in the greenhouse experiments (pH 5.5; N-P-K 12-14-24; 0.8 kg/m³). At the seedling, first true leaf, and transfer stages, plants were grown in the tray. At the transfer stage, plants were transplanted into pots to grow on to the mature stages (vegetative, 50% flowering, seed-forming, and mature-seed stages). Plants were harvested at different growth stages and separated into different parts (cotyledon, hypocotyl, leaf, stem, flower, young pod including seed, pod, root, and mature seed) depending on the growth stage (Table 1) to quantify the four active diterpenoids. Plant materials were washed and then dried in an oven at 35–45°C [27]. The dried plant materials were ground into a powder using a blender (Waring Commercial, CT, USA), and kept at room temperature until extraction and analysis.

Field experiment

Seeds were sown in the field at Ratchaburi Province, Thailand. Plants were harvested at different growth stages; transfer, vegetative, 50% flowering, seed-forming, and mature-seed stages (2, 3.5, 5, 6, and 6.5 months old, respectively) to quantify the four active diterpenoids. Sun-dried plant materials from the field were separated into two parts (leaves and aerial part). The dried leaves and aerial parts were ground into powder, and kept at room temperature until extraction and analysis.

Sample preparation

Greenhouse experiment

Dried powder (0.1 g) of *A. paniculata* from different plant organs at different growth stages was accurately

weighed (three replicates per sample, 20 plants per replicate at seedling, first true leaf, and transfer stages, two plants per replicate at other maturity stages). The plant powder was extracted with 4 mL methanol and rotated with a rugged rotator (Glas-Col, IN, USA) for 5 min, and then centrifuged (Sorvall, Munich, Germany) at 1876 × g for 10 min. The supernatant was transferred into a 10 mL volumetric flask and the crude powder was re-extracted twice with 3 mL methanol. All extracts were combined, the volume was adjusted, and then the solution was filtered through a 0.45-µm nylon membrane (Chrom Tech, MN, USA) before HPLC analysis. The extraction method was modified from our previous study [25].

Field experiment

Powdered plant material (leaves or aerial part) (0.1 g) of *A. paniculata* at different growth stages was accurately weighed (three replicates per sample, 20 plants per replicate at transfer stage, two plants per replicate at other maturity stages) and then extracted as described above for the greenhouse experiment.

HPLC-DAD

The four compounds in the methanolic extract of *A. paniculata* were analyzed simultaneously by HPLC-DAD (Agilent Technologies, Germany) on a reverse phase column (Zorbax SB-C18; 4.6 × 75 mm, 3.5 µm) connected to a cartridge guard column (Agilent Technologies, USA). The simple mobile phase consisted of 28% acetonitrile in water with a flow rate of 1.2 mL/min. The temperature of the column was controlled at 25°C and the detection with the diode array detector (DAD; Model G1315A, Agilent Technologies, Germany) at 205 nm. The injection volume was 5 µL. A standard mixture containing AP₁, AP₃, AP₄, and AP₆ in methanol was prepared in the range of 0.5–1000 µg/mL. The peak

Table 1 *Andrographis paniculata* materials from plants at various developmental stages in greenhouse experiment in two growing seasons, September 2006, and January 2007

Stages	Harvesting time (days)		Plant organs
	September 2006	January 2007	
Tray			
1. Seedling	7	7	Cotyledon, hypocotyl, root
2. First true leaf	14	14	Cotyledon, young leaf, stem, root
3. Transfer	39	45	Cotyledon, leaf, stem, root
Pot			
4. Vegetative	108	114	Leaf, stem, root
5. 50% Flowering	128	141	Leaf, stem, root, flower, young pod (including seed)
6. Seed forming	133	149	Leaf, stem, root, flower, young pod (including seed)
7. Mature seed	161	206	Leaf, stem, root, pod, and seed

area of each compound was plotted against the concentration.

Statistical analysis

Data are expressed as mean \pm standard deviation (SD, $n = 3$). Linear regression analysis was used to compare DAD responses and concentrations using ChemStation software version A.10.02 (Agilent Technologies, Germany). Accuracy was assessed as percentage recovery using the following formula: $\left(\frac{(C_{\text{Spiked}} - C_{\text{Sample}})}{C_{\text{Standard added}}} \times 100 \right)$. Precision was expressed as the percent relative standard deviation; %RSD ($\%RSD = \frac{SD}{X} \times 100$), which was calculated with Excel 2010 software (Microsoft, USA).

Results and discussion

Method validation

Xu et al. [28] determined diterpenoids in *A. paniculata* leaves by silver ion complexation in the mobile phase to separate AP₃ and AP₆ with a 25-min analysis time. The analysis time in the present study was less than 18 min. The chromatograms in Figure 2 showed good separation of these four active diterpenoids; a chromatogram of the standard solution is shown, compared with samples containing the highest contents of AP₆ and AP₁ at the transfer and mature-seed stages, respectively. The calibration curves showed good linearity in the range of 0.5 to 1000 $\mu\text{g/mL}$, and the correlation coefficients were 0.9999 for AP₁, AP₃, AP₄, and AP₆. Accuracy was determined (expressed as percentage recovery) by adding three known amounts of standard solutions to the plant samples, 10 replicates per concentration, and then extracting the samples and analyzing by the HPLC method described above. The mean recoveries of the four active diterpenoids, AP₁, AP₃, AP₄, and AP₆, were 99.66 \pm 2.22%, 98.91 \pm 2.29%, 100.20 \pm 1.85%, and 97.98 \pm 1.90%, respectively, indicating good accuracy of the method. To access the precision of the method, we analyzed the extracted sample solutions 10 times on the same day and 10 times on 3 consecutive days. The results showed acceptable intra- and inter-day precision, as represented by the percent relative standard deviation (%RSD) ranging from 0.80% to 1.02% (content) and from 0.41% to 0.66% (retention time) for intra-day precision, and from 3.42% to 4.92% (content) and from 0.39% to 0.74% (retention time) for inter-day precision. The limit of detection (LOD) and limit of quantitation (LOQ) were measured based on the signal to noise ratio of 3:1 and 10:1, respectively. The LOD of AP₁, AP₃, AP₄, and AP₆ were 0.10, 0.25, 0.25, and 0.25 $\mu\text{g/mL}$, respectively, and the LOQ were 0.25, 0.5, 0.5, and 0.5 $\mu\text{g/mL}$, respectively. These results showed that our analytical method met acceptable criteria for all

analytes, and could be used for routine analysis for the four active diterpenoids in *A. paniculata*.

Greenhouse experiment

There were large variations in contents of the four active diterpenoids in different plant organs at different growth stages. We studied the patterns of accumulation of these four active diterpenoids in two cultivation periods: September 2006 (Figure 3) and January 2007 (Figure 4). The contents of four active diterpenoids in seeds and roots of *A. paniculata* were lower than limit of detection (LOD). From the seedling to the first true leaf stages, the highest content of AP₆ was in the cotyledons (Figure 3A and B). At the transfer stage, the highest content of AP₆ was in leaves both in 2006 and 2007 (28.02 \pm 0.15 and 36.05 \pm 0.69 mg/g, respectively; Figures 3C and 4C). In 2006, the highest content of AP₁ was in leaves at the mature-seed stage (24.02 \pm 1.18 mg/g) (Figure 3G), whereas in 2007, the highest content of AP₁ was at the seed-forming stage (24.72 \pm 1.89 mg/g) (Figure 4F). Flowers at the 50% flowering stage also contained high contents of AP₁ in both 2006 and 2007 (21.82 \pm 1.61 and 21.42 \pm 3.74 mg/g, respectively) (Figures 3E and 4E). In contrast, flowers contained low contents of the other three active diterpenoids (\leq 0.9 mg/g). There were low levels of AP₃ in all plant organs at all growth stages (\leq 1.4 mg/g). The highest AP₃ content was at the mature-seed stage in 2006 (1.4 \pm 0.12 mg/g) and at the vegetative stage in 2007 (0.77 \pm 0.09 mg/g) (Figures 3G and 4D). The highest content of AP₄ was in cotyledons at the transfer stage in 2006 and 2007 (3.28 \pm 0.23 and 16.65 \pm 4.48 mg/g, respectively) (Figures 3C and 4C). The range of AP₄ in leaves was from 0.22 mg/g to 3.72 mg/g and the content increased as plants matured. The highest content of AP₄ in leaves was at the mature-seed stage in 2006 and 2007 (3.72 \pm 1.31 and 1.88 \pm 0.15 mg/g, respectively) (Figures 3G and 4G). The AP₆ content in leaves was lower in 2006 (28.02 \pm 0.15 mg/g) than in 2007 (36.05 \pm 0.69 mg/g), while the contents of the other three active diterpenoids were similar in both cultivation times. The present results also indicated that the young plants at seedling and first true leaf stages (harvesting time \leq 14 days) contained high contents of these four active diterpenoids.

Field experiment

In greenhouse and field experiments, the changes in the contents of the four active diterpenoids at different growth stages were similar in leaves and the aerial part (Figures 5 and 6). However, leaves contained higher contents of four active diterpenoids than the aerial part. In the greenhouse experiment, the highest AP₁ content in leaves (24.02 \pm 1.18 mg/g) was about three times higher than that in the aerial part (8.64 \pm 0.25 mg/g) at the




Figure 2 HPLC chromatograms of four diterpenoids in *Andrographis paniculata*. **A.** Four standard diterpenoids; **B.** Leaf sample of *Andrographis paniculata* at transfer stage; and **C.** Leaf sample of *Andrographis paniculata* at mature-seed stage.




Figure 3 Contents of four diterpenoids in various tissues of *Andrographis paniculata* at different growth stages (A–G) in greenhouse experiment (September 2006). Stages were as follows: A. Seedling (7 days); B. First true leaf (14 days); C. Transfer (39 days); D. Vegetative (108 days); E. 50% Flowering (128 days); F. Seed-forming (133 days); and G. Mature seed (161 days).




Figure 4 (See legend on next page.)

(See figure on previous page.)

Figure 4 Contents of four diterpenoids in various tissues of *Andrographis paniculata* at different growth stages (A–G) in greenhouse experiment (January 2007). Stages were as follows: **A.** Seedling (7 days); **B.** First true leaf (14 days); **C.** Transfer (45 days); **D.** Vegetative (114 days); **E.** 50% Flowering (141 days); **F.** Seed-forming (149 days); and **G.** Mature seed (206 days).

mature-seed stage in 2006 (Figures 5A and 6A). Similarly, in 2007, the highest AP₁ content in leaves was four times higher than that in the aerial part at the seed-forming stage (24.72 ± 1.89 mg/g in leaves, 5.71 ± 0.65 mg/g in aerial part) (Figures 5B and 6B). In the field experiment, the highest AP₁ content in leaves was only twice that in the aerial part at the vegetative stage (43.16 ± 0.92 mg/g in leaves, 24.31 ± 1.68 mg/g in aerial part; Figures 5C and 6C). In greenhouse (2006 and 2007) and field experiments, there was a drastic increase of AP₁ in leaves from the transfer stage (7.78 ± 0.03 , 9.09 ± 0.47 , and 18.43 ± 0.54 mg/g, respectively) to the vegetative stage (19.61 ± 0.28 , 15.39 ± 0.06 , and 43.16 ± 0.92 mg/g, respectively; Figure 5). In the field experiment, the highest content of AP₁ was at the early growth stage (vegetative stage) whereas it was at the mature-seed and seed-forming stages in the greenhouse experiment in 2006 and 2007, respectively. The highest content of AP₆ in leaves was at the transfer stage in the field experiment (30.59 ± 1.39 mg/g) (Figure 5C) and in the greenhouse experiment (28.02 ± 0.15 and 36.05 ± 0.69 mg/g in 2006 and 2007, respectively) (Figure 5A, 5B). In the greenhouse experiment, the AP₆ content in leaves and the aerial part increased markedly from the first true leaf to the transfer stages and then decreased at maturity. The pattern of AP₆ accumulation in plants was similar in the field and greenhouse experiments (Figures 5 and 6). Moreover, there were low levels of AP₃ and AP₄ in leaves and the aerial part at all growth stages in both the greenhouse and field experiments. *A. paniculata* grown in field conditions contained higher levels of the four active diterpenoids than plants grown in greenhouse conditions. For example, AP₁ contents in leaves at all growth stages (transfer to mature-seed stages) were higher in field-grown plants than in greenhouse-grown plants. Because *A. paniculata* was grown under natural conditions in the field experiment, it is difficult to control cultivation conditions that may affect the levels of these active diterpenoids and quality of this medicinal plant. However, these results showed that the pattern of accumulation of the four active diterpenoids was similar in the field and greenhouse experiments.

To our knowledge, this is the first report to evaluate the contents of these four active diterpenoids in different plant organs during plant development in greenhouse and field experiments. *A. paniculata* is normally harvested at the 50–60% flowering stage, since it was proposed that plants at this stage contain the highest levels

of AP₁ [22,23]. Among the active compounds extracted from *A. paniculata*, AP₁ is generally reported as the major active compound [2]. Matsuda *et al.* [26] analyzed whole *A. paniculata* plants and reported the following diterpenoids contents on a dry weight basis: AP₁ (0.6%), AP₃ (0.06%), AP₄ (0.005%), and AP₆ (0.02%). Sharma *et al.* [29] reported that the leaves of *A. paniculata* contained the highest content of AP₁ (2.39%), similar to our results (2.4%) in 2006 and 2007 in the greenhouse experiment. However, the highest content of AP₁ in our field experiment (4.3%) was higher than that reported previously. A new finding in our study was that the most abundant diterpenoid was not AP₁ but AP₆ (in 2006 and 2007 at the seedling to vegetative stages in greenhouse experiments, and at transfer stage in the field experiment). The highest content of AP₆ was found in leaves at the transfer stage; it was four times higher than AP₁ in 2006 (28.02 ± 0.15 and 7.78 ± 0.03 mg/g, respectively) and 2007 (36.05 ± 0.69 and 9.09 ± 0.47 mg/g, respectively). *A. paniculata* at the transfer stage with short harvesting time (39–45 days) contained the highest level of AP₆, which showed bioactivity as an effective vasorelaxant [16].

Since the four active diterpenoids of *A. paniculata* have different pharmacological properties, knowledge of their different patterns of accumulation among various plant organs and growth stages will be helpful to select plant materials for particular purposes or disease treatments. The patterns of accumulation of AP₁ and AP₆ were similar to those reported by Bhan *et al.* [22]. AP₁ was continuously produced during plant growth while AP₆ content in leaves peaked at the transfer stage and then decreased as the plants matured. In a previous report, harvesting at 100 days after transplantation was recommended to obtain plants with the highest AP₁ content [22]. Prathanturarug *et al.* [23] reported levels of AP₁ and AP₃ similar to those detected at 50% flowering stage in the present study, although the *A. paniculata* plants were harvested at the 60% flowering stage in their study. Recently, Parasher *et al.* [24] analyzed the AP₁ content in leaves at different growth stages, and found the highest levels of AP₁ in leaves at 120 days of maturity, similar to our results (highest AP₁ content was in leaves at mature-seed and seed-forming stages). The content of AP₃ was lower than that of other diterpenoids in these experiments. However, in our previous study, we observed that the AP₃ increased in plants and plant products stored for a period of time [25]. Furthermore, the stability of the amorphous form of AP₁ was




Figure 5 Contents of four diterpenoids in *Andrographis paniculata* leaves at different growth stages. **A.** Greenhouse experiment (September 2006); **B.** Greenhouse experiment (January 2007); **C.** Field experiment.




Figure 6 Contents of four diterpenoids in *Andrographis paniculata* aerial part at different growth stages. **A.** Greenhouse experiment (September 2006); **B.** Greenhouse experiment (January 2007); **C.** Field experiment.

temperature-dependent, and it could be converted to AP₃ [30]. Fresh mature plants with low AP₃ content and high AP₁ content should be used to treat the common cold, rather than products that have been stored for a period of time, which may exert cardiovascular side effects [31]. The highest content of AP₄ was at the transfer stage, similar to AP₆, but in cotyledons rather than leaves. The AP₄ content in leaves reported in this study was low and slightly increased with maturity. Bhan et al. [22] reported that the simultaneous dehydration and glycosylation of AP₁ formed andrographoside and AP₄. However, the AP₄ content in our study was lower than that reported in their study, which may be because of genetic variations and/or cultivation conditions.

Conclusion

In *A. paniculata*, AP₆ was at the highest level in leaves at transfer stage in both greenhouse and field experiments. AP₁ was at the highest level in leaves at vegetative stage and seed-forming stage in field and greenhouse experiments, respectively. The contents of AP₃ and AP₄ in leaves were low at all growth stages.

Abbreviations

AP₁: Andrographolide; AP₃: 14-deoxy-11,12-didehydroandrographolide; AP₄: Neoandrographolide; AP₆: 14-deoxyandrographolide; HPLC-DAD: High performance liquid chromatography-diode array detector; TLC: Thin-layer chromatography; UV: Ultraviolet; MP: Melting point; IR: Infrared; MS: Mass spectrometer; NMR: Nuclear magnetic resonance; CRI: Chulabhorn Research Institute; %RSD: The percent relative standard deviation; LOD: Limit of detection; LOQ: Limit of quantitation.

Competing interests

The authors declare that they have no competing interests.

Authors' contributions

JS conceived the idea for the study. JS, NR, and NP designed the experiments. NP and NR performed the experiments. NP analyzed the data. JSH cultivated plants in the greenhouse experiment and conducted extractions. SR separated, purified, and identified the four diterpenoids reference standards. NP, NR, and JS wrote the manuscript. All authors read and approved the final version of the manuscript.

Acknowledgements

The authors kindly acknowledge Mr. Likhit Suchikran for providing seed materials and for assistance with the field experiment. We also thank Mrs. Sujanya Rueangveerayut, Laboratory of Biotechnology, CRI, for assistance with greenhouse experiments.

Some results from this work were also applied for patent in Thailand. (Request Number: 1203001500, Date: December 27, 2012, Status: Pending)

Author details

¹Laboratory of Pharmacology, Chulabhorn Research Institute (CRI), Kamphaeng Phet 6 Road, Laksi, Bangkok 10210, Thailand. ²Chulabhorn Graduate Institute (CGI), Kamphaeng Phet 6 Road, Laksi, Bangkok 10210, Thailand. ³Center of Excellence on Environmental Health and Toxicology, Office of the Higher Education Commission, Ministry of Education, Bangkok 10400, Thailand.

Received: 8 August 2012 Accepted: 19 December 2012

Published: 15 January 2013

References

- Mishra SK, Sangwan NS, Sangwan RS: *Andrographis paniculata* (Kalmegh): A review. *Pharmacogn Rev* 2007, 1:283–298.
- Chao WW, Lin BF: Isolation and identification of bioactive compounds in *Andrographis paniculata* (*Chuanxinlian*). *Chin Med* 2010, 5:1–15.
- Jarukamjorn K, Nemoto N: Pharmacological aspects of *Andrographis paniculata* on health and its major diterpenoid constituent andrographolide. *J Health Sci* 2008, 54:370–380.
- Niranjan A, Tewari SK, Lehri A: Biological activities of Kalmegh (*Andrographis paniculata* Nees) and its active principles-a review. *Indian J Nat Prod Resour* 2010, 1:125–135.
- Abu-Ghefreh AA, Canatan H, Ezeamuzie CI: *In vitro* and *in vivo* anti-inflammatory effects of andrographolide. *Int Immunopharmacol* 2009, 9:313–318.
- Kumar RA, Sridevi K, Kumar NV, Nanduri S, Rajagopal S: Anticancer and immunostimulatory compounds from *Andrographis paniculata*. *J Ethnopharmacol* 2004, 92:291–295.
- Woo AY, Waye MM, Tsui SK, Yeung ST, Cheng CH: Andrographolide up-regulates cellular-reduced glutathione level and protects cardiomyocytes against hypoxia/reoxygenation injury. *J Pharmacol Exp Ther* 2008, 325:226–235.
- Chao CY, Lii CK, Tsai IT, Li CC, Liu KL, Tsai CW, Chen HW: Andrographolide inhibits ICAM-1 expression and NF-κappaB activation in TNF-alpha-treated EA.hy926 cells. *J Agric Food Chem* 2011, 59:5263–5271.
- Yoopan N, Thisoda P, Rangkadilok N, Sahasitwat S, Pholphana N, Ruchirawat S, Satayavivad J: Cardiovascular effects of 14-deoxy-11, 12-didehydroandrographolide and *Andrographis paniculata* extracts. *Planta Med* 2007, 73:503–511.
- Thisoda P, Rangkadilok N, Pholphana N, Worasuttiayangkurn L, Ruchirawat S, Satayavivad J: Inhibitory effect of *Andrographis paniculata* extract and its active diterpenoids on platelet aggregation. *Eur J Pharmacol* 2006, 553:39–45.
- Kadem RE, Sang S, Ho CT: Mechanism of the superoxide scavenging activity of neoandrographolide - a natural product from *Andrographis paniculata* Nees. *J Agric Food Chem* 2002, 50:4662–4665.
- Batkhuu J, Hattori K, Takano F, Fushiya S, Oshiman K, Fujimiya Y: Suppression of NO production in activated macrophages *in vitro* and *ex vivo* by neoandrographolide isolated from *Andrographis paniculata*. *Biol Pharm Bull* 2002, 25:1169–1174.
- Misra P, Pal NL, Guru PY, Katiyar JC, Srivastava V, Tandon JS: Antimalarial activity of *Andrographis paniculata* (Kalmegh) against *Plasmodium berghei* NK 65 in *Mastomys natalensis*. *Pharm Biol* 1992, 30:263–274.
- Kapil A, Koul IB, Banerjee SK, Gupta BD: Antihepatotoxic effects of major diterpenoid constituents of *Andrographis paniculata*. *Biochem Pharmacol* 1993, 46:182–185.
- Burgos RA, Hidalgo MA, Monsalve J, LaBranche TP, Eyre P, Hancke JL: 14-deoxyandrographolide as a platelet activating factor antagonist in bovine neutrophils. *Planta Med* 2005, 71:604–608.
- Zhang CY, Tan BKH: Vasorelaxation of rat thoracic aorta caused by 14-deoxyandrographolide. *Clin Exp Pharmacol Physiol* 1998, 25:424–429.
- Burgos RA, Loyola M, Hidalgo MA, LaBranche TP, Hancke JL: Effect of 14-deoxyandrographolide on calcium-mediated rat uterine smooth muscle contractility. *Phytother Res* 2003, 17:1011–1015.
- Sharma A, Shanker C, Tyagi LK, Singh M, Rao CV: Herbal medicine for market potential in India: An Overview. *Academic J Plant Sci* 2008, 1:26–36.
- Kanjilal PB, Bordoloi S, Kalita R, Burman P, Singh R: *Cultivation practices for Kalmegh (*Andrographis paniculata*) and Spiderling (*Boerhaavia diffusa*) in Assam India*. USA: Studium Press LLS; 2002.
- Kumar B, Verma SK, Singh HP: Effect of temperature on seed germination parameters in Kalmegh (*Andrographis paniculata* Wall. ex Nees.). *Ind Crop Prod* 2011, 34:1241–1244.
- Anuradha VE, Jaleel CA, Salem MA, Gomathinayagam M, Panneerselvam R: Plant growth regulators induced changes in antioxidant potential and andrographolide content in *Andrographis paniculata* Wall. ex Nees. *Pestic Biochem Physiol* 2010, 98:312–316.
- Bhan MK, Dhar AK, Khan S, Lattoo SK, Gupta KK, Choudhary DK: Screening and optimization of *Andrographis paniculata* (Burm.f.) Nees for total andrographolide content, yield and its components. *Sci Hortic* 2006, 107:386–391.
- Prathanturarug S, Soonthornchareonnon N, Chuakul W, Saralamp P: Variation in growth and diterpene lactones among field-cultivated *Andrographis paniculata*. *J Nat Med* 2007, 61:159–163.

24. Parasher R, Upadhyay A, Khan NA, Dwivedi SK: Biochemical estimation and quantitative determination of medicinally important andrographolide in *Andrographis paniculata* at different growth stages. *EJAFChe* 2011, **10**:2479–2486.
25. Pholphana N, Rangkadilok N, Thongnest S, Ruchirawat S, Ruchirawat M, Satayavivad J: Determination and variation of three active diterpenoids in *Andrographis paniculata* (Burm.f.) Nees. *Phytochem Anal* 2004, **15**:365–371.
26. Matsuda T, Kuroyanagi M, Sugiyama S, Umehara K, Ueno A, Nishi K: Cell differentiation-inducing diterpenes from *Andrographis paniculata* NEES. *Chem Pharm Bull* 1994, **42**:1216–1225.
27. Department of Medical Sciences, Ministry of Public Health: *Thai Herbal Pharmacopoeia. Volume I*. Bangkok: Prachachon Publishing; 1995:24–31.
28. Xu T, Pan J, Zhao L: Simultaneous determination of four andrographolides in *Andrographis paniculata* Nees by silver ion reversed-phase high-performance liquid chromatography. *J Chromatogr Sci* 2008, **46**:747–750.
29. Sharma A, Lal K, Handa SS: Standardization of the Indian crude drug kalmegh by high pressure liquid chromatographic determination of andrographolide. *Phytochem Anal* 1992, **3**:129–131.
30. Lomlim L, Jirayupong N, Plubrukarn A: Heat-accelerated degradation of solid-state andrographolide. *Chem Pharm Bull* 2003, **51**:24–26.
31. Zhang CY, Kuroyangi M, Tan BKH: Cardiovascular activity of 14-deoxy-11,12-didehydroandrographolide in the anaesthetized rat and isolated right atria. *Pharmac Res* 1998, **38**:413–417.

doi:10.1186/1749-8546-8-2

Cite this article as: Pholphana et al.: Changes in the contents of four active diterpenoids at different growth stages in *Andrographis paniculata* (Burm.f.) Nees (*Chuanxinlian*). *Chinese Medicine* 2013 **8**:2.

Submit your next manuscript to BioMed Central and take full advantage of:

- Convenient online submission
- Thorough peer review
- No space constraints or color figure charges
- Immediate publication on acceptance
- Inclusion in PubMed, CAS, Scopus and Google Scholar
- Research which is freely available for redistribution

Submit your manuscript at
www.biomedcentral.com/submit

