

(R)- β -Lysine-modified Elongation Factor P Functions in Translation Elongation*

Received for publication, November 21, 2012, and in revised form, December 27, 2012. Published, JBC Papers in Press, December 31, 2012, DOI 10.1074/jbc.M112.438879

Tammy J. Bullwinkle[‡], S. Betty Zou^{§1}, Andrei Rajkovic[¶], Steven J. Hersch^{§1}, Sara Elgamal[‡], Nathaniel Robinson[‡], David Smil[¶], Yuri Bolshan[¶], William Wiley Navarre^{§2}, and Michael Ibba^{‡¶13}

From the [‡]Department of Microbiology and [¶]Molecular, Cell, and Developmental Biology Program, Ohio State University, Columbus, Ohio 43210 and the [§]Department of Molecular Genetics and [¶]Structural Genomics Consortium, University of Toronto, Toronto, Ontario M5S 1A8, Canada

Background: Post-translational modification activates bacterial elongation factor P (EF-P) in several Gram-negative bacteria.

Results: The addition of β -lysine alone is sufficient for activation of EF-P to function in translation.

Conclusion: Modified EF-P acts by regulating translation of a subset of mRNAs.

Significance: EF-P can post-transcriptionally regulate gene expression by controlling translation elongation.

Post-translational modification of bacterial elongation factor P (EF-P) with (R)- β -lysine at a conserved lysine residue activates the protein *in vivo* and increases puromycin reactivity of the ribosome *in vitro*. The additional hydroxylation of EF-P at the same lysine residue by the YfcM protein has also recently been described. The roles of modified and unmodified EF-P during different steps in translation, and how this correlates to its physiological role in the cell, have recently been linked to the synthesis of polyproline stretches in proteins. Polysome analysis indicated that EF-P functions in translation elongation, rather than initiation as proposed previously. This was further supported by the inability of EF-P to enhance the rate of formation of fMet-Lys or fMet-Phe, indicating that the role of EF-P is not to specifically stimulate formation of the first peptide bond. Investigation of hydroxyl-(β)-lysyl-EF-P showed 30% increased puromycin reactivity but no differences in dipeptide synthesis rates when compared with the β -lysylated form. Unlike disruption of the other genes required for EF-P modification, deletion of *yfcM* had no phenotypic consequences in *Salmonella*. Taken together, our findings indicate that EF-P functions in translation elongation, a role critically dependent on post-translational β -lysylation but not hydroxylation.

Elongation factor P (EF-P)⁴ is a highly conserved bacterial protein that can increase the efficiency of translation *in vitro* but is not required for ribosome activity and is dispensable for viability in *Escherichia coli* and *Salmonella enterica* (1–3). Structural analyses showed that EF-P can bind to the ribosome

between the P and E sites while making several interactions with the 3' acceptor stem of a P-site-bound initiator tRNA (4). Based on these structural data and other functional studies, it was proposed that EF-P specifically promotes formation of the first peptide bond during protein synthesis. However, two recent studies also indicated that EF-P could act during elongation by facilitating the synthesis of proteins containing stretches of consecutive proline residues (5, 6). EF-P is similar in size and shape to a tRNA and can be post-translationally modified by a lysyl-tRNA synthetase paralog, PoxA (7, 8). In a reaction analogous to tRNA aminoacylation, PoxA activates (R)- β -lysine and subsequently transfers it to a conserved residue (Lys-34) in EF-P (9). Attachment of a β -lysyl moiety is necessary for EF-P functionality both *in vivo* and *in vitro* (2, 8). Recent mass spectroscopy analyses identified a second post-translational modification of EF-P involving hydroxylation of the C4(γ) or C5(δ) of β -lysyl-Lys-34 by YfcM (10). However, the role of hydroxylation in EF-P function remains unclear.

In *S. enterica*, disruptions of the genes encoding EF-P or the β -lysyl modification pathway result in a wide range of phenotypic changes including increased sensitivity to several antibiotics, loss of motility, attenuated virulence, and sensitivity to detergents and low osmolarity conditions (8). Many of these phenotypes are consistent with increased membrane permeability, suggesting a role for EF-P in regulating expression of outer membrane proteins, such as KdgM (3). Proteomic analyses indicated that expression of a subset of genes is under post-transcriptional regulation by EF-P and that this activity is dependent on the β -Lys modification (3).

EF-P is a homolog of eukaryotic initiation factor 5A (eIF5A), which is also post-translationally modified at a conserved lysine residue analogous to Lys-34 of EF-P. eIF5A is modified with a hypusine residue through a two-step pathway different from the bacterial EF-P modification pathway. Hypusinated eIF5A is essential in yeast and is required for post-transcriptional regulation of a subset of genes, including those encoding factors involved in cell cycle progression and growth under stress conditions (11). The detection of eIF5A in polysomes parallels other elongation factors, leading to the inference of a role for

* This work was supported, in whole or in part, by National Institutes of Health Grant GM 65183.

¹ Funded by Vanier Canada Graduate Scholarships.

² Supported by an Operating Grant and a New Investigator Award from the Canada Institutes for Health Research (MOP-86683 and MSH-87729) and a Discovery Grant from the Natural Sciences and Engineering Research Council of Canada (RGPIN 386286-10).

³ To whom correspondence should be addressed: Dept. of Microbiology, Ohio State University, 484 W. 12th Ave., Columbus, OH 43210. Tel.: 614-292-2120; Fax: 614-292-8120; E-mail:ibba.1@osu.edu.

⁴ The abbreviations used are: EF-P, elongation factor P; 70SIC, 70 S initiation complexes; PM, polysome/monosome.

this factor in translation elongation. Further evidence for a role in translation elongation comes from the ability of eIF5A to promote *in vitro* di- and tripeptide synthesis (12). To date, the effects of EF-P and its β -lysine modification on translation have largely been investigated using a puromycin reactivity assay, which does not accurately represent a physiologically relevant peptide synthesis reaction (2, 13, 14). We now show that EF-P functions in translation elongation, with activity dependent on β -lysylation, whereas the additional hydroxyl modification is dispensable. The possible role of YfcM-catalyzed hydroxylation in modulating EF-P activity is discussed.

EXPERIMENTAL PROCEDURES

Strains, Plasmids, and General Methods—EF-Tu, EF-G, and translation initiation factors were prepared as described previously (15). *E. coli* BL21(DE3)/pYTB11-*efp*, WT and K34R, and *E. coli* BL21(DE3)/pYTB11-*poxA* WT were prepared as described (8). Intein-tagged proteins were purified on a chitin affinity column (New England Biolabs) and stored in 25 mM Tris-HCl, pH 8.0, 150 mM NaCl, 4 mM 2-mercaptoethanol, 20% glycerol. *E. coli* BL21(DE3)/pQE31-*Ec-lysS* and *E. coli* BL21(DE3)/pQE31-*Ec-pheS-pheT* were used to prepare lysyl-tRNA synthetase and phenylalanyl-tRNA synthetase, respectively (16, 17). Native tRNAs were purchased from Chemical Block (Moscow, Russia). [³⁵S]fMet-tRNA^{fMet} was prepared as described previously (18). Wild type BW25113 and Δ *efp* *E. coli* strains were obtained from the Keio collection, and kanamycin cassettes were removed via pCP20-encoded FLP recombinase (19, 20). The *S. enterica* Δ *yfcM* mutant was constructed according to the protocol of Datsenko and Wanner (20) using the *red-gam* recombinase as described previously. Briefly, the *yfcM* gene was replaced by a kanamycin resistance gene cassette amplified from the plasmid pKD4 using primers WNp658 (5'-CCACGGACAGGAGATCCTCCACTGGTTGGGGATGATTAAGTGTAGGCTGGAGCTGCTTC) and WNp659 (5'-CGTCAATTAATGCCAGAATGCGTGATTCAAACCTCCGCGATCATATGAATATCCTCCTTAG). The null allele was moved into a fresh 14028s background by transduction using phage P22 HT105/1 *int-201* prior to downstream analyses. Verification of the knock-out strain was carried out by PCR amplification using primers flanking the *yfcM* region.

Ribosome Preparation and Analysis—Strains were grown at 37 °C, 250 rpm to $A_{600\text{ nm}}$ 0.4–0.5, and as indicated, 100 μ g/ml chloramphenicol was added to the culture, and the culture was incubated for 2 min. Cells were harvested at 5,000 \times g for 5 min over blocked ice or blocked ice supplemented with 100 μ g/ml chloramphenicol as appropriate. Cell pellets were immediately suspended in lysis buffer (20 mM Tris-HCl, pH 8.0, 10.5 mM MgCl₂, 40 units/ml RNase inhibitor (Roche Applied Science), and 100 units/ml Turbo[®] DNase (New England Biolabs)) and lysed by freeze/thaw. Lysate was loaded onto 10/40% sucrose gradients and separated by ultracentrifugation at 35,000 rpm (151,000 \times g) for 2.5 h at 4 °C. Fractions were harvested from the gradient, and ribosomes were detected by monitoring $A_{254\text{ nm}}$.

For detection of EF-P via Western blotting, ribosomal fractions were first normalized by $A_{260\text{ nm}}$, and equal amounts of ribosomes were then analyzed by SDS-PAGE. Proteins were

transferred to nitrocellulose and probed with anti-EF-P polyclonal antibodies (ProSci) followed by visualization using anti-rabbit-HRP secondary antibody.

Tightly coupled 70 S ribosomes were prepared as described previously (21). *E. coli* strain BW25113 Δ *efp* was harvested and lysed, and ribosomal fractions were separated by ultracentrifugation on a 10–40% sucrose gradient. 70 S fractions were pooled and stored in ribosome buffer A (20 mM Tris-HCl, pH 7.5, 10.5 mM MgCl₂, 100 mM NH₄Cl₂, 0.5 mM EDTA, and 6 mM 2-mercaptoethanol). 30 S and 50 S subunits were purified similarly with a dialysis step in 30/50 buffer (20 mM Tris-HCl, pH 7.5, 1 mM MgCl₂, 100 mM NH₄Cl₂, 0.5 mM EDTA, and 6 mM 2-mercaptoethanol) for 3 h prior to loading a 10–30% sucrose gradient.

Modification of EF-P with β -Lysine—Purified EF-P (40 μ M) was incubated with 200 μ M β -lysine (9), 10 mM ATP, 1 \times buffer (100 mM glycine, pH 9.0, 30 mM KCl, 10 mM MgCl₂, 1 mM DTT), and 5 μ M PoxA. Reactions were performed at 37 °C for 1 h. The reaction mixture was then dialyzed against 20 mM Tris-HCl, 100 mM KCl, 2 mM 2-mercaptoethanol, and 10% glycerol. The degree of EF-P aminoacylation (routinely >99%) was monitored by one-dimensional isoelectric focusing followed by Western blotting with anti-EF-P as described above.

Native EF-P Purification—Purification of native EF-P was adapted from Refs. 2, 10, and 22. Wild type *E. coli* MRE600 cells were grown in autoinduction medium (0.5% glycerol, 0.25% glucose, 0.33% ammonium sulfate, 0.68% potassium dihydrogen phosphate, and 0.71% sodium phosphate dibasic) overnight at 37 °C. Cells were harvested by centrifugation at 6,000 rpm (3,600 \times g) for 10 min. The cell pellet was resuspended in buffer A (25 mM Tris-HCl, pH 8, 150 mM NaCl, 10% glycerol, 1 mM 2-mercaptoethanol, 0.75 mM PMSF) and passed through a French press. Lysate was centrifuged at 75,000 \times g for 75 min, and the resulting supernatant was retained. Proteins were precipitated from the supernatant at 35–55% ammonium sulfate, dissolved, and then dialyzed overnight against buffer B (25 mM Tris-HCl, pH 8, 50 mM NaCl, 10% glycerol, 1 mM 2-mercaptoethanol). Following dialysis, the sample was loaded onto a HiPrep XK26 Sepharose Q column, washed with buffer C (25 mM Tris-HCl, pH 8, 50 mM NaCl, 1 mM 2-mercaptoethanol), and developed with a linear gradient of 50 mM to 1 M NaCl. Fractions containing EF-P, as assessed by immunoblotting, were pooled and applied to a HiLoad 26/600 Superdex 200 prep grade column and developed in buffer C. EF-P-containing fractions were pooled, dialyzed against buffer B, and then loaded onto a MonoQ column, which was then developed with a linear gradient of 50 mM to 1 M NaCl. EF-P-containing fractions were pooled, concentrated with an Amicon Ultra centrifugal filter (molecular weight cut-off 10,000), loaded onto a Superose 12 HR 10/30 size exclusion column, and eluted in buffer C. Fractions containing EF-P were pooled, concentrated, and dialyzed against buffer D (25 mM Tris-HCl, pH 8, 150 mM NaCl, 1 mM 2-mercaptoethanol, 20% glycerol). Aliquots of EF-P were flash-frozen in liquid nitrogen and stored at –80 °C.

Mass Spectroscopy—Native EF-P protein was separated on a 13% SDS-PAGE gel and then fixed in a solution of 50% (v/v) ethanol and 10% (v/v) acetic acid for 1 h. Solvent was removed, and the gel was then washed in 50% (v/v) methanol and 10%

EF-P Functions in Translation Elongation

(v/v) acetic acid overnight followed by staining for 4 h at room temperature with 0.1% (w/v) Coomassie Blue, 20% (v/v) methanol, and 10% (v/v) acetic acid. The gel was destained with 50% (v/v) methanol and 10% (v/v) acetic acid, and the EF-P containing band was excised and stored in 5% (v/v) acetic acid. The sample was then analyzed using ultrahigh resolution TOF liquid chromatography tandem mass spectrometry at the Ohio State University Campus Chemical Instrument Center (CCIC) Mass Spectrometry and Proteomics Facility. Analysis of modifications on natively purified *E. coli* EF-P revealed that Lys-34 was 100% β -lysylated and >98% hydroxylated.

Subunit Docking—30 S preinitiation complexes were formed by incubating 0.1 μ M 30 S ribosome subunits, 0.2 μ M fMet-tRNA^{fMet}, 0.5 μ M *poxB* mRNA, 0.2 μ M initiation factors, and 0.1 mM GTP in ribosome buffer A for 20 min at 37 °C. 0.3 μ M 50 S ribosome subunits were preincubated with 1 μ M EF-P in ribosome buffer A for 10 min at 21 °C and then rapidly mixed at equal volumes with the preinitiation complex in an SX20 stopped-flow apparatus at 21 °C. Light scattering over time was monitored without a cutoff filter at an angle perpendicular to the excitation beam (430 nm).

Poly(U)-directed Poly-Phe Synthesis—Poly-Phe synthesis was adapted from a previous protocol (23). Briefly, 0.2 μ M 70 S ribosomes, 1 μ M EF-G, 5 μ M activated EF-Tu, and 0.3 μ g/ μ l polyuridine RNA were incubated in poly-Phe buffer (50 mM Tris-HCl, pH 7.5, 5 mM MgCl₂, 20 mM NH₄Cl, 1 mM DTT, 30 mM DTT, 0.5 mM GTP) in the presence or absence of EF-P at 21 °C for 10 min. To initiate the reaction, 1 μ M [¹⁴C]Phe-tRNA^{Phe} was added. Time points were quenched on 10% TCA-soaked filter paper and incubated for 20 min before washing two times in 5% TCA at 90 °C. Filter papers were then washed once with ethanol, dried, and quantified by scintillation counting.

fMet-Puromycin Reactivity Assay—70 S initiation complexes (70SIC) were formed as follows. 70 S ribosomes (2 μ M) were incubated with 2 μ M [³⁵S]fMet-tRNA^{fMet}, 4 μ M *poxB* upstream mRNA, 5'-CAGGAGAUGGAGAACCAUGAAACAA-3' (start codon underlined), 2 μ M each of IF1, IF2, IF3, and 1 mM GTP in polymix buffer (5 mM KPO₄, pH 7.5, 1 mM DTT, 5 mM Mg(OAc)₂, 0.5 mM CaCl₂, 95 mM KCl, 5 mM NH₄Cl, 8 mM putrescine, 1 mM spermidine) for 30 min at 37 °C.

70SIC (0.2 μ M) were incubated with 2 mM EF-P (or equal mass BSA) in polymix buffer for 10 min at 21 °C. Equal volume of EF-P-bound 70SIC (0.1 μ M final) was added to puromycin (1 μ M final) to initiate the reaction. The reaction was quenched at given time points in 0.5 mM KOH. Unreacted [³⁵S]fMet-tRNA^{fMet} was deacylated by incubating quenched samples at 37 °C for 20 min. The [³⁵S]fMet-puromycin dipeptide was separated from unreacted [³⁵S]fMet using silica TLC (running buffer of 4:1:1 butanol:acetic acid:H₂O).

fMet-Lys and fMet-Phe Dipeptide Formation—Aminoacylation of tRNA^{Lys} or tRNA^{Phe} was performed in 0.1 M Na-HEPES, pH 7.2, 30 mM KCl, 10 mM MgCl₂, 10 mM DTT, and 8 mM ATP. 1–10 μ M native tRNA was incubated in above conditions with 100 μ M lysine or phenylalanine and 1–2 μ M lysyl-tRNA synthetase or phenylalanyl-tRNA synthetase for 20 min at 37 °C. Aminoacyl-tRNA was purified by phenol:chloroform extraction followed by passage over a G-25 spin column pre-equilibrated with 5 mM sodium acetate (pH 5.2).

EF-Tu was first activated with 50 μ g/ml pyruvate kinase in a buffer containing 50 mM Tris-HCl, pH 7.5, 5 mM MgCl₂, 20 mM NH₄Cl, 1 mM DTT in the presence of 5 mM fresh P-enolpyruvate and 0.5 mM GTP for 30 min at 37 °C. Activated EF-Tu (10 μ M) was incubated with 2 μ M Lys-tRNA^{Lys} for 20 min on ice to form the GTP-EF-Tu-aminoacyl-tRNA ternary complex. 70SIC (0.2 μ M) were incubated with 2 μ M EF-P (or equal mass BSA) in 1 \times polymix buffer for 10 min at 21 °C. Using a KinTek rapid quench apparatus, an equal volume of EF-P-bound 70SIC (0.1 μ M final) was added to ternary complex (1 μ M final) to initiate the reaction. The reaction was quenched at given time points in 0.5 M KOH. Unreacted [³⁵S]fMet-tRNA was deacylated by incubating quenched samples at 37 °C for 20 min. The [³⁵S]fMet-Lys dipeptide was separated from unreacted [³⁵S]fMet using electrophoretic TLC in pyridine/acetic acid buffer.

To improve product signal, fMet-Phe dipeptide formation was also performed under pseudo-first order conditions, but with limiting amounts of radiolabeled acceptor tRNA in complex with EF-Tu (0.1 μ M) and excess 70SIC with cold fMet-tRNA^{fMet} (1 μ M). Prior to aminoacylation, tRNA^{Phe} was 3'-labeled with α [³²P]ATP using snake venom phosphodiesterase (Sigma) and tRNA nucleotidyl transferase (24) or tRNA nucleotidyl transferase alone (25). After dipeptide formation and rapid quench (0.5 M sodium acetate, pH 4.5), the fMet-Phe-[³²P]tRNA^{Phe} product was cleaved using S1 endonuclease (Promega) to release fMet-Phe-[³²P]AMP, which was separated from unreacted Phe-[³²P]AMP on cellulose TLC (75:15:15 isopropyl alcohol:HCl:H₂O).

Growth Assays—Sensitivity to hypoosmolarity, gentamicin, lauryl sulfobetaine, and *S*-nitrosoglutathione was assessed as described previously (3). The migratory ability of the $\Delta yfcM$ mutant was determined using soft agar assays, and accumulation of 1-*N*-phenyl-naphthylamine was conducted to measure membrane permeability, both as described previously (3).

RESULTS

Polysome Profiles Indicate a Role for EF-P in Translation Elongation—Polysomes were isolated and fractionated from *E. coli* wild type and Δefp strains to investigate at which stages in translation this factor might function. Polysome/monosome (P/M) ratios were then compared, similar to previous studies involving the eukaryotic EF-P homolog eIF5a (12). We reasoned that if EF-P functions to promote the elongation reaction, then the P/M ratio would increase in the Δefp strain relative to wild type because ribosomes would move more slowly and accumulate to higher levels on mRNA. Chloramphenicol, an inhibitor of peptide synthesis that increases polysome retention, was used as a positive control. In the absence of chloramphenicol, the Δefp strain showed a higher P/M ratio when compared with wild type consistent with a reduced rate of translation elongation (Fig. 1). The addition of chloramphenicol to the wild type culture also resulted in an increase in the P/M ratio relative to polysomes isolated in the absence of the inhibitor. For the Δefp strain, the addition of chloramphenicol did not further increase the P/M ratio. The translational effects of hydroxylation or β -lysylation of EF-P were also investigated by comparing P/M ratios from ribosomes isolated from $\Delta yfcM$ and $\Delta poxA$ strains in the presence and absence of chloram-




FIGURE 1. **Deletion of *efp* results in increased retention of polysomes.** Shown is the quantification of representative A_{254} profiles of ribosomal fractions isolated from BW25113 (WT) or Δefp , $\Delta poxA$, or $\Delta yfcM$ *E. coli* strains and separated via sucrose gradient centrifugation. Cultures were grown to mid-log and in some cases (+) exposed to 100 $\mu\text{g/ml}$ chloramphenicol (CAM) for 2 min prior to harvesting. The P/M ratio for each fractionation is calculated from three separate experiments.

phenicol. Although some differences were observed in the appearance of the fractionated ribosome profiles (Fig. 1), the overall level of polysome retention for the $\Delta yfcM$ strain in the absence of chloramphenicol was only slightly higher than for wild type *E. coli* and increased in the presence of the antibiotic. The $\Delta poxA$ strain exhibited increased polysome retention relative to $\Delta yfcM$, with P/M ratios comparable with the Δefp strain in the presence and absence of chloramphenicol (Fig. 1). These results suggest that the role of EF-P in translation elongation requires β -lysylation by PoxA but is not critically dependent on YfcM-mediated hydroxylation.

In contrast to the data above, previous studies (4) suggested that EF-P functions during translation initiation by stabilizing initiator tRNA on the ribosome. To further examine possible roles of EF-P in translation initiation, the rate of 30 S and 50 S subunit joining was measured during formation of the 70 S initiation complex. Subunit joining was measured using a 30 S preinitiation complex loaded with the initiation region of *poxB*, a gene previously shown to be regulated by EF-P (8), mRNA, and fMet-tRNA^{fMet}. No difference in the rate of subunit joining was observed in the presence or absence of unmodified EF-P or β -Lys-EF-P, suggesting that EF-P does not function in initiation complex formation (Fig. 2).

EF-P Does Not Function as an Essential Translation Factor—The above analyses of polysome profiles, and previous investigations of puromycin reactivity (2, 22), support a role for modified EF-P in translation elongation. To further analyze the effect of EF-P β -lysylation and hydroxylation on translation, puromycin reactivity was measured in the presence of modified and unmodified EF-P and a K34R variant that mimics a conserved replacement seen in numerous bacterial EF-Ps (26). β -Lys-modified EF-P was generated by *in vitro* aminoacylation, whereas the hydroxylated form was purified in its native form from *E. coli*. Neither the addition of unmodified EF-P nor the addition of the K34R variant had any effect on puromycin reactivity (Fig. 3). The k_{obs} for fMet-puromycin formation increased 3-fold in the presence of β -Lys-EF-P and 4-fold when Lys-34 was also hydroxylated (Fig. 3). These data support a role for modified EF-P in translation elongation and indicate that hydroxylation is important but not critical for this function.




FIGURE 2. **Initiation of translation of *poxB* mRNA in the presence of β -Lys-EF-P.** Representative graphs of 50 S and 30 S preinitiation complex joining are shown as light scattering over time. Rates were measured in the presence or absence of unmodified EF-P or β -Lys-EF-P and fit to the single exponential equation ($y = m1 \times \exp(-m0 \times m2) + m3 + m4 \times m0$). $k_{\text{obs}} = 0.0088 \pm 0.0013 \text{ s}^{-1}$ (BSA), $0.0089 \pm 0.0009 \text{ s}^{-1}$ (EF-P), and $0.0082 \pm 0.0027 \text{ s}^{-1}$ (β -Lys-EF-P).

The role of EF-P modification in translation was further investigated using *in vitro* dipeptide and polypeptide synthesis assays. The effects of EF-P on dipeptide formation were measured using a ribosomal 70 S initiation complex containing the initiation region of *poxB* mRNA and fMet-tRNA^{fMet} in the P-site. The second codon (in the ribosomal A-site) was AAA, which encodes lysine. Excess EF-Tu-bound Lys-tRNA^{Lys} was then added, and the formation of the fMet-Lys dipeptide was monitored. Neither the modified nor the unmodified EF-P variants had any significant effect on the rate of fMet-Lys dipeptide formation (Fig. 4). It had previously been proposed that the size of the acceptor amino acid determines the extent of the effect of EF-P on peptide bond formation (32). To test the role of the acceptor amino acid, the second codon of the *poxB* message was changed to UUA (Phe), and the rate of fMet-Phe synthesis measured. No effects of modified or unmodified EF-P on the

EF-P Functions in Translation Elongation




FIGURE 3. [³⁵S]fMet-puromycin formation by the ribosome in the presence of β -Lys-EF-P and hydroxyl- β -Lys-EF-P. Puromycin reactions were performed under pseudo-first order conditions using 70SIC preincubated with variants of EF-P or BSA. Error bars represent the S.D. of three separate experiments.




FIGURE 4. fMet-Lys dipeptide synthesis in the presence of β -Lys-EF-P. Dipeptide formation assays were performed under pseudo-first order conditions using 70SIC preincubated with variants of EF-P or BSA. Error bars represent the S.D. of three separate experiments.

rate of fMet-Phe synthesis were observed (data not shown). To further investigate the role of EF-P as a translation factor, its impact on polypeptide synthesis was measured using a poly(U) RNA template. As with dipeptide synthesis, neither unmodified or modified EF-P had any effect on the rate of poly-Phe formation (Fig. 5). Taken together, these studies on di- and polypeptide synthesis indicate that EF-P is not an essential translation elongation factor and, as suggested previously from proteomic data, may instead only affect the translation of a particular subset of mRNAs (27).

YfcM Is Dispensable for EF-P-dependent Growth Phenotypes—Previous studies have shown that the pathway encoded by the genes *efp*, *poxA*, and *yjeK* is linked to a number of growth phenotypes in *E. coli* and *S. enterica* (3, 8). To test the effects of




FIGURE 5. Elongation of a poly-Phe peptide in the presence of modified EF-P. The rate of poly(U)-directed poly-Phe synthesis by 70S ribosomes was measured by the incorporation of [¹⁴C]Phe. Reactions were performed in the presence or absence of unmodified EF-P (●), β -Lys-EF-P (▲), or BSA (■), preincubated with 70S ribosomes. Error bars represent the S.D. of three separate experiments.

hydroxylation on EF-P activity *in vivo*, an *S. enterica* $\Delta yfcM$ strain was characterized with respect to phenotypes previously described for Δefp , $\Delta poxA$, and $\Delta yjeK$ strains (3, 8). The phenotypes of the $\Delta yfcM$ mutant were very similar to those of wild type *Salmonella* when grown on the hypoosmolar antibiotic medium 2 agar and when exposed to gentamicin, lauryl sulfo-betaine, or *S*-nitrosoglutathione (Fig. 6, A–D). Unlike the Δefp , $\Delta poxA$, and $\Delta yjeK$ mutants, the $\Delta yfcM$ mutant did not exhibit a migration defect on a soft agar assay, nor any change in membrane permeability (Fig. 6, E and F). Similarly, characterization of *E. coli* mutants from the Keio collection showed no difference in growth between the $\Delta yfcM$ strain and wild type BW25113. In contrast, both the Δefp and the $\Delta poxA$ strains grew more slowly than their wild type counterparts (Fig. 7). These results indicate that YfcM-catalyzed hydroxylation has no effect on cellular functions known to be regulated by EF-P.

DISCUSSION

The β -lysylation of EF-P by the PoxA/YjeK pathway leads to stimulation of puromycin reactivity *in vitro* and post-transcriptional regulation of certain genes *in vivo* (2, 8). These studies and structural investigations (4) supported the hypothesis that EF-P acts by stabilizing initiator tRNA in the ribosomal P-site and promoting formation of the first peptide bond. However, there has been no other biochemical evidence to directly support a role for EF-P in initiation, whereas it was recently shown that EF-P facilitates synthesis of stretches of proline residues during translation elongation (5, 6). Consistent with this last point, our results also suggest that although modified EF-P does increase puromycin reactivity, it does not directly affect initiation, nor does it globally promote the rate of translation elongation. Although EF-P is not an essential translation factor, analyses of polysome profiles isolated in the presence and




FIGURE 6. *Salmonella* $\Delta yfcM$ mutants do not phenocopy $\Delta poxA$, $\Delta yjeK$, and Δefp mutants. *Salmonella* mutants deficient in *yfcM* exhibit phenotypes similar to that of wild type *Salmonella* 14028s when grown under hypoosmolar conditions (antibiotic medium 2 agar plates (A)) or when exposed to gentamicin (B), lauryl sulfobetaine (C), or *S*-nitrosoglutathione (GSNO) (D). $\Delta yfcM$ mutants also do not exhibit a migration defect on soft agar plates (E) or a change in 1-*N*-phenyl-naphthylamine accumulation (NPN) (F). Data points represent the average of three independent experiments. Error bars represent S.E. For 1-*N*-phenyl-naphthylamine accumulation, a strain lacking the AcrAB multidrug efflux pump served as a positive control, and standard errors are shown only at 0, 2.5, 5, and 9.5 min for clarity. AU, arbitrary fluorescence units.

absence of chloramphenicol indicate that EF-P is likely to be playing a role at some stage of elongation. Chloramphenicol binds the ribosomal A-site in the same place as the aminoacyl moiety of the A-site tRNA and blocks peptidyl transferase activity, leading to increased polysome retention (28). Disruption of the *efp* gene also led to increased retention of polysomes, indicating that EF-P functions by increasing translation elongation rates. Chloramphenicol has also been shown to inhibit the effect of EF-P on puromycin reactivity, further supporting the proposed role for the factor in translation elongation (29). The $\Delta poxA$ strain displayed similar changes when compared with the Δefp strain with respect to polysome retention and effects of chloramphenicol on the P/M ratio. However, less pronounced effects for $\Delta poxA$ when compared with Δefp strains were observed for several growth phenotypes (Figs. 6 and 7), suggesting that β -lysylation of EF-P is not required for the function of EF-P in translation elongation in all situations.

The exact mechanism by which EF-P affects translation elongation was previously unclear, in part due to the inherent limitations of puromycin reactivity assays to investigate ribosome peptidyl transferase function. Puromycin resembles an amino-

acyl-tRNA 3' end (tyrosyl-adenosine) and binds the A-site of the 50 S subunit. Although it is able to form a peptide bond with the P-site tRNA, translocation does not occur, and the ribosome is trapped in a "pretranslocation" state (30). Any effect EF-P may have on translocation or on the transition of the ribosome-bound tRNAs to the hybrid states would not be seen by monitoring puromycin reactivity. Furthermore, puromycin does not contain the critical tRNA residues needed for the rate-limiting accommodation step in the A-site (31). It was previously shown that EF-P also affects the rate of reaction between a P-site tRNA and CA-amino acids, which mimic the 3' end of an aminoacylated tRNA, but only in the case of small amino acids (32). The effects of EF-P on peptidyl transfer observed with these analogs were not observed with full-length aminoacyl-tRNA (Fig. 5), consistent with previous observations that extensive tRNA-A-site interactions during accommodation compensate for differences in the size of amino acids (33). Taken in the context of these structural and functional studies of bacterial ribosomes, the data described here suggest that EF-P does not act as a factor that generally promotes accommodation or translocation during translation elongation. Instead, the




FIGURE 7. Growth of $\Delta yfcM$, Δefp and $\Delta poxA$ *E. coli* strains. Growth in LB at 37 °C was monitored by A_{600} over time for BW25113 (WT) (■), $\Delta yfcM$ (▲), Δefp (●), and $\Delta poxA$ (★) *E. coli* strains. OD, optical density.

extensive phenotypic changes observed in Δefp strains and the accompanying changes in post-transcriptional control of gene expression agree with the recent proposal that EF-P is required for translation elongation for a particular subset of mRNAs encoding stretches of consecutive prolines (5, 6). Whether other sequences or structures within either the mRNA or the nascent peptide chain are also specifically targeted by EF-P during translation elongation is currently unclear.

Recent structural studies of EF-P identified an additional hydroxylation modification catalyzed by YfcM at the conserved Lys-34 residue of EF-P (10). Whether this modification is independent of β -lysylation of Lys-34, or is an integral part of the PoxA/YjeK/EF-P pathway, is unknown. Our data show that the absence of YfcM does not phenocopy loss of EF-P, PoxA, or YjeK. These results strongly suggest that YfcM is not a critical component of the modification pathway necessary for EF-P activity *in vivo* (8, 9). Although not essential for EF-P activation, hydroxylation further stimulated puromycin reactivity, suggesting that it may function to modulate EF-P activity. More extensive proteomic analyses, coupled with global profiling of actively translating ribosomes, are now warranted to investigate the exact mechanisms by which different modified forms of EF-P act as post-transcriptional regulators of gene expression.

Acknowledgments—We thank Dr. Kurt Fredrick and the members of the Fredrick laboratory for helpful advice and discussions as well as strains for the initiation factors. The Structural Genomics Consortium is a registered charity (Number 1097737) that receives funds from the Canadian Institutes for Health Research, Genome Canada, the Canadian Foundation for Innovation, the Ontario Innovation Trust, the Ontario Ministry for Research and Innovation, the Wellcome Trust, AbbVie, Eli Lilly Canada, GlaxoSmithKline, Novartis, Pfizer, and Takeda.

REFERENCES

- Aoki, H., Xu, J., Emili, A., Chosay, J. G., Golshani, A., and Ganoza, M. C. (2008) Interactions of elongation factor EF-P with the *Escherichia coli* ribosome. *FEBS J.* **275**, 671–681
- Park, J. H., Johansson, H. E., Aoki, H., Huang, B. X., Kim, H. Y., Ganoza, M. C., and Park, M. H. (2012) Post-translational modification by β -lysylation is required for activity of *Escherichia coli* elongation factor P (EF-P). *J. Biol. Chem.* **287**, 2579–2590
- Zou, S. B., Hersch, S. J., Roy, H., Wiggers, J. B., Leung, A. S., Buranyi, S., Xie, J. L., Dare, K., Ibba, M., and Navarre, W. W. (2012) Loss of elongation factor P disrupts bacterial outer membrane integrity. *J. Bacteriol.* **194**, 413–425
- Blaha, G., Stanley, R. E., and Steitz, T. A. (2009) Formation of the first peptide bond: the structure of EF-P bound to the 70S ribosome. *Science* **325**, 966–970
- Doerfel, L. K., Wohlgemuth, I., Kothe, C., Peske, F., Urlaub, H., and Rodnina, M. V. (2013) EF-P is essential for rapid synthesis of proteins containing consecutive proline residues. *Science* **339**, 85–88
- Ude, S., Lassak, J., Starosta, A. L., Kraxenberger, T., Wilson, D. N., and Jung, K. (2013) Translation elongation factor EF-P alleviates ribosome stalling at polyproline stretches. *Science* **339**, 82–85
- Yanagisawa, T., Sumida, T., Ishii, R., Takemoto, C., and Yokoyama, S. (2010) A paralog of lysyl-tRNA synthetase aminoacylates a conserved lysine residue in translation elongation factor P. *Nat. Struct. Mol. Biol.* **17**, 1136–1143
- Navarre, W. W., Zou, S. B., Roy, H., Xie, J. L., Savchenko, A., Singer, A., Edvokimova, E., Prost, L. R., Kumar, R., Ibba, M., and Fang, F. C. (2010) PoxA, YjeK, and elongation factor P coordinately modulate virulence and drug resistance in *Salmonella enterica*. *Mol. Cell* **39**, 209–221
- Roy, H., Zou, S. B., Bullwinkle, T. J., Wolfe, B. S., Gilreath, M. S., Forsyth, C. J., Navarre, W. W., and Ibba, M. (2011) The tRNA synthetase paralog PoxA modifies elongation factor-P with (*R*)- β -lysine. *Nat. Chem. Biol.* **7**, 667–669
- Peil, L., Starosta, A. L., Virumäe, K., Atkinson, G. C., Tenson, T., Remme, J., and Wilson, D. N. (2012) Lys34 of translation elongation factor EF-P is hydroxylated by YfcM. *Nat. Chem. Biol.* **8**, 695–697
- Park, M. H., Nishimura, K., Zanelli, C. F., and Valentini, S. R. (2010) Functional significance of eIF5A and its hypusine modification in eukaryotes. *Amino Acids* **38**, 491–500
- Saini, P., Eyler, D. E., Green, R., and Dever, T. E. (2009) Hypusine-containing protein eIF5A promotes translation elongation. *Nature* **459**, 118–121
- Ganoza, M. C., Zahid, N., and Baxter, R. M. (1985) Stimulation of peptidyltransferase reactions by a soluble protein. *Eur. J. Biochem.* **146**, 287–294
- Aoki, H., Adams, S. L., Turner, M. A., and Ganoza, M. C. (1997) Molecular characterization of the prokaryotic *efp* gene product involved in a peptidyltransferase reaction. *Biochimie* **79**, 7–11
- Ling, J., Yadavalli, S. S., and Ibba, M. (2007) Phenylalanyl-tRNA synthetase editing defects result in efficient mistranslation of phenylalanine codons as tyrosine. *RNA* **13**, 1881–1886
- Levengood, J., Ataide, S. F., Roy, H., and Ibba, M. (2004) Divergence in noncognate amino acid recognition between class I and class II lysyl-tRNA synthetases. *J. Biol. Chem.* **279**, 17707–17714
- Roy, H., Ling, J., Irnov, M., and Ibba, M. (2004) Post-transfer editing *in vitro* and *in vivo* by the β subunit of phenylalanyl-tRNA synthetase. *EMBO J.* **23**, 4639–4648
- Walker, S. E., and Fredrick, K. (2008) Preparation and evaluation of acylated tRNAs. *Methods* **44**, 81–86
- Baba, T., Ara, T., Hasegawa, M., Takai, Y., Okumura, Y., Baba, M., Datsenko, K. A., Tomita, M., Wanner, B. L., and Mori, H. (2006) Construction of *Escherichia coli* K-12 in-frame, single-gene knockout mutants: the Keio collection. *Mol. Syst. Biol.* **2**, 2006.0008
- Datsenko, K. A., and Wanner, B. L. (2000) One-step inactivation of chromosomal genes in *Escherichia coli* K-12 using PCR products. *Proc. Natl. Acad. Sci. U.S.A.* **97**, 6640–6645
- McGarry, K. G., Walker, S. E., Wang, H., and Fredrick, K. (2005) Destabilization of the P site codon-anticodon helix results from movement of

- tRNA into the P/E hybrid state within the ribosome. *Mol. Cell* **20**, 613–622
22. Aoki, H., Adams, S. L., Chung, D. G., Yaguchi, M., Chuang, S. E., and Ganoza, M. C. (1991) Cloning, sequencing, and overexpression of the gene for prokaryotic factor EF-P involved in peptide bond synthesis. *Nucleic Acids Res.* **19**, 6215–6220
23. Ling, J., So, B. R., Yadavalli, S. S., Roy, H., Shoji, S., Fredrick, K., Musier-Forsyth, K., and Ibba, M. (2009) Resampling and editing of mischarged tRNA prior to translation elongation. *Mol. Cell* **33**, 654–660
24. Gilreath, M. S., Roy, H., Bullwinkle, T. J., Katz, A., Navarre, W. W., and Ibba, M. (2011) β -Lysine discrimination by lysyl-tRNA synthetase. *FEBS Lett.* **585**, 3284–3288
25. Ledoux, S., and Uhlenbeck, O. C. (2008) [3'-32P]-labeling tRNA with nucleotidyltransferase for assaying aminoacylation and peptide bond formation. *Methods* **44**, 74–80
26. Bailly, M., and de Crécy-Lagard, V. (2010) Predicting the pathway involved in post-translational modification of elongation factor P in a subset of bacterial species. *Biol. Direct.* **5**, 3
27. Zou, S. B., Roy, H., Ibba, M., and Navarre, W. W. (2011) Elongation factor P mediates a novel post-transcriptional regulatory pathway critical for bacterial virulence. *Virulence* **2**, 147–151
28. Bulkley, D., Innis, C. A., Blaha, G., and Steitz, T. A. (2010) Revisiting the structures of several antibiotics bound to the bacterial ribosome. *Proc. Natl. Acad. Sci. U.S.A.* **107**, 17158–17163
29. Swaney, S., McCroskey, M., Shinabarger, D., Wang, Z., Turner, B. A., and Parker, C. N. (2006) Characterization of a high-throughput screening assay for inhibitors of elongation factor P and ribosomal peptidyl transferase activity. *J. Biomol. Screen.* **11**, 736–742
30. Schmeing, T. M., Seila, A. C., Hansen, J. L., Freeborn, B., Soukup, J. K., Scaringe, S. A., Strobel, S. A., Moore, P. B., and Steitz, T. A. (2002) A pre-translocational intermediate in protein synthesis observed in crystals of enzymatically active 50S subunits. *Nat. Struct. Biol.* **9**, 225–230
31. Beringer, M., and Rodnina, M. V. (2007) Importance of tRNA interactions with 23S rRNA for peptide bond formation on the ribosome: studies with substrate analogs. *Biol. Chem.* **388**, 687–691
32. Ganoza, M. C., and Aoki, H. (2000) Peptide bond synthesis: function of the *efp* gene product. *Biol. Chem.* **381**, 553–559
33. Dale, T., Fahlman, R. P., Olejniczak, M., and Uhlenbeck, O. C. (2009) Specificity of the ribosomal A site for aminoacyl-tRNAs. *Nucleic Acids Res.* **37**, 1202–1210