

Published in final edited form as:

Accid Anal Prev. 2013 March ; 51: 185–191. doi:10.1016/j.aap.2012.11.016.

ADOLESCENT EXPOSURE TO DRINK DRIVING AS A PREDICTOR OF YOUNG ADULTS' DRINK DRIVING

Tracy J. Evans-Whipp^{a,b,* , ^}, Stephanie M. Plenty^{a, ^}, John W. Toumbourou^{a,c}, Craig Olsson^{a,c}, Bosco Rowland^c, and Sheryl A. Hemphill^d

Tracy J. Evans-Whipp: tracy.evanswhipp@mcri.edu.au; Stephanie M. Plenty: stephanie.plenty@mcri.edu.au; John W. Toumbourou: john.toumbourou@deakin.edu.au; Craig Olsson: craig.olsson@rch.org.au; Bosco Rowland: bosco.rowland@deakin.edu.au; Sheryl A. Hemphill: sheryl.hemphill@acu.edu.au

^aCentre for Adolescent Health, Royal Children's Hospital & Adolescent Health Research, Murdoch Children's Research Institute, Parkville, Victoria, 3052, Australia

^bThe University of Melbourne Department of Paediatrics, Royal Children's Hospital, Parkville, Victoria, 3052, Australia

^cPrevention Sciences, School of Psychology and Centre for Mental Health and Wellbeing Research, Deakin University, Waterfront Campus, Geelong Victoria, 3217, Australia

^dSchool of Psychology, Australian Catholic University, Fitzroy, Victoria, 3065, Australia

Abstract

The purpose of this study was to investigate the influence of exposure to others' drink driving during adolescence on self-reported driving under the influence (DUI) of alcohol in young adulthood. Data were drawn from 1,956 participants with a driving license enrolled in the International Youth Development Study from Victoria, Australia. During 2003 and 2004, adolescents in Grades 7, 9 and 10 (aged 12 – 17) completed questionnaires examining whether they had ridden in a vehicle with a driver who had been drinking, as well as other demographic, individual, peer and family risk factors for DUI. In 2010, the same participants (aged 18 to 24) then reported on their own DUI behaviour. 18% of young adults with a driving license reported DUI in the past 12 months. Exposure to others' drink driving during adolescence was associated with an increased likelihood of DUI as a young adult (OR=2.13, 95% CI 1.68 – 2.69). This association remained after accounting for the effects of other potential confounding factors from the individual, peer and family domains (OR=1.62, 95% CI 1.23 – 2.13). Observing the drink driving behaviours of others during adolescence may increase the likelihood of DUI as a young adult. Strategies to reduce youth exposure to drink driving are warranted.

Keywords

drink driving; driving under the influence; adolescence; alcohol; riding with drinking driver

© 2012 Elsevier Ltd. All rights reserved.

*Author to whom correspondence should be addressed; tracy.evanswhipp@mcri.edu.au; Tel.: +613-9345-6677; Fax: +613-9345-6343.

[^]Tracy Evans-Whipp and Stephanie Plenty are joint first authors

Publisher's Disclaimer: This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final citable form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

1. Introduction

Traffic accidents and drink driving are serious health concerns for the young adult population. Road fatality is the leading cause of death among young adults in Australia and the US (Australian Institute of Health and Welfare, 2008; Australian Medical Association, 2009; Australian Transport Council, 2008; Hoyert *et al.*, 2005). For example, traffic accidents account for 31% of all deaths in young Australians aged 15 to 24 years (Australian Bureau of Statistics, 2008). This is higher than the road fatality rate in any other age group (Australian Institute of Health and Welfare, 2008; Transport Accident Commission of Victoria, 2003). Alcohol plays a key role in many fatal accidents (Siskind *et al.*, 2011), with up to 50% of U.S. road fatalities amongst 18 to 24 year olds involving alcohol (Hingson *et al.*, 2009; National Highway Traffic Safety Administration, 2004). Approximately 25% of all drivers killed on Australian roads have a blood alcohol concentration (BAC) above the legal limit (0.05%) (Australian Transport Council, 2008; Office of Road Safety, 2012). Additionally, young drivers are at increased risk of road accidents compared to older drivers with the same BAC (Cavallo and Triggs, 1996; Engström *et al.*, 2003; Palamara *et al.*, 2001).

Young adult drink driving is prevalent, with self-report estimates ranging from 20 to 25% in Australian and US studies (Beck *et al.*, 2010; Davey *et al.*, 2005; Hingson *et al.*, 2009; LaBrie *et al.*, 2011; Vassallo *et al.*, 2010). Given this high prevalence and that young adults are over-represented in road accidents there is an important need to identify approaches that can help reduce driving under the influence (DUI) of alcohol in this population.

Previous research has identified a number of key risk factors associated with DUI. These factors span the demographic, family, peer, personality and community domains. A key risk factor for DUI is frequent and intense (binge) alcohol use (Bingham *et al.*, 2007; LaBrie *et al.*, 2010; Tin Tin *et al.*, 2008; Vassallo *et al.*, 2010). Earlier alcohol initiation has also been shown to increase the likelihood of DUI (Hingson *et al.*, 2009; Tomas Dols *et al.*, 2010). Demographic risk factors include being male (Beck *et al.*, 2010; Bina *et al.*, 2006; LaBrie *et al.*, 2011; Vassallo *et al.*, 2007) and lower socio-economic status (Morrison *et al.*, 2002); (Vaez and Laflamme, 2005). Other individual characteristics associated with DUI include greater levels of aggression (Bingham *et al.*, 2007), delinquency (Bingham *et al.*, 2007), impulsivity (Pedersen and McCarthy, 2008) and sensation seeking (Fernandes *et al.*, 2007; Jonah *et al.*, 2001; Pedersen and McCarthy, 2008; Zakletskaia *et al.*, 2009). The family context is also relevant as poor family relationships (Tomas Dols *et al.*, 2010) and parent alcohol use (LaBrie *et al.*, 2010; Maldonado-Molina *et al.*, 2011) have been shown to predict DUI. Family factors can also play a protective role. For example, adolescent perceptions of supportive relationships with their parents and schools may protect against DUI (Tin Tin *et al.*, 2008). Within the broader community context, perceiving that others in the community hold permissive attitudes towards DUI has been shown to be associated with a greater tendency to drink drive (Bingham *et al.*, 2007; Chen *et al.*, 2008).

The attitudes others hold towards drink driving can be expected to exert a particularly strong influence during adolescence as attitudes and expectations of acceptable behaviour are developing. Children and adolescents rely on lifts from others and many high school students accept rides from drivers who have been drinking alcohol (Centers for Disease Control and Prevention, 2010). In this sense, the importance of other drivers as role models in the socialization process should not be underestimated. Social Learning Theory proposes that normative values and behaviours are learnt by observing others' actions (Bandura, 1977, 1986). There is a body of evidence that supports the social modelling effect in many areas of development, such as the relationship between parent alcohol use and a child's alcohol use (Ward *et al.*, 2010), as well as the accident driving records shared between

parents and their children (Ferguson *et al.*, 2001). The current study proposes that exposure during adolescence to others' drink driving is an important potentially modifiable risk factor worthy of investigation.

Some previous studies have explored the link between exposure to others' drink driving and DUI. Cross-sectional data from the U.S. National Survey on Drug Use and Health indicated that 16 and 17 year olds were more likely to drink drive if they lived with a mother or father who also had driven under the influence in the past year (Center for Behavioral Health Statistics and Quality, 2011). In a cross-sectional study of Canadian high school students, Leadbeater and colleagues (Leadbeater *et al.*, 2008) found that having ridden with a drink driving peer predicted students' DUI. This effect was stronger if they had more frequent exposure to adults' drink driving. In a longitudinal New Zealand study, Gulliver & Begg (2004) investigated the influence of riding with an alcohol impaired driver during adolescence on DUI at age 21. A path analysis showed that, for males, exposure to adult DUI at age 15 predicted travelling with an alcohol impaired peer at age 18, which in turn increased the chance of DUI at age 21. These studies highlight the possible influence of observing drink driving during adolescence. However, they have some limitations in terms of study design and consideration of potential confounders. The first two are cross-sectional and the third utilized a relatively small sample of only male young adults. Furthermore, none of the studies accounted for a range of important confounding factors. Therefore it is uncertain if the relationship between riding with a drinking driver and DUI would remain above and beyond the influence of other risk factors (such as sensation seeking). Furthermore, some other previous studies have combined measures of exposure to DUI with an individual's own DUI (e.g. Finken *et al.*, 1998; McCarthy and Pedersen, 2009), making it impossible to disentangle the effects of exposure compared to a young adult's prior behaviour. No single study has used a longitudinal design with a large sample of young adults whilst accounting for a range of risk factors for drink driving.

The purpose of this study is to investigate the influence of adolescent exposure to others' drink driving on later self-reported DUI in young adulthood. The study is based on a representative sample of Victorian Australian school students followed from adolescence to young adulthood. It is hypothesised that adolescent exposure to others' drink driving will increase the likelihood of DUI as a young adult. It is also expected that this effect will remain after accounting for the influence of other important confounding factors in the demographic, family, peer and individual domains.

2. Material and methods

2.1 Participants and procedure

Data for this paper come from a larger study, the International Youth Development Study (IYDS). The IYDS is a multi-cohort longitudinal study that investigates the etiology of substance use and associated problems for adolescents from Victoria, Australia and Washington State, U.S. In 2002 the IYDS used a two-stage cluster sampling approach to recruit statewide representative samples of students from three age cohorts, Grades 5, 7 and 9. Participants came from 152 schools in Victoria and 153 schools in Washington State. Across the three cohorts 2,885 (74.8%) of eligible Washington State parents and students consented to participate and 2,884 (73.5%) of eligible Victorian parents and students consented to participate. See (McMorris *et al.*, 2007) for more details regarding recruitment.

Data from the current study are drawn from Victorian participants. At time 1, a total of 2,821 participants in three cohorts were surveyed when they were in Grades 7, 9 or 10. The Grade 10 cohort (n=907) was surveyed in 2003 and the Grades 7 and 9 cohorts (n=955 and n=959 respectively) were surveyed in 2004. At time 1 participants were aged 12 to 17

(median age = 15.0; interquartile range 13.3 to 15.7). At time 2 (2010), 2,397 (85% retention rate) participants completed surveys again as young adults when they were 18 to 24 years of age (median age 21.0; interquartile range 19.3 to 22.6). Of these, 83% reported that they had a vehicle license. The current study includes a total of 1,956 young adults with a vehicle license who had complete data available for both time-points.

Ethics approval was first gained through the Ethics in Human Research Office at the Royal Children's Hospital in Victoria. At time 1 permission was then gained from the Department of Education and Training for government schools and the Catholic Education Office for some private schools and then by school principals. Parents provided written consent for their child to participate in the study and students provided assent to complete the survey. Students completed surveys during a 50- to 60-minute class period. Students absent on the day of testing completed the survey under the supervision of trained school personnel, or in a small percentage of cases (approximately 4%), over the telephone with study staff. At time 2, as young adults, participants provided consent for their own participation. They either completed a questionnaire online (77.5%), over the telephone with trained study staff (1.5%), or returned a postal survey (less than 5%).

2.2 Measures

2.2.1. Outcome variable

Young adult drink driving (DUI): At time 2, as young adults, participants were asked how often they had driven when probably affected by alcohol since gaining their license (adapted from the Australian Temperament Project, see (Prior *et al.*, 2000; Vassallo *et al.*, 2002). Response options ranged from *never* to *whenever I can* on a 7-point scale. Responses were recoded to generate a dichotomous variable with responses of *never* versus *at least once*. Less than 2% of respondents selected more frequently than the second response option of *occasionally*.

2.2.2. Exposure variable

Riding with a drinking driver during adolescence: At time 1, students were asked how many times during the past year they had ridden in a car or other motor vehicle driven by someone who had been drinking alcohol (adapted from the Youth Risk Behavior Survey (YRBS) High School Questionnaire, (Brener *et al.*, 2002; Centers for Disease Control and Prevention, 2010). Response options ranged from *Never*, *1 or 2 times*, to *40 or more times* on an eight-point scale. Responses were skewed with relatively few respondents selecting 3 or more times (13%). Responses were recoded to generate a dichotomous variable with responses reflecting *never during the past year* versus *at least once during the past year*.

2.2.3. Confounding factors measured during adolescence

Demographics: Information on students' gender, age and family socio-economic status was also collected. A single composite measure of family socioeconomic status (SES) was calculated from responses to questions on maternal and paternal education status and family income. This information was provided in a separate telephone interview with a parent/guardian of each student in the first year of the study (2002) as described in Evans-Whipp and colleagues (Evans-Whipp *et al.*, 2007). The SES variable was a continuous measure with a higher score indicating a higher level of SES.

The following factors were measured at time 1 during adolescence.

Poor family management: This scale came from the Communities that Care Youth Survey (Arthur *et al.*, 2002; Glaser *et al.*, 2005). It asked participants if their parents: would know if

they did not come home on time, know who they are with when they are not home, ask if they have done their homework, have clear rules in the family, have clear rules about alcohol and drug use, want them to call if they are going to be home late, would find out if they drank alcohol, carried a weapon or skipped school without their permission. Response options were on a four-point scale ranging from *definitely no* to *definitely yes* (Cronbach's Alpha = .83).

Family history of antisocial behaviour: This scale also came from the Communities that Care Youth Survey (Arthur *et al.*, 2002; Glaser *et al.*, 2005). It asked participants if any of their siblings had ever smoked cigarettes, drunk alcohol, used marijuana, taken a weapon to school, been suspended from school and if anyone in the family had ever had a severe alcohol or drug problem. Participants indicated *yes* or *no*. They were also asked how many adults they had known in the past year who had gotten drunk or high, used marijuana, sold drugs or done other things that could get them in trouble with the police. Response options ranged from *none* to *5 or more adults* on a five-point scale (Cronbach's Alpha = .76).

Binge alcohol use reflected students who indicated that they had drunk five or more alcoholic drinks in a row at least once during the last two weeks compared to those who had not. This measure was adapted from the Communities that Care Survey (Arthur *et al.*, 2002; Glaser *et al.*, 2005).

Sensation seeking: Participants were asked how many times they have: done crazy things even if they are a little dangerous, done something dangerous because someone dared you to do it and done what feels good no matter what. These items were adapted from the Seattle Social Development Project (SSDP) Sensation-Seeking/Disinhibition scale (Hawkins *et al.*, 2003). Response options ranged from *never* to *once a week or more* on a six-point scale (Cronbach's Alpha = .80).

Rebelliousness: Participants responded to the following three statements from the Personal Experiences Inventory (Winters and Henly, 1989). I do the opposite of what people tell me, just to get them mad, I ignore rules that get in my way and I like to see how much I can get away with. Response options were on a four-point scale ranging from *definitely no* to *definitely yes* (Cronbach's Alpha = .81).

Friends' use of drugs was measured by asking how many of the student's friends in the past year had: smoked cigarettes, tried alcohol, used marijuana or used other illegal drugs. Items were adapted from the SSDP Deviant Peer Involvement and Interaction scale (Hawkins *et al.*, 2003). Response options ranged from *none of my friends* to *4 of my friends* on a 5 point scale (Cronbach's Alpha = .77).

The mean of responses to items within each subscale formed measures of poor family management, family history of antisocial behaviour, sensation seeking, rebelliousness, and friends' use of drugs.

2.3 Statistical analysis

The following analyses are based on the sample of 1,956 respondents from 3 grade cohorts described above. To compare the frequencies and means of exposure and potential confounder variables in drink driving and non-drink driving groups, chi square tests for categorical variables and *t*-tests for continuous variables were performed. The unadjusted association between riding with a drinking driver during adolescence and young adult drink driving was then examined in a bivariate logistic regression. A series of logistic regressions was also performed to identify associations between each potential confounder with young adult DUI. A multivariate logistic regression predicting young adult drink driving was then

performed to test the predictive value of riding with a drinking driver during adolescence while accounting for the influence of potential confounding risk factors. To identify differential effects between subgroups within the sample, interactions between gender, age and cohort with adolescent exposure to riding with a drinking driver were also tested. However, none were significant and so the results are pooled across all participants.

3. Results

3.1 Descriptive statistics

As shown in Table 1, 34% of adolescents reported riding with a drinking driver at least once in the past year. As young adults, 18% of participants reported DUI in the past year. A statistically significantly higher proportion of young adults who had ridden with a drinking driver during adolescence reported DUI (compared to those who reported DUI but had not ridden with a drinking driver in adolescence (26% versus 14%, $\chi^2=41.33$, $p < .001$).

Descriptive statistics for potential confounding factors are also presented in Table 1. These showed differences on all factors between self-reported young adult drink drivers and non-drink drivers. A greater proportion of young adult drink drivers reported binge drinking during adolescence and were male compared to non-drink drivers. Drink drivers also reported higher rates of other individual and peer level risk factors during adolescence than non-drink drivers. They also showed higher rates of family history of antisocial behaviour, poor family management and higher family SES.

3.2 Bivariate associations between riding with a drinking driver during adolescence, confounding factors and young adult DUI

A series of logistic regressions were performed for riding with a drinking driver during adolescence and each potential confounder with young adult DUI (see Table 2). These analyses showed that adolescents exposed to others' drink driving were approximately twice as likely to drive under the influence of alcohol as a young adult. Males were three times more likely to DUI than females. All confounding factors measured also showed statistically significant associations, increasing the likelihood of young adult DUI. No statistically significant interactions between gender, age and cohort with riding with a drinking driver during adolescence were observed.

3.3 Multivariate associations between riding with a drinking driver during adolescence, confounding factors and DUI

Next, a multivariate logistic regression was performed to examine the main effects of riding with a drinking driver during adolescence on young adult DUI after the effects of key confounding factors were taken into account. Pairwise correlation analyses indicated weak to moderate correlations amongst the predictor variables (r ranged from $-.12$ to $.49$), indicating no signs of multicollinearity amongst these risk factors. As shown in Table 2, in the multivariate model, riding with a drinking driver during adolescence was associated with young adult DUI, increasing the likelihood by 62%. Other significant risk factors included being male, older in age, sensation seeking and higher family SES. Males were two and a half times more likely to DUI than females.

4. Discussion

This large longitudinal study examined the predictive relationship between self-reported DUI in young adulthood and riding with a drinking driver during adolescence in a large community sample in Victoria, Australia. As hypothesised, it found that exposure to drink driving during adolescence is a risk factor in young adulthood. Even when taking into

account other key influences on DUI there was a greater than 60 percent increase in the likelihood of DUI for those who reported riding with a drinking driver at least once in the past year in adolescence. Other variables found to be related to DUI were being male, increased age, sensation-seeking and higher SES.

The finding of the link between adolescent exposure to DUI and subsequent young adult DUI adds to the small body of research using longitudinal studies to understand the influence of personal, family and social characteristics on the development of risky driving behaviours (Gulliver and Begg, 2004; Reeder *et al.*, 1998; Vassallo *et al.*, 2007) and extends previous cross-sectional findings (Leadbeater *et al.*, 2008). Consistent with major social learning frameworks of behaviour (Akers, 1985; Bandura, 1977, 1986; Catalano and Hawkins, 1996) it is hypothesised this effect largely occurs through the influence of observing adult role models on the development of normative attitudes to DUI in adolescence.

It is apparent that many adolescents are being exposed to others' drink driving. Over one third of the adolescent sample in this study reported having ridden in a car with a drinking driver in the past year. This prevalence falls within the range reported in other studies: a Canadian study reported that 54% of Grade 10 and 12 students had ever ridden with an adult who had been drinking alcohol (Leadbeater *et al.*, 2008); another Canadian study reported that 23% of 9th to 12th graders had ridden in a car with a driver who had had too much to drink in the past year (Poulin *et al.*, 2007); a U.S. study found 45% of 15–20 year olds in California had ridden with a drinking driver in the past year (Chen *et al.*, 2008); and a national U.S. study found 28% of 9th to 12th graders had rode in a vehicle driven by someone who had been drinking alcohol in the past 30 days (Centers for Disease Control and Prevention, 2010). The risk to these adolescent passengers in terms of immediate safety has been well established but the negative influence of observing others' drink driving on their future driving behaviours is identified here as a further risk.

This study confirms previous findings that being male and sensation seeking may also be risk factors for DUI. Other potential risk factors for DUI, such as binge alcohol use, friends' alcohol use and family factors, whilst showing bivariate relationships with DUI, became statistically non-significant in the multivariate model. These findings suggest that adolescent exposure to others' drink driving is highly associated with the social contextual factors of family history of antisocial behaviour, poor family management and peer drug use, as well as individual behaviours of binge drinking and rebelliousness. The finding that higher family SES is associated with increased DUI is in contrast to some previous reports (Morrison *et al.*, 2002; Vaez and Laflamme, 2005). It is possible that the observed positive relationship with SES relates to increased access to alcohol and motor vehicles and less vigilance from police in the form of random breath testing (RBT) in high SES areas. The observed increased risk with age may be reflective of older participants having greater access to alcohol and motor vehicles, possibly from increased income. This finding may also reflect the nature of the measurement used, as older participants would have had a longer time period since gaining their license during which to DUI.

4.1 Strengths and limitations

As with all studies, the results of this study should be interpreted in light of its potential limitations. The self-report measure of DUI might have led to underestimations due to the potentially illegal nature of the behaviour. However, respondents were assured of the confidential nature of their responses and mostly completed the survey online which would have provided a greater sense of privacy and confidentiality. Also, it has been argued that self-report is the most appropriate method of measuring antisocial behaviours that may not be detected by others and therefore underestimated in objective evaluations (Jolliffe *et al.*,

2003). The rate of self-reported DUI (18%) in this study is consistent with previous reports based on Australian youth (Davey *et al.*, 2005; Vassallo *et al.*, 2010) and is considerably higher than rates ascertained using RBT (Wundersitz *et al.*, 2009).

The wording of the outcome variable (“driven when probably affected by alcohol”) rather than other measures such as the number of drinks consumed in a specified period before driving or BAC level was a deliberate attempt to measure self-perceived impaired driving rather than actual intoxication. This approach captures the development of attitudes towards ‘safe’ driving behaviour because participants indicated having driven despite feeling the effects of alcohol. However, it should be acknowledged that perceptions of intoxication are likely to vary between individuals. Further investigations using alternative measures of DUI would be beneficial. Nevertheless, perceptions of intoxication are important because even being slightly above the BAC limit greatly increases the risk of fatal accidents (Phillips and Brewer, 2011).

The exposure variable used in this study did not provide information on who the drinking driver was (parent, sibling, friend or other adult) or how much they had drunk. Further studies could investigate the differences in influences between different types of driver and whether this influence changes at different stages of adolescence. Measurement of the amount of alcohol consumed might help determination of the modeling and influence of views that “a little over the legal limit” is safe and acceptable driving behaviour. This is an important area of concern since casualty crash risk doubles when driving with a BAC just above the legal limit, and the risk of a fatal crash rises more sharply (Office of Road Safety, 2012).

Parent alcohol use is a known predictor of young adult DUI (Maldonado-Molina *et al.*, 2011) but was not measured directly in this study. The measure of family history of antisocial behaviour, which incorporates sibling and adult drug and alcohol use as well as other types of antisocial behaviour, was used as an alternative measure. This broader measure was considered more relevant given the exposure variable used in this study did not define the relationship of the drinking driver to the respondent.

Some studies have identified a relationship between rural (non-metropolitan) living status and DUI (Leadbeater *et al.*, 2008). The current study did not collect location information in young adulthood and so could not account for any possible effects of rural living status on DUI. Location information was available for the school of each adolescent participant at the time of the first survey. A possible link was explored but no association was found between time 1 location as measured on the Australian Remoteness Index for Areas (ARIA) (Australian Bureau of Statistics, 2003) and young adult DUI.

The exposure variable was measured 6 to 7 years before the outcome variable and so it is possible that other factors over the intervening time could have modified the effect on the outcome variable. Furthermore, the current outcome variable required participants to report on any DUI behavior ‘since getting a license’, rather than within the past year, as the exposure variable did. The findings of this study suggest there is a link, particularly evidenced by the association remaining when the influence of a range of important risk factors were controlled for. Nevertheless, future research should consider the use of other statistical techniques such as “growth modeling” to help better understand the association between exposure to riding with a drinking driver and cumulative DUI across an extended period of time.

Despite these limitations, this study has extended the findings of previous investigations of the personal and socio-contextual influences on DUI behaviour. Whilst previous studies have been limited by their sample design or the range of potential risk factors for which they

accounted, this study has utilised a large community sample to prospectively measure the influence of a range of potential factors in mid to late adolescence on later DUI. Retention rates across the 6 to 7 years of the study were high so negative effects due to sample attrition are minimal.

4.2 Implications

The finding that exposure in adolescence to others' drink driving increases the risk of self-reported DUI has important implications for prevention efforts. Most of the current measures aimed at preventing youth drink driving are based on structural, legal and policy changes. For example, higher minimum drinking ages (Wagenaar and Toomey, 2002), graduated licensing schemes (Russell *et al.*, 2011; Shope, 2007) and increased policing and enforcement of drink driving laws have all been shown to be effective. Some have argued, however, that further gains could be made by targeting key aspects of the individual and social context within which DUI behaviours develop (Chen *et al.*, 2008; Shults *et al.*, 2009; Vassallo *et al.*, 2007). The findings of this study suggest that adult role modeling of DUI to adolescents is one influence that could be focused upon.

Efforts to reduce adolescents riding with a drinking driver might target either the adult driver or the adolescent passenger. In the former case adult education via social media campaigns on the harmful effect of role modeling DUI behaviour would be warranted. One group that might be particularly susceptible to such messages is parents of younger aged children. It has been estimated that children in New South Wales, Australia are subject to 16 years of watching their parents drive before they embark on driver training themselves (Papakosmas and Noble, 2011). Awareness campaigns may also involve broader efforts to change community attitudes on the acceptability of drink driving and improve knowledge of the relationship between alcohol intake and impairment. Similar efforts have been made in relation to the harmful effect of child exposure to environmental tobacco smoke in cars. In this case legislation specifically targeting this behaviour has been successfully introduced and may be a method of conveying the seriousness of poor role modeling and exposing children to risk. The second form of prevention campaign involves downstream approaches that provide adolescents with tools and strategies for avoiding riding with a drinking driver. Many determinants are out of the control of younger adolescents but as they get older they are able to exert more influence over decisions. Harm reduction programs could educate on safe alcohol consumption rates for driving and refusal skills to help adolescents say no to alcohol impaired adults (which during later adolescence age may include parents, peers or siblings) offering to drive them. A number of school-based programs have been implemented with some evidence of effectiveness for reducing riding with drinking drivers (Elder *et al.*, 2005).

4.3 Conclusion

Rather than aiming to develop a broad based predictive model, the purpose of the current study was to specifically examine the relationship between riding with a drinking driver in the teens and subsequent DUI in young adulthood, net of a comprehensive range of potential confounding factors. The findings have important implications for adult drivers in the community who are acting as role models for adolescents riding in vehicles with them. The implications of DUI extend beyond the immediate risk to the driver of police enforcement, accident and injury to influence on young people exposed to their risky behaviour and the behaviour of future drivers. Incorporation of this message into safe drink drive media campaigns targeting parents and other adults may be an effective addition to existing enforcement policies which strive to reduce the unacceptable death and injury rate related to DUI.

Acknowledgments

We would like to express our gratitude to the participants of the International Youth Development Study. Further information is available from the IYDS website (<http://www.iyds.org>). The authors are grateful for the financial support of the National Institute on Drug Abuse (R01-DA012140) for the International Youth Development Study initial data collection and the National Institute on Alcoholism and Alcohol Abuse (R01AA017188-01) for analysis of the data. Continued data collection in Victoria, Australia has been supported by three Australian Research Council Discovery Projects (DPO663371, DPO877359, and DP1095744) and an Australian National Health and Medical Research Council grant (NHMRC; project number 594793). The IYDS was also supported by the Victorian Government's Operational Infrastructure Support program. The content is solely the responsibility of the authors and does not necessarily represent the official views of the sponsors.

References

- Akers, RL. *Deviant Behavior: A Social Learning Approach*. Wadsworth; Belmont, CA: 1985.
- Arthur MW, Hawkins JD, Pollard JA, Catalano RF, Baglioni AJ. Measuring risk and protective factors for substance use, delinquency, and other adolescent problem behaviors: The Communities That Care youth survey. *Eval Rev.* 2002; 26:575–601. [PubMed: 12465571]
- Australian Bureau of Statistics. ASGC remoteness classification: Purpose and use (census paper no 03/01). Canberra: 2003.
- Australian Bureau of Statistics. Australian social trends. ABS catalogue no. 4102.0. Australian Bureau of Statistics; 2008. Available at: <http://www.abs.gov.au/AUSSTATS/abs@.nsf/Lookup/4102.0Main+Features62008?OpenDocument>
- Australian Institute of Health and Welfare. AIHW bulletin series no. 60. Cat. no. AUS 102. AIHW; Canberra: 2008. Injury among young Australians.
- Australian Medical Association. Alcohol use and harms in Australia. Australian Medical Association; 2009. Available at: <http://ama.com.au/node/4762>
- Australian Transport Council. National road safety action plan 2009 and 2010. Australian Government Publishing Service; Canberra: 2008.
- Bandura, A. *Social Learning Theory*. Prentice-Hall; Englewood Cliffs, NJ: 1977.
- Bandura, A. *Social Foundations of Thought and Action: A Social Cognitive Theory*. Prentice-Hall; Englewood Cliffs, NJ: 1986.
- Beck KH, Kasperski SJ, Caldeira KM, Vincent KB, O'Grady KE, Arria AM. Trends in alcohol-related traffic risk behaviors among college students. *Alcohol Clin Exp Res.* 2010; 34 (8):1472–1478. [PubMed: 20528819]
- Bina M, Graziano F, Bonino S. Risky driving and lifestyle in adolescence. *Accid Anal Prev.* 2006; 38:472–481. [PubMed: 16375844]
- Bingham CR, Elliott MR, Shope JT. Social and behavioral characteristics of young adult drink/drivers adjusted for level of alcohol use. *Alcohol Clin Exp Res.* 2007; 31 (4):655–664. [PubMed: 17374045]
- Brener ND, Kann L, McManus T, Kinchen SA, Sundberg EC, Ross JG. Reliability of the 1999 Youth Risk Behavior Survey questionnaire. *J Adolesc Health.* 2002; 31:336–342. [PubMed: 12359379]
- Catalano, RF.; Hawkins, JD. The Social Development Model: A Theory of Antisocial Behaviour. In: Hawkins, JD., editor. *Delinquency and Crime: Current Theories*. Cambridge University Press; New York: 1996. p. 149-197.
- Cavallo, A.; Triggs, TJ. Directions for improving young driver safety within Victoria: A discussion paper. Monash University Accident Research Centre; Melbourne, Vic: 1996.
- Center for Behavioral Health Statistics and Quality. Data Spotlight: December 6, 2011. Substance Abuse and Mental Health Services Administration (SAMHSA); 2011. Adolescents living with a parent who drives under the influence are at increased risk for driving under the influence themselves.
- Centers for Disease Control and Prevention. Youth risk behavior surveillance —United States, 2009 surveillance summaries, [June]. *MMWR* 2010. 2010; 59(SS-5)
- Chen MJ, Grube JW, Nygaard P, Miller BA. Identifying social mechanisms for the prevention of adolescent drinking and driving. *Accid Anal Prev.* 2008; 40:576–585. [PubMed: 18329409]

- Davey JD, Davey T, Obst PL. Drug and drink driving by university students: An exploration of the influence of attitudes. *Traffic Inj Prev.* 2005; 6 (1):44–52. [PubMed: 15823874]
- Elder RW, Nichols JL, Shults RA, Sleet DA, Barrios LC, Compton R. Effectiveness of school-based programs for reducing drinking and driving and riding with drinking drivers. A systematic review. *Am J Prev Med.* 2005; 28 (Supplement):288–304. [PubMed: 15894162]
- Engström, I.; Gregersen, NP.; Hernetkoski, K.; Keskinen, E.; Nyberg, A. Young novice drivers, driver education and training: Literature review. Swedish National Road and Transport Research Institute; Linköping, Sweden: 2003.
- Evans-Whipp TJ, Bond L, Toumbourou JW, Catalano RF. School, parent, and student perspectives of school drug policies. *J Sch Health.* 2007; 77 (3):138–146. [PubMed: 17302856]
- Ferguson SA, Williams AF, Chapline JF, Reinfurt DW, De Leonardis DM. Relationship of parent driving records to the driving records of their children. *Accid Anal Prev.* 2001; 33:229–234. [PubMed: 11204894]
- Fernandes R, Job RFS, Hatfield J. A challenge to the assumed generalizability of prediction and countermeasure for risky driving: Different factors predict different risky driving behaviors. *J Safety Res.* 2007; 38 (1):59–70. [PubMed: 17275028]
- Finken LL, Jacobs JE, Laguna KD. Risky drinking and driving/riding decisions: The role of previous experience. *J Youth Adolescence.* 1998; 27 (4):493–511.
- Glaser RR, Van Horn ML, Arthur MW, Hawkins JD, Catalano RF. Measurement properties of the Communities That Care youth survey across demographic groups. *J Quant Criminol.* 2005; 21:73–102.
- Gulliver P, Begg D. Influences during adolescence on perceptions and behaviour related to alcohol use and unsafe driving as young adults. *Accid Anal Prev.* 2004; 36:773–781. [PubMed: 15203354]
- Hawkins, JD.; Smith, BL.; Hill, KG.; Kosterman, R.; Catalano, RF.; Abbott, RD. Understanding and Preventing Crime and Violence: Findings from the Seattle Social Development Project. In: Thornberry, TP.; Krohn, MD., editors. *Taking Stock of Delinquency: An Overview of Findings from Contemporary Longitudinal Studies.* Kluwer Academic/Plenum Publishers; New York: 2003. p. 255-312.
- Hingson RW, Zha W, Weitzman ER. Magnitude of and trends in alcohol-related mortality and morbidity among U.S. College students ages 18–24, 1998–2005. *J Stud Alcohol Drugs Suppl.* 2009; (16):12–20. [PubMed: 19538908]
- Hoyert, DL.; Kung, HC.; Smith, BL. National vital statistics reports. Vol. 53. National Center for Health Statistics; Hyattsville, MD: 2005. Deaths: Preliminary data for 2003.
- Jolliffe D, Farrington DP, Hawkins JD, Catalano RF, Hill KG, Kosterman R. Predictive, concurrent, prospective and retrospective validity of self-reported delinquency. *Crim Behav Ment Health.* 2003; 13 (3):179–197. [PubMed: 14654870]
- Jonah BA, Thiessen R, Au-Yeung E. Sensation seeking, risky driving and behavioral adaptation. *Accid Anal Prev.* 2001; 33 (5):679–684. [PubMed: 11491249]
- Labrie JW, Kenney SR, Mirza T, Lac A. Identifying factors that increase the likelihood of driving after drinking among college students. *Accid Anal Prev.* 2011; 43:1371–1377. [PubMed: 21545868]
- Labrie JW, Migliuri S, Kenney SR, Lac A. Family history of alcohol abuse associated with problematic drinking among college students. *Addict Behav.* 2010; 35:721–725. [PubMed: 20359831]
- Leadbeater BJ, Foran K, Grove-White A. How much can you drink before driving? The influence of riding with impaired adults and peers on the driving behaviors of urban and rural youth. *Addiction.* 2008; 103 (4):629–637. [PubMed: 18339107]
- Maldonado-Molina MM, Reingle JM, Delcher C, Branchini J. The role of parental alcohol consumption on driving under the influence of alcohol: Results from a longitudinal, nationally representative sample. *Accid Anal Prev.* 2011; 43 (6):2182–2187. [PubMed: 21819850]
- McCarthy DM, Pedersen SL. Reciprocal associations between drinking-and-driving behavior and cognitions in adolescents. *J Stud Alcohol Drugs.* 2009; 70 (4):536–542. [PubMed: 19515293]
- McMorris BJ, Hemphill SA, Toumbourou JW, Catalano RF, Patton GC. Prevalence of substance use and delinquent behavior in adolescents from Victoria, Australia and Washington State, United States. *Health Educ Behav.* 2007; 34 (4):634–50. [PubMed: 16740513]

- Morrison L, Begg DJ, Langley JD. Personal and situational influences on drink driving and sober driving among a cohort of young adults. *Inj Prev*. 2002; 8 (2):111–115. [PubMed: 12120828]
- National Highway Traffic Safety Administration. Traffic safety facts 2003 annual report: Early edition. U.S. Department of Transportation; Washington, DC: 2004.
- Office of Road Safety. Drink driving fact sheet. Government of Western Australia; 2012. Available at: <http://www.ors.wa.gov.au/Documents/DrinkDriving/drinkdriving-factsheet-may2012.aspx>
- Palamara, PG.; Legge, M.; Stevenson, MR. An investigation of the relationship between years of licensing, traffic offences, and crash involvement: A comparison of first year drivers with drivers licensed for ten years and five years. Injury Research Centre, University of Western Australia; Crawley, Australia: 2001.
- Papakosmas MF, Noble G. Parents and young drivers: The role of learning, behaviour modelling, communication and social marketing. *Journal of the Australasian College of Road Safety*. 2011; 22 (1):45–52.
- Pedersen SL, McCarthy DM. Person-environment transactions in youth drinking and driving. *Psychol Addict Behav*. 2008; 22 (3):340–348. [PubMed: 18778127]
- Phillips DP, Brewer KM. The relationship between serious injury and blood alcohol concentration (BAC) in fatal motor vehicle accidents: BAC = 0.01% is associated with significantly more dangerous accidents than BAC = 0.00%. *Addiction*. 2011; 106 (9):1614–1622. [PubMed: 21689195]
- Poulin C, Boudreau B, Asbridge M. Adolescent passengers of drunk drivers: A multi-level exploration into the inequities of risk and safety. *Addiction*. 2007; 102 (1):51–61. [PubMed: 17207123]
- Prior, M.; Sanson, A.; Smart, D.; Oberklaid, F. Pathways from infancy to adolescence: Australian Temperament Project 1983–2000. Australian Institute for Family Studies; Melbourne, Australia: 2000. 0642394784
- Reeder AI, Alsop JC, Begg DJ, Nada-Raja S, McLaren RL. A longitudinal investigation of psychological and social predictors of traffic convictions among young New Zealand drivers. *Transportation Research Part F Traffic Psychology and Behaviour*. 1998; 1 (1):25–46.
- Russell K, Vandermeer B, Hartling L. Graduated driver licensing for reducing motor vehicle crashes among young drivers. *Cochrane Database Syst Rev*. 2011; 10:CD003300. [PubMed: 21975738]
- Shope JT. Graduated driver licensing: Review of evaluation results since 2002. *J Safety Res*. 2007; 38:165–175. [PubMed: 17478187]
- Shults RA, Elder RW, Nichols JL, Sleet DA, Compton R, Chattopadhyay SK. Guide to community preventive services: Effectiveness of multicomponent programs with community mobilization for reducing alcohol-impaired driving. *Am J Prev Med*. 2009; 37:360–371. [PubMed: 19765509]
- Siskind V, Steinhardt D, Sheehan M, O'Connor T, Hanks H. Risk factors for fatal crashes in rural Australia. *Accid Anal Prev*. 2011; 43:1082–1088. [PubMed: 21376905]
- Tin Tin S, Ameratunga S, Robinson E, Crengle S, Schaaf D, Watson P. Drink driving and the patterns and context of drinking among New Zealand adolescents. *Acta Paediatr*. 2008; 97 (10):1433–1437. [PubMed: 18624990]
- Tomas Dols S, González FJÁ, Aleixandre NL, Vidal-Infer A, Rodrigo MJT, Valderrama-Zurián JC. Predictors of driving after alcohol and drug use among adolescents in Valencia (Spain). *Accid Anal Prev*. 2010; 42 (6):2024–2029. [PubMed: 20728658]
- Transport Accident Commission of Victoria. Young driver statistics (2002). 2003. Available at: <http://www.tacsafety.com.au/jsp/content/NavigationController.do?areaID=13&tierID=1&navID=AF3476DD&navLink=null&pageID=209>
- Vaez M, Laflamme L. Impaired driving and motor vehicle crashes among Swedish youth: An investigation into drivers' sociodemographic characteristics. *Accid Anal Prev*. 2005; 37:605–611. [PubMed: 15949450]
- Vassallo, S.; Smart, D.; Cockfield, S.; Gunatillake, T.; Harris, A.; Harrison, W. Research Report No 17. Australian Institute of Family Studies; Melbourne: 2010. In the driver's seat ii: Beyond the early driving years.
- Vassallo, S.; Smart, D.; Sanson, A.; Dussuyer, I.; McKendry, B.; Toumbourou, JW. Patterns and precursors of adolescent antisocial behaviour: The first report December 2002. Crime Prevention Victoria; Melbourne: 2002.

- Vassallo S, Smart D, Sanson A, Harrison W, Harris A, Cockfield S, McIntyre A. Risky driving among young Australian drivers: Trends, precursors and correlates. *Accid Anal Prev.* 2007; 39 (3):444–458. [PubMed: 17359925]
- Wagenaar AC, Toomey TL. Effects of minimum drinking age laws: Review and analyses of the literature from 1960–2000. *J Stud Alcohol Suppl.* 2002; 14:2006–225.
- Ward, B.; Snow, P.; James, E.; Griffith, J. The influence of parents and siblings on children's and adolescents' alcohol use: A critical review of the literature. *Drinkwise Australia*; 2010. Available at: <http://www.drinkwise.org.au/free-resources/research/the-influence-of-parents-and-siblings-on-childrens-and-adolescents-attitudes-and-behaviours-towards-alcohol-a-critical-review-of-the-literature/>
- Winters, KC.; Henly, GA. *Personal experiences inventory test & test manual.* Western Psychological Services; Los Angeles, CA: 1989.
- Wundersitz, LN.; Hiranandani, K.; Baldock, MRJ. Annual performance indicators of enforced driver behaviours in South Australia, 2007. Centre for Automotive Safety Research, University of Adelaide; 2009. Available at: <http://casr.adelaide.edu.au/publications/researchreports/casr058.pdf>
- Zakletskaia LI, Mundt MP, Balousek SL, Wilson EL, Fleming MF. Alcohol-impaired driving behavior and sensation-seeking disposition in a college population receiving routine care at campus health service centers. *Accid Anal Prev.* 2009; 41:380–386. [PubMed: 19393782]

Highlights

- We investigated role modelling of drink driving to adolescents
- Adolescent exposure to drink driving increased risk of drink driving in adulthood
- This effect remained when accounting for other known risk factors for drink driving
- Drink drive prevention efforts should target adult role models in the community

Table 1

Means and prevalence of key variables for young people with a driver's license

Adolescent Risk Factor	Whole Sample (N= 1,956)	Non-Drink Drivers (N=1,599)	Drink Drivers (N=357)
Adolescent exposure to DUI	34.15%	14.21%	26.05% ***
Gender - Male	45.81%	41.03%	67.23% ***
Age	14.76 (1.23)	14.67 (1.24)	15.17 (1.09) ***
Binge Drinking	27.90%	25.31%	39.55% ***
Sensation Seeking	2.55 (1.32)	2.43 (1.27)	3.09 (1.41) ***
Rebelliousness	1.99 (.67)	1.96 (.67)	2.12 (.64) ***
Family History of ASB	1.93 (.81)	1.88 (.80)	2.13 (.84) ***
Poor Family Management	1.80 (.53)	1.77 (.53)	1.94 (.50) ***
Family SES	1.96 (.50)	1.95 (.50)	2.02 (.52) *
Friends' Drug Use	2.08 (1.01)	2.01 (.99)	2.39 (1.03) ***

Note.

* $p < .05$,

** $p < .01$,

*** $p < .001$.

Family History of ASB = Family history of antisocial behaviour; Standard deviations for continuous variables are in parenthesis.

Table 2

Logistic regression models predicting young adult DUI

	Bivariate Models		Multivariate Model	
	OR	95% CI	OR	95% CI
Adolescent exposure to DUI	2.13 ^{***}	1.68–2.69	1.62 ^{**}	1.23–2.13
Gender - Male	2.95 ^{***}	2.31–3.76	2.64 ^{***}	2.02–3.43
Age	1.33 ^{***}	1.24–1.44	1.32 ^{***}	1.16–1.50
Sensation seeking	1.42 ^{***}	1.31–1.54	1.24 ^{***}	1.10–1.39
Rebelliousness	1.43 ^{***}	1.20–1.69	.90	.72–1.13
Binge drinking	1.93 ^{***}	1.52–2.46	1.10	.79–1.53
Family History of ASB	1.42 ^{***}	1.24–1.62	1.11	.92–1.35
Poor Family Management	1.79 ^{***}	1.45–2.22	1.00	.75–1.33
Family SES	1.32 [*]	1.04–1.66	1.41 ^{**}	1.09–1.81
Friends' drug use	1.42 ^{***}	1.27–1.59	.99	.83–1.17

*Note.**
p < .05,**
p < .01,***
p < .001.

Family History of ASB = Family history of antisocial behaviour. Multivariate models control for all variables in the table.