



Published in final edited form as:

Addiction. 2013 March ; 108(3): 584–591. doi:10.1111/add.12014.

Egocentric Social Network Analysis of Pathological Gambling

Matthew K. Meisel¹, Allan D. Clifton², James MacKillop^{1,3}, Joshua D. Miller¹, W. Keith Campbell¹, and Adam S. Goodie¹

¹Department of Psychology, University of Georgia, Athens, GA 30602-3013

²Department of Psychology, Vassar College, 124 Raymond Ave., Box 713, Poughkeepsie, NY 12604-0713

³Department of Behavioral and Social Sciences, Brown University, Providence, RI 02912

Abstract

Aims—To apply social network analysis (SNA) to investigate whether frequency and severity of gambling problems were associated with different network characteristics among friends, family, and co-workers. is an innovative way to look at relationships among individuals; the current study was the first to our knowledge to apply SNA to gambling behaviors.

Design—Egocentric social network analysis was used to formally characterize the relationships between social network characteristics and gambling pathology.

Setting—Laboratory-based questionnaire and interview administration.

Participants—Forty frequent gamblers (22 non-pathological gamblers, 18 pathological gamblers) were recruited from the community.

Findings—The SNA revealed significant social network compositional differences between the two groups: pathological gamblers (PGs) had more gamblers, smokers, and drinkers in their social networks than did nonpathological gamblers (NPGs). PGs had more individuals in their network with whom they personally gambled, smoked, and drank with than those with who were NPG. Network ties were closer to individuals in their networks who gambled, smoked, and drank more frequently. Associations between gambling severity and structural network characteristics were not significant.

Conclusions—Pathological gambling is associated with compositional but not structural differences in social networks. Pathological gamblers differ from non-pathological gamblers in the number of gamblers, smokers, and drinkers in their social networks. Homophily within the networks also indicates that gamblers tend to be closer with other gamblers. This homophily may serve to reinforce addictive behaviors, and may suggest avenues for future study or intervention.

Keywords

Pathological gambling; social network analysis; egocentric; alcohol; tobacco

Corresponding author: Adam S. Goodie, Department of Psychology, University of Georgia, Athens, GA 30602-3013, goodie@uga.edu, Tel 706-542-6624, Fax 706-542-3275.

Declaration of interest: Dr. MacKillop receives funding from the National Institutes of Health and the Institute for Research on Gambling Disorders. In the past, he has received research grants from the Russell Sage Foundation, the Robert Wood Johnson Foundation; the Pfizer, Inc., Global Research Advances in Nicotine Dependence program, and the Alcoholic Beverage Medical Research Foundation. Dr. Miller and Dr. Campbell receive funding from the Institute for Research on Gambling Disorders. Dr. Goodie receives funding from the Institute for Research on Gambling Disorders and the Army Research Office. In the past, he has received research grants from the National Institutes of Health, Air Force Office of Sponsored Research, and the Ontario Problem Gambling Research Centre. None of these sources constitutes a conflict of interest with this study.

Social factors contribute to the initiation and maintenance of gambling behavior. For example, the most frequent reason for gambling among older adults reported was to socialize with friends (1). In a college-aged sample, social factors were the third most cited motivation to gamble (2). Based on Becker's early studies on the initiation of drug use (3, 4), Reith and Dobbie (5) argue that the social environment interacts with the individual, such that an individual learns how and where to gamble from his or her social network. Recreational gamblers and pathological gamblers (PGs) who were introduced to gambling in early life were at the greatest risk of developing gambling problems (5). Further, as adolescents age and their gambling involvement increases, they spent more time with their gambling friends, resulting in fewer close relationships with non-gambling friends (6), which may result in a pernicious cycle of a social network that reinforces gambling, which in turn results in spending more time with gambling friends. Social factors, as well as perceptions of social norms, are also implicated during gambling. For example, participants who believe that others are gambling and winning, play for longer periods resulting in greater losses (7). In the presence of onlookers, people place smaller bets (8), suggesting that social factors can have a considerable impact on gambling play. When students perceive that important others approve of gambling, they gamble more frequently (9). Friends' and families' approval of gambling are also strong predictors of past year gambling frequency and severity (10).

Social network analysis

The current study utilized an established method that has only recently been applied to gambling and other addictive behaviors. Social network analysis (SNA) is an innovative technique for understanding group prevalence and structure. The current study utilized egocentric network analysis, in which the participant (referred to as "ego") lists his or her closest friends, family members, and co-workers (referred to as "alters"), and assesses the relationships among the alters. (In a sociocentric network analysis, by contrast, information is gathered from each person, about each person, in a relatively closed network.)

A frequent focus of SNA studies is homophily, or the tendency of individuals who are similar in their beliefs, attitudes, and behaviors to be more frequently and more closely linked in social networks than those who are dissimilar (11). In his classic housing study, Festinger (12) found evidence of homophily based on propinquity, the tendency of people who live close together to be more connected. Social network analysis is also used to examine the structural characteristics of social networks. One structural characteristic that may affect addictive behavior is network density, which reflects how connected the members of a network are to each other. Dense networks make it easier for egos to observe and to replicate the behaviors of alters in their network (13).

SNA and addictive behaviors

Within the DSM-IV, pathological gambling (PG) is categorized as an impulse control disorder defined by symptoms including loss of control of gambling, preoccupation with gambling, and persistence despite negative consequences (14). The DSM-5 will most likely categorize PG under Substance Use and Addictive Disorders, reduce the diagnostic threshold from 5 to 4 symptoms and eliminate the criterion of illegal activities (15). SNA has been successfully utilized to study substance use and abuse. For example, the proportion of drinkers and heavy drinkers in an individual's network is positively related to participants' drinking (16). In contrast, the proportion of family members in a drinker's network is negatively related to the participants' drinking.

Homophily has been observed in the addiction domain. For example, drinkers prefer friends with the same drinking and smoking behavior (17, 18). We therefore posit that individuals who gamble, smoke and drink, will be more frequently and more closely connected to others who gamble, smoke and drink, respectively. Peer group substance use has also been examined in several studies utilizing sociocentric SNA applied to samples of middle and high school students. Fang and colleagues (19) found a negative relationship between network density and substance abuse among isolates, (those who are unconnected to peers; 20), whereas Henry and Kobus (21) found greater substance abuse among those who link otherwise unconnected groups (or “liaisons”). Liaisons have been found to smoke more than others, but are less affected by the prevalence of smoking in their networks (22). Surprisingly, there is no effect of network position on alcohol use, but alcohol use is related to the proportion of network peers who use alcohol. As the prevalence of alcohol and marijuana use increases in peer networks, so does the frequency of an individual engaging in that behavior (22).

The present study

The primary aim of the present study was to apply SNA to PG for the first time, investigating the role of social networks in PG, in a comparison of recreational gamblers and problem gamblers. We hypothesized that, compared to nonpathological gamblers (NPGs), PGs would have social networks that were denser with gamblers and also differed structurally. However, in the absence of previous studies, no a priori hypotheses were made for specific structural indices. A second aim of the study was to investigate substance use comorbidity in PGs’ and NPGs’ social networks. Based on the comorbidity literature (23, 24), we hypothesized that PGs’ network members would gamble, drink, and smoke more often than NPGs’ network members. We also hypothesized that PGs would engage in all of these behaviors more often than NPGs with their network members. As friends have been found to be a primary reason to gamble for older adults, we also hypothesize that they will have significant impact on gambling, smoking, and drinking behavior.

Methods

Participants

Forty frequent-gambling adult participants (75% male) were recruited from the Athens, GA community. All participants were recruited through advertisements in newspapers and buses as well as word-of-mouth. Exclusion criteria were gambling less than weekly, currently living with another participant, inability to use a computer, self-reported symptoms of psychosis, or age greater than 65 years. Participants were an average age of 35.25 years (SD=11.09). 67.5% earned less than \$15,000 pre-tax in the past year, and 17.5% earned between \$15,000–\$30,000. Most participants were African American (72.5%) followed by Caucasian (25%) and mixed race (2.5%). Participants were compensated \$20. Based on the DSM-IV Structured Clinical Interview for Pathological Gambling (SCI-PG; 25), 18 participants met criteria for PG and 22 participants did not.

Measures

We used an egocentric network analysis approach, in which the participant (“ego”) listed his or her 30 closest social associates including friends, family members, present/past romantic partners, and co-workers (“alters”). The amount of structural information gained about a network increases as the number of alters increases, but begins to plateau around 25 alters, with 35 alters providing virtually identical information as 45 alters (26). Participants did not report difficulty listing the 30 alters, although tests of order effects revealed some significant

differences in gambling or substance use between later- and earlier-named alters (reported below).

Participants indicated the sex and race of each alter, how long he or she knew each alter, how often he or she spent time with each alter, how close they were, whether they ever lived together, and whether they ever were in a romantic relationship with one another. Participants also indicated how frequently each alter gambled, smoked, and drank, and how often the participant gambled, smoked, and drank with each alter. Each of these behaviors was assessed on a 6-point Likert frequency scale that included the following levels: 1) Not in the past year, 2) Less than once a month, 3) Once a month, 4) Once a week, 5) Multiple times a week, 6) Daily.

Participants additionally answered questions about the relationships among the alters. Each alter pairing was rated on a scale ranging from *very close* (5) to *they have never met* (1). Assessment was conducted using EgoNet, a program designed for the collection of egocentric social network data (27).

Social networks were structurally characterized using the validated SNA indices of network density and betweenness centrality. Network density is the proportion of the number of actual connections relative to the number of possible connections in a network. Dense networks have many strong connections between members whereas a less dense network has fewer and weaker connections. We also calculated the betweenness centrality of each alter, which assesses how well-connected and integral each individual is to his/her network. Betweenness centrality is the degree to which the shortest paths between any pair of people in the network pass through a particular alter (28).

Data analysis

A Jonckheere-Tepstra test (29) was used to analyze differences in gambling, smoking and drinking frequency between the social networks of PGs and NPGs, as well as the frequency of joint engagement in these behaviors by ego and alter together. These use median values, with lower numbers representing higher frequencies. We dichotomized alters' gambling, drinking, and smoking frequency as less than once a month or at least once a month (30). We then used Mann-Whitney U tests to compare these two categories between the networks of PGs and NPGs.

We also used a Mann-Whitney U test to examine differences in network density. For other tests, we used multilevel models with a "one-with-many" design (31), which allowed for multiple ratings of alters by a single participant. We used these to account for nonindependence of alters within a participant's network and interactions between the individual and the social network. We also conducted a multilevel model with the participant's diagnostic status (PG or NPG) and the participant's gambling as fixed effects predicting homophily, and with each alter's gambling, smoking, or drinking frequency as fixed effects predicting closeness or centrality. Data analysis was conducted on SPSS 19.0, and UCINET (32) was used to generate the structural aspects of the participants' social networks. All non-dichotomized independent variables were grand mean centered.

Results

Compositional social network characteristics

The number of networks members who were friends, family members, co-workers, and present/past romantic partner was associated with PG status, with PGs having significantly more family members and fewer coworkers in their self-reported networks than is expected

by random proportional assignment ($\chi^2=21.01$, $df=4$, $p<.001$). See Table 1 for full descriptive statistics.

Overall activity of alters

With each of 40 participants naming 30 members of his or her social network, we accumulated data on $40 \times 30=1200$ alters (660 alters named in NPGs' network and 540 alters named in PGs' network). The gambling frequency of PGs' network members (Mdn=2; *less than once a month*) were significantly higher than NPGs' network members (Mdn=1; *not in the past year*, $Z=4.98$, $p<.001$). For example, 19% of people listed in the PGs' networks gambled daily, whereas 11% of people listed in the NPGs' networks gambled daily. The PGs' networks included more alters who gambled at least once a month ($U=202620$, $p<.001$). We also found significant differences in the networks' distribution of smoking ($Z=2.80$, $p<.01$) and drinking ($Z=3.42$, $p<.001$) behavior. For both comparisons, the PGs' median scores were 2 (*less than once a month*), whereas the NPGs' median scores were 1 (*not in the past year*). As revealed in Table 2, the networks of PGs had frequency distributions that were more weighted to frequent engagement in all three behaviors.

Figure 1 presents examples of PG and NPG networks, selected to be maximally illustrative of the effects in question. A line between two nodes represents a connection between alters, and darker and larger nodes represent more frequent gambling, ranging from black (daily) to white (not in the past year). Panels A and B reflect gambling in the alters of an NPG and PG participant, respectively; Panels C and D depict smoking in the alters of the same NPG and PG participants; and Panels E and F depict drinking in the alters of the same NPG and PG participants. In each case, the networks reveal the greater occurrence of gamblers, smokers, and drinkers for the PG participant; in contrast, the NPG participant exhibits a network in which the addictive behaviors are restricted to more distinct subgroups of associates.

In addition to effects in their overall behavior, consistent with our hypotheses, we also found significant differences in alters' frequency of gambling ($Z=3.84$, $p<.001$), smoking ($Z=4.42$, $p<.001$), and drinking ($Z=3.74$, $p<.001$) with ego, although the medians for both groups on all three behaviors were 1 (*not in the past year*), reflecting the fact that these are generally low-frequency behaviors. PGs gambled with 37% of their network members at least once a month compared with 27% in the NPGs' networks ($U=196350$, $p<.001$). PGs smoked with 42% of their network members at least once a month compared with 29% of NPGs' ($U=202470$, $p<.001$). PG's drank with 41% of the individuals in their networks at least once a month, compared with 33% of the members of NPGs' ($U=192540$, $p<.01$).

In general, the effects appeared stronger among friends as opposed to all network members, as floor effects on frequency were attenuated. Alters described as friends gambled ($U=64856$, $p<.001$), smoked ($U=62772$, $p=.001$), and drank ($U=63254$, $p=.001$) more in PGs' networks than in NPGs' networks (Mdns=3 and 1 for gambling, 4 and 1 for smoking, and 5 and 3 for drinking, respectively). The friends in PGs' networks also gambled (Mdns both=1; $U=64845$, $p<.001$), smoked (Mdns=3 and 1; $U=67133$, $p<.001$) and drank (Mdns=3 and 1; $U=65027$, $p<.001$) significantly more with the participant than the friends in NPGs' networks.

Structural social network characteristics

We next examined the structural characteristics of both groups' networks. There were no significant differences in density between the networks of PGs' (1.95, $SD=0.77$) and NPGs' (2.10, $SD=1.00$; $t(38)=.54$). Likewise, no significant relationships were observed between centrality and alter gambling, drinking or smoking (all t 's <1.16 , all p 's $>.24$). Similarly, when using dichotomized behaviors (less than once a month vs. once a month or more), no

significant relationships between centrality and alter gambling ($t(949) = -.560, p = .576$), drinking ($t(881) = .031, p = .975$) or smoking ($t(1065) = -1.085, p = .278$) were found. These findings suggest that the organizational structure of the participants' social networks do not significantly differ by PG status.

We also tested the relationship between subjective closeness and gambling severity. Participants felt subjectively closer to alters who gambled ($F(1, 1179) = 31.27, p < .001$), smoked ($F(1, 1195) = 6.32, p = .01$), and drank more frequently ($F(1, 1192) = 6.43, p < .05$). There were no interactions between diagnostic severity and alters' gambling ($F(1, 1179) = 0.97, p = .32$) or smoking ($F(1, 1195) = 1.02, p = .32$) in predicting closeness, although NPGs felt subjectively closer to the drinkers in their networks to a greater degree than PGs did ($F(1, 1192) = 6.49, p < .05$). Furthermore, there were no differences in friend subjective closeness ($U = 54582, p = .76$) and entire network subjective closeness ($U = 170782, p = .195$) with the ego between the networks of PGs and NPGs. The relationship between closeness and frequency of gambling, smoking and drinking is further borne out by associations between order of identification and all three addictive behaviors. The first 10 listed alters gambled ($U = 71695, p < .01$), smoked ($U = 73373, p < .05$), and drank ($U = 74185, p = .05$) more frequently with the ego than did the second 10 alters listed, who in turn did not differ from the third 10 alters listed.

Homophily was observed in the gambling behavior of alters, with a positive correlation between an alters gambling score and the average gambling scores of the other alters to whom that alter was connected ($r(977) = .61, p < .001$). There was no evidence of differential homophily in the networks of PGs ($r(438) = .60, p < .001$) compared with the NPGs group ($r(539) = .61, p < .001$; $F = 1.77, p = .18$).

Discussion

To our knowledge, the current study constitutes the first formal social network analysis of pathological and recreational gamblers. This is a particularly promising methodology for gambling studies, both insofar as SNA has made significant strides in other addictive behaviors (19, 21), and as social factors are known to contribute substantially to PG (1, 2).

Consistent with the literature on comorbidity (23, 24), PGs had not only more gamblers, but also more smokers and drinkers in their networks, who gambled, smoked, and drank more frequently than those in a NPG's network. We also found that PGs gambled, smoked and drank more frequently with members of their networks than did NPGs. At a correlational level, as an individual's gambling severity increases, so may the importance and frequency of gambling, smoking, and drinking in the network. The members of a PG's network may reinforce the addictive behavior.

Consistent with our hypotheses, PGs were found to gamble, smoke and drink alcohol significantly more often with their friends than NPGs did. We also found that in the networks of PGs, their friends gambled, smoked and drank significantly more than the friends in the NPG's network. There are two prominent theories on why social affiliates engage in similar behaviors: socialization and selection. In the former, friends' attitudes and behaviors affect the individual (conformity), while in the latter the person seeks out peers with similar beliefs and behaviors (33, 34). Research suggests that socialization is associated with closed, tight networks (e.g., military class) while selection is associated with open, looser networks (e.g., high schools; 35). As the current study is cross-sectional, it cannot differentially support either of the two theories, but it clearly represents a methodology that, applied across time, could clarify whether individuals with PG seek out similarly affected people or whether social groups directly confer risk for developing PG.

Participants felt significantly closer to alters who gambled, smoked and drank more frequently. Surprisingly, we found that this effect was virtually identical for the PG and NPG groups in their subjective feelings of closeness to the gamblers and smokers in their networks. Furthermore, when comparing the networks of PGs and NPGs, there were no differences in subjective closeness for friends and the entire network. These results suggest that PG status does not directly affect closeness and that closeness may be defined by several other factors besides mutual interests.

We found evidence of homophily in the networks of both PGs and NPGs. Alters who gambled were more connected to others who gamble, and those who did not gamble were more connected to others who did not gamble. Consistent with these results pertaining to gambling, homophily is also found in drug using networks (16, 17). This suggests that the networks of gamblers are similar to those of substance abusers. The absence of significant differential homophily and network density may have been due to issues of range restriction arising from the entire sample being comprised of gamblers. In a study examining heroin injectors and non-injectors, the authors found that although injectors had more friends and a larger network size, there was not a significant difference in network density between the two groups (36). Similarly, the main differences between these networks in our data were compositional, not structural. That is, taken together, the most salient social network factors observed for PG participants were significantly more gamblers in the network, more frequent gambling among those gamblers, and significantly greater joint gambling with network members. The lack of difference in density, which reflects closeness among alters and not between ego and any alters, is independent of the low social support that is associated with greater gambling severity (37).

Strengths of this study include the systematic application of an SNA approach to PG and a well characterized sample with considerable diversity. However, limitations include that the participants reported the behavior of others in their network, possibly resulting in a false consensus effect, an inherent limitation of egocentric SNA wherein participants project their own behavior onto others (38). This possibility is diminished by the fact that homophily was observed across networks and not just in alters' relationships with ego. Future research would benefit from utilizing a sociocentric network design and a longitudinal design that addresses the causal role of social influence and selection on addictive behaviors. Another limitation of this study was its relatively modest sample size, which may reduce the generalizability of the findings. We also cannot eliminate the possibility of overlapping networks, as alters were kept anonymous. Future research will be needed to establish whether the correlational effects reported here are attributable to gambling problems per se or to gambling frequency. Finally, the current study included higher proportions of African American and low-income individuals than is reflective of the broader US population, likely due to these demographic characteristics being more prevalent in the recruitment catchment area.

These caveats notwithstanding, the current study advances understanding of the role of the social network in addictive behavior by providing the first formal SNA of pathological gambling. Distinct and theoretically relevant differences were observed in the composition of PGs' and NPGs' networks, in the absence of structural differences. Pathological gamblers had more gamblers, smokers and drinkers in their networks in general and more individuals with whom they personally gambled, smoked, and drank alcohol. These compositional differences may provide important insights into the causes and maintaining factors in PG and, ultimately, may also be leveraged to enhance treatment.

Acknowledgments


This research was partially supported by grants from the Institute for Research on Gambling Disorders (AC, JM, JDM, WKC and ASG) and the National Institutes of Health (K23 AA016936 – JM; P30 DA027827 – ASG, JM).

References


1. Hope J, Havir L. You bet they're having fun! Older Americans and casino gambling. *Journal of Aging Studies*. 2002; 16:177–197.
2. Neighbors C, Lostutter TW, Crouse JM, Larimer ME. Exploring college student gambling motivation. *Journal of Gambling Studies*. 2002; 18(4):361–370. [PubMed: 12514915]
3. Becker H. Becoming a marijuana user. *American Journal of Sociology*. 1953; 59:235–242.
4. Becker, H. *Outsiders: Studies in the sociology of deviance*. New York: Free Press; 1963.
5. Reith G, Dobbie F. Beginning gambling: The role of social networks and environment. *Addiction Research and Theory*. 2011; 19:483–493.
6. Gupta R, Derevensky JL. Adolescents with gambling problems: from research to treatment. *Journal of Gambling Studies*. 2000; 16:315–342. [PubMed: 14634318]
7. Rockloff MJ, Dyer V. An experiment on the social facilitation of gambling behavior. *Journal of Gambling Studies*. 2007; 23:1–12. [PubMed: 17191146]
8. Rockloff M, Greer N, Fay C. The social contagion of gambling: How venue size contributes to player losses'. *Journal of Gambling Studies*. 2010; 27:487–497. [PubMed: 20872055]
9. Larimer ME, Neighbors C. Normative misperception and the impact of descriptive and injunctive norms on college student gambling. *Psychology of Addictive Behaviors*. 2003; 3:235–243. [PubMed: 14498818]
10. Welte JW, Wieczorek WF, Barnes GM, Tidwell MO. Multiple risk factors for frequent and problem gambling: Individual, social, and ecological. *Journal of Applied Social Psychology*. 2006; 36:1548–1568.
11. McPherson M, Smith-Lovin L, Cook JM. Birds of a feather: Homophily in social networks. *Annual Review of Sociology*. 2001; 27:415–444.
12. Festinger, L.; Schachter, S.; Back, KW. *Social pressures in informal groups: A study of human factors in housing*. Stanford, California: Stanford University Press; 1950.
13. Latané B. Dynamic Social Impact: The Creation of Culture by Communication. *Journal of Communication*. 1996; 46:13–25.
14. American Psychiatric Association. *Diagnostic and Statistical Manual of Mental Disorders*. 4. American Psychiatric Press; Washington, DC: 1994.
15. American Psychiatric Association. DSM-5 development: Proposed revision of pathological gambling. 2012. Retrieved June 16, 2012, from <http://www.dsm5.org/proposedrevision/Pages/SubstanceUseandAddictiveDisorders.aspx>
16. Fondacaro MR, Heller K. Social support factors and drinking among college student males. *Journal of Youth and Adolescence*. 1983; 12(4):285–299.
17. Christakis NA, Fowler JH. The collective dynamics of smoking in a large social network. *The New England Journal of Medicine*. 2008; 358:2249–2258. [PubMed: 18499567]
18. Rosenquist JN, Murabito J, Fowler JH, Christakis NA. The spread of alcohol consumption in a large social network. *Annals of Internal Medicine*. 2010; 152(7):426–433.
19. Fang X, Li X, Stanton B, Dong Q. Social network positions and smoking experimentation among Chinese adolescents. *American Journal of Health Behavior*. 2003; 27(3):257–267. [PubMed: 12751622]
20. Hansell S, Tackaberry SN, Slavin RE. Cooperation, competition, and the structure of student peer groups. *Representative Research in Social Psychology*. 1981; 12(1–2):46–61.
21. Henry DB, Kobus K. Early adolescent social networks and substance use. *The Journal of Early Adolescence*. 2007; 27:346–362.
22. Kobus K, Henry DB. Interplay of network position and peer substance use in early adolescent cigarette, alcohol, and marijuana use. *The Journal of Early Adolescence*. 2010; 30:225–245.

23. Cunningham-Williams RM, Cottler LB, Compton WM III, Spitznagel EL. Taking chances: Problem gamblers and mental health disorders-Results from the St. Louis epidemiologic catchment area study. *American Journal of Public Health*. 1998; 88(7):1093–1096. [PubMed: 9663161]
24. Kessler RC, Hwang I, LaBrie R, Petukhova M, Sampson NA, Winters KC, Shaffer HJ. DSM-IV pathological gambling in the National Comorbidity Survey Replication. *Psychological Medicine*. 2008; 38:1351–1360. [PubMed: 18257941]
25. Grant JE, Steinberg MA, Kim SW, Rounsaville BJ, Potenza MN. Preliminary validity and reliability testing of a structured clinical interview for pathological gambling. *Psychiatry Research*. 2004; 128:79–88. [PubMed: 15450917]
26. McCarty C, Killworth PD, Rennell J. Impact of methods for reducing respondent burden on personal network structural measures. *Social networks*. 2007; 29:300–315.
27. McCarty, C. Egonet (Version 2.0 Beta 6). Gainesville, FL: Endless Loop Software; 2004.
28. Freeman LC. Centrality in social networks: conceptual clarification. *Social Networks*. 1979; 1:215–239.
29. Jonckheere AR. A distribution-free k-sample test against ordered alternatives. *Biometrika*. 1954; 41:133–145.
30. Weinstock J, Ledgerwood DM, Petry NM. Association between posttreatment gambling behavior and harm in pathological gamblers. *Psychology of Addictive Behaviors*. 2007; 21:185–193. [PubMed: 17563138]
31. Kenny, DA.; Kashy, DA.; Cook, WL. *Dyadic data analysis*. New York: The Guilford Press; 2006.
32. Borgatti, SP.; Everett, MG.; Freeman, LC. *Ucinet 6 for windows: Software for social network analysis*. Harvard, MA: Analytic Technologies; 2002.
33. Cohen JM. Sources of peer group homogeneity. *Sociology of Education*. 1977; 50:227–241.
34. Kandel DB. Homophily, selection and socialization in adolescent friendships. *American Journal of Sociology*. 1978; 84:4217–436.
35. de Klepper M, Sleenbos E, van de Bunt G, Agneessens F. Similarity in friendship networks: Selection or influence? The effect of constraining contexts and non-visible individual attributes. *Social Networks*. 2010; 32:82–90.
36. Koram N, Liu H, Li J, Li H, Luo J, Nield J. Role of social network dimensions in the transition to injection drug use: Actions speak louder than words. *AIDS and Behavior*. 2011; 15:1579–1588. [PubMed: 21431412]
37. Petry NM, Weiss LM. Social support is associated with gambling treatment outcomes in pathological gamblers. *The American Journal of Addictions*. 2009; 18:402–408.
38. Ross L, Greene D, House P. The ‘False Consensus Effect’: An egocentric bias in social perception and attribution processes. *Journal of Experimental Social Psychology*. 1977; 13:279–301.


A




B



C



D






Figure 1. Structural social networks of gambling, alcohol use, and tobacco use in two illustrative participants. (A) NPG network's gambling; (B) PG network's gambling; (C) NPG network's smoking; (D) PG network's smoking; (E) NPG network's drinking; (F) PG network's drinking. The depictions are of the same two participants' networks in each case. The participant is not shown in the graphs. Darker colors and larger nodes reflect more frequent gambling, drinking, or smoking behavior. In each case, the networks reveal the significantly greater occurrence of gamblers, smokers, and drinkers for the PG participant; in contrast, the NPG participant exhibits a network in which the addictive behaviors are restricted to more distinct subgroups of associates.

Table 1

Distribution of network members among relationship types, by PG status.

Relationship	Diagnostic Severity			Total
	Count	NPG	PG	
Friend	Count	375	295	670
	Expected count	368.50	301.50	670
Current romantic partner	Count	20	23	43
	Expected count	23.60	19.40	43
Past romantic partner	Count	41	32	73
	Expected count	40.20	32.80	73
Family member	Count	172	176	348
	Expected count	191.40	156.60	348
Co-worker	Count	52	14	66
	Expected count	36.30	29.70	66
Total	Count	660	540	1200
	Expected count	660	540	1200

Table 2

Distribution of alters' overall gambling, smoking and drinking frequency by PG status.

Frequency	Gamble			Smoke			Drink		
	NPG	PG	NPG	NPG	PG	NPG	NPG	PG	PG
Daily	11.06	19.07	27.73	31.85	11.82	23.52			
Multiple times a week	8.33	13.15	6.67	6.85	16.36	11.11			
Once a week	8.33	10.56	1.82	4.26	10.00	10.19			
Once a month	7.27	5.93	2.58	4.44	5.76	6.85			
Less than once a month	7.42	5.00	3.64	4.63	7.42	6.48			
Not in the past year	57.58	46.3	57.58	47.06	48.64	41.85			

Frequency	Gamble			Smoke			Drink		
	NPG	PG	NPG	NPG	PG	NPG	NPG	PG	PG
Daily	11.06(73)	19.07(103)	27.73(183)	31.85(172)	11.82 (78)	23.52(127)			
Multiple times a week	8.33(55)	13.15(71)	6.67 (44)	6.85 (37)	16.36(108)	11.11 (60)			
Once a week	8.33(55)	10.56(57)	1.82 (12)	4.26 (23)	10 (66)	10.19 (55)			
Once a month	7.27(48)	5.93(32)	2.58 (17)	4.44 (24)	5.76 (38)	6.85 (37)			
Less than once a month	7.42(49)	5(27)	3.64 (24)	4.63 (25)	7.42 (49)	6.48 (35)			
Not in the past year	57.58(380)	46.3(250)	57.58(380)	47.06(259)	48.64(321)	41.85(226)			

All values are percentages. Total N's are given in parentheses.

Table 3

Distribution of alters' gambling, smoking and drinking frequency with ego, by PG status.

Frequency	Gamble with Ego		Smoke with Ego		Drink with Ego	
	NPG	PG	NPG	PG	NPG	PG
Daily	8.03	12.41	13.79	20.93	6.06	17.96
Multiple times a week	5.15	7.96	7.88	7.96	8.64	9.81
Once a week	5.61	9.63	2.88	8.70	11.06	6.11
Once a month	7.88	6.85	4.24	4.81	7.12	7.04
Less than once a month	6.82	5.93	5.45	3.70	8.48	6.85
Not in the past year	66.52	57.22	65.76	53.89	58.64	52.22

Frequency	Gamble with Ego		Smoke with Ego		Drink with Ego	
	NPG	PG	NPG	PG	NPG	PG
Daily	8.03 (53)	12.41 (67)	13.79 (91)	20.93(113)	6.06 (40)	17.96 (97)
Multiple times a week	5.15 (34)	7.96 (43)	7.88 (52)	7.96 (43)	8.64 (57)	9.81 (53)
Once a week	5.61 (37)	9.63 (52)	2.88 (19)	8.7 (47)	11.06 (73)	6.11 (33)
Once a month	7.88 (52)	6.85 (37)	4.24 (28)	4.81 (26)	7.12 (47)	7.04 (38)
Less than once a month	6.82 (45)	5.93 (32)	5.45 (36)	3.7 (20)	8.48 (56)	6.85 (37)
Not in the past year	66.52(439)	57.22(309)	65.76(434)	53.89(291)	58.64(387)	52.22(282)

All values are percentages. Total N's are given in parentheses.