Correction to: Foy KC, Miller MJ, Moldovan N, Carson III WE, Kaumaya PTP. Combined vaccination with HER-2 peptide followed by therapy with VEGF peptide mimics exerts effective anti-tumor and anti-angiogenic effects in vitro and in vivo. OncoImmunology 2012; 1:1048–60

Kevin C. Foy,[1,2] Megan J. Miller,[1,2] Nicanor Moldovan,[3] William E. Carson III[4] and Pravin P.T. Kaumaya[1,2,4]

[1]Department of Microbiology; The Ohio State University; Columbus, OH USA; [2]Department of Obstetrics and Gynecology; The Ohio State University; Columbus, OH USA; [3]Department of Internal Medicine; Division of Cardivascular Medicine; The Ohio State University; Columbus, OH USA; [4]James Cancer Hospital and Solove Research Institute and the Comprehensive Cancer Center; The Ohio State University; Columbus, OH USA

In our research paper published in Issue 1, Volume 7 of *OncoImmunology*, a panel from Figure 8 was accidently submitted as part of both Figure 8 and Figure 7. The correct version of **Figure** 7, with its legend, is presented on the following page.

Figure 7. Combination treatment decreases the number of actively dividing cells in a transplantable mouse model of breast cancer. Quantification of the number of actively dividing cells in tumor sections using Ki-67 staining. (**A**) Tissue sections show the amount of positive cells. The stain is specific for cells that are actively dividing. (**B**) Quantification of the staining using Image J software. Data represent mean values from four different fields and error bars represent SD of the mean.