

Marek Szpalski
Robert Gunzburg
Bernard Sztern

An overview of blood-sparing techniques used in spine surgery during the perioperative period

Received: 4 May 2004
Accepted: 7 May 2004
Published online: 15 June 2004
© Springer-Verlag 2004

M. Szpalski (✉)
Department of Orthopedic Surgery,
IRIS South Teaching Hospitals,
142 rue Marconi, 1190 Brussels, Belgium
e-mail: mszp@win.be

R. Gunzburg
Department of Orthopedic Surgery,
Euwfeestkliniek, Antwerp, Belgium

B. Sztern
Department of Internal Medicine,
IRIS South Teaching Hospitals,
Brussels, Belgium

Abstract The problems linked to blood loss and blood-sparing techniques in spine surgery have been less studied than in other fields of orthopedics, such as joint-replacement procedures. Decreasing bleeding is not only important for keeping the patient's hemodynamic equilibrium but also for allowing a better view of the surgical field. In spine surgery the latter aspect is especially important because of the vicinity of major and highly fragile neurologic structures. The techniques and agents used for hemostasis and blood sparing in spinal procedures are mostly similar to those used elsewhere in surgery. Their use is modulated by the specific aspects of spinal approach and its relation to the contents of the spinal canal. Blood-sparing techniques can be divided into two categories based on their goals:

either they are aimed at decreasing the bleeding itself, or they are aimed at decreasing the need for homologous transfusion. Various hemodynamic techniques, as well as systemic and local drugs and agents, can be used separately or in combination, and their use in the field of spine surgery is reported. The level of evidence for the efficacy of many of those methods in surgery as a whole is limited, and there is a lack of evidence for most of them in spine surgery. However, several blood-saving procedures and drugs, as well as promising new agents, appear to be efficient, although their efficacy has yet to be assessed by proper randomized controlled trials.

Keywords Hemostasis · Spine surgery · Bleeding · Blood sparing · Prevention

Introduction

Blood loss in spine surgery is an important issue, even though it appears underestimated, or at least understudied, compared to hip and knee arthroplasty surgery. Blood loss, however, may be an acute problem not only in major deformity surgery but also in less extensive fusion procedures. Decreasing bleeding is important for maintaining a patient's hemodynamic equilibrium and allowing a better view of the surgical field. In spine surgery, the latter aspect is especially important because of the vicinity of major and highly fragile neurologic structures. The surgeon's comfort shortens surgery time, which further decreases bleeding.

In this article we will summarize the main techniques used to spare blood in spine surgery during the perioperative period: from the immediate pre-op to the post-op period. Most of those modalities will be described in greater detail elsewhere in this issue.

Blood-sparing techniques

Blood-sparing techniques can be divided into two groups based on their goals: they are aimed either at decreasing the bleeding itself, or at decreasing the need for homologous transfusion.

Methods aimed at decreasing bleeding

Hemodynamic:

- Controlled hypotension
- Local vasoconstrictors
- Epidural blockade

Chemical/biological:

- Systemic:
 - Desmopressin
 - Aprotinin
 - Tranexamic acid
 - Epsilon-aminocaproic acid
 - Oestrogens
 - Other
- Local:
 - Bone wax
 - Hemostatic “sponges” (gelatin, collagen, cellulose)
 - Fibrin sealants
 - Other

Methods aimed at decreasing homologous transfusion needs

Hemodynamic:

- Acute hemodilution
- Planned autologous transfusion
- Blood saving:
 - Perioperative: cell saving systems
 - Postoperative: drainage recovery systems or cell saving

Chemical/biologic:

- Erythropoietin (EPO)
- Substitutive oxygen carriers

Planned preoperative autologous donation and erythropoietin treatment will not be discussed here, as they take place many weeks prior to surgery and, therefore, are not truly perioperative. This paper will discuss all blood-sparing techniques reported in orthopedics, even if they have not (yet) been published in the context of spine surgery.

Controlled hypotensive anesthesia

Controlled hypotension has been used with success since the 1950s in orthopedic surgery [52]. It is widely applied to spine surgery, and good results have been published since the early 1970s [44, 66, 72, 84, 87, 91, 109].

However, Lennon et al. [77] report in a retrospective study that hypotensive anesthesia did not decrease transfusion requirements compared with normotensive narcosis in scoliosis surgery. The goal of hypotensive anesthe-

sia is to reach a systolic blood pressure of around 60–80 mm Hg. Many different drugs have been used over time, such as anesthetic gases [13], Ca channel blockers [75], beta-blockers [80], nitroglycerin, nitroprusside [59, 50], opiates [21] and anesthetic drugs.

The mechanism by which controlled hypotension decreases blood loss is still unclear and goes beyond the simple explanation that lower blood pressure induces lower blood extravasation. Some authors have hypothesized the existence of an ischemic wound during hypotensive anesthesia, but few studies have really tried to measure blood flow through scientific measures such as flowmetry. Lee et al. measured blood flow in the paraspinal muscles during spine surgery with two different hypotensive drugs, reaching a similar degree of hypotension. They found widely different values for local blood flow [73], although the blood losses were not very different. This indicates that the effect on local blood flow is not the only factor involved. The effect on blood flow in the epidural venous plexuses has also been hypothesized [74], and blood pressure by itself seems to be an important variable influencing blood loss [100].

In the context of spinal fusions, some authors report that, since bleeding is mainly linked to bone decortication and is, therefore, essentially venous, blood loss will not be influenced by a decrease in arterial pressure [15]. In their series of Jehovah’s Witnesses, Brodsky et al. [15] found that blood flow was correlated to surgery duration more than to blood pressure. Kakiushi [62] measured intraosseous pressure in thoracic vertebral bodies during surgery and found that the intraoperative blood loss correlated with intraosseous pressure but that the latter was not correlated to the arterial pressure.

Hypotensive anesthesia may lead to complications [81] and is contraindicated in some cases, mainly in patients with hypertension or ischemic disorders: coronary, cerebral or peripheral.

The possibility of neurological damage at the level of already compressed and compromised nerve roots, where hypotension could add to the suffering of the root, was raised by Krengel et al. [69]. The effect of hypotension on spinal cord function during scoliosis surgery has also been questioned [44]. However, although evoked potential monitoring may show temporary alterations, it does not appear that hypotensive anesthesia increases the risk of neurologic damage [45].

Acute normovolemic hemodilution

In this technique venous blood is collected at the beginning of the procedure, after the induction of anesthesia, in order to diminish hematocrit to a level around 30. In children even lower values can safely be attained (under 20) [33, 101]. The lost volume is compensated with synthetic colloids. Hemodilution can be combined with hypotensive anesthesia.

Despite the hemoglobin loss, tissue oxygenation is maintained through increased cardiac output and better venous return due to reduced viscosity, shear and sludge effect. Furthermore, the loss of red blood cells is decreased, as the blood volume lost during surgery will contain fewer cells. The collected blood can then be retransfused according to need.

Acute normovolemic hemodilution is widely used in spine surgery with good results in fusion [55, 60, 88], as well as in scoliosis surgery [11, 24, 30]. The patient can be kept in hemodilution beyond the surgical-procedure duration by delaying transfusion until the next day, with transfusion then being performed based on clinical judgment [55]. The main contraindications to this technique are ischemic disorders and hemoglobinopathies.

Local vasoconstrictors

Local infiltration of paraspinal muscles with vasoconstrictors (epinephrine, ornipressin) is widely used in spine surgery. This is based on the common belief that vasoconstriction will decrease blood loss. However, the literature is very scarce, and there is little evidence of true efficacy [11]. Also, blood loss does not appear to be related to the dose injected [47].

Epidural blockade

Epidural blockade with normotensive anesthesia has been described as reducing blood loss [61]. It induces vasodilatation in the pelvis and lower limbs with a reactive vasoconstriction above the blocked level and, therefore, can only be used at the lumbar level.

Postoperative wound instillation

Bianconi et al. [12] reported that postoperative wound instillation with ropivacaine was effective in controlling pain and decreasing postoperative blood loss.

Lost blood salvage

Perioperative

With this technique, blood lost during surgery is recuperated and processed through a pump system (Cell Saver, Haemonetics), then transfused back to the patient. In this case it is scavenged blood that returns to the patient. It does not contain platelets or coagulation factors. Therefore, in the case of significant loss and return, a supplementation with fresh-frozen plasma is required [70]. This means that the need for homologous blood products is not

entirely eliminated in this method. It is estimated that about half of the lost red blood cells can be salvaged [37].

The main complication is that a dilutional or disseminated coagulopathy can occur, and there is also a question about the complete elimination of tissue residues. Cell saving is therefore contraindicated in the presence of coagulopathies. Other rare complications include pulmonary injuries probably linked to leukoagglutinins [113] and transient hemoglobinuria [37]. This technique has been reported to be effective in spine surgery [10, 77, 84]. However, Copley et al. [24] did not show an efficacy in cell saving in conjunction with hypotensive anesthesia or with hemodilution in adolescents undergoing surgery for idiopathic scoliosis, and concludes that, in these cases, the cost exceeds the benefit. Shulman et al. compared acute normovolemic hemodilution and hemapheresis for blood-component sequestration with salvage alone. They found that the first technique was more cost effective, decreasing allogeneic as well as autologous blood need [99]. In a meta-analysis, Huet et al. [53] conclude that cell salvage in orthopedic surgery decreases the frequency of allogeneic transfusions, and a recent Cochrane review on cell salvage in surgery suggests that it is efficient in reducing need for allogeneic transfusion, despite the poor methodological quality of most studies [18, 19].

Postoperative

Postoperative collection and transfusion of either washed (cell saving) or unwashed blood has been described in many surgical fields. The most frequent technique consists of re-infusing filtered but unwashed blood from the wound-drainage systems. Many questions remain about the safety and real efficacy of this technique; e.g., the viability of the recovered erythrocytes is doubtful. The nephrotoxicity of free hemoglobin, originating from the hemolyzed red blood cells, poses a safety concern. Furthermore, the efficacy of the filtering system on all tissue residues and metabolites is not entirely clear. However, it seems that, although serum levels of fibrin-degradation products, free hemoglobin and some enzymes [97] increase, they return to previous levels after 24 h. Minor transient transfusion reactions may occur: chills, tachycardia and temperature-increase [103]. Infection and malignancy are contraindications.

This technique is very widely used in hip and knee arthroplasties. There are a few publications in the field of the spine, and these claim efficacy in reducing homologous transfusion requirements [10, 84]. Sebastian et al. [97] showed a reduction of 30% in transfusion needs after re-infusion of drainage-salvaged blood. A combination of intraoperative cell saving with the postoperative infusion of unwashed shed blood enabled Behrman to decrease transfusion requirements by 68% in spinal procedures [10].

Systemic drugs

Epsilon-aminocaproic acid (EACA):
Hemocaprol, Amicar (Lederle) and others

This is an antifibrinolytic agent that prevents plasmin from binding to fibrin. Some rare complications have been described: deep venous thrombosis, pulmonary embolism, renal failure and severe bradycardia.

In the field of spine surgery there are scarce and conflicting results published. In a controlled but not blinded study [35] and then in a proper randomized controlled study against placebo [36], Florentino-Pineda et al. showed a decrease in blood loss and transfusion requirements in patients undergoing idiopathic scoliosis surgery. In fusions in adults, Urban et al. showed limited effect, EACA being only marginally more effective than placebo but much less effective than aprotinin [110]. Amar et al. [5] used it in major orthopedic surgery (including spine) in patients with malignancy and found no efficacy. In a Cochrane review on antifibrinolytic drugs in surgery, Henry et al. [48] found a trend in favor of the use of EACA, but this was marred by poor methodology and biases in the reviewed articles. In other reviews Kovesi et al. [68] did not find any data to support routine use of EACA in orthopedic surgery, and Erstadt [32] came to the same conclusions concerning all types of surgery (including orthopedics).

Aprotinin: Trasylol (Bayer)

The complete mechanism of action of this substance remains unclear. It appears to decrease fibrinolysis by inhibiting plasmin, trypsin and kallikrein. It also avoids pathologic platelet activation by stabilizing the platelet membrane. Finally, it seems to decrease the inflammatory response by inhibiting bradykinin, interleukin and TNF.

Numerous problems are linked to its use: hypercoagulation, thrombus formation and effect on renal function. There also is a high risk of anaphylactic reaction after previous administration. This is especially important, as this drug is very widely used in cardiac surgery. This risk could preclude the drug's use in heart surgery in those patients having previously received it for other procedures such as spine surgery. In a multicenter randomized controlled trial, Samama et al. [94] have demonstrated a decrease of blood loss and transfusion with aprotinin in major orthopedic surgery. A Cochrane review [48] of antifibrinolytic drugs in surgery found existing evidence for the use of aprotinin, despite biases in some of the reviewed studies.

The use of aprotinin in spine surgery has been documented in a few controlled studies. Urban showed a decrease in blood loss and transfusion needs in adult fusion, and Cole et al. [23] showed similar results in deformity surgery on children and adolescents. Lentschener et al.

[78] demonstrated dramatic reduction of intraoperative and 24-h blood loss but did not notice a significant reduction in homologous transfusion needs. This study demonstrated the decrease on intraoperative fibrinolysis through intraoperative dosage of D-dimmeder. Amar et al. used aprotinin for major orthopedic surgery (including spine) in patients with malignancy and found no efficacy [5] in reducing blood loss. There are conflicting results as to the dose regimen to be used, with some reporting high dosage [78, 94] and others half-dosage efficacy [110].

Deamino-8-d-arginine-vasopressin
(DDVAP or desmopressin): Minirin (Ferring)

This is an analog to L-arginine-vasopressin or antidiuretic hormone (ADH). Its mechanism of action lies principally in an increase of the secretion of factor VIIIc. It also increases the secretion of von Willebrand factor (vWF) and has a paradoxical effect on the increase of plasminogen activator. There are conflicting reports about the use of desmopressin in scoliosis surgery. In a controlled trial, Kobrinsky et al. [67] found decrease in blood loss, transfusion requirements and use of analgesic agents in the postoperative period. The latter might be due to the decrease of bleeding in the wound. Other studies, however, do not confirm such findings. Alanay et al. [2] conducted a randomized trial against placebo and did not find any significant effect on blood loss in idiopathic or congenital scoliosis. Guay et al. [46] reached the same conclusion.

In neuromuscular scoliosis, where coagulation abnormalities are often present [63, 89], leading to severe blood loss [31], Theroux et al. [104], in a randomized controlled trial, showed that although administration of DDVAP significantly increased factor VIIIc and vWF, it didn't decrease blood loss compared to placebo. Similarly, Lett's et al. [79] conclude that DDVAP decreases bleeding in some patients while others show no response. They propose testing bleeding time after a test dose of DDAVP a few days before surgery.

Several reviews do not find arguments for a wide use of DDAVP in order to reduce surgical bleeding. Kovesi [68] concluded that it can be efficient in patients with a defect in platelet function but found no evidence for use in routine elective orthopedic surgery. A recent Cochrane review of DDAVP in surgery does not find any evidence for surgical use outside patients with congenital bleeding disorders [18]. Desmopressin might be well indicated in patients with von Willebrand disease, acquired platelet disorders, renal failure, cirrhosis or long-term salicylate treatment.

Oestrogens: estriol, conjugated oestrogens

Oestrogens have been widely used in the past for hemostasis. However, their apparent lack of efficacy has made

their use scarce [32]. The mode of action is unclear. It seems that they decrease the permeability and increase the resistance of capillary walls as well as improve the platelet/wall interaction. They also have an antifibrinolytic activity by altering levels of factor V and decreasing the antithrombin activity of plasma.

In a controlled study McCall et al. [86] have reported the efficacy of conjugated oestrogens (Premarin, Wyeth-Ayerst) in decreasing the postoperative drainage volume in adolescents having undergone deformity surgery. There were no side effects. In a review of systemic hemostatic drugs, Erstadt [32] found only limited efficacy of conjugated oestrogens in the reduction of blood loss in surgery.

Tranexamic acid: Exacyl (Sanofi-Synthelabo),
Cyklokapron (Pfizer) and others

Tranexamic acid decreases fibrinolysis by inhibiting transformation of plasminogen into plasmin. Recent studies show efficacy in reduction of bleeding during hip- [76] and knee-replacement [56] procedures. Other studies, however, seem to show that tranexamic acid does not decrease hidden losses [40]. Meta-analysis of the use of tranexamic acid in surgery shows either specific efficacy in knee arthroplasty [32, 51] or a trend towards effectiveness in other procedures, but results are marred by methodological flaws in the existing publications [48]. In the field of spine surgery, only one double-blind, placebo-controlled study in children and adolescents undergoing scoliosis surgery showed a reduction in transfusion needs without added complications [90].

Recombinant factor VIIa/RhFVIIa
(NovoSeven, Novo Nordisk)

This drug is an anti-hemophilic agent. Its efficacy on hemostasis in non-hemophilic subjects during surgery [4], neurosurgery [64] and trauma [43] has been reported. A double-blind randomized control trial has shown good results in prostatic surgery [39]. In spine surgery there are some anecdotal reports [106] but there is currently a lack of evidence, due to a paucity of available studies [68]. This agent seems promising but true evidence of its efficacy in reducing bleeding during spine surgery has yet to be collected.

Other systemic means

Amino-acid infusion that induces thermogenesis prevented a drop in body temperature and decreased blood loss in hip arthroplasty [114]. Likewise, aggressive warming to keep body temperature at 36.5°C appears to decrease blood loss in hip replacement [115]. There are no

reports on the effect of warming during spine surgery. Etamsylate (Dycinone, Sanofi-Synthelabo), a drug marketed as hemostatic, was used without success in hip replacement [65], and no data on its use in spine surgery is available.

Substitutive hemoglobin carriers (hemoglobin raffimers)

Substitutive oxygen carriers are an elegant solution for the future [6]. They are often incorrectly called “artificial hemoglobins,” whereas the products in clinical experimentation are made from human or animal stabilized acellular hemoglobin. Hemopure (Biopure) is already marketed in South Africa, although the ethical issues regarding the early commercial human use of such products are discussed [99]. It is a bovine stabilized hemoglobin for which an FDA regulatory filing phase III trial has been completed in orthopedic surgery (including spine). This study showed an important decrease in transfusion needs. There do not appear to be major safety issues [102].

Hemolink (Hemosol) is a similar product but based on human hemoglobin. Results of phase II trials in cardiac surgery have been published showing decreased allogeneic transfusion needs with good safety [20]. A clinical phase III study, along the same lines, shows a marked reduction in the need for transfusions [42]. Phase II trials in orthopedic surgery are on the way in Canada and the US.

Local agents

Bone wax

Bone wax is a mixture of beeswax (70%) and petroleum jelly (30%). It acts in a purely mechanical way, by covering and filling bleeding bone surfaces. Its best known use is probably the hemostasis of sternotomy surfaces. There have been few reports in the field of spine surgery. Abumi et al. [1] describe an injury of the vertebral artery during cervical pedicular-screw insertion that was subsequently stopped with bone wax. Use of bone wax to stop bleeding of the bony surface in the spine is hampered by the fear of leaving a foreign body that can intrude into the spinal canal. Indeed, Cirak and Unal describe a case of tetraplegia following use of bone wax [22].

Hemostatic “sponges”

*Gelatin-based (Gelfoam (Pharmacia),
Surgifoam (Johnson & Johnson), and others*

Gelatin-based, local hemostatic agents have been used in surgery for decades. They can be of bovine, porcine or equine origin and are available in multiple presentations:

sheets, powder, foam, etc. They can be used alone or soaked with thrombin. Their mechanism of action is still not completely clear but appear to be more physical, by “surface effect”, than through any action on the blood-clotting mechanism. Gelatin-based devices have been reported to induce a better quality clot than collagen-based products [28]. The substance has been widely used in spine surgery, as it is considered safe to leave in the canal because it does not swell. Some authors have even suggested that gelatin reduces scar adhesion [71].

However, several publications report severe neurological consequences including cauda equina syndromes linked to the use of gelatin products in the spinal canal [3, 9, 38, 49]. Allergic reaction to gelatin after spinal use has also been reported [92].

*Collagen-based: Instat (Johnson & Johnson),
Lyostrict (B-Braun), Hemocol (Pilling-Weck), and others*

Collagen based hemostatic products are from bovine, porcine or equine origin, and also exist in different forms: sheets, powder, aggregates, etc. They act on platelet aggregation and activate Hageman factor (F XII). They should not be left in the spinal canal as they provoke adhesion formation and foreign body reactions.

A more elaborate product is Tachocomb (Nycomed). It consists of a patch of collagen coated with fibrinogen and thrombin. In contact with liquids the components dissolve and the last phase of coagulation is launched, resulting in a fibrin-clot formation. That mode of action is similar to that of the fibrin sealants reviewed later. It also contains aprotinin in order to inhibit clot fibrinolysis. It is mostly used in thoracic and abdominal surgery, but repair of lacerations to the dural sac with this product have been reported [58] and experimental models have shown effectiveness in avoiding epidural fibrosis [73]. The use in spine surgery appears to raise the potential danger of leaving collagen-based agents in the spinal canal.

*Oxidized cellulose based: Surgicel (Johnson & Johnson),
Curacel (Curaspon) and others*

These products also exist in multiple forms, such as sheets, gauze and powder. Their action is essentially a surface effect, which activates the initial coagulation phase. They also induce a moderate acceleration of fibrinogen polymerization. They should not be left in the canal, as they swell and cause foreign-body reactions. Neurological complications have been widely reported [8, 14, 57, 82]. Epidural migration, causing severe, and sometimes permanent, neurological lesions after use of cellulose hemostatic agents for thoracotomy have also been reported [98, 112]. The foreign-body reactions provoked by cellulose may induce granulomas and pseudo tumors [16, 95]. These reactions

may also result in misleading pseudo-compression images on MRI examinations [7].

Fibrin sealants

Fibrin sealants reproduce the last phase of coagulation: the formation of a fibrin clot. There are two main families of fibrin sealant. The first combines two components: a fibrinogen component and a thrombin solution. A more recent type of sealant uses the fibrinogen of the bleeding source itself.

*Bi-component fibrin sealant: Tissucol/Tisseel (Baxter),
Beriplast (Behring), Hemaseel (Haemacure),
CoStasis (Cohesion Tech)*

All the available sealants in this category are used by mixing a fibrinogen component with a thrombin solution to form a fibrin clot. They usually contain factor XII to speed the cross linking of the clot. Some may contain aprotinin to inhibit fibrinolysis and even plasminogen to control the reaction. They are applied with a syringe, but a sprayable form exists to cover larger surfaces. However, they adhere poorly on wet and bleeding surfaces. Content of the sealant solution is especially important in spine surgery because of possible contact with major neurological structures. Quixil (Omrix, Brussels, Belgium) contains tranexamic acid instead of aprotinin as antifibrinolytic adjuvant. It appears that tranexamic acid is neurotoxic. Fatal neurotoxic reactions have been reported after use in neurosurgery [85].

Fibrin sealants have been reported to be effective in spine surgery [107] and to diminish scar formation [111]. A Cochrane review on the use of fibrin sealants in surgery suggests efficacy, but this conclusion is hampered by the small number of trials and their mostly dubious methodology [17].

The use of bovine fibrinogen may pose a safety problem in the context of spongiform encephalitis and the marketing of one fibrin glue, Biocol, had to be stopped. CoStasis uses the patient’s centrifuged plasma with bovine thrombin. It is useful for avoiding the need for bovine fibrinogen (although bovine thrombin remains), but it seems expensive, cumbersome and time consuming. It has been reported to be effective in spine surgery [29, 105].

*Fibrin sealant using wound fibrinogen:
FloSeal (Baxter, formerly Proceed, by Centerpulse)*

This second family of sealants contains thrombin but relies on the wound’s fibrinogen. The bi-component medium contains a collagen/thrombin component and a gelatin matrix. The swelling of the collagen granules will restrict bleeding through a tamponing mechanism, while the gelatin

matrix will provide structural integrity to remain in situ. The main advantage of that form is that it can more easily be used on wet and bleeding tissues. Good results have been reported in spinal procedures [93].

N-butyl-cyanoacrylate: Histoacryl (B-Braun)

Apart from its cutaneous applications, this acrylic glue is used classically in gastroenterology to control gastro-esophageal bleeding. In spine surgery it has been used for injection of vertebral hemangiomas or malignant tumors prior to surgery in order to reduce peroperative bleeding [25, 27, 108].

Conclusions: where is the evidence?

The range of techniques available for sparing blood in spine procedures, as in other areas of surgery, is very wide and the variety of concepts is large. However, the evidence, aside from the anecdotal, is often lacking or conflicting. Controlled studies are scarce, and we do not really know if we can adapt, to spinal surgery, conclusions stemming from other surgical fields, mainly cardiac surgery, as to the efficacy of hemostatic means.

There seems to be adequate evidence showing the efficacy of most hemodynamic methods, such as hypotensive anesthesia and normovolemic hemodilution. The systemic hemostatic agents show fragmented and conflicting evi-

dence as to their effectiveness in spine surgery. A Cochrane review of antifibrinolytic drugs in elective surgery has found good evidence to support the use of aprotinin in cardiac surgery and found trends for the effectiveness of tranexamic acid and epsilon aminocaproic acid, without, however, finding conclusive evidence, due to the heterogeneity and biases of many studies [48]. In elective orthopedic surgery, some reviews state that no hard data exist to support use of any fibrinolytic or other systemic drug [68]. RhFVIIa seems promising in the field of spine surgery, in light of results in other fields, but conclusive evidence should be demonstrated in spine surgery by proper randomized controlled trials. Alternative hemoglobin carriers are undoubtedly experimental products. As far as the local agents are concerned, once again, conclusive evidence is scarce and reports are mainly anecdotal. However, the realization of true randomized trials for those kinds of techniques is very difficult.

The daily, routine use of all these methods is also quite limited at the present time. Several surveys have shown that, outside of cardiac surgery, regular use of any blood-sparing method is infrequent [41, 54]. Lack of familiarity is the first reason quoted for their infrequent use [54]. Techniques (hemodilution, cell salvage, etc.) are more frequently used than pharmaceuticals, and there is a considerable variation among different countries [34].

Blood sparing in spine surgery is important, and it clearly appears that, contrary to other orthopedic and surgical fields, it has been understudied, with the current practice being more based on beliefs than on evidence.

References

1. Abumi K, Shono Y, Ito M, Taneichi H, Kotani Y, Kaneda K (2000) Complications of pedicle screw fixation in reconstructive surgery of the cervical spine. *Spine* 15:962–969
2. Alanay A, Acaroglu E, Ozdemir O, Ercelen O, Bulutcu E, Surat A (1999) Effects of deamino-8-D-arginin vasopressin on blood loss and coagulation factors in scoliosis surgery. A double-blind randomized clinical trial *Spine* 24:877–882
3. Alander DH, Stauffer ES (1995). Gelfoam-induced acute quadriplegia after cervical decompression and fusion. *Spine* 20:970–971
4. Aldouri M (2002) The use of recombinant factor VIIa in controlling surgical bleeding in non-haemophilic patients. *Pathophysiol Haemost Thromb* 32 [Suppl 1]:41–46
5. Amar D, Grant FM, Zhang H, Boland PJ, Leung DH, Healey JA (2003) Antifibrinolytic therapy and perioperative blood loss in cancer patients undergoing major orthopedic surgery. *Anesthesiology* 98:337
6. Arnoldo BD, Minei JP (2001) Potential of hemoglobin-based oxygen carriers in trauma patients. *Curr Opin Crit Care* 7:431–436
7. Awwad EE, Smith KR (1999) MRI of marked dural sac compression by surgical in the immediately postoperative period after uncomplicated lumbar laminectomy. *J Comput Assist Tomogr* 23:969–975
8. Banerjee T, Goldschmidt K (1998) "Surgiceloma" manifested as cauda equina syndrome. *South Med J* 91:481–483
9. Bellen P (1992) Prevention of peridural fibrosis following laminectomy. Apropos of a case of monoradicular paralysis due to an intracanalicular hematoma on Gelfoam. *Acta Orthop Belg* 58:236–239
10. Behrman MJ, Keim HA (1992) Perioperative red blood cell salvage in spine surgery. A prospective analysis. *Clin Orthop* 278:51–57
11. Berniere J, Scemama MP, Granados M, Beurrier C, Bercovy M (1982) Value of deliberate normovolemic hemodilution associated with a capillary vasoconstrictor, ornipressin, in surgery of the spine. *Ann Fr Anesth Reanim* 1:419–423
12. Bianconi M, Ferraro L, Ricci R, Zanoli G, Antonelli T, Giulia B, Guberti A, Massari L (2004) The pharmacokinetics and efficacy of ropivacaine continuous wound instillation after spine fusion surgery. *Anesth Analg* 98:166–172
13. Bourreli B, Pinaud M, Passuti N, Gunst JP, Drouet JC, Remi JP (1988) Additive effects of dihyalazine during enflurane or isoflurane hypotensive anaesthesia for spinal fusion. *Can J Anaesth* 35:242–248
14. Brodbelt AR, Miles JB, Foy PM, Broome JC (2002) Intraspinal oxidized cellulose (Surgicel) causing delayed paraplegia after thoracotomy – a report of three cases. *Ann R Coll Surg Engl* 84:97–99

15. Brodsky JW, Dickson JH, Erwin WD, Rossi CD (1991) Hypotensive anesthesia for scoliosis surgery in Jehovah's Witnesses. *Spine* 16:304–306
16. Buckley SC, Broome JC (1995) *Br J Neurosurg* 9:561–563
17. Carless PA, Anthony DM, Henry DA (2002) Systematic review of the use of fibrin sealant to minimize perioperative allogeneic blood transfusion. *Br J Surg* 89:695–703
18. Carless P, Henry D, Moxey A, O'Connell D, McClelland B, Henderson K, Sly K, Laupacis A, Fergusson D (2004) Desmopressin for minimising perioperative allogeneic blood transfusion. *Cochrane Database Syst Rev* 1:CD001884
19. Carless PA, Henry DA, Moxey AJ, O'Connell DL, Ferguson DA (2004) Cell salvage for minimising perioperative allogeneic blood transfusion. IN: *The Cochrane Library, Issue 1*. Wiley, London
20. Cheng DC, Mazer CD, Martineau R, Ralph-Edwards A, Karski J, Robblee J, Finegan B, Hall RI, Latimer R, Vuylsteke A (2004) A phase II dose-response study of hemoglobin rafi-fimer (Hemolink) in elective coronary artery bypass surgery. *J Thorac Cardiovasc Surg* 127:79–86
21. Chillemi S, Sinardi D, Marino A, Mantarro G, Campisi R (2002) The use of remifentanyl for bloodless surgical field during vertebral disc resection. *Minerva Anestesiologica* 68:645–649
22. Cirak B, Unal O (2000) Iatrogenic quadriplegia and bone wax. Case illustration. *J Neurosurg* 92 [Suppl 2]: 248
23. Cole JW, Murray DJ, Snider RJ, Bassett GS, Bridwell KH, Lenke LG (2003) Aprotinin reduces blood loss during spinal surgery in children. *Spine* 28:2482–2485
24. Copley LA, Richards BS, Safavi FZ, Newton PO (1999) Hemodilution as a method to reduce transfusion requirements in adolescent spine fusion surgery. *Spine* 24:219–222
25. Cortet B, Cotten A, Deprez X, Deramond H, Lejeune JP, Leclerc X, Chastanet P, Duquesnoy B, Delcambre B (1994) Value of vertebroplasty combined with surgical decompression in the treatment of aggressive spinal angioma. Apropos of 3 cases. *Rev Rhum Ed Fr* 61:16–22
26. CoStasis Multi-Center Collaborative Writing Committee (2001) A novel collagen-based composite offers effective hemostasis for multiple surgical indications: Results of a randomized controlled trial. *Surgery* 129: 445–450
27. Cotten A, Deramond H, Cortet B, Lejeune JP, Leclerc X, Chastanet P, Clarisse J (1996) Preoperative percutaneous injection of methyl methacrylate and n-butyl cyanoacrylate in vertebral hemangiomas. *Am J Neuro-radiol* 17:137–142
28. Dehen M, Niederdelmann H, Lachner J (1990) Mechanical properties of collagen or gelatine-stabilized blood clots. *Dtsch Z Mund Kiefer Gesichtschir* 14:229–231
29. Dunne JR et al (2003) A novel approach to the treatment of gunshot injuries to the sacrum. *Am Surg* 69: 91–94
30. Du Toit G, Relton JE, Gillespie R (1978) Acute haemodilutional autotransfusion in the surgical management of scoliosis. *J Bone Joint Surg Br* 60:178–180
31. Edler A, Murray DJ, Forbes RB (2003) Blood loss during posterior spinal fusion surgery in patients with neuromuscular disease: is there an increased risk? *Paediatr Anaesth* 13: 818–822
32. Erstad BL (2001) Systemic hemostatic medications for reducing surgical blood loss. *Ann Pharmacother* 35: 925–934
33. Feldman JM, Roth JV, Bjoraker DG (1995) Maximum blood savings by acute normovolemic hemodilution. *Anesth Analg* 80:108–113
34. Fergusson D, Blair A, Henry D, Hisashige A, Huet C, Koopman-van Gemert A, Katz E, McClelland B, Sigmund H, Laupacis A (1999) Technologies to minimize blood transfusion in cardiac and orthopedic surgery. Results of a practice variation survey in nine countries. *International Study of Peri-operative Transfusion (ISPOT) Investigators. Int J Technol Assess Health Care.* 15:717–728
35. Florentino-Pineda I, Blakemore LC, Thompson GH, Poe-Kochert C, Adler P, Tripi P (2001) The Effect of epsilon-aminocaproic acid on perioperative blood loss in patients with idiopathic scoliosis undergoing posterior spinal fusion: a preliminary prospective study. *Spine* 26:1147–1151
36. Florentino-Pineda I, Thompson GH, Poe-Kochert C, Huang RP, Haber LL, Blakemore LCT (2004) The effect of amicar on perioperative blood loss in idiopathic scoliosis: the results of a prospective, randomized double-blind study. *Spine*. 29:233–238
37. Flynn JC, Metzger CR, Csencsitz TA (1982) Intraoperative autotransfusion (IAT) in spinal surgery. *Spine* 7:432–435
38. Friedman J, Whitecloud TS (2001) Lumbar cauda equina syndrome associated with the use of gelfoam: case report. *Spine* 26:E485–487
39. Friedrich PW, Henny CP, Messelink EJ, Geerdink MG, Keller T, Kurth KH, Buller HR, Levi M (2003) Effect of recombinant activated factor VII on perioperative blood loss in patients undergoing retropubic prostatectomy: a double-blind placebo-controlled randomised trial. *Lancet* 361(9353): 201–205
40. Good L, Peterson E, Lisander B (2003) Tranexamic acid decreases external blood loss but not hidden blood loss in total knee replacement. *Br J Anaesth* 90:596–599
41. Graham ID, Fergusson D, McAuley L, Laupacis (2000) The use of technologies to minimize exposure to perioperative allogeneic blood transfusion in elective surgery. A survey of Canadian hospitals. *Int J Technol Assess Health Care* 16:228–241
42. Greenburg AG, Kim HW, Hemolink Study Group (2004) Use of an oxygen therapeutic as an adjunct to intraoperative autologous donation to reduce transfusion requirements in patients undergoing coronary artery bypass graft surgery. *J Am Coll Surg* 198:373–383
43. Grounds M (2003) Recombinant factor VIIa (rFVIIa) and its use in severe bleeding in surgery and trauma: a review. *Blood Rev* 17 [Suppl 1]:S11–21
44. Grundy BL, Nash CL Jr, Brown RH (1981) Arterial pressure manipulation alters spinal cord function during correction of scoliosis. *Anesthesiology* 54:249–253
45. Grundy BL, Nash CL Jr, Brown RH (1982) Deliberate hypotension for spinal fusion: prospective randomized study with evoked potential monitoring. *Can Anaesth Soc J* 29:452–462
46. Guay J, Reinberg C, Poitras B, David M, Mathews S, Lortie L, Rivard GE (1992) A trial of desmopressin to reduce blood loss in patients undergoing spinal fusion for idiopathic scoliosis. *Anesth Analg* 75:317–318
47. Guay J, Haig M, Lortie L, Guertin MC, Poitras B (1994) Predicting blood loss in surgery for idiopathic scoliosis. *Can J Anaesth* 41:775–781
48. Henry DA, Moxey AJ, Carless PA, O'Connell D, McClelland B, Henderson KM, Sly K, Laupacis A, Fergusson D (2001) Anti-fibrinolytic use for minimizing perioperative allogeneic blood transfusion. *Cochrane Database Syst Rev* CD001886
49. Herndon JH, Grillo HC, Riseborough EJ, Rich JC Jr (1972) Compression of the brain and spinal cord following use of gelfoam. *Arch Surg* 104:107

50. Hersey SL, O'Dell NE, Lowe S, Rasmussen G, Tobias JD, Deshpande JK, Mencia G, Green N (1997) Nicardipine versus nitroprusside for controlled hypotension during spinal surgery in adolescents. *Anesth Analg* 84:1239–1244
51. Ho KM, Ismail H (2003) Use of intravenous tranexamic acid to reduce allogeneic blood transfusion in total hip and knee arthroplasty: a meta-analysis. *Anaesth Intensive Care* 31:529–537
52. Holmes F (1956) Induced hypotension in orthopaedic surgery; hexamethonium bromide (C6) in 407 orthopaedic operations. *J Bone Joint Surg Br* 38:846–854
53. Huet C, Salmi LR, Fergusson D, Koopman-van Gemert AW, Rubens F, Laupacis A (1999) A meta-analysis of the effectiveness of cell salvage to minimize perioperative allogeneic blood transfusion in cardiac and orthopedic surgery. *International Study of Perioperative Transfusion (ISPOT) Investigators. Anesth Analg* 89:861–869
54. Hutchinson AB, Fergusson D, Graham ID, Laupacis A, Herrin J, Hillyer CD (2001) Utilization of technologies to reduce allogeneic blood transfusion in the United States. *Transfus Med* 11:79–85
55. Hur SR, Huizenga BA, Major M (1992) Acute normovolemic hemodilution combined with hypotensive anesthesia and other techniques to avoid homologous transfusion in spinal fusion surgery. *Spine* 17:867–873
56. Hynes M, Calder P, Scott G (2003) The use of tranexamic acid to reduce blood loss during total knee arthroplasty. *Knee* 10:375–377
57. Iwabuchi S, Koike K, Okabe T, Tago S, Murakami T (1997) Iatrogenic paraplegia caused by surgicel used for hemostasis during a thoracotomy: report of a case. *Surg Today* 27:969–970
58. Kaczmarczyk R, Markiewicz P, Trojanowski T (2001) The application and evaluation of TachoComb in repair of lacerations of dural sac. *Neurol Neurochir Pol* 35 [Suppl 5]:64–69
59. Kadam PP, Saksena SG, Jagtap SR, Pantavaidya SM (1993) Hypotensive anaesthesia for spine surgery – nitroglycerin vs halothane. *J Postgrad Med* 39:26–28
60. Kafer ER (1986) *Anesth Analg* 65: S76
61. Kakiuchi M (1997) Reduction of blood loss during spinal surgery by epidural blockade under normotensive general anesthesia. *Spine* 22:889–894
62. Kakiuchi M (2002) Intraoperative blood loss during cervical laminoplasty correlates with the vertebral intrasosseous pressure. *J Bone Joint Surg Br* 84:518–520
63. Kannan S, Meert KL, Mooney JF, Hillman-Wiseman C, Warriar I (2002) Bleeding and coagulation changes during spinal fusion surgery: a comparison of neuromuscular and idiopathic scoliosis patients. *Pediatr Crit Care Med* 3:364–369
64. Karadimov D, Binev K, Nachkov Y, Platikanov V (2003) Use of activated recombinant Factor VII (NovoSeven) during neurosurgery. *J Neurosurg Anesthesiol* 15:330–332
65. Keith I (1979) Ethamsylate and blood loss in total hip replacement. *Anaesthesia* 34:666–670
66. Khambatta HJ, Stone JG, Matteo RS, Michelsen WJ (1978) Hypotensive anesthesia for spinal fusion with sodium nitroprusside. *Spine* 3:171–174
67. Kobrinsky NL, Letts RM, Patel LR, Israels ED, Monson RC, Schwetz N, Cheang MS (1987) 1-Desamino-8-D-arginine vasopressin (desmopressin) decreases operative blood loss in patients having Harrington rod spinal fusion surgery. A randomized, double-blinded, controlled trial. *Ann Intern Med* 107:446–450
68. Kovesi T, Royston D (2003) Pharmacological approaches to reducing allogeneic blood exposure. *Vox Sang* 84: 2–10
69. Krengel WF 3rd, Robinson LR, Schneider VA (1993) Combined effects of compression and hypotension on nerve root function. A clinical case. *Spine* 18:306–309
70. Kruger LM, Colbert JM (1985) Intraoperative autologous transfusion in children undergoing spinal surgery. *J Pediatr Orthop* 5:330–332
71. LaRocca H, McNab I (1974) The laminectomy membrane. Studies in its evolution, characteristics, effects and prophylaxis in dogs. *J Bone Joint Surg Br* 56:545–550
72. Lawhon SM, Kahn A 3rd, Crawford AH, Brinker MS (1984) Controlled hypotensive anesthesia during spinal surgery. A retrospective study. *Spine* 9:450–453
73. Lee JY, Ebel H, Friese M, Schillinger G, Schroder R, Klug N (2003) Influence of TachoComb in comparison to local hemostyptic agents on epidural fibrosis in a rat laminectomy model. *Minim Invasive Neurosurg* 46:106–109
74. Lee TC, Yang LC, Chen HJ (1998) Effect of patient position and hypotensive anesthesia on inferior vena caval pressure. *Spine* 23:941–948
75. Lee TC, Buerkle H, Wang CJ, Liang CL, Lu K, Huang PL, Cheng CH, Yang LC (2001) Effect of isoflurane versus nicardipine on blood flow of lumbar paraspinal muscles during controlled hypotension for spinal surgery. *Spine* 26:105–109
76. Lemay E, Guay J, Cote C, Roy A (2004) Tranexamic acid reduces the need for allogeneic red blood cell transfusions in patients undergoing total hip replacement. *Can J Anaesth* 51:31–37
77. Lennon RL, Hosking MP, Gray JR, Klassen RA, Popovsky MA, Warner MA (1987) The effects of intraoperative blood salvage and induced hypotension on transfusion requirements during spinal surgical procedures. *Mayo Clin Proc* 62:1090–1094
78. Lentschener C, Cottin P, Bouaziz H, Mercier FJ, Wolf M, Aljabi Y, Boyer-Neumann C, Benhamou D (1999) Reduction of blood loss and transfusion requirement by aprotinin in posterior lumbar spine fusion. *Anesth Analg* 89:590–597
79. Letts M, Pang E, D'Astous J, Jarvis J, Lawton L, Luke B, Rhine E, Menard E (1998) The influence of desmopressin on blood loss during spinal fusion surgery in neuromuscular patients. *Spine* 23:475–478
80. Lim YJ, Kim CS, Bahk JH, Ham BM, Do SH (2003) Clinical trial of esmolol-induced controlled hypotension with or without acute normovolemic hemodilution in spinal surgery. *Acta Anaesthesiol Scand* 47:74–78
81. Lindop MJ (1975) Complications and morbidity of controlled hypotension. *Br J Anaesth* 47:799–803
82. Lovstad RZ, Steen PA, Forsman M (1999) Paraplegia after thoracotomy – not caused by the epidural catheter. *Acta Anaesthesiol Scand* 43:230–232
83. Malcolm-Smith NA, McMaster M (1983) The use of induced hypotension to control bleeding during posterior fusion for scoliosis. *J Bone Joint Surg Br* 65:255–258
84. Mandel RJ, Brown MD, McCollough NC 3rd, Pallares V, Varlotta R (1981) Hypotensive anesthesia and autotransfusion in spinal surgery *Clin Orthop* 154:27–33
85. MCA/CSM (1999) Current problems in pharmacovigilance. 25:19
86. McCall RE, Bilderback KK (1997) Use of intravenous Premarin to decrease postoperative blood loss after pediatric scoliosis surgery. *Spine* 22: 1394–1397
87. McNeill TW, DeWald RL, Kuo KN, Bennett EJ, Salem MR (1974) Controlled hypotensive anesthesia in scoliosis surgery. *J Bone Joint Surg Am* 56:1167–1172

88. Messmer K (1975) Hemodilution. *Surg Clin North Am* 55:659–675
89. Meert KL, Kannan S, Mooney JF (2002) Predictors of red cell transfusion in children and adolescents undergoing spinal fusion surgery. *Spine* 27:2137–2142
90. Neilipovitz DT, Murto K, Hall L, Barrowman NJ, Splinter WM (2001) A randomized trial of tranexamic acid to reduce blood transfusion for scoliosis surgery. *Anesth Analg* 93:82–87
91. Patel NJ, Patel BS, Paskin S, Laufer S (1985) Induced moderate hypotensive anesthesia for spinal fusion and Harrington-rod instrumentation. *J Bone Joint Surg Am* 67:1384–1387
92. Purrello-D'Ambrosio F, Gangemi S, La Rosa G, Merendino RA, Tomasello F (2000) Allergy to gelatin. *Allergy* 55:414–415
93. Renkens KL Jr, Payner TD, Leipzig TJ, Feuer H, Morone MA, Koers JM, Lawson KJ, Lentz R, Shuey H Jr, Conaway GL, Andersson GB, An HS, Hickey M, Rondinone JF, Shargill NS (2001) A multicenter, prospective, randomized trial evaluating a new hemostatic agent for spinal surgery. *Spine* 26:1645–1650
94. Samama CM, Langeron O, Rosencher N, Capdevila X, Rouche P, Pegoix M, Berniere J, Coriat P (2002) Aprotinin versus placebo in major orthopedic surgery: a randomized, double-blinded, dose-ranging study. *Anesth Analg* 95:287–293
95. Sandhu GS, Elexpuru-Camiruaga JA, Buckley S (1996) Oxidized cellulose (Surgicel) granulomata mimicking tumour recurrence. *Br J Neurosurg* 10: 617–619
96. Schutte AE (2002) Ethical considerations concerning South Africa's approval of a blood substitute. *Ethics Med* 18:55–62
97. Sebastian C, Romero R, Olalla E, Ferrer C, Garcia-Vallejo JJ, Munoz M (2000) Postoperative blood salvage and reinfusion in spinal surgery: blood quality, effectiveness and impact on patient blood parameters. *Eur Spine J* 9:458–465
98. Short HD (1990) Paraplegia associated with the use of oxidized cellulose in posterolateral thoracotomy incisions. *Ann Thorac Surg* 50:288–289
99. Shulman G, Solanki DR, Nicodemus CL, Flores IM, Hadjipavlou AG (1998) Audit of autotransfusion in spine surgery. *Int Orthop* 22:303–307
100. Sivarajan M, Amory DW, Everett GB, Buffington C (1980) Blood pressure, not cardiac output, determines blood loss during induced hypotension. *Anesth Analg* 59:203–206
101. Singler RC, Furman EB (1980) Hemodilution: How low a minimum hematocrit? *Anesthesiology* 53:S72
102. Sprung J, Kindscher JD, Wahr JA, Levy JH, Monk TG, Moritz MW, O'Hara PJ (2002) Hemopure in surgical patients: results of a multicenter, randomized, single-blinded trial. *Anesth Analg* 94:799–808
103. Tate DE Jr, Friedman RJ (1992) Blood conservation in spinal surgery. Review of current techniques. *Spine* 17:1450–1456
104. Theroux MC, Corddry DH, Tietz AE, Miller F, Peoples JD, Kettrick RG (1997) A study of desmopressin and blood loss during spinal fusion for neuromuscular scoliosis: a randomized, controlled, double-blinded study. *Anesthesiology* 87:260–267
105. Thibodeaux KT, Lorio MP, Block JE (2003) Intraoperative hemostasis during spinal reconstructive procedures. *Orthopedics* 26:413–414
106. Tobias JD (2002) Synthetic factor VIIa to treat dilutional coagulopathy during posterior spinal fusion in two children. *Anesthesiology* 96:1522–1525
107. Tredwell SJ, Sawatzky B (1990) The use of fibrin sealant to reduce blood loss during Cotrel-Dubousset instrumentation for idiopathic scoliosis. *Spine* 15:913–915
108. Trystram D, Aymard A, Godon Hardy S, Cioloca C, Fredy D, Meder J (2000) Preoperative devascularization of a vertebral metastasis with a spinal artery at the same level. *J Radiol* 81:250–253
109. Ullrich PF Jr, Keene JS, Hogan KJ, Roecker EB (1990) Results of hypotensive anesthesia in operative treatment of thoracolumbar fractures. *J Spinal Disord* 3:329–333
110. Urban MK, Beckman J, Gordon M, Urquhart B, Boachie-Adjei O (2001) The efficacy of antifibrinolytics in the reduction of blood loss during complex adult reconstructive spine surgery. *Spine* 10:1152–1157
111. Vaquero J et al (1993) Effect of fibrin glue on postlaminectomy scar formation. *Acta Neurochir (Wien)* 120: 159–163
112. Wada E, Yonenobu K, Ebara S, Kuwahara O, Ono K (1993) Epidural migration of hemostatic agents as a cause of postthoracotomy paraplegia. Report of two cases. *J Neurosurg* 78: 658–660
113. Walker R (1987) Special report: transfusion risks. *Am J Clin Pathol* 88:374–378
114. Widman J, Hammarqvist F, Sellden E (2002) Amino acid infusion induces thermogenesis and reduces blood loss during hip arthroplasty under spinal anesthesia. *Anesth Analg* 95:1757–1762
115. Winkler M, Akca O, Birkenberg B, Hetz H, Scheck T, Arkilic CF, Kabon B, Marker E, Grubl A, Czepan R, Greher M, Goll V, Gottsauner-Wolf F, Kurz A, Sessler DI (2000) Aggressive warming reduces blood loss during hip arthroplasty. *Anesth Analg* 91:978–984