

MOZ increases p53 acetylation and premature senescence through its complex formation with PML

Susumu Rokudai^a, Oleg Laptenko^a, Suzzette M. Arnal^a, Yoichi Taya^b, Issay Kitabayashi^c, and Carol Prives^{a,1}

^aDepartment of Biological Sciences, Columbia University, New York, NY 10027; ^bCancer Science Institute of Singapore and Department of Biochemistry, Center for Life Sciences #02-07, National University of Singapore, Singapore 117456; and ^cMolecular Oncology Division, National Cancer Center Research Institute, Chuo-ku, Tokyo 104-0045, Japan

Contributed by Carol Prives, January 18, 2013 (sent for review September 18, 2012)

Monocytic leukemia zinc finger (MOZ)/KAT6A is a MOZ, Ybf2/Sas3, Sas2, Tip60 (MYST)-type histone acetyltransferase that functions as a coactivator for acute myeloid leukemia 1 protein (AML1)- and Ets family transcription factor PU.1-dependent transcription. We previously reported that MOZ directly interacts with p53 and is essential for p53-dependent selective regulation of p21 expression. We show here that MOZ is an acetyltransferase of p53 at K120 and K382 and colocalizes with p53 in promyelocytic leukemia (PML) nuclear bodies following cellular stress. The MOZ–PML–p53 interaction enhances MOZ-mediated acetylation of p53, and this ternary complex enhances p53-dependent p21 expression. Moreover, we identified an Akt/protein kinase B recognition sequence in the PML-binding domain of MOZ protein. Akt-mediated phosphorylation of MOZ at T369 has a negative effect on complex formation between PML and MOZ. As a result of PML-mediated suppression of Akt, the increased PML–MOZ interaction enhances p21 expression and induces p53-dependent premature senescence upon forced PML expression. Our research demonstrates that MOZ controls p53 acetylation and transcriptional activity via association with PML.

protein modification | DNA damage

The p53 protein functions as a key regulator of pathways mediating cellular responses by inducing myriad target genes that regulate diverse cellular processes including cell-cycle arrest, apoptosis, and genomic stability (1–4). Regulation of p53 transcriptional activities is crucial for genotoxic stress because of the varieties of cellular responses that are mediated by p53, which, in some cases, can be mutually exclusive (e.g., arrest and apoptosis) (5). p53 has been detected in discrete nuclear speckles, known as promyelocytic leukemia nuclear bodies (PML-NBs), in which CREB binding protein (CBP)/p300, Tip60, and pRB are also found (6–8).

PML was originally identified as a *t*(15, 17) chromosomal translocation partner with the retinoic acid receptor- α (RAR α) in acute promyelocytic leukemia, in which fusion genes encoding the PML–RAR α fusion protein are generated (9–11). The ability of PML to interact with activators such as CBP/p300 in the nuclear body suggests that PML could modulate transcription through its ability to stabilize complexes of cofactors (12, 13). Overexpression of PML, γ -irradiation of cells, or oncogenic signals such as Ras overexpression can recruit p53 into PML-NBs (14, 15). The resulting ternary p53–PML–CBP complex then promotes the acetylation of p53 and harmonically coordinates critical tumor-suppressive functions such as apoptosis, senescence, and growth arrest (16, 17). Unlike bona fide coactivators such as CBP and p300, PML does not possess intrinsic histone acetylase activity. PML-mediated recruitment of both coactivators promotes p53 modification, such as acetylation and phosphorylation (16, 17). Acetylation levels of p53 are known to be significantly enhanced in response to stress and are involved in p53 activation and stabilization (18, 19). Given that p300^{-/-} mouse embryonic fibroblasts (MEFs) retain the ability to respond to UV irradiation by stabilization of p53 and induction of p21, it is possible that other cofactors may compensate for p300 function in p300^{-/-} MEFs (20).

Recent studies have shown PML to be a component of the PI3K-signaling network, negatively regulating the nuclear content

of phosphorylated Akt/protein kinase B via its dephosphorylation by protein phosphatase 2A (PP2A) (21). The PI3K–Akt pathway has also been well documented to delay the p53-mediated response (22). The central role of Akt in this process is illustrated by the physical association of Akt with murine double minute 2 (MDM2), which inhibits p21 expression through its phosphorylation and activation of MDM2 and subsequent MDM2-mediated ubiquitination of p53 (23, 24). Recent studies suggest that Mdm2 mediates transcriptional repression by forming a protein complex with p53 on the promoter of specific p53-responsive genes (25), whereas accumulating observations suggest that degradation-independent mechanisms are also crucial for MDM2 in controlling p53 activities (26). Regardless, it is clear that the molecular mechanisms by which p53 activity is controlled are complex and how such mechanisms function in opposition to the Akt-signaling network remain to be elucidated.

Monocytic leukemia zinc finger (MOZ) protein is a MYST [MOZ, Ybf2/Something about silencing protein 3 (Sas3), Something about silencing protein 2 (Sas2), Tip60/KAT5]-type histone acetyltransferase (HAT) that functions as a coactivator for AML1- and p53-dependent transcription (27–29). MOZ^{-/-} mice die at around embryonic day 15 (30). In serially passaged MOZ^{-/-} MEFs, DNA damage-induced expression of p21 and ensuing cell-cycle arrest is profoundly impaired, suggesting that the p53 pathway may be altered in MOZ^{-/-} MEFs (27). MOZ is involved in leukemia-associated chromosome rearrangements such as *t*(8, 16)(p11;p13) (31), *t*(8, 22) (32), and *inv*(8) (33), which result in fusion to the transcription coactivators CBP, p300, and transcriptional intermediary factor 2 (TIF2), respectively. Recently, we reported that MOZ directly interacts with p53 through its modification and selective activation of p21 but not Bcl-2 associated x protein (Bax) (27). Although MOZ stimulates p53-mediated transcription of the p21 gene selectively to induce cell-cycle arrest, neither the precise mechanisms of p53 transcriptional activation by MOZ nor the interactions between MOZ and p53 are fully understood.

In this report, we demonstrate that MOZ interacts with PML and recruits it into PML-NBs. A ternary p53–PML–MOZ interaction promotes the acetylation of p53 at K120 and K382 and subsequent p53-induced transcription of the p21 gene. Moreover, the MOZ amino acid sequence contains an Akt substrate motif, and phosphorylation of MOZ by Akt at T369 has a negative effect on its complex formation with PML. PML–MOZ complex formation, which is increased through PML-mediated Akt kinase activity suppression, induces acetylation of p53 and, consequently, p21 expression and p53-dependent premature senescence. These findings reveal a second mechanism that is distinct from Akt phosphorylation and activation of Mdm2, by which the Akt pathway counteracts p53 and prevents it from activating its downstream target genes.

Author contributions: S.R. and C.P. designed research; S.R. and S.M.A. performed research; O.L., Y.T., and I.K. contributed new reagents/analytic tools; S.R. and C.P. analyzed data; and S.R. and C.P. wrote the paper.

The authors declare no conflict of interest.

¹To whom correspondence should be addressed. E-mail: clp3@columbia.edu.

This article contains supporting information online at www.pnas.org/lookup/suppl/doi:10.1073/pnas.1300490110/-DCSupplemental.

Results

MOZ Interacts with PML in PML-NBs. PML is critical for the proper activation of p53 transcriptional activity (15, 16). Upon exposure to DNA damage (e.g., Ras activation, UV radiation), p53 becomes associated with PML-NBs. To investigate whether MOZ colocalizes with PML in PML-NBs, we exposed MCF-7 (breast cancer cell line) cells to UV radiation. In untreated cells, MOZ exhibited both speckled and diffuse nuclear staining in both the nucleoplasm and the nucleolus. When cells were exposed to UV radiation, both PML and MOZ showed a high degree of colocalization in PML-NBs (Fig. 1A and Fig. S1A). Coimmunoprecipitation analysis confirmed that both PML–MOZ and p53–MOZ interactions were increased after UV radiation, but not after treatment with As₂O₃ (Fig. 1B and Fig. S1B and C). These results suggest that UV radiation enhances p53–PML–MOZ ternary complex formation.

To identify the PML-interacting domain in MOZ, the interactions between various deletion mutants of MOZ and PML were examined by immunoprecipitation (IP) immunoblotting. Coimmunoprecipitation experiments using these deletion mutants identified two PML-interacting regions (amino acids spanning 144–664 and 1517–1741) in MOZ (Fig. 1C and D and Fig. S2A). To determine the region of PML required for binding to MOZ, a pull-down assay using anti-FLAG epitope peptide antibody (see *SI Materials and Methods*) was performed using a series of PML deletion mutants. Results from these studies suggest that PML possesses one MOZ-interacting coiled-coil domain within amino acids 229–360 (Fig. 1E and Fig. S2B).

To confirm that the interaction between PML and MOZ is direct, we performed a Far-Western blot analysis. In brief, full-length PML protein was translated in reticulocyte lysates, and then proteins were separated by SDS/PAGE, followed by transfer to nitrocellulose. Following denaturation and renaturation, proteins on the membrane were incubated with sf9 insect cells-purified MOZ proteins, and bound MOZ was detected by immunoblotting analysis. As shown in Fig. 1F, PML directly bound to MOZ in vitro.

As negative controls, BSA and the polypeptides present in reticulocyte extracts did not interact with MOZ. Therefore, we conclude that PML is a bona fide interacting partner for MOZ.

PML-MOZ Association Enhances MOZ-Mediated p53 Acetylation and p21 Expression. PML directly interacts with HATs, including CBP, p300, and Tip60, and PML-NBs serve as sites for posttranslational modifications of p53 (17, 34). MOZ is a MYST-type HAT, like Tip60 and MOF (35, 36), and the MYST domain of MOZ has 69% homology with that of Tip60. We therefore hypothesized that MOZ may play an important role in acetylation of p53.

In fact, an in vitro acetylation assay showed that MOZ could acetylate p53 at K120 and K382 (Fig. 2A and Fig. S3). Furthermore, knockdown of MOZ expression partially inhibited irradiation-dependent acetylation of p53 at K120 and K382, but not at K373 (Fig. 2B). Therefore, MOZ displays aspects of both Tip60 and p300 in acetylation of p53. To further confirm that MOZ acetylates p53 at K120 and K382, we generated an acetylase-deficient MOZ variant (AD-MOZ: Q654E/G657E), in which the acetyl-CoA-binding sequences are mutated to glutamine, mimicking the MYST domain of a previously characterized acetylase-deficient mutant of Tip60 (37) (Fig. 2C). Immunoblotting showed that wild-type (WT) MOZ, but not AD-MOZ, could acetylate itself and p53 at K382 (Fig. S4A and B). Because we observed that MOZ is recruited into PML-NBs like CBP/p300, we wanted to determine whether the PML–MOZ interaction could also be enhanced by MOZ-mediated acetylation of p53. To this end, we performed immunoblotting with antibodies that specifically recognize acetylated-p53 at K120 and K382 and showed that PML promotes MOZ-mediated p53 acetylation (Fig. 2D).

To test whether PML also affects transcriptional activation of the p53–MOZ complex, we used reporter plasmids under the control of promoters for p53 target genes such as p21 and bax in MOZ^{-/-} MEFs. Whereas PML stimulated p53-mediated activation of p21-luc in wild-type MEFs, p21-luc induction was




Fig. 1. MOZ is recruited into promyelocytic leukemia nuclear bodies after DNA damage. (A) Colocalization of MOZ with PML. MCF-7 cells were irradiated with UV (30 J/m²) and then processed for immunofluorescence. (B) MOZ interacts with PML and p53. MCF7 cells were cotransfected with HA-epitope tagged PML and FLAG-epitope tagged MOZ and then subjected to UV irradiation (30 J/m²) followed by harvesting at indicated times. The cell lysates were immunoprecipitated with anti-FLAG beads and subjected to immunoblotting. (C) Schematic representation of the structure of MOZ deletion mutants. The plant homeo domain (PHD) finger domain (P), the basic domain (B), the HAT domain, the serine-rich region (S), the methionine-rich region (M), and the regions required for interaction with PML (bars) and for strong interaction with PML (dotted bar) are indicated. (D) BOSC23 cells were transfected with HA-MOZ deletion mutants along with FLAG-PML. Cell lysates (CL) were immunoprecipitated with anti-FLAG beads and were subjected to immunoblotting. (E) Schematic representation of the structure of PML deletion mutants. The region required for interaction with MOZ is indicated (bar). (F) PML can interact with MOZ in vitro as indicated by Far-Western analysis. FLAG-MOZ protein purified from sf9 (Right) and HAPML immunoprecipitated from reticulocyte lysate (Left) were subjected to SDS/PAGE and stained by Coomassie blue. Mock (T7) and PML (T7-PML) proteins were resolved and transferred to a nitrocellulose membrane, which was then incubated with 1 μg/mL of purified FLAG-MOZ (Center) and detected by immunoblotting.




Fig. 2. MOZ acetylates p53 at K382 and K120. (A) In vitro acetylation of p53 by MOZ, Tip60, and p300. Baculovirus-expressed p53, MOZ, Tip60, and p300 were assayed for acetylation activity of p53. The acetylation samples were detected by immunoblotting with antibodies against Acp53, AcK382, AcK120, and p53 (Upper). MOZ, Tip60, and p300 proteins were visualized by silver staining (Lower). (B) Inhibition of MOZ blocks p53 acetylation. The siRNA-treated MCF-7 cells were irradiated with 30 J/m² of UV for indicated times and were subjected to immunoblotting. (C) MYST domain sequences of MOZ and Tip60. (D) PML enhances MOZ acetylation of p53. BOSC23 cells were transfected with indicated FLAG-p53 along with either HA-MOZ or PML and were subjected to immunoblotting. (E) PML enhances MOZ-mediated p21-luc but not Bax-luc reporter activity. Either wild-type (WT) or MOZ^{-/-} MEFs were transfected with p21-luc (Left) or Bax-luc (Right), p53, PML, and pRL-CMV as a transfection control. (F) PML enhances MOZ-mediated p21 expression in a p53-dependent manner. Either HCT116 p53^{+/+} or p53^{-/-} cells were cotransfected with FLAG-MOZ along with HA-PML and were subjected to immunoblotting.

suppressed in MOZ^{-/-} MEFs. On the other hand, induction of bax-luc was slightly induced in MOZ^{-/-} MEFs compared with wild-type MEFs (Fig. 2E). This supports the likelihood that MOZ is a key component of the regulation of p53 by PML. Although further luciferase and immunoblotting demonstrated that wild-type MOZ strongly enhanced p21 gene transcription and p21 expression in the presence of PML, AD-MOZ partially suppressed the expression of p21 compared with wild-type MOZ, suggesting that MOZ-mediated acetylation of p53 is important but not sufficient for MOZ-mediated expression of p21 (Fig. 2F and Fig. S5). A MOZ-CBP fusion gene, produced by the t(8, 16) translocation, is associated with acute monocytic leukemia. We tested the effects of the MOZ-CBP chimera and found that MOZ-CBP suppressed p53-dependent transcriptional activation of p21 (Fig. S5). These results indicate that MOZ associates with PML and that the PML-MOZ complex induces p21 expression.

Akt Phosphorylates MOZ and Inhibits Its Interaction with PML. Analysis of the human MOZ sequence revealed one site (T369) that conforms to the Akt phosphorylation consensus site (Fig. 3A).

The Akt phosphorylation motif (R-X-R-X-X-S/T-B, where “X” represents any amino acid and “B” represents a hydrophobic residue), has been refined to include amino acids that contribute to its 3D structure (38). We found that the Akt phosphorylation motif is also conserved in human-, mouse-, rat-MOZ, and RXXRXT in human MORF/MYST4, suggesting its possible functional importance. The presence of putative Akt phosphorylation sites in MOZ led us to determine whether MOZ is a substrate for Akt.

In fact, we observed that recombinant Akt was sufficient for inducing phosphorylation of wild-type MOZ in vitro, but not T369A-MOZ (Fig. 3B). We also confirmed that wild-type Akt, but not dominant negative (DN) Akt, was capable of inducing MOZ phosphorylation in vitro (Fig. 3C). To further examine whether Akt can phosphorylate MOZ at T369, we generated a polyclonal antibody that specifically recognizes MOZ phosphorylated at that site (pT369-MOZ). By immunoblotting, this antibody detected MOZ that had been phosphorylated at T369 by myristoylated active Akt (Myr) while it partially cross-reacted with nonphosphorylated MOZ in a manner similar to a pAkt-substrate antibody that recognized phosphorylated MOZ (Fig. 3D). These results support the possibility that Akt mediates MOZ phosphorylation at T369 in vivo and in vitro.

Because T369 of MOZ is located within the PML-binding domain, we further tested whether this phosphorylation affects binding between MOZ and PML or p53 (Fig. 1C). Coimmunoprecipitation




Fig. 3. Akt-mediated phosphorylation of MOZ suppresses the interaction between PML and MOZ. (A) Structure of the consensus Akt phosphorylation motif in MOZ. Sequences of human-, mouse-, rat-MOZ, and MORF are shown for comparison. (B) Recombinant Akt phosphorylates MOZ in vitro. FLAG-tagged wild-type (WT) and T369A (TA) of MOZ proteins were immunoprecipitated with anti-FLAG beads from transfected BOSC23 cells. MOZ proteins were incubated with recombinant Akt (Rec. Akt) in the presence of [³²P]ATP. (C) FLAG-tagged WT and dominant negative (DN) Akt and MOZ proteins were immunoprecipitated with anti-FLAG beads from transfected BOSC23 cells. MOZ proteins were incubated with WT- or DN-Akt in the presence of [³²P]ATP. (D) Akt phosphorylates MOZ at T369. BOSC23 cells were cotransfected with FLAG-tagged WT- or TA-MOZ along with myristoylated (Myr)- or DN-Akt. Cell lysates were immunoprecipitated with anti-FLAG beads and were subjected to immunoblotting using antibodies for phospho-T369 (pT369-MOZ), phospho-Akt substrate (pAkt-Sub), FLAG (MOZ), Akt, and β -actin. (E and F) Akt-mediated phosphorylation of MOZ suppresses the PML-MOZ interaction but not the p53-MOZ interaction. BOSC23 cells were cotransfected with FLAG-tagged WT, AD, T369A, and T369D of MOZ along with HA-PML (E) or HA-p53 (F). Cell lysates were immunoprecipitated with anti-FLAG beads and subjected to immunoblotting. Ten percent of input lysates were used for input control.

analysis using phosphorylation-deficient MOZ (T369A) demonstrated that it had increased interaction with PML (Fig. 3E). Interestingly, these mutations did not affect MOZ-p53 binding (Fig. 3F). These results indicate that Akt phosphorylates T369 of MOZ to inhibit its interaction with PML.

Phosphorylation of MOZ at T369 Is Important for Negative Regulation of p53 Acetylation. Because the interaction between PML and MOZ is likely dependent on the phosphorylation state of MOZ at T369, we determined whether Akt-mediated phosphorylation of MOZ inhibits its acetylation activity. An IP immunoblotting showed that T369A-MOZ strongly acetylates p53 at K382 and K120 in transfected cells, unlike the phosphorylation-mimicking mutant T369D-MOZ (Fig. 4A). To examine the T369A-MOZ acetylation activity per se, we next performed an *in vitro* HAT activity assay using histone peptides as substrates. In contrast to Fig. 4A, T369A-MOZ showed only slightly increased acetylation of histone H3 and H4 peptides compared with T369D-MOZ (Fig. 4B). These results are consistent with results showing that wild-type or T369 mutant forms of MOZ displayed equivalent abilities to self-acetylate and that the acetylation activity of MOZ was enhanced in the presence of PML (Fig. 4C). Thus, our data indicate that phosphorylation of MOZ at T369 is important for

the negative regulation of acetylation of p53 in the presence of PML, whereas such phosphorylation is not important for its acetylation activity per se.

Further luciferase analysis and real-time PCR analysis upon coexpression of MOZ with PML revealed that T369A-MOZ strongly enhanced the expression of p21, whereas T369D-MOZ partially suppressed the expression of p21 compared with T369A-MOZ (Fig. 4D and E). These data are consistent with p21 expression analyzed by immunoblotting (Fig. 4F). Taken together, these findings indicate that suppression of Akt-mediated phosphorylation at T369 of MOZ is critical for MOZ recruitment into PML-NBs and its acetylation of p53. We conclude that the PI3K-Akt pathway-mediated suppression of p21 is not only through Akt phosphorylation and stabilization of MDM2, but also through blocking the recruitment of MOZ into PML-NBs and the acetylation of p53.

Loss of MOZ Contributes to Resistance to PML-Induced Senescence.

To characterize PML expression-induced cell-cycle arrest and senescence, we infected wild-type and MOZ^{-/-} primary mouse embryonic fibroblasts (pMEFs) with a pLNCX retroviral vector expressing PML and Ha-Ras Ha-Ras (v-Ha-ras Harvey rat sarcoma viral oncogene homolog, V12), cultured them for 3 d under puromycin selection, and counted cells for the next 5 d. Wild-type pMEFs expressing PML as well as oncogenic Ha-Ras (V12) ceased to proliferate at subconfluent densities (Fig. 5A). Conversely, in MOZ^{-/-} pMEFs, ectopic PML expression did not lead to cessation of proliferation, although RasV12-expressing MOZ^{-/-} MEFs exhibited slowed proliferation (Fig. 5B). The morphology of MOZ^{-/-} pMEFs differed from senescent wild-type pMEFs, even after infection of these cells with retroviruses containing PML or Ras (Fig. S6).

We next examined the cell-cycle distribution of pMEFs arrested by retroviral-mediated PML expression by following BrdU incorporation. Wild-type pMEFs did not incorporate BrdU (Fig. 5C and D) and became positive for senescence-associated β -galactosidase (SA β -gal) staining, a marker of cellular senescence (Fig. 5E and Fig. S7). Consistent with Fig. 5B, forced PML expression in MOZ^{-/-} pMEFs did not alter BrdU incorporation (Fig. 5C and D). The frequency of SA β -gal-positive cells was markedly lower in PML-expressing MOZ^{-/-} pMEF cells than in wild-type pMEFs (Fig. 5E and Fig. S7). These results indicate that loss of MOZ contributes to resistance to PML-induced premature senescence.

Based on the above findings, we examined p53 modifications and p21 expression in WT and MOZ^{-/-} MEFs expressing either RasV12 or PML. Although both p53 and p16 are vital components of the intracellular senescence pathway (39), PML does not affect p16 expression (17), consistent with our observed results (Fig. 5F). Although ectopically expressed Ras or PML induced equivalent levels of p53 in both wild-type and MOZ^{-/-} MEFs, their expression led to the appearance of more nonphosphorylated MOZ in wild-type cells, which corresponded to increased acetylation of p53 at K117 and K379 (equivalent to human p53 K120 and K382, respectively). In line with this, p21 expression was induced in wild-type pMEFs, but not in MOZ^{-/-} pMEFs after PML expression despite similar p53 transcriptional levels in wild-type and MOZ^{-/-} pMEFs (Fig. 5F and Fig. S8). This suggests that, through PML-mediated suppression of the Akt pathway, increased PML-MOZ complex formation enhances p21 expression, which mediates p53-dependent premature senescence upon PML overexpression. In short, MOZ is crucial for p53-mediated p21 expression induced by PML expression and the ensuing PML-induced senescence.

Discussion

Several factors, such as Tip60, hematopoietic zinc finger (Hzf), and human cellular apoptosis susceptibility protein (hCAS/CSE1L) are involved in the selection of p53 target genes (35, 40, 41). However, the molecular mechanisms by which p53 "chooses" cell-cycle arrest or apoptosis are not fully understood. Here we discovered that the MYST-type HAT MOZ interacts with PML and is recruited into PML-NBs. Significantly, we confirmed that the




Fig. 4. Phosphorylation of MOZ is important for negative regulation of p53 acetylation. (A) BOSC23 cells were cotransfected with HA-tagged WT, AD, T369A or T369D along with FLAG-p53. Cell lysates were purified with anti-FLAG beads and were subjected to immunoblotting using antibodies for AcK382, AcK120, FLAG (f-p53), HA (MOZ), and β -actin. (B) Phosphorylation of MOZ has no effect on HAT activity. FLAG-tagged WT, AD, T369A, and T369D MOZ were purified with anti-FLAG beads from transfected BOSC23 cells. The acetylase activities of mutant MOZ were measured as described in *Materials and Methods*. (C) BOSC23 cells were cotransfected with FLAG-tagged WT, AD, T369A, or T369D of MOZ with HA-PML. Cell lysates were immunoprecipitated with anti-FLAG beads and subjected to immunoblotting. (D) The PML-MOZ interaction stimulates p53-mediated p21 reporter activity. H1299 cells were cotransfected with p21-luc, p53, and PML along with WT, AD, T369A, or T369D of MOZ. (E and F) The PML-MOZ interaction stimulates p53-dependent p21 induction and expression. H1299 cells were cotransfected with p53 and PML along with HA-tagged WT, AD, T369A, and T369D of MOZ. The harvested cells were subjected to real-time PCR (E) and to immunoblotting.




Fig. 5. MOZ-knockout cells display resistance to PML-induced senescence. (A and B) Growth curves of wild-type (A) and MOZ^{-/-} (B) pMEFs infected with vector (Mock), RasV12, or PML. Day 0 is the first day after puromycin selection. Error bars represent the standard deviation for three independent littermate experiments. **P* < 0.02, ***P* > 0.03. (C) Flow cytometry analysis of pMEFs. pMEFs in growth phase were stained with an anti-BrdU antibody and propidium iodide at 5 d after selection. (D) Histogram comparing BrdU incorporation of RasV12- and PML-infected pMEFs at 1 or 5 d after selection. (E) SA β -gal staining of WT or MOZ^{-/-} pMEFs infected with pLNCX (Mock), RasV12, or PML at 5 d after selection. (Scale bars: 50 μ m.) (F) MOZ^{-/-} cells are resistant to PML-induced p21 expression. The infected WT or MOZ^{-/-} pMEFs were harvested at 5 d after selection. The cells were partially subjected to immunoprecipitation with anti-p53 (IP: p53) and anti-MOZ (IP: MOZ) and were subjected to immunoblotting using indicated antibodies.

PML–MOZ interaction promotes the MOZ-mediated acetylation of p53 at K120 and K382, which makes MOZ distinct from other KATs that can target only one of these residues and p53-mediated expression of p21 (Fig. S9).

It has been reported that p53 can be acetylated at K373, K382, K164 (catalyzed by p300/CBP), K320 (PCAF/KAT2B), and K120 (Tip60) (42, 43). Although p53 mice engineered to lack acetylatable lysines within their C termini are viable and phenotypically normal, p53-mediated transcriptional activation upon DNA damage is partially impaired in the ES cells and thymocytes of these mice (44). In addition, the p53 7KR mutation, where the seven C-terminal lysines were changed to arginine, significantly contributes to hematopoietic stem cell homeostasis and mouse radiosensitivity (45). Thus, it is possible that other cofactors or additional acetylation sites of p53 may compensate for the loss of p53 acetylation at the C terminus. In this report, we found that MOZ is an acetylation regulator of p53. In contrast to normal MOZ and CBP, the leukemia-associated MOZ–CBP fusion protein inhibited p53-mediated transcription (Fig. S5). These results suggest that inhibition of MOZ–p53-mediated transcription might be involved in pathogenesis of tumors and leukemia.

Our previous study revealed that MOZ-deficient MEFs exhibit impaired p21 expression and fail to arrest in G1 phase in response to DNA damage (27). Evidence of the significance of lysine modifications in the functions of the DNA-binding region comes from studies showing that Tip60/KAT5 and hMOF/KAT8, members of the MYST family of acetyltransferases, acetylate K120 of p53 followed by its accumulation on Puma and Bax promoters (35) and that K120/K164R double-mutant mice display reduced induction of *p21* and *puma* (43). In addition, K120 acetylation of p53

exhibited specific DNA binding and discriminated among response elements at effective physiological salt concentration (46). The ability of MOZ to simultaneously acetylate both K120 and K382 might be responsible for the difference of target genes from previous reports. It might be also possible that MOZ acetylation of p53 occurs in a different subcellular compartment, such as PML–NBs, from that of Tip60/hMOF. Thus, the DNA-binding domain acetylation by MOZ appears to play a role in selective gene regulation.

Akt kinase activity is frequently elevated in several high-grade, late-stage cancers (47), and a somatic constitutively active Akt mutant has been identified in human breast, colorectal, and ovarian cancers (48). In fact, the role of Akt in the p53-mediated response was shown to involve Akt suppression of p21 expression through phosphorylation and activation of MDM2 and subsequent MDM2-mediated ubiquitination of p53 (23, 24, 49). Here we demonstrate that Akt works by a second process in which Akt phosphorylation of MOZ at T369 within its PML interaction region blocks its complex formation with PML (Fig. 3A). A mutant MOZ (T369A) displays enhanced acetylation of p53 at K120 and K382 with ensuing increased expression of p21, indicating that phosphorylation at T369 of MOZ is important for the acetylation of p53, but not for its acetylation activity per se (Fig. 4A and C). Suppressing Akt-mediated phosphorylation of MOZ at T369 is likely very important for the recruitment of MOZ into PML–NBs and the subsequent acetylation of p53.

In this report, we demonstrate that forced PML expression in MOZ^{-/-} primary MEFs somewhat suppresses either G1 arrest or senescence (Fig. 5B). PML expression in MOZ^{-/-} MEFs only slightly alters BrdU incorporation (Fig. 5C and D), but the frequency of SA β -gal-positive senescent cells is markedly lower (Fig. 5E and Fig. S7). These results indicate that loss of MOZ contrib-

utes to resistance to PML-induced G1 arrest and premature senescence. In contrast, MOZ^{-/-} primary MEFs exhibit a high proportion of cells in S phase but lower BrdU incorporation after forced PML expression. This is also true for serially passaged MEFs after DNA damage, suggesting that MOZ might be involved in DNA replication. These results are consistent with a report that depletion of inhibitor of growth family, member 5 (ING5), which comprises a subunit of the ING5-MOZ stoichiometric HAT complex, renders cells unable to complete S phase, allowing only a few cells to proceed to the G2/M phase. Together, these results suggest that ING5-MOZ complexes are essential for DNA replication, not only for initiation but also for replication fork movement (50). Recently, it has been reported that Pten-loss-induced cellular senescence, where the activation of the PI3K-Akt pathway is the key mechanism triggering a p53-dependent senescence, represents a senescence response that is distinct from oncogene-induced senescence (51). Further studies are required to investigate whether the sequential phenomenon—down-regulation of Akt activity and MOZ acetylation of p53—occurs in the PML-NB subcellular compartment.

Materials and Methods

Plasmids expressing FLAG-tagged and HA-tagged human MOZ, CBP, MOZ-CBP, and p300 were cloned into the pLNCX vector (Clontech). Human cDNA encoding FLAG- and HA-tagged PML isoform IV and HA-tagged V12 Ras were cloned into pLPCX retrovirus vectors. Myristoylated active, FLAG-tagged, wild-type, or dominant negative Akt1 were described previously (52). Details of other methods used in this article, including cell culture, antisense oligonucleotides, baculovirus protein expression and purification, flow cytometry, SA β -gal assay, immunofluorescence, immunoprecipitation, immunoblotting, Far-Western blotting, real-time PCR, luciferase assay, in vitro acetylation assay, and in vitro kinase assay are described in *SI Materials and Methods*.

ACKNOWLEDGMENTS. Baculovirus expressing His₆-tagged p300 was a generous gift from Dr. W. L. Kraus. We thank Ella Freulich for expert technical assistance. This work was supported in part by Grants-in-Aid for Scientific Research from the Ministry of Health, Labor, and Welfare; the Ministry of Education, Culture, Sports, Science, and Technology (to I.K. and S.R.); The Mochida Memorial Foundation for Medical and Pharmaceutical Research (to S.R.); and by National Institutes of Health Grant CA77742 (to C.P.).

- Vogelstein B, Lane D, Levine AJ (2000) Surfing the p53 network. *Nature* 408(6810):307–310.
- Vousden KH, Lu X (2002) Live or let die: The cell's response to p53. *Nat Rev Cancer* 2(8):594–604.
- Laptenko O, Prives C (2006) Transcriptional regulation by p53: One protein, many possibilities. *Cell Death Differ* 13(6):951–961.
- Harris SL, Levine AJ (2005) The p53 pathway: Positive and negative feedback loops. *Oncogene* 24(17):2899–2908.
- Vousden KH, Prives C (2009) Blinded by the light: The growing complexity of p53. *Cell* 137(3):413–431.
- Alcalay M, et al. (1998) The promyelocytic leukemia gene product (PML) forms stable complexes with the retinoblastoma protein. *Mol Cell Biol* 18(2):1084–1093.
- Vallian S, et al. (1998) Modulation of Fos-mediated AP-1 transcription by the promyelocytic leukemia protein. *Oncogene* 16(22):2843–2853.
- Zhang Y, Xiong Y (1999) Mutations in human ARF exon 2 disrupt its nucleolar localization and impair its ability to block nuclear export of MDM2 and p53. *Mol Cell* 3(5):579–591.
- de Thé H, et al. (1991) The PML-RAR alpha fusion mRNA generated by the t(15;17) translocation in acute promyelocytic leukemia encodes a functionally altered RAR. *Cell* 66(4):675–684.
- Pandolfi PP, et al. (1991) Structure and origin of the acute promyelocytic leukemia myl/RAR alpha cDNA and characterization of its retinoid-binding and transactivation properties. *Oncogene* 6(7):1285–1292.
- Piazza F, Gurrieri C, Pandolfi PP (2001) The theory of APL. *Oncogene* 20(49):7216–7222.
- LaMorte VJ, Dyck JA, Ochs RL, Evans RM (1998) Localization of nascent RNA and CREB binding protein with the PML-containing nuclear body. *Proc Natl Acad Sci USA* 95(9):4991–4996.
- Zhong S, et al. (2000) Role of SUMO-1-modified PML in nuclear body formation. *Blood* 95(9):2748–2752.
- Ferbeyre G, et al. (2000) PML is induced by oncogenic ras and promotes premature senescence. *Genes Dev* 14(16):2015–2027.
- Fogal V, et al. (2000) Regulation of p53 activity in nuclear bodies by a specific PML isoform. *EMBO J* 19(22):6185–6195.
- Guo A, et al. (2000) The function of PML in p53-dependent apoptosis. *Nat Cell Biol* 2(10):730–736.
- Pearson M, et al. (2000) PML regulates p53 acetylation and premature senescence induced by oncogenic Ras. *Nature* 406(6792):207–210.
- Barlev NA, et al. (2001) Acetylation of p53 activates transcription through recruitment of coactivators/histone acetyltransferases. *Mol Cell* 8(6):1243–1254.
- Ito A, et al. (2001) p300/CBP-mediated p53 acetylation is commonly induced by p53-activating agents and inhibited by MDM2. *EMBO J* 20(6):1331–1340.
- Yao TP, et al. (1998) Gene dosage-dependent embryonic development and proliferation defects in mice lacking the transcriptional integrator p300. *Cell* 93(3):361–372.
- Trotman LC, et al. (2006) Identification of a tumour suppressor network opposing nuclear Akt function. *Nature* 441(7092):523–527.
- Sabbatini P, McCormick F (1999) Phosphoinositide 3-OH kinase (PI3K) and PKB/Akt delay the onset of p53-mediated, transcriptionally dependent apoptosis. *J Biol Chem* 274(34):24263–24269.
- Mayo LD, Donner DB (2001) A phosphatidylinositol 3-kinase/Akt pathway promotes translocation of Mdm2 from the cytoplasm to the nucleus. *Proc Natl Acad Sci USA* 98(20):11598–11603.
- Zhou BP, et al. (2001) HER-2/neu induces p53 ubiquitination via Akt-mediated MDM2 phosphorylation. *Nat Cell Biol* 3(11):973–982.
- Minsky N, Oren M (2004) The RING domain of Mdm2 mediates histone ubiquitylation and transcriptional repression. *Mol Cell* 16(4):631–639.
- Pant V, Xiong S, Iwakuma T, Quintás-Cardama A, Lozano G (2011) Heterodimerization of Mdm2 and Mdm4 is critical for regulating p53 activity during embryogenesis but dispensable for p53 and Mdm2 stability. *Proc Natl Acad Sci USA* 108(29):11995–12000.
- Rokudai S, et al. (2009) Monocytic leukemia zinc finger (MOZ) interacts with p53 to induce p21 expression and cell-cycle arrest. *J Biol Chem* 284(1):237–244.
- Champagne N, Pelletier N, Yang XJ (2001) The monocytic leukemia zinc finger protein MOZ is a histone acetyltransferase. *Oncogene* 20(3):404–409.
- Kitabayashi I, Aikawa Y, Nguyen LA, Yokoyama A, Ohki M (2001) Activation of AML1-mediated transcription by MOZ and inhibition by the MOZ-CBP fusion protein. *EMBO J* 20(24):7184–7196.
- Katsumoto T, et al. (2006) MOZ is essential for maintenance of hematopoietic stem cells. *Genes Dev* 20(10):1321–1330.
- Borrow J, et al. (1996) The translocation t(8;16)(p11;p13) of acute myeloid leukaemia fuses a putative acetyltransferase to the CREB-binding protein. *Nat Genet* 14(1):33–41.
- Chaffanet M, et al. (2000) MOZ is fused to p300 in an acute monocytic leukemia with t(8;22). *Genes Chromosomes Cancer* 28(2):138–144.
- Carapeti M, Aguiar RC, Goldman JM, Cross NC (1998) A novel fusion between MOZ and the nuclear receptor coactivator TIF2 in acute myeloid leukemia. *Blood* 91(9):3127–3133.
- Wu Q, et al. (2009) PML3 orchestrates the nuclear dynamics and function of TIP60. *J Biol Chem* 284(13):8747–8759.
- Sykes SM, et al. (2006) Acetylation of the p53 DNA-binding domain regulates apoptosis induction. *Mol Cell* 24(6):841–851.
- Tang Y, Luo J, Zhang W, Gu W (2006) Tip60-dependent acetylation of p53 modulates the decision between cell-cycle arrest and apoptosis. *Mol Cell* 24(6):827–839.
- Ikura T, et al. (2000) Involvement of the TIP60 histone acetylase complex in DNA repair and apoptosis. *Cell* 102(4):463–473.
- Alessi DR, Caudwell FB, Andjelkovic M, Hemmings BA, Cohen P (1996) Molecular basis for the substrate specificity of protein kinase B: Comparison with MAPKAP kinase-1 and p70 S6 kinase. *FEBS Lett* 399(3):333–338.
- Lin AW, et al. (1998) Premature senescence involving p53 and p16 is activated in response to constitutive MEK/MAPK mitogenic signaling. *Genes Dev* 12(19):3008–3019.
- Das S, et al. (2007) Hzf determines cell survival upon genotoxic stress by modulating p53 transactivation. *Cell* 130(4):624–637.
- Tanaka T, Ohkubo S, Tatsuno I, Prives C (2007) hCAS/CSE1L associates with chromatin and regulates expression of select p53 target genes. *Cell* 130(4):638–650.
- Gu W, Roeder RG (1997) Activation of p53 sequence-specific DNA binding by acetylation of the p53 C-terminal domain. *Cell* 90(4):595–606.
- Tang Y, Zhao WH, Chen Y, Zhao Y, Gu W (2008) Acetylation is indispensable for p53 activation. *Cell* 133(4):612–626.
- Krummel KA, Lee CJ, Toledo F, Wahl GM (2005) The C-terminal lysines fine-tune P53 stress responses in a mouse model but are not required for stability control or transactivation. *Proc Natl Acad Sci USA* 102(29):10188–10193.
- Wang YV, et al. (2011) Fine-tuning p53 activity through C-terminal modification significantly contributes to HSC homeostasis and mouse radiosensitivity. *Genes Dev* 25(13):1426–1438.
- Arbely E, et al. (2011) Acetylation of lysine 120 of p53 endows DNA-binding specificity at effective physiological salt concentration. *Proc Natl Acad Sci USA* 108(20):8251–8256.
- Sun M, et al. (2001) AKT1/PKBalpha kinase is frequently elevated in human cancers and its constitutive activation is required for oncogenic transformation in NIH3T3 cells. *Am J Pathol* 159(2):431–437.
- Carpenter JD, et al. (2007) A transforming mutation in the pleckstrin homology domain of AKT1 in cancer. *Nature* 448(7152):439–444.
- Michael D, Oren M (2002) The p53 and Mdm2 families in cancer. *Curr Opin Genet Dev* 12(1):53–59.
- Doyon Y, et al. (2006) ING tumor suppressor proteins are critical regulators of chromatin acetylation required for genome expression and perpetuation. *Mol Cell* 21(1):51–64.
- Alimonti A, et al. (2010) A novel type of cellular senescence that can be enhanced in mouse models and human tumor xenografts to suppress prostate tumorigenesis. *J Clin Invest* 120(3):681–693.
- Rokudai S, Fujita N, Kitahara O, Nakamura Y, Tsuruo T (2002) Involvement of FKHR-dependent TRADD expression in chemotherapeutic drug-induced apoptosis. *Mol Cell Biol* 22(24):8695–8708.