

Endophthalmitis caused by *Acinetobacter baumannii*: a case series

R Roy¹, P Panigrahi¹, J Malathi², SS Pal¹, K Nandi¹, A Patil¹, E Nigam¹ and V Arora¹

Abstract

Purpose To profile the etiology, clinical outcomes and drug sensitivity patterns in endophthalmitis caused by *Acinetobacter baumannii*.

Methods Retrospective analysis of all the cases of *Acinetobacter baumannii* endophthalmitis presenting to tertiary referral care ophthalmic hospital in Eastern India from January 2009 to December 2011 were done.

Results A total of four cases were included in the study. Out of the four cases one was post traumatic and the rest were post cataract surgery. All the cases underwent vitreoretinal surgical intervention followed by intravitreal antibiotics. *A. Baumannii* was isolated from vitreous in all the cases. Among all the drugs tested bacteria were found sensitive to ciprofloxacin (100 %) whereas all tested resistant to ceftazidime. Out of the four cases one had to be eviscerated, another developed retinal detachment post vitrectomy, one was phthisical at final followup, and only one patient achieved a vision of 20/200 with clear media and attached retina at final visit.

Conclusion *A. Baumannii* is a very rare cause of endophthalmitis with poor visual and anatomical outcomes. Ciprofloxacin should be considered as first the line intravitreal antibiotic.

Eye (2013) 27, 450–452; doi:10.1038/eye.2012.277; published online 11 January 2013

Keywords: retina; endophthalmitis; *Acinetobacter*

Introduction

Endophthalmitis is a visually devastating condition having poor clinical outcome in majority of the cases. Amongst other factors, the nature of the infective strain has got a profound bearing on the final result. *Acinetobacter baumannii* is a rare cause of endophthalmitis

with only a single report available in literature.¹ Purpose of this study is to report the clinical profile and drug sensitivity patterns of *A. baumannii* endophthalmitis.

Case reports

Case 1

A 60-year-old lady presented with pain, redness, and loss of vision in her right eye 2 days after phacoemulsification. Visual acuity was light perception (PL). Examination revealed corneal abscess with exudates in anterior chamber. Ultrasound revealed dense vitreous echoes with attached retina. She underwent vitrectomy with intraocular lens (IOL) removal and intravitreal antibiotics (vancomycin: 1 mg/0.1 ml, ceftazidime: 2.25 mg/0.1 ml, and dexamethasone: 0.4 mg/0.1 ml). In spite of surgery, her condition worsened and she developed panophthalmitis. Finally she underwent evisceration.

Case 2

A 48-year-old male patient presented with pain, redness, and loss of vision in his left eye 3 days after cataract surgery. At presentation, vision was PL. Examination revealed superior scleral wound dehiscence with iris prolapse. Cornea was oedematous with hypopyon and membrane over the IOL. Ultrasound revealed attached retina with dense vitreous echoes. He underwent wound resuturing, excision of the prolapsed iris, removal of IOL, vitrectomy, and intravitreal antibiotics. He developed retinal detachment at the 7th post-operative day. He was advised resurgery, which he declined.

Case 3

A 74-year-old lady presented with pain and redness in her right eye 15 days after cataract

¹Vitreo Retina Services, Aditya Birla Sankara Nethralaya, Kolkata, West Bengal, India

²Department of Microbiology, L&T Microbiology Research Centre, Sankara Nethralaya, Chennai, India

Correspondence: R Roy, Vitreo Retina Services, Aditya Birla Sankara Nethralaya, 147, E.M.Bypass, Kolkata 700099, West Bengal, India.
 Tel: +91 33 4401 3000;
 Fax: +91 33 4401 3199.
 E-mail: rayrupak@gmail.com

Received: 1 August 2012
 Accepted in revised form: 19 November 2012
 Published online: 11 January 2013

Table 1 Salient clinical details

Case no	Age/gender	Aetiology	Presentation	BCVAI	Intervention	BCVAF
1	60/F	Post cat sx	Fulminant	PL	V + intravit Ab	No PL
2	48/M	Post cat sx	Fulminant	PL	V + intravit Ab	PL
3	74/F	Post cat sx	Acute	PL	V + intravit Ab	20/200
4	7/F	Post traumatic	Acute	PL	L + V + intravit Ab	No PL

Abbreviations: Ab, antibiotics; Acute, >4 days and <6 weeks; BCVAF, final best corrected visual acuity; BCVAI, best corrected visual acuity at presentation; Cat Sx, cataract surgery; F, Female; Fulminant, <4 days, L=lensectomy; M, male; PL, Perception of light; V, vitrectomy.

Table 2 Microbiological details and drug sensitivity patterns

Case no	Aspirate	Gm stain	PCR	Culture	Cefta	Ami	Cipro	Genta	Cepho
1	V	N	Eubac	<i>A. baumannii</i>	R	S	S	S	S
2	V	N	Eubac	<i>A. baumannii</i>	R	S	S	S	R
3	V	N	Eubac	<i>A. baumannii</i>	R	R	S	R	S
4	V	Gm negative bacilli	Eubac	<i>A. baumannii</i>	R	S	S	S	R

Abbreviations: Ami, amikacin; Cefta, ceftazidime; Cepho, cephotaxime; Cipro, ciprofloxacin; Eubac, eubacterial genome positive; Genta, gentamycin; Gm, gram; N, no result; PCR, polymerase chain reaction; R, resistant; S, sensitive; V, vitreous.

surgery. Visual acuity was PL. Cornea was oedematous with hypopyon and exudative membrane over IOL. Ultrasound showed dense vitreous echoes with attached retina. She underwent vitrectomy with IOL removal and intravitreal antibiotics. Intraoperatively, a subretinal abscess was noted at the macular area. In the post-operative period, the patient was on alternate day regimen of intravitreal antibiotics. At final followup, disc was pale with a scar at the macular area. Visual acuity improved to 20/200.

Case 4

A 7-year-old girl presented 19 days after penetrating trauma with needle to her right eye. Examination revealed a sutured corneal tear with exudates in the pupillary area. Ultrasound revealed dense vitreous echoes with attached retina. She underwent lensectomy, vitrectomy, and intravitreal antibiotics. Intraoperatively, retina was found to be ischaemic with a pale disc. At final followup, the eye was phthisical.

Salient details of the cases are summarised in Table 1.

Discussion

Acinetobacter are nonmotile, oxidase negative gram negative bacilli classified under family Moraxcellaceae.² *A. baumannii* is an important member of this group, which causes a variety of human clinical infections involving skin, soft tissue, and bone.³ Recently, it has emerged as an important cause of community-acquired infections. It has

gained notoriety for its ability to upregulate and acquire resistance to antibiotics.² Endophthalmitis caused by *A. baumannii* is rare. English language literature review reveals only a single report of two cases of *A. baumannii* endophthalmitis. To the best of our knowledge, this is the second report on *A. baumannii* endophthalmitis. In the present series, the organism *A. baumannii* was isolated from all cases in more than one media. The bacteria were identified to species level by application of standard biochemical reactions. Antibiotic-disc diffusion test was carried out according to CSLI guidelines. Chen *et al*¹ has reported two cases of *A. baumannii* endophthalmitis. Out of those two cases, one was endogenous and the other was post keratoplasty. Though *A. baumannii* is an important cause of nosocomial infections, none of our cases were endogenous. All our cases had an early presentation, indicating the virulence of the organism. In spite of aggressive therapy, all but one had very poor outcome. Similarly, Chen *et al*¹ reports visual acuity outcomes of 20/60 and no LP in their two cases. Antimicrobial resistance of *Acinetobacter* is of serious concern. Among all the drugs tested, bacteria were found sensitive to ciprofloxacin (100%), whereas all tested resistant to ceftazidime (Table 2). Gopal *et al*⁴ has also reported high level of sensitivity to ciprofloxacin exhibited by *Acinetobacter* species. Chen *et al*¹ has reported sensitivity to imipenem of their isolates, however, due to logistical reasons, we were unable to test sensitivity to imipenem. In non-responding cases and where resistance to ceftazidime is seen, ciprofloxacin can be used as an intravitreal antibiotic. To conclude, *A. baumannii* is a rare cause of endophthalmitis with mostly poor visual outcome.

Summary

What was known before

- *Acinetobacter baumannii* is a rare cause of endophthalmitis.

What this study adds

- Adds to the body of literature of endophthalmitis caused by this organism. Provides drug sensitivity results that can be used for treatment.
-

Conflict of interest

The authors declare no conflict of interest.

Acknowledgements

This research received no specific grant from any funding agency in the public, commercial or not-for-profit sectors.

References

- 1 C Kuan-Jen, H Chiun-Ho, S Ming-Hui, L Chi-Chun, S Chi-Chin, H Ching-Hsi. Endophthalmitis caused by *Acinetobacter baumannii*: report of two cases. *J Clin Microbiol* 2008; **46**(3): 1148–1150.
- 2 Peleg Anton Y, S Harald, LP David. *Acinetobacter baumannii*: emergence of a successful pathogen. *Clin Microbiol Rev* 2008; **21**(3): 538–582.
- 3 Hsueh PR, Teng LJ, Chen CY, Chen WH, Yu CJ, Ho SW *et al*. Pandrug-resistant *Acinetobacter baumannii* causing nosocomial infections in a university hospital. *Taiwan Emerg Infect Dis* 2002; **8**: 827–832.
- 4 Gopal L, Ramaswamy AA, Madhavan HN, Battu RR, Sharma T, Shanmugam MP *et al*. Endophthalmitis caused by *Acinetobacter calcoaceticus*: a profile. *Indian J Ophthalmol* 2003; **51**: 335–340.