

RESEARCH ARTICLE

Open Access

# Carotenoid biosynthesis and overproduction in *Corynebacterium glutamicum*

Sabine A E Heider, Petra Peters-Wendisch and Volker F Wendisch\*

## Abstract

**Background:** *Corynebacterium glutamicum* contains the glycosylated C50 carotenoid decaprenoxanthin as yellow pigment. Starting from isopentenyl pyrophosphate, which is generated in the non-mevalonate pathway, decaprenoxanthin is synthesized via the intermediates farnesyl pyrophosphate, geranylgeranyl pyrophosphate, lycopene and flavuxanthin.

**Results:** Here, we showed that the genes of the carotenoid gene cluster *crtE-cg0722-crtBIY<sub>e</sub>Y<sub>f</sub>Eb* are co-transcribed and characterized defined gene deletion mutants. Gene deletion analysis revealed that *crtI*, *crtEb*, and *crtY<sub>e</sub>Y<sub>f</sub>*, respectively, code for the only phytoene desaturase, lycopene elongase, and carotenoid C45/C50  $\epsilon$ -cyclase, respectively. However, the genome of *C. glutamicum* also encodes a second carotenoid gene cluster comprising *crtB2I2-1/2* shown to be co-transcribed, as well. Ectopic expression of *crtB2* could compensate for the lack of phytoene synthase CrtB in *C. glutamicum*  $\Delta$ *crtB*, thus, *C. glutamicum* possesses two functional phytoene synthases, namely CrtB and CrtB2. Genetic evidence for a *crtI2-1/2* encoded phytoene desaturase could not be obtained since plasmid-borne expression of *crtI2-1/2* did not compensate for the lack of phytoene desaturase CrtI in *C. glutamicum*  $\Delta$ *crtI*. The potential of *C. glutamicum* to overproduce carotenoids was estimated with lycopene as example. Deletion of the gene *crtEb* prevented conversion of lycopene to decaprenoxanthin and entailed accumulation of lycopene to  $0.03 \pm 0.01$  mg/g cell dry weight (CDW). When the genes *crtE*, *crtB* and *crtI* for conversion of geranylgeranyl pyrophosphate to lycopene were overexpressed in *C. glutamicum*  $\Delta$ *crtEb* intensely red-pigmented cells and an 80 fold increased lycopene content of  $2.4 \pm 0.3$  mg/g CDW were obtained.

**Conclusion:** *C. glutamicum* possesses a certain degree of redundancy in the biosynthesis of the C50 carotenoid decaprenoxanthin as it possesses two functional phytoene synthase genes. Already metabolic engineering of only the terminal reactions leading to lycopene resulted in considerable lycopene production indicating that *C. glutamicum* may serve as a potential host for carotenoid production.

**Keywords:** *Corynebacterium*, Carotenoid production, Lycopene production, Phytoene synthase, Phytoene desaturase, Lycopene elongase, Carotenoid epsilon-cyclase, Decaprenoxanthin

## Background

Carotenoids are yellow to red colored pigments originating from the terpenoid biosynthetic pathway. They are very abundant in plants and microorganisms, where they can have diverse functions such as photo protection or light harvesting molecules or as membrane stabilizers [1,2]. In the biosynthetic pathways of certain hormones (like retinoic acid, a hormone regulating the epidermal growth of mammals) they serve as precursors [3]. Carotenoids are also proposed to prevent cancer and reduce

the risk of cardiovascular and Alzheimer disease due to their antioxidative properties [4-6]. Traditionally, terpenoids have been used in the feed, food and nutraceutical industries [1]. As the large-scale chemical synthesis of terpenoids is often difficult and/or costly due to their structural complexity [7] and as their isolation from natural sources usually does not yield sufficient quantities [8], microbial production processes offer a promising alternative.

Carotenoids are derived from the universal precursor isopentenyl pyrophosphate (IPP) and its isomer dimethylallyl pyrophosphate (DMPP) [9]. Enhancing cellular metabolic flux toward IPP and DMAPP is one strategy to

\* Correspondence: volker.wendisch@uni-bielefeld.de  
Chair of Genetics of Prokaryotes, Faculty of Biology & CeBITec, University of Bielefeld, P.O. Box 100131, D-33501 Bielefeld, Germany

improve rates and yield of microbial isoprenoid production [10,11]. There are two independent pathways leading to IPP: the mevalonic acid (MVA) pathway and the methylerythritol phosphate (MEP) pathway. The MVA pathway is found in eukaryotes (mammals, fungi, in the cytoplasm of plant cells), archaea, and a limited number of bacteria. Most bacteria as well as plant plastids synthesize IPP through the MEP pathway [1,12,13]. The MVA pathway requires acetyl-CoA as the primary educt, whereas the MEP pathway starts by condensation of pyruvate and glyceraldehyde 3-phosphate (GAP) [14,15].

*Corynebacterium glutamicum* is used commercially for the annual production of more than 3,000,000 tons of amino acids (Ajinomoto, Food Products Business. Available from World Wide Web: <http://www.ajinomoto.com/ir/pdf/Food-Oct2010.pdf>. 2010, cited 20 April 2012). The predominant carotenoids in *C. glutamicum* are the C50-terpene decaprenoxanthin and its glucosides [16]. To date, only three different C50 carotenoid biosynthetic pathways have been described: the biosynthetic pathways of the  $\epsilon$ -cyclic C50 carotenoid decaprenoxanthin in *C. glutamicum* [17,18], the  $\beta$ -cyclic C50 carotenoid C.p. 450 in *Dietzia* sp. CQ4 [19] and the  $\gamma$ -cyclic C50 carotenoid sarcinaxanthin in *Micrococcus luteus* NCTC2665 [20]. In addition, only a few other corynebacteria have been identified to contain carotenoid pigments i.e. *C. michiganense* [21], *C. erythrogenes* [22], *C. fascians* [23] and *C. poinsettiae* [24]. *C. poinsettiae* (*Curtobacterium flaccumfaciens*) e.g. is known to produce the C50 carotenoids bacterioruberin, bisanhydrobacterioruberin and C.p. 450 [2]. The genome of *C. glutamicum* encodes the enzymes of the MEP pathway [2,25]. Based on transposon mutant analysis and biochemical evidence *C. glutamicum* possesses a carotenogenic gene cluster encoding the responsible enzymes for the entire decaprenoxanthin biosynthesis starting from DMPP [17,18]. The immediate precursors of C30 and C40 carotenoids, farnesyl pyrophosphate (FPP, C15) and geranylgeranyl pyrophosphate (GGPP, C20), are generated from DMPP by prenyl transferase CrtE [18]. Subsequently, phytoene synthase (CrtB) condenses two GGPP molecules yielding the colorless carotenoid phytoene. Four subsequent desaturation reactions by phytoene desaturase (CrtI) yield the red-colored lycopene [17,18]. The elongation of lycopene with DMPP to the acyclic C50 carotenoid flavuxanthin is catalyzed by the *crtEb* gene product lycopene elongase. The cyclization of flavuxanthin to decaprenoxanthin is catalyzed by heterodimeric carotenoid  $\epsilon$ -cyclase, encoded by *crtY<sub>e</sub>* and *crtY<sub>f</sub>* [16,20,26]. While mono- and diglucosylated decaprenoxanthin can be found in *C. glutamicum*, the genes and enzymes for glucosylation of decaprenoxanthin are still unknown [20].

In this study, gene-directed deletion mutagenesis was employed to decipher the functions of the genes present in the main carotenogenic gene cluster of *C. glutamicum*

and in a second cluster encoding putative phytoene synthase and phytoene desaturase paralogs. Moreover, the potential of *C. glutamicum* to produce carotenoids was estimated by metabolic engineering of the conversion of GGPP to lycopene.

## Results


### Bioinformatical analysis of the carotenogenic genes

The genome of *C. glutamicum* ATCC 13032 (wild type; WT) encodes genes showing homology to carotenoid biosynthesis genes in two gene clusters that are separated by almost 2 Mbp. The larger cluster is composed of seven genes, *crtE* (cg0723), cg0722 (encoding a putative membrane protein), *crtB* (cg0721), *crtI* (cg0720), *crtY<sub>e</sub>*, *crtY<sub>f</sub>* (cg01719/18) and *crtEb* (cg0717) (Figure 1 and 2). The second cluster consists of a gene putatively encoding phytoene synthase (here named *crtB2*, cg2672) and two genes with similarity to an N-terminal fragment (*crtI2-1*, cg2670) and a C-terminal fragment (*crtI2-2*, cg2668) of phytoene desaturase/dehydrogenase (Figure 1).

The cluster *crtB2/crtI2-1/crtI2-2* has not yet been analyzed. While CrtB and CrtB2 share 49% identity, CrtI2-1 shares 49% identical amino acids with the 364 N-terminal amino acids of CrtI and CrtI2-2 63% identical amino acids with the 104 C-terminal amino acids of CrtI. Thus, it is not clear whether CrtI2-1 and CrtI2-2 function as a two-subunit phytoene desaturase or whether a frameshift mutation disrupted the gene downstream of *crtB2*.

A comparison of the sequenced genomes of corynebacteria (Figure 1, Additional file 1: Table S1) revealed that *C. glutamicum* WT is the only species possessing two *crtB* and *crtI* like genes, while the organization of the large gene cluster is comparable in *C. glutamicum* WT, *C. glutamicum* R (and ATCC 14067 and S9114) and *C. efficiens* YS-314. In *C. glutamicum* R, no *crtY<sub>e</sub>Y<sub>f</sub>* is annotated as likely a G->T mutation at position 814478 of the *C. glutamicum* R genome altered the start codon of an open reading frame coding for a protein with 99% amino acid identity to *crtY<sub>e</sub>Y<sub>f</sub>* of *C. glutamicum* WT to a leucine codon.

A second group of corynebacterial species (e.g. *C. diptheriae*, *C. aurimucosum* and *C. pseudotuberculosis*) only possess the clustered genes *crtB* and *crtI* (50 to 55% amino acid identity to the *C. glutamicum* enzymes; Additional file 1: Table S1). An intermediate situation is found in *C. lipophiloflavum*, which possesses a gene cluster with *crtB*, *crtI*, *crtY<sub>ef</sub>* and *crtEb*, as well as in *C. genitalium* possessing *crtB*, *crtI* and *crtY<sub>ef</sub>* but lacking *crtEb* (Additional file 1: Table S1). Members of a third group (*C. kroppenstedtii*, *C. jeikeium*, *C. urealyticum* as examples) also lack *crtY<sub>ef</sub>* and *crtEb* orthologs, but possess *crtB* and *crtI*, however not clustered. Although the overall amino acid sequence identities of the *crtB* and *crtI* gene products are below 50% as compared to the respective CrtB and CrtI from *C. glutamicum* WT,



**Figure 1** Genomic organization of the putative and characterized carotenogenic genes in different corynebacteria.

their domain structure includes the *crtI* domain (TIGR02734) as well as an N-terminal NAD(P)-binding Rossmann-like domain (NCBI Domain structure). As an exception, *C. variabile* only possesses CrI with an amino acid identity to CrI from *C. glutamicum* WT of 58%.


The phylogeny of the *crtI* gene product (Additional file 2: Figure S1), which is present in all analysed corynebacteria, is congruent to the grouping of corynebacterial species with respect to occurrence and clustering of *crt* genes as shown in Figure 1 and Additional file 1: Table S1.

#### Analysis of the transcriptional organization of the carotenogenic gene clusters

Annotation of the carotenogenic gene cluster of the *C. glutamicum* WT for the biosynthesis of decaprenoxanthin from the precursor GGPP suggests co-transcription of *crtB*,

*crtI*, *crtY<sub>e</sub>* and *crtY<sub>f</sub>* and *crtEb*, while the upstream GGPP synthase gene *crtE* appears to be monocistronic. To characterize the transcriptional organization of this cluster RT-PCR experiments have been carried out. PCR analysis of cDNA synthesized from total RNA of the *C. glutamicum* WT using primer *crtEb-rv* (see Additional file 3: Table S2) revealed that the entire gene cluster is co-transcribed since fragments overlapping adjacent genes could be amplified in each case. A cDNA preparation without the addition of reverse transcriptase served as a negative control (Figure 3).


Similarly, RT-PCR analysis of the small gene cluster revealed that *crtB2*, *crtI2-1* and *crtI2-2* are co-transcribed. Figure 3B displays the amplificate of a fragment overlapping *crtB2* and *crtI2-1* based on cDNA generated by reverse transcription using the *crtI-rv* primer.



To determine the transcriptional start point (TSP) of *crtE* and *crtB2*, respectively, RNA was isolated from *C. glutamicum* WT grown in LB complex medium. By use of 5' RACE\_PCR, the TSP of *crtE* was identified as a guanosine 114 nucleotides upstream of the first nucleotide of the ATG start codon. The three most conserved nucleotides of the consensus -10 hexamer of *C. glutamicum* promoters [27] can be found in the -15 to -10 region. The -39 to -34 region contains a sequence motif sharing four identical nucleotides to -35 consensus. The TSP of *crtB2* was determined as a guanosine thirteen nucleotides upstream of the first nucleotide of the start codon GTG. The hexamer TAAAGT at position -13 to -8 relative to the TSP matches the three most conserved bases of the TANANT consensus sequence of the -10 region of *C. glutamicum* promoters [27]. At position -32 to -27 the hexamer TTGTCT was found, which resembles the key recognition motif for the -35 region of *C. glutamicum* promoters TTGNCA [27].

#### Gene deletion and complementation analysis of the carotenogenic gene clusters in *C. glutamicum*

Gene-directed deletion mutants of *C. glutamicum* WT lacking *crtB*, *crtI*, *crtEb*, or *crtY<sub>e</sub>Y<sub>f</sub>* were constructed and characterized regarding carotenoid production. Besides the single deletion mutants, strain *C. glutamicum*  $\Delta\Delta$  lacking *crtB*, *crtI*, *crtEb*, and *crtY<sub>e</sub>Y<sub>f</sub>* as well as the putative paralogs *crtB2*, *crtI2-1* and *crtI2-2* was constructed. All strains showed growth rates of about 0.35 h<sup>-1</sup> in CGXII minimal medium with 100 mM glucose as carbon source. Thus, growth was comparable to *C. glutamicum* WT. However, pigment accumulation differed between the various strains (Figure 2). The different composition of carotenoids in the cell extracts could be demonstrated by HPLC analyses (Additional file 4: Figure S2, Additional file 5: Figure S3, Additional file 6: S4 and data not shown). The spectrophotometric analysis of the methanolic cell extracts of the *C. glutamicum* WT showed the characteristic absorption maxima at 415, 440, and 470 nm for the yellow



decaprenoxanthin, whereas the spectra of the pale red-colored *C. glutamicum* strains  $\Delta$ *crtEb* and  $\Delta$ *crtY* showed absorption maxima at 445, 470 and 500 nm (Additional file 4: Figure S2).


The multiple deletion strain *C. glutamicum*  $\Delta\Delta$  (Additional file 3: Table S2) was used for stepwise reconstruction of the decaprenoxanthin biosynthetic pathway. Expression of *crtB* and *crtI* in the white strain *C. glutamicum*  $\Delta\Delta$  entailed a pale pink cell color and accumulation of lycopene was observed in cell extracts. Additional expression of *crtEb* entailed an orange cell color and accumulation of flavuxanthin. When *crtY<sub>e</sub>Y<sub>f</sub>* was expressed additionally, a color comparable to that of the wild type was observed and the HPLC chromatograms of the cell extracts were comparable to those of the wild type. Thus, expression of *crtB*, *crtI*, *crtEb*, *crtY<sub>e</sub>* and *crtY<sub>f</sub>* in the multiple deletion strain was sufficient to allow for decaprenoxanthin biosynthesis.

This finding was supported by analysis of the single gene deletion strains. Each deletion mutant could be complemented by ectopic expression of the respective gene deleted in the chromosome (Figure 2). The mutant  $\Delta$ *crtY* lacking the final reaction in the synthesis of decaprenoxanthin, i.e. introduction of two  $\epsilon$ -ionone groups into the acyclic flavuxanthin catalyzed by gene products of *crtY<sub>e</sub>Y<sub>f</sub>*, accumulated flavuxanthin and exhibited a pale orange to red color. In the absence of the penultimate enzyme reaction of decaprenoxanthin biosynthesis, i.e. prenylation of lycopene to flavuxanthin by lycopene elongase, in the mutant  $\Delta$ *crtEb*, lycopene accumulated

and neither flavuxanthin nor decaprenoxanthin were observed (HPLC analysis of cell extracts not shown). Accordingly, mutants  $\Delta$ *crtB* lacking phytoene synthase and  $\Delta$ *crtI* lacking phytoene desaturase showed white cell color and  $\Delta$ *crtI* accumulated phytoene, which absorbs light at wavelengths below 300 nm. Taken together, our gene deletion and complementation analysis corroborates previous biochemical and transposon mutagenesis data and results from heterologous gene expression regarding the functions of the enzymes encoded by *crtB*, *crtI*, *crtEb*, *crtY<sub>e</sub>* and *crtY<sub>f</sub>*.

The function of the putative *crtB* paralogous gene *crtB2* and of the putative *crtI* paralogous genes *crtI2-1* and *crtI2-2* has not yet been analyzed. As hardly any phytoene was detectable in  $\Delta$ *crtB*, but faint quantities of other carotenogenic intermediates were observed, *CrtB* appears to be the major phytoene synthase active under the chosen conditions. Similarly, the lack of the red chromophore lycopene in  $\Delta$ *crtI* indicated that *CrtI* is the only active phytoene desaturase. By contrast, a deletion mutant lacking the paralogous genes *crtB2*, *crtI2-1* and *crtI2-2* showed the same yellow phenotype as *C. glutamicum* WT and the cell extracts showed the identical elution pattern in the HPLC analysis. Moreover, deletion mutants lacking *crtB2**I2-1**I2-2* and either *crtB* or *crtI* grew like wild type and showed the same white phenotype as the *crtB* and *crtI* single deletion mutants. Thus, the paralogous genes annotated as *crtB2* and *crtI2-1* and *crtI2-2* are either not functional or not expressed (enough) under the chosen conditions.





**Figure 4 Lycopene production in *C. glutamicum*  $\Delta$ crtEb overexpressing carotenogenic genes. (A)** Cell pellets of cultures grown in glucose CGXII minimal medium after consumption of the carbon source. By the overexpression of the indicated carotenogenic genes the intensity of the red color was enhanced. **(B)** Lycopene concentrations of the cells depicted in A as determined by HPLC analyses of cell extracts.

synthesized. On the other hand, *C. michiganense* [21], *C. erythrogenes* [22], *C. fascians* [23] and *C. poinsettiae* [24] are known to synthesize carotenoids, but their genome sequences are unknown.

In this study it could be shown that the genes of the carotenoid gene cluster of *C. glutamicum* ATCC 13032 *crtE-cg0722-crtBIY<sub>e</sub>Y<sub>f</sub>Eb* are co-transcribed. Similarly, also the second cluster is transcriptionally organized as an operon. Transcriptional regulation of both operons has not yet been reported. The *in vivo* activity of the *crtB2* gene product appears low due either to low expression levels or to low catalytic activity as plasmid-

borne overexpression was required to complement the phenotype of the deletion mutant lacking the paralog *crtB*. Currently, it remains unknown whether *crtB2* expression is affected by environmental stimuli and if/how the function of the two paralogs is regulated.

The potential of *C. glutamicum* for overproduction of carotenoids is to our knowledge described here for the first time. The interest in production of carotenoids, which find application in a wide variety of products due to their antioxidative properties and their colors, by cost-effective, environmentally friendly microbial fermentation processes is steadily increasing.

The carotenogenic *C. glutamicum* is generally recognized as safe (GRAS), can readily be metabolically engineered and has been safely used in the million-ton-scale production of food-additives since more than 50 years [28]. Lycopene was chosen as a test carotenoid product as it may serve as a platform intermediate and as its red color serves as a simple read out. Lycopene is a commercial product obtained by fermentation with the fungus *Blakeslea trispora* [29] (Vitatene, Leon, Spain). Here we show that *C. glutamicum* overproduces lycopene if *crtEb* is deleted and that additional overexpression of the carotenogenic genes *crtE*, *crtB* and *crtI* boosted lycopene production 80 fold. The achieved lycopene concentration of 2.4 mg/g CDW is already comparable to that obtained with other popular biotechnological hosts like *E. coli*, for which e.g. a lycopene yield of 1.8 mg/g CDW was reported when the *crtE*, *crtB* and *crtI* genes of the plant pathogen *Pantoea ananatis* were overexpressed [20]. A higher lycopene concentration (6.6 mg/g CDW) could only be achieved in an *E. coli* strain overexpressing genes for IPP synthesis and carotenogenesis after a systematic screen identified three gene knock-outs in the central carbon metabolism [30]. In *E. coli* harboring multiple modifications, i.e. carrying a plasmid with genes of the lycopene biosynthetic pathway (*crtE*, *crtB* and *crtI*) and a plasmid containing the entire heterologous MVA pathway as well as the IPP isomerase gene, *idi*, and overexpressing the endogenous *dxs* gene, a lycopene concentration of  $6.8 \pm 0.4$  mg/g was obtained in batch culture [31]. Process engineering as fed-batch process with glycerol with 10 g/l glucose and 7.5 g/l arabinose boosted the lycopene concentration to 32 mg/g CDW [31].

The very good lycopene concentration obtained by *C. glutamicum* after engineering only the final three enzymatic steps of lycopene synthesis can likely be further enhanced by additional metabolic engineering of (a) IPP synthesis using the endogenous MEP pathway and/or the heterologous MVA pathway, (b) genome-based or computational approaches to identify target genes in the central metabolism or its regulation and (c) by process engineering using e.g. fed-batch protocols. Thus, *C. glutamicum* may serve as a suitable production host for lycopene and related carotenoids.

In addition, *C. glutamicum* is a natural producer of the relatively rare group of C50 carotenoids that feature strong antioxidative properties due to the multiple conjugated double bonds and the hydroxyl group [32-34]. The pharmaceutical potential of these C50 carotenoids is not yet well studied [35]. It is imaginable that decaprenoxanthin, its direct precursor flavuxanthin or the C50 carotenoid of *Micrococcus luteus*, sarcinaxanthin, could be of commercial interest. Notably, genes of *C. glutamicum* and

of *M. luteus* have been used to engineer *E. coli* for the production of sarcinaxanthin [20]. Thus, the product range of structurally diverse C50 carotenoids could be accessible by engineered hosts including *C. glutamicum*.

## Conclusion

The genes of the carotenoid gene cluster of *C. glutamicum* ATCC 13032 *crtE-cg0722-crtBIY<sub>e</sub>Y<sub>f</sub>Eb* are co-transcribed and encode the enzymes involved in the biosynthesis of the C50 carotenoid decaprenoxanthin. An alternative, functionally active phytoene synthase is encoded in the *crtB2/crtI2-1/crtI2-2* operon leading to a certain degree of redundancy in carotenoid synthesis in *C. glutamicum*. The potential of *C. glutamicum* as production host for terpenoids in general was demonstrated by considerable lycopene production after engineering the terminal reactions leading to lycopene.

## Methods

### Bacterial strains, media and growth conditions

The strains and plasmids used in this work are listed in Additional file 3: Table S2. *C. glutamicum* ATCC 13032 was used as wild type (WT). Precultivation of *C. glutamicum* strains was performed in BHI or LB medium. For cultivation in CGXII medium [36] precultivated cells were washed once with CGXII medium without carbon source and inoculated to an initial OD<sub>600</sub> of 1. Glucose was added as carbon and energy source to a concentration of 100 mM. Standard cultivations of *C. glutamicum* were performed at 30°C in a volume of 50 ml in 500 ml flasks with two baffles shaking at 120 rpm. The OD<sub>600</sub> was measured in dilutions using a Shimadzu UV-1202 spectrophotometer (Duisburg, Germany). For cloning, *E. coli* DH5α was used as host and cultivated in LB medium at 37°C. When appropriate kanamycin or spectinomycin were added to concentrations of 25 µg/ml and 100 µg/ml, respectively. Gene expression was induced by adding 1 mM IPTG.

### Recombinant DNA work

Plasmids were constructed in *E. coli* DH5α from PCR-generated fragments (KOD, Novagen, Darmstadt, Germany) and isolated with the QIAprep spin miniprep kit (QIAGEN, Hilden, Germany). Oligonucleotides used in this study were obtained from Eurofins MWG Operon (Ebersberg, Germany) and are listed in Additional file 3: Table S1. Standard reactions like restriction, ligation and PCR were performed as described previously [37]. If applicable, PCR products were purified using the PCR purification kit or MinElute PCR purification kit (QIAGEN, Hilden, Germany). For transformation of *E. coli* the RbCl method was used [38] and *C. glutamicum* was transformed via electroporation [39] at



2.5 kV, 200  $\Omega$  and 25  $\mu$ F. All cloned DNA fragments were shown to be correct by sequencing.

#### Determination of the transcriptional start point of *crtE* and *crtB2*

Total RNA was isolated from an exponentially growing culture of *C. glutamicum* WT as described previously [40]. Purified RNA was analyzed by UV-spectrometry in regard to quantity and quality and was stored at  $-20^{\circ}\text{C}$  until use. 2  $\mu$ g of total RNA were used to perform 5' rapid amplification of cDNA ends-PCR (5' RACE-PCR) basically as described previously [41] with use of *crtE*-rv and *crtB2*-rv primers, respectively, for reverse transcription. Both, individual C tailing and A tailing were performed and analyzed. RACE-PCR was performed with primers *crtE*-RACE and *crtB2*-RACE and either OligoT or OligoG. Sequencing of the generated PCR fragments was accomplished using the suitable RACE primers and gave identical results for C tailing and A tailing reactions.

#### Reverse transcription (RT) for the analysis of transcription units

Total RNA was isolated from an exponentially growing culture of *C. glutamicum* WT as described previously [40]. Purified RNA was analyzed by UV-spectrometry in regard to quantity and quality and was stored at  $-20^{\circ}\text{C}$  until use. 2  $\mu$ g of total RNA were used to perform reverse transcription to generate cDNA that was subsequently used as template for PCRs applying primer that bind at adjacent genes. The reverse transcription reactions were performed using SuperScript<sup>™</sup> II reverse transcriptase (Invitrogen, Karlsruhe, Germany), and the remaining RNA was removed by the use of RNase H (MBI Fermentas GmbH, St. Leon-Rot).

#### Overexpression of carotenogenic genes from *C. glutamicum*

Plasmids harboring a carotenogenic gene allowed its IPTG-inducible overexpression and were based on pEKEx3 [42] or pVWEx1 [43], respectively. Amplification of a carotenogenic gene by polymerase chain reaction (PCR) from genomic DNA of *C. glutamicum* WT, which was prepared as described [44], was carried out using the respective primers (Additional file 3: Table S1). The amplified products were cloned into the appropriately restricted pEKEx3 or pVWEx1 plasmid DNA.

#### Deletion of carotenogenic genes in *C. glutamicum* WT

For deletion of a carotenogenic gene, the suicide vector pK19*mobsacB* was used [36]. Genomic regions flanking a carotenogenic gene were amplified from genomic DNA of *C. glutamicum* WT using the respective primer pairs A/B and C/D as indicated in Additional file 3:

Table S1. The PCR products were purified and linked by crossover PCR using the respective primer pair A/D (Additional file 3: Table S1). The either *Sma*I or *Bam*HI restricted purified PCR product was cloned into pK19*mobsacB* resulting in the construction of the respective deletion vector (Additional file 3: Table S1). Targeted deletion of a carotenogenic gene via two-step homologous recombination using the respective deletion vector was carried out as described previously [26]. For the first recombination event integration of the vector into one of the flanking regions was selected via kanamycin resistance. Integration of the deletion vector into the genome results in a sucrose sensitivity because of the *sacB* gene product levansucrase. Selection for clones that have excised the deletion vector in a second recombination event was carried out via sucrose-resistance. Deletion of a carotenogenic gene was verified by PCR analysis of the constructed mutant using the respective primer pair E/F (Additional file 3: Table S1).

#### Extraction of carotenoids from bacterial cell cultures

To extract carotenoids from the *C. glutamicum* strains 20 ml aliquots of the cell cultures were centrifuged at  $10,000 \times g$  for 15 min, and the pellets were washed with deionized  $\text{H}_2\text{O}$ . The pigments were extracted with 10 ml methanol:acetone mixture (7:3) at  $60^{\circ}\text{C}$  for 80 min with thorough vortexing every 20 min. When necessary, several extraction cycles were performed to remove all visible colors from the cell pellet.

#### Analysis of carotenoids

The extraction mixture was centrifuged  $10,000 \times g$  for 15 min and the methanol supernatant was transferred to a new tube. The absorption spectra of the various extracts were measured at wavelengths between 400 and 800 nm using the UV-1202 spectrophotometer (Shimadzu, Duisburg, Germany). High performance liquid chromatography (HPLC) analyses of the *C. glutamicum* extracts were performed on an Agilent 1200 series HPLC system (Agilent Technologies Sales & Services GmbH & Co. KG, Waldbronn), including a diode array detector (DAD) for UV/visible (Vis) spectrum recording. Quantification of carotenoids was performed using the extracted wavelength chromatogram at peak  $\lambda_{\text{max}}$ , 450 nm for decaprenoxanthin and carotenoids with corresponding UV/Vis profiles and 470 nm for lycopene and corresponding carotenoids. Lycopene from tomato (Sigma, Steinheim, Germany) was used as standard. It was dissolved in chloroform according to its solubility and diluted in methanol. The HPLC protocol comprised isocratic elution for 25 min using a flow rate of 1.4 ml/min, a mobile phase composition of methanol/methyl tert-butyl ether/ethyl acetate (5:4:1) and a column system consisting of a precolumn (10x4 mm MultoHigh 100 RP18-5, CS Chromatographie

Service GmbH, Langerwehe, Germany) and a main column (ProntoSIL 200–5 C30, 250x4 mm, CS Chromatographie Service GmbH, Langerwehe, Germany). 100 µl extract was injected.

### Computational analysis

To compare the *crt* genes from *C. glutamicum* ATCC 13032 with other sequenced corynebacteria, the amino acid sequences of the respective genes were obtained from CoryneRegNet database (<http://www.coryneregnet.de/>). Sequence comparisons were carried out using BLASTP ([www.ncbi.nlm.nih.gov/blast/Blast.cgi](http://www.ncbi.nlm.nih.gov/blast/Blast.cgi), [45]) and CLUSTALW [46] and the identification of potential orthologs and paralogs to *C. glutamicum* ATCC 13032 proteins was achieved by pairwise reciprocal BLAST analysis [47]. Species names, gene identifier and accession numbers are given in Additional file 1: Table S2. A phylogenetic tree was constructed using the neighbor joining method [48] with 1,000 bootstrap replicates.

### Additional files

**Additional file 1: Table S1.** Comparison of the *crt* genes from different corynebacteria, for which genome sequence information is available in the database (NCBI). The rows show the gene identifier (top) and accession number (middle) for each *crt* gene of the corresponding species and the amino acid identity to the respective *crt* gene product from *C. glutamicum* ATCC 13032 (bottom).

**Additional file 2: Figure S1.** Phylogenetic tree of phytoene desaturase from different corynebacteria. Numbers at the nodes represent bootstrap values. Gene identifiers are given in Additional file 1: Table S2.

**Additional file 3: Table S2.** Bacterial strains, plasmids and oligonucleotides [38,40,47,49,50].

**Additional file 4: Figure S2.** HPLC chromatograms of carotenoids extracted from *C. glutamicum*  $\Delta crtB$  strains (A) and  $\Delta crtI$  (B). Detection by absorption at 470 nm. (A) Elution profiles of carotenoids extracted from *C. glutamicum* WT (blue),  $\Delta crtB$ (pEKEx3) (red),  $\Delta crtB$ (pEKEx3-*crtB*) (green),  $\Delta crtB$ (pEKEx3-*crtB2*) (pink). (B) Elution profiles of carotenoids extracted from *C. glutamicum* WT (blue),  $\Delta crtI$ (pEKEx3) (red),  $\Delta crtI$ (pEKEx3-*crtI*) (green),  $\Delta crtI$ (pEKEx3-*crtI2-1/2*) (pink).

**Additional file 5: Figure S3.** Absorption spectra of cell extracts of *C. glutamicum* WT and *crt* deletion strains. The extract of the strains *C. glutamicum*  $\Delta crtEb$  (black line) and  $\Delta crtY$  (grey line) show an additional absorption maximum at about 500 nm compared to the wild type (red line). *C. glutamicum*  $\Delta crtB$  (dotted line) and  $\Delta crtI$  (dashed line) show no absorption.

**Additional file 6: Figure S4.** HPLC elution profiles of carotenoids extracted from *C. glutamicum*  $\Delta\Delta$  strains. Detection by absorption at 470 nm. (A) Elution profiles of carotenoids extracted from *C. glutamicum*  $\Delta\Delta$ (pEKEx3/pVWEx1) (blue),  $\Delta\Delta$ (pEKEx3-*crtB*/pVWEx1-*crtI*) (red),  $\Delta\Delta$ (pEKEx3-*crtB2*/pVWEx1-*crtI*) (green). The red and green chromatograms show the accumulation of carotenoids including lycopene (L) which elutes after about 18 min. (B) Elution profiles of carotenoids extracted from *C. glutamicum*  $\Delta\Delta$ (pEKEx3/pVWEx1) (blue) and  $\Delta\Delta$ (pEKEx3-*crtI2-1/2*/pVWEx1-*crtB2*) (red).

### Abbreviations

DAD: Diode array detector; DMPP: Dimethylallyl pyrophosphate; EV: Empty vector; FPP: Farnesyl pyrophosphate; GAP: Glyceraldehyde 3-phosphate; GGPP: Geranylgeranyl pyrophosphate; GPP: Geranyl pyrophosphate; HPLC: High performance liquid chromatography; IPP: Isopentenyl pyrophosphate; IPTG: Isopropyl  $\beta$ -D-1-thiogalactopyranoside;

MEP: Methylerythritol phosphate; MVA: Mevalonate; RACE: Rapid amplification of cDNA ends; RT: Reverse transcription; TSP: Transcriptional start point; WT: Wild type.

### Competing interests

The authors do not declare competing interests.

### Authors' contributions

All authors contributed to designing the study. SAEH constructed and characterized the recombinant strains. VFW and PPW supervised the experiments. SAEH and PPW were responsible for the draft of the manuscript. All authors contributed to writing and approved the final manuscript.

### Acknowledgements

The authors thank Maria Metzler for experimental support and acknowledge support of the publication fee by Deutsche Forschungsgemeinschaft and the Open Access Publication Funds of Bielefeld University.

Received: 10 July 2012 Accepted: 5 September 2012

Published: 10 September 2012

### References

1. Lee PC, Schmidt-Dannert C: **Metabolic engineering towards biotechnological production of carotenoids in microorganisms.** *Appl Microbiol Biotechnol* 2002, **60**:1–11.
2. Sandmann G, Yukawa H: **Vitamin synthesis: carotenoids, biotin and pantothenate.** In *Handbook of Corynebacterium glutamicum*. Edited by Eggeling L, Bott M. Boca Raton: CRC Press; 2005:399–417.
3. Vershinin A: **Biological functions of carotenoids—diversity and evolution.** *Biofactors* 1999, **10**:99–104.
4. Kirsh VA, Mayne ST, Peters U, Chatterjee N, Leitzmann MF, Dixon LB, Urban DA, Crawford ED, Hayes RB: **A prospective study of lycopene and tomato product intake and risk of prostate cancer.** *Cancer Epidemiol Biomarkers Prev* 2006, **15**:92–98.
5. Mayne ST: **Beta-carotene, carotenoids, and disease prevention in humans.** *FASEB J* 1996, **10**:690–701.
6. Wang W, Shinto L, Connor WE, Quinn JF: **Nutritional biomarkers in Alzheimer's disease: the association between carotenoids, n-3 fatty acids, and dementia severity.** *J Alzheimers Dis* 2008, **13**:31–38.
7. Misawa N: **Pathway engineering for functional isoprenoids.** *Curr Opin Biotechnol* 2011, **22**:627–633.
8. Kim SW, Keasling JD: **Metabolic engineering of the nonmevalonate isopentenyl diphosphate synthesis pathway in *Escherichia coli* enhances lycopene production.** *Biotechnol Bioeng* 2001, **72**:408–415.
9. Rodriguez-Villalon A, Perez-Gil J, Rodriguez-Concepcion M: **Carotenoid accumulation in bacteria with enhanced supply of isoprenoid precursors by upregulation of exogenous or endogenous pathways.** *J Biotechnol* 2008, **135**:78–84.
10. Martin VJ, Pitera DJ, Withers ST, Newman JD, Keasling JD: **Engineering a mevalonate pathway in *Escherichia coli* for production of terpenoids.** *Nat Biotechnol* 2003, **21**:796–802.
11. Leonard E, Ajikumar PK, Thayer K, Xiao WH, Mo JD, Tidor B, Stephanopoulos G, Prather KL: **Combining metabolic and protein engineering of a terpenoid biosynthetic pathway for overproduction and selectivity control.** *Proc Natl Acad Sci USA* 2010, **107**:13654–13659.
12. Rohmer M: **The discovery of a mevalonate-independent pathway for isoprenoid biosynthesis in bacteria, algae and higher plants.** *Nat Prod Rep* 1999, **16**:565–574.
13. Lange BM, Rujan T, Martin W, Croteau R: **Isoprenoid biosynthesis: the evolution of two ancient and distinct pathways across genomes.** *Proc Natl Acad Sci USA* 2000, **97**:13172–13177.
14. Daum M, Herrmann S, Wilkinson B, Bechthold A: **Genes and enzymes involved in bacterial isoprenoid biosynthesis.** *Curr Opin Chem Biol* 2009, **13**:180–188.
15. Kirby J, Keasling JD: **Biosynthesis of plant isoprenoids: perspectives for microbial engineering.** *Annu Rev Plant Biol* 2009, **60**:335–355.
16. Krubasik P, Takaichi S, Maoka T, Kobayashi M, Masamoto K, Sandmann G: **Detailed biosynthetic pathway to decaprenoxanthin diglucoside in *Corynebacterium glutamicum* and identification of novel intermediates.** *Arch Microbiol* 2001, **176**:217–223.

17. Krubasik P, Kobayashi M, Sandmann G: Expression and functional analysis of a gene cluster involved in the synthesis of decaprenoxanthin reveals the mechanisms for C50 carotenoid formation. *Eur J Biochem* 2001, **268**:3702–3708.
18. Krubasik P, Sandmann G: A carotenogenic gene cluster from *Brevibacterium linens* with novel lycopene cyclase genes involved in the synthesis of aromatic carotenoids. *Mol Gen Genet* 2000, **263**:423–432.
19. Tao L, Yao H, Cheng Q: Genes from a *Dietzia* sp. for synthesis of C40 and C50 beta-cyclic carotenoids. *Gene* 2007, **386**:90–97.
20. Netzer R, Stafsnes MH, Andreassen T, Goksoyr A, Bruheim P, Brautaset T: Biosynthetic pathway for gamma-cyclic sarcinaxanthin in *Micrococcus luteus*: heterologous expression and evidence for diverse and multiple catalytic functions of C(50) carotenoid cyclases. *J Bacteriol* 2010, **192**:5688–5699.
21. Saperstein S, Starr MP: The ketonic carotenoid canthaxanthin isolated from a colour mutant of *Corynebacterium michiganense*. *Biochem J* 1954, **57**:273–275.
22. Hodgkiss W, Liston J, Goodwin TW, Jamikorn M: The Isolation and Description of 2 Marine Micro-Organisms with Special Reference to Their Pigment Production. *J Gen Microbiol* 1954, **11**:438–450.
23. Pebble J: The Carotenoids of *Corynebacterium fascians* Strain 2 Y. *J Gen Microbiol* June 1968, **52**:15–24. doi:10.1099/00221287-52-1-15.
24. Starr MP, Saperstein S: Thiamine and the carotenoid pigments of *Corynebacterium poinsettiae*. *Arch Biochem Biophys* 1953, **43**:157–168.
25. Kalinowski J, Bathe B, Bartels D, Bischoff N, Bott M, Burkovski A, Dusch N, Eggeling L, Eikmanns BJ, Gaigalat L, et al: The complete *Corynebacterium glutamicum* ATCC 13032 genome sequence and its impact on the production of L-aspartate-derived amino acids and vitamins. *J Biotechnol* 2003, **104**:5–25.
26. Eggeling L, Bott M (Eds): *Handbook of Corynebacterium glutamicum*. Boca Raton: CRC Press; 2005. ISBN 978-0-8493-1821-4.
27. Patek M, Nesvera J: Sigma factors and promoters in *Corynebacterium glutamicum*. *J Biotechnol* 2011, **154**:101–113.
28. Wendisch VF, Bott M, Eikmanns BJ: Metabolic engineering of *Escherichia coli* and *Corynebacterium glutamicum* for biotechnological production of organic acids and amino acids. *Curr Opin Microbiol* 2006, **9**:268–274.
29. Choudhari SM, Ananthanarayan L, Singhal RS: Use of metabolic stimulators and inhibitors for enhanced production of beta-carotene and lycopene by *Blakeslea trispora* NRRL 2895 and 2896. *Bioresour Technol* 2008, **99**:3166–3173.
30. Alper H, Jin YS, Moxley JF, Stephanopoulos G: Identifying gene targets for the metabolic engineering of lycopene biosynthesis in *Escherichia coli*. *Metab Eng* 2005, **7**:155–164.
31. Kim YS, Lee JH, Kim NH, Yeom SJ, Kim SW, Oh DK: Increase of lycopene production by supplementing auxiliary carbon sources in metabolically engineered *Escherichia coli*. *Appl Microbiol Biotechnol* 2011, **90**:489–497.
32. Jackson H, Braun CL, Ernst H: The chemistry of novel xanthophyll carotenoids. *Am J Cardiol* 2008, **101**:50D–57D.
33. Naguib YM: Antioxidant activities of astaxanthin and related carotenoids. *J Agric Food Chem* 2000, **48**:1150–1154.
34. Miller NJ, Sampson J, Candeias LP, Bramley PM, Rice-Evans CA: Antioxidant activities of carotenes and xanthophylls. *FEBS Lett* 1996, **384**:240–242.
35. Osawa A, Ishii Y, Sasamura N, Morita M, Kasai H, Maoka T, Shindo K: Characterization and antioxidative activities of rare C(50) carotenoids-sarcinaxanthin, sarcinaxanthin monoglucoside, and sarcinaxanthin diglucoside-obtained from *Micrococcus yunnanensis*. *J Oleo Sci* 2010, **59**:653–659.
36. Eggeling L, Reyes O: Experiments. In *Handbook of Corynebacterium glutamicum*. Edited by Eggeling L, Bott M. Boca Raton: CRC Press; 2005:3535–566.
37. Sambrook J, Russell D: *Molecular Cloning. A Laboratory Manual*. 3rd edition. Cold Spring Harbor: Cold Spring Harbor Laboratory Press; 2001.
38. Hanahan D: Studies on transformation of *Escherichia coli* with plasmids. *J Mol Biol* 1983, **166**:557–580.
39. van der Rest ME, Lange C, Molenaar D: A heat shock following electroporation induces highly efficient transformation of *Corynebacterium glutamicum* with xenogeneic plasmid DNA. *Appl Microbiol Biotechnol* 1999, **52**:541–545.
40. Netzer R, Krause M, Rittmann D, Peters-Wendisch PG, Eggeling L, Wendisch VF, Sahn H: Roles of pyruvate kinase and malic enzyme in *Corynebacterium glutamicum* for growth on carbon sources requiring gluconeogenesis. *Arch Microbiol* 2004, **182**:354–363.
41. Youn JW, Jolkver E, Kramer R, Marin K, Wendisch VF: Identification and characterization of the dicarboxylate uptake system DccT in *Corynebacterium glutamicum*. *J Bacteriol* 2008, **190**:6458–6466.
42. Stansen C, Uy D, Delaunay S, Eggeling L, Goergen JL, Wendisch VF: Characterization of a *Corynebacterium glutamicum* lactate utilization operon induced during temperature-triggered glutamate production. *Appl Environ Microbiol* 2005, **71**:5920–5928.
43. Peters-Wendisch PG, Schiel B, Wendisch VF, Katsoulidis E, Mockel B, Sahn H, Eikmanns BJ: Pyruvate carboxylase is a major bottleneck for glutamate and lysine production by *Corynebacterium glutamicum*. *J Mol Microbiol Biotechnol* 2001, **3**:295–300.
44. Eikmanns BJ, Rittmann D, Sahn H: Cloning, sequence analysis, expression, and inactivation of the *Corynebacterium glutamicum* *icd* gene encoding isocitrate dehydrogenase and biochemical characterization of the enzyme. *J Bacteriol* 1995, **177**:774–782.
45. Altschul S, et al: Basic local alignment search tool. *J Mol Biol* 1990, **215**:403–410.
46. Thompson JD, Higgins DG, Gibson TJ: CLUSTALW: improving the sensitivity of progressive multiple sequence alignment through sequence weighting, position-specific gap penalties and weight matrix choice. *Nucleic Acids Res* 1994, **22**:4673–4680.
47. Kohl TA, Tauch A: The GlxR regulon of the amino acid producer *Corynebacterium glutamicum*: Detection of the corynebacterial core regulon and integration into the transcriptional regulatory network model. *J Biotechnol* 2009, **143**:239–246.
48. Saitou N, Nei M: The neighbor-joining method: a new method for reconstructing phylogenetic trees. *Mol Biol Evol* 1987, **4**:406–425.
49. Abe S, Takayama K, Kinoshita S: Taxonomical studies on glutamic acid producing bacteria. *J Gen Appl Microbiol* 1967, **13**:279–301.
50. Schäfer A, Tauch A, Jäger W, Kalinowski J, Thierbach G, Puhler A: Small mobilizable multi-purpose cloning vectors derived from the *Escherichia coli* plasmids pK18 and pK19: selection of defined deletions in the chromosome of *Corynebacterium glutamicum*. *Gene* 1994, **145**:69–73.

doi:10.1186/1471-2180-12-198

Cite this article as: Heider et al.: Carotenoid biosynthesis and overproduction in *Corynebacterium glutamicum*. *BMC Microbiology* 2012 12:198.

Submit your next manuscript to BioMed Central and take full advantage of:

- Convenient online submission
- Thorough peer review
- No space constraints or color figure charges
- Immediate publication on acceptance
- Inclusion in PubMed, CAS, Scopus and Google Scholar
- Research which is freely available for redistribution

Submit your manuscript at  
www.biomedcentral.com/submit

