

DATABASE

Open Access

PASmiR: a literature-curated database for miRNA molecular regulation in plant response to abiotic stress

Shihua Zhang^{1,2†}, Yi Yue^{3†}, Liang Sheng^{2†}, Yunzhi Wu³, Guohua Fan³, Ao Li⁴, Xiaoyi Hu², Mingzhu ShangGuan² and Chaoling Wei^{2*}

Abstract

Background: Over 200 published studies of more than 30 plant species have reported a role for miRNAs in regulating responses to abiotic stresses. However, data from these individual reports has not been collected into a single database. The lack of a curated database of stress-related miRNAs limits research in this field, and thus a cohesive database system should necessarily be constructed for data deposit and further application.

Description: *PASmiR*, a literature-curated and web-accessible database, was developed to provide detailed, searchable descriptions of miRNA molecular regulation in different plant abiotic stresses. *PASmiR* currently includes data from ~200 published studies, representing 1038 regulatory relationships between 682 miRNAs and 35 abiotic stresses in 33 plant species. *PASmiR*'s interface allows users to retrieve miRNA-stress regulatory entries by keyword search using plant species, abiotic stress, and miRNA identifier. Each entry upon keyword query contains detailed regulation information for a specific miRNA, including species name, miRNA identifier, stress name, miRNA expression pattern, detection method for miRNA expression, a reference literature, and target gene(s) of the miRNA extracted from the corresponding reference or miRBase. Users can also contribute novel regulatory entries by using a web-based submission page. The *PASmiR* database is freely accessible from the two URLs of <http://hi.ustc.edu.cn:8080/PASmiR>, and <http://pcsb.ahau.edu.cn:8080/PASmiR>.

Conclusion: The *PASmiR* database provides a solid platform for collection, standardization, and searching of miRNA-abiotic stress regulation data in plants. As such this database will be a comprehensive repository for miRNA regulatory mechanisms involved in plant response to abiotic stresses for the plant stress physiology community.

Background

MicroRNAs (miRNAs) are endogenous, single-stranded, small (~21-23nt), non-coding, regulatory RNA molecules that manipulate messenger RNA (mRNA) degradation and translation repression by binding complementary sites in the protein-encoding, or 3'-untranslated regions, of target mRNAs [1-3]. Contemporary research has revealed numerous miRNAs in diverse plant species, such as *Arabidopsis thaliana* (mouse-ear cress), *Oryza sativa* (Asian rice), *Chlamydomonas reinhardtii* (flagellated green algae), and

Populus euphratica (subtropical poplar tree) [4-7]. An abundance of recent evidence suggests that plant miRNAs are critical to several essential biological processes in plant physiology, such as floral organ identity, leaf morphogenesis, cellular signaling, and stress responses [8-11]. Because of the diverse and important functions of miRNAs in plants, the dysfunction, mutation, and dysregulation of specific miRNAs has the potential to negatively influence plant growth and productivity.

Initially discovered as regulators of developmental timing in *Caenorhabditis elegans* (transparent nematode), miRNAs have already been demonstrated to play important regulatory roles, particularly in the complex network of gene regulatory pathways. During abiotic stress to plants, such as drought, salinity, wounding, and high temperature, miRNAs act at the posttranscriptional

* Correspondence: weichl@ahau.edu.cn

†Equal contributors

²Key laboratory of Tea Biochemistry and Biotechnology, Ministry of Education, Ministry of Agriculture, Anhui Agricultural University, Hefei 230036, China

Full list of author information is available at the end of the article

level in gene regulatory networks associated with stress adaptation and tolerance. Under adverse stress conditions, gene expression can be elaborately regulated to facilitate the action of these tolerance mechanisms [12-14], providing further evidence of the importance of miRNAs in regulation of plant responses to abiotic stress conditions.

In the past decade, the regulatory mechanisms of miRNAs involved in plant stress responses have garnered increasing attention. For example, miRNAs have been identified to regulate plant response to drought [15,16], salinity [17,18], low temperature [19,20], and nutrient deficiency [21-23]. More recently, a set of conserved and non-conserved miRNAs from *Medicago truncatula* (barrel clover) [24], *Oryza sativa* (Asian rice) [25], *Brassica napus* (rapeseed) [26], and *Arabidopsis thaliana* (mouse-ear cress) [27] that act to regulate plant response to heavy metal exposure, have been identified. These findings indicated that certain stress conditions stimulate specific plant species to produce miRNAs involved in species-specific regulatory processes that vary highly between different species, potentially accounting for differences in plant tolerances to specific stress conditions. Additionally, these results indicate that miRNAs could represent a discriminative evolutionary footprint associated with abiotic stress regulation, relationships that can only be determined by comprehensive organization of miRNA regulation data from many plant species.

To date, more than one thousand miRNAs have been discovered in diverse plant species [28]. Many of these miRNAs have confirmed involvement in regulatory networks associated with plant stress response. Despite notable recent progress in identifying miRNA regulation during stress conditions, detailed data of miRNA molecular regulation in different plant abiotic stresses, such as miRNA expression pattern and target genes of miRNAs, remains widely distributed in published literature. In the current stage, a cohesive database system is sorely needed for data deposit and further application. Therefore, development of the literature-curated database system "*PASmiR*" (miRNA molecular regulation in Plant Abiotic Stress) for the storage of regulatory relationships between miRNAs and plant abiotic stresses will be a valuable resource for contemporary researchers. While these immediate, tangible benefits are apparent, the *PASmiR* database may also serve as a valuable ongoing program for the study of miRNA regulatory mechanisms associated with plant responses to abiotic stress as additional information becomes available in this field.

Construction and content

Collection of regulatory relationships between miRNA and plant abiotic stress

We searched the databases of the PubMed, Scopus, Google Scholar, and Ovid [29-32], by using keywords,

such as 'stress microRNA' and 'stress miRNA', to retrieve publications that described miRNA-stress regulatory relationships. *PASmiR* currently encompasses data from approximately 200 published studies, representing 1038 regulatory relationships between 682 miRNAs and 35 abiotic stresses in 33 plant species. For each of these regulatory relationships, we collected detailed information for miRNA molecular regulation in abiotic stress response, which contains the plant species name, a miRNA identifier, the abiotic stress name, miRNA expression pattern (up-regulated or down-regulated) during stress condition, detection method for miRNA expression (e.g., miRNA microarray, qRT-PCR, northern blot, and deep sequencing, etc.), a reference literature, the target gene(s) of miRNA extracted from the corresponding reference, and validated (predicted) target gene(s) derived from miRBase [33]. In *PASmiR*, convenient links are provided to other external databases, including miRNA sequence and annotation in miRBase, and species taxonomic description from UniProt taxonomy [34]. In the original publications, different members of a miRNA family may not be stated properly due to limited experimental design. Thus, a family-like miRNA identifier (e.g., *osa-miR172*) documented with expression pattern assignment will denote the regulatory pattern of an unspecified family member.

Nomenclature standardization

The published studies used to create *PASmiR* entries use varied standards for miRNA and abiotic stress presentation, differing widely between individual journals and researchers. As a result, variant miRNA and abiotic stress spellings and descriptions exist, making literature curation a complicated task. For example, two forms of *osa-miR171* and *miR-171* that appear in different publications may actually refer to the same miRNA in *Oryza sativa* (Asian rice) [19,35].

Establishment of the *PASmiR* database necessitated the development of a curated standard nomenclature for miRNAs and abiotic stresses. Each miRNA inputted into the *PASmiR* database was converted to a standard terminology by referring to miRBase [36], as follows: a prefix of species abbreviation based on Latin name was followed by a dash, "miR", and a family number. For instance, "*osa-miR172*" would be correct terminology for *Oryza sativa* miRNA from family 172.

A total of 35 different plant abiotic stresses have been identified. For the purposes of inclusion in *PASmiR*, each inputted abiotic stress was assigned unique names and manually classified into one of 11 abiotic stress classes according to physiological affection (Figure 1). Figure 2 illustrates the distribution of miRNAs and plant species involved in different abiotic stresses. Examination of the prevalent abiotic stresses represented in *PASmiR*

Abiotic stress	Abiotic stress type
Abscisic acid (ABA)	Hormone
Auxin	
Ethylene	
Gibberellic acid (GA)	
Cadmium (Cd)	Heavy metal
Manganese (Mn)	
Mercury (Hg)	
Compression	Mechanical damage
Cropping	
Tension	
Topping	
Wounding	Nutritional starvation
Carbon (C) deficiency	
Nitrogen (N) deficiency	
Phosphorus (P) deficiency	
Sulphur (S) deficiency	Oxygenous stress
Hypoxia	
Ozone	Radiational stress
High light	
Radiation	
Ultraviolet B (UV-B)	Saline, alkaline
High osmotic	
High alkali	
High salt	Temperature stress
Heat	
Low temperature	Trace element
Aluminum (Al) excess	
Copper (Cu) deficiency	
Copper (Cu) excess	
Ferrum (Fe) deficiency	Drought, waterlogging
Drought (water deficit, dehydration)	
Waterlogging (submergence)	Unclassified
Arsenite treatment	
Glyphosate	
Sucrose	

Figure 1 Eleven abiotic stress classes generated by manual curation.

demonstrated miRNA involvement was most often reported in stresses of drought, low temperature, high salt, and phosphate (Pi) deficiency.

Database construction


The *PASmiR* database was constructed using freely available and open source frameworks, such as Apache, JavaServer Pages (JSP), MySQL, Struts2, Spring, Hibernate and so on. The curated miRNA-stress regulatory relationships were stored in a MySQL Server and accessed using Structured Query Language (SQL) queries. The web platform is based on Apache Tomcat server and its pages are generated via a combination of Java language and JSP scripts. In addition,

Hibernate modules were applied to manipulate data and convert data formats.

Utility

Keyword-based searching

The web-accessible *PASmiR* database allows for curated miRNA-stress regulatory relationships to be clearly and simply browsed, searched, downloaded, and updated (Figure 3). Using the search page, remote users can search the database by keywords. Three search fields are present that may be used together or separately: 1) plant species, 2) abiotic stress, and 3) miRNA. Logical operators (AND and OR) were configured between two of these fields to allow rapid, targeted access to specific entries of interest.



Fuzzy search capabilities

Users can remotely access the web-based user interface to conveniently generate queries on the database using logical, standardized keywords. *PASmiR* also offers a fuzzy search engine, allowing querying of miRNA-stress regulatory information by plant species, abiotic stress or miRNA, even when the exact name is unknown. Notably, multiple hits may result from input of single queries containing few or vague keywords, allowing for manual selection of data presented by relevance to the search term(s).

User submit to *PASmiR*

PASmiR provides a submit page that allows non-affiliated researchers to independently contribute novel miRNA-stress regulatory relationships as they become available. For user submissions, primary required fields are: 1) plant species, 2) abiotic stress, 3) miRNA and 4) reference title, as indicated by the conventional red star symbol. The curator(s) at our site conducts manual verification of the original publication(s) for data validation purposes to maintain the quality and integrity of the database. Submissions that pass this review process are then approved for entry into the *PASmiR* database.

Notably, all new submission will be made available in coming *PASmiR* versions on a monthly release schedule.

Discussion

Contemporary exploration of miRNA molecular regulation associated with plant abiotic stresses has demonstrated that miRNAs act as important regulators in plant physiological adaption mechanisms during response to unfavorable stresses. Unfortunately, data information on specific miRNAs involved in regulation of plant responses to abiotic stress is widely dispersed throughout current scientific literature. To provide a comprehensive repository for collection, standardization, and searching of these miRNA-abiotic stress regulatory relationships, the literature-curated and web-accessible *PASmiR* database was established.

Notably, the number of miRNAs involved in particular abiotic stress responses is highly dependent on the experimental approaches proposed in the original investigations. For example, more miRNAs will be observed using a microarray platform than other low-throughput technologies [37,38]. In different plant species, miRNAs may be defined as functionally conserved if their homologs behave as regulators of specific abiotic stress responses in more than one other plant species. For example, the miRNA




Figure 3 Homepage of the PASMIR database. The web-accessible PASMIR database allows for curated miRNA-stress regulatory entries to be clearly and simply browsed, searched, downloaded, and updated.

homologs *peu-miR164* and *mtr-miR164* are functionally conserved in plant responses to drought stress in both *Populus euphratica* (Euphrates Poplar) [6] and *Medicago truncatula* (Barrel Clover) [39]. Of 682 total plant miRNAs, 242 (35.5%) represent functional conserved miRNAs, and 16 (6.6%) of which are functionally conserved in more than 3 abiotic stresses response in different plant species (see Additional file 1). Investigation of these conserved miRNAs using a structured database approach, such as that provided by PASMIR, will facilitate improvements in the understanding of stress-focused miRNA functional evolutionarily relationship in plant species. This data has potential applications for experimental biology aiming to improve plant crop performance and breeding characteristics.

Current efforts have demonstrated that miRNAs play critical roles in regulating plant responses to abiotic stresses in a complicated network manner [40-43]. Thus, it is possible that analysis of the miRNA-stress regulation data archived in PASMIR may be able to yield novel information about the network regulatory mechanisms of miRNAs involved in stress response. In the staple food crop *Oryza sativa* (Asian rice), a bipartite network can be used to describe the 162 regulatory relationships between 113 miRNAs and 13 abiotic stresses known in this species (Figure 4). Within the network, drought, arsenite treatment, cadmium (Cd), low temperature, aluminum (Al) excess, and high salt demonstrate the highest connectivity by involving 35, 25, 25, 22, 19, and 13 regulatory miRNAs in the stress response, respectively.

Several miRNAs, such as *osa-miR156*, *osa-miR393*, and *osa-miR528*, have been demonstrated to play regulatory roles in more than two types of abiotic stress responses. Among these miRNAs, a small portion (solid purple circles) exhibit opposite expression patterns in different abiotic stresses response. The miRNA *osa-miR156*, for example, has a high-expression tendency when subjected to radiation exposure, but exhibits the opposite tendency when exposed to either low temperature or drought, indicating that, as a regulator of stress, *osa-miR156* exhibits unique behaviors under variant stress conditions. Cumulatively, these findings indicate the complex network regulatory relationships between miRNAs and abiotic stresses. By using PASMIR, researchers will have access to a comprehensive knowledge base for assembling bipartite networks for plant species documented, allowing the completion of novel derivations for detailed stress-driven miRNA molecular regulatory mechanisms.

The PASMIR database project, reported herein, provides the initial groundwork for an ongoing program for cataloging and utilizing miRNA-abiotic stress regulatory information in plants. Notably, the current version is not as comprehensive as we envision future versions to be. Ongoing collection and curation of new publications related to miRNA-stress response regulation is essential to this program's success. New versions of the databases are currently scheduled to be released on a monthly basis, incorporating all new submissions to the database over the preceding month. While this work is currently time and




Figure 4 (See legend on next page.)

(See figure on previous page.)

Figure 4 Bipartite network visualization of the regulatory relationship between miRNAs and abiotic stresses in the commercial crop *Oryza sativa* (Asian rice). In the network, squares and circles correspond to stresses and miRNAs, respectively. A link was placed between stresses and miRNAs indicating miRNA involvement in the regulation of a specific abiotic stress response. The outlined red, outlined blue, and solid purple of the circle denote different miRNA expression pattern (up-regulated, down-regulated and combined, respectively) in different stress conditions, respectively. Notably, a family-like miRNA identifier (e.g., osa-miR172) with expression pattern assignment will denote the regulatory pattern of an unspecified family member.

resource intensive, the pending employment of a natural language processing (NLP) technique to allow more rapid abstract prescreening will facilitate the addition of many new entries to the database in the near future. This strategy is expected to increase comprehensiveness and accessibility of the database, promoting broader interest from researchers worldwide.

Conclusion

We constructed the first comprehensive database for miRNA molecular regulation in plant abiotic stress responses. The web-based program was created by manually reviewing widely scattered scientific literature. This database, however, also provides a convenient submission interface for contribution of novel miRNA-stress regulatory relationship by independent researchers. *PASmiR* provides comprehensive, keyword-searchable miRNA-abiotic regulatory relationships focused on a specific plant species, abiotic stress, and miRNA family. Thus, *PASmiR* will aid in rapid and complete exploration of the complex network of miRNA regulatory mechanisms involved in plant abiotic stress responses, serving as a valuable resource for future investigators in experimental biology concerned with commercial crop breeding and characteristics.

Availability and requirements

Project name: *PASmiR*: A literature-curated database for miRNA molecular regulation in plant response to abiotic stress

Project home pages: <http://hi.ustc.edu.cn:8080/PASmiR>

<http://pcsb.ahau.edu.cn:8080/PASmiR>

Operating system(s): Platform independent

Programming languages: JSP, MySQL, HTML and JavaScript

License: Not required.

Any restrictions to use by non-academics: None

Additional file

Additional file 1: Distribution of functionally conserved miRNAs in 35 different abiotic stresses. The expression patterns of miRNAs are indicated as red and blue colors according to their up-regulated and down-regulated patterns, respectively.

Abbreviations

PASmiR: miRNA molecular regulation in Plant Abiotic Stress;
microRNA: miRNA.

Competing interests

The authors declare that they have no competing interests.

Authors' contributions

SZ, YY and LS performed literature mining, constructed the database platform, and wrote the manuscript. YW, GF, AL, XH and MS helped with the design of database platform and update of the database, and provided scientific suggestions and criticisms for improving the manuscript and website. SZ and CW participated in the design, helped write the manuscript and supervised the whole project. All authors read and approved the final manuscript.

Acknowledgements

The research was supported by grants from the National "Twelfth Five-Year" basic research program of China (973, 2012CB722903), the National Natural Science Foundation of China (Grant Nos. 31171608, 31270714, and 31270729), the National "Twelfth Five-Year" Technology Support Program of China (2011BAD01B01), and the Youth Research Foundation of Anhui Agricultural University (2012zd010). Funding to pay the Open Access publication charges for this article was provided by the above grants. Many thanks will be given to the editor Michael K Deyholos and the anonymous reviewers for their instructive suggestions. We are also grateful to the unknown editor at the MedSci editing service for the English polishing.

Author details

¹School of life sciences, Anhui Agricultural University, Hefei 230036, China.

²Key laboratory of Tea Biochemistry and Biotechnology, Ministry of Education, Ministry of Agriculture, Anhui Agricultural University, Hefei 230036, China. ³College of information and computer science, Anhui Agricultural University, Hefei 230036, China. ⁴Department of Electronic Science and Technology, University of Science and Technology of China, Hefei, China.

Received: 9 September 2012 Accepted: 21 February 2013

Published: 1 March 2013

References

1. Bartel DP: **MicroRNAs: genomics, biogenesis, mechanism, and function.** *Cell* 2004, **116**(2):281–297.
2. Meister G, Tuschl T: **Mechanisms of gene silencing by double-stranded RNA.** *Nature* 2004, **431**(7006):343–349.
3. Zhang B, Stellwag EJ, Pan X: **Large-scale genome analysis reveals unique features of microRNAs.** *Gene* 2009, **443**(1–2):100–109.
4. Molnar A, Schwach F, Studholme DJ, Thuenemann EC, Baulcombe DC: **miRNAs control gene expression in the single-cell alga *Chlamydomonas reinhardtii*.** *Nature* 2007, **447**(7148):1126–1129.
5. Jian X, Zhang L, Li G, Wang X, Cao X, Fang X, Chen F: **Identification of novel stress-regulated microRNAs from *Oryza sativa* L.** *Genomics* 2010, **95**(1):47–55.
6. Li B, Qin Y, Duan H, Yin W, Xia X: **Genome-wide characterization of new and drought stress responsive microRNAs in *Populus euphratica*.** *J Exp Bot* 2011, **62**(11):3765–3779.
7. Sunkar R, Zhu JK: **Novel and stress-regulated microRNAs and other small RNAs from *Arabidopsis*.** *Plant Cell* 2004, **16**(8):2001–2019.
8. Aukerman MJ, Sakai H: **Regulation of flowering time and floral organ identity by a MicroRNA and its APETALA2-like target genes.** *Plant Cell* 2003, **15**(11):2730–2741.
9. Chiou TJ, Lin SJ: **Signaling network in sensing phosphate availability in plants.** *Annu Rev Plant Biol* 2011, **62**:185–206.
10. Lu S, Sun YH, Chiang VL: **Stress-responsive microRNAs in *Populus*.** *Plant J* 2008, **55**(1):131–151.

11. Palatnik JF, Allen E, Wu X, Schommer C, Schwab R, Carrington JC, Weigel D: **Control of leaf morphogenesis by microRNAs.** *Nature* 2003, **425**(6955):257–263.
12. Covarrubias AA, Reyes JL: **Post-transcriptional gene regulation of salinity and drought responses by plant microRNAs.** *Plant Cell Environ* 2010, **33**(4):481–489.
13. Jia X, Ren L, Chen QJ, Li R, Tang G: **UV-B-responsive microRNAs in *Populus tremula*.** *J Plant Physiol* 2009, **166**(18):2046–2057.
14. Zuo J, Wang Y, Liu H, Ma Y, Ju Z, Zhai B, Fu D, Zhu Y, Luo Y, Zhu B: **MicroRNAs in tomato plants.** *Sci China Life Sci* 2011, **54**(7):599–605.
15. Sunkar R, Jagadeeswaran G: **In silico identification of conserved microRNAs in large number of diverse plant species.** *BMC Plant Biol* 2008, **8**:37.
16. Zhao B, Liang R, Ge L, Li W, Xiao H, Lin H, Ruan K, Jin Y: **Identification of drought-induced microRNAs in rice.** *Biochem Biophys Res Commun* 2007, **354**(2):585–590.
17. Ding D, Zhang L, Wang H, Liu Z, Zhang Z, Zheng Y: **Differential expression of miRNAs in response to salt stress in maize roots.** *Ann Bot* 2009, **103**(1):29–38.
18. Liu HH, Tian X, Li YJ, Wu CA, Zheng CC: **Microarray-based analysis of stress-regulated microRNAs in *Arabidopsis thaliana*.** *RNA* 2008, **14**(5):836–843.
19. Lv DK, Bai X, Li Y, Ding XD, Ge Y, Cai H, Ji W, Wu N, Zhu YM: **Profiling of cold-stress-responsive miRNAs in rice by microarrays.** *Gene* 2010, **459**(1–2):39–47.
20. An FM, Hsiao SR, Chan MT: **Sequencing-based approaches reveal low ambient temperature-responsive and tissue-specific microRNAs in phalaenopsis orchid.** *PLoS One* 2011, **6**(5):e18937.
21. Bari R, Datt Pant B, Stitt M, Scheible WR: **PHO2, microRNA399, and PHR1 define a phosphate-signaling pathway in plants.** *Plant Physiol* 2006, **141**(3):988–999.
22. Chiou TJ, Aung K, Lin SJ, Wu CC, Chiang SF, Su CL: **Regulation of phosphate homeostasis by MicroRNA in *Arabidopsis*.** *Plant Cell* 2006, **18**(2):412–421.
23. Fujii H, Chiou TJ, Lin SJ, Aung K, Zhu JK: **A miRNA involved in phosphate-starvation response in *Arabidopsis*.** *Curr Biol* 2005, **15**(22):2038–2043.
24. Zhou ZS, Huang SQ, Yang ZM: **Bioinformatic identification and expression analysis of new microRNAs from *Medicago truncatula*.** *Biochem Biophys Res Commun* 2008, **374**(3):538–542.
25. Huang SQ, Peng J, Qiu CX, Yang ZM: **Heavy metal-regulated new microRNAs from rice.** *J Inorg Biochem* 2009, **103**(2):282–287.
26. Xie FL, Huang SQ, Guo K, Xiang AL, Zhu YY, Nie L, Yang ZM: **Computational identification of novel microRNAs and targets in *Brassica napus*.** *FEBS Lett* 2007, **581**(7):1464–1474.
27. Ding YF, Zhu C: **The role of microRNAs in copper and cadmium homeostasis.** *Biochem Biophys Res Commun* 2009, **386**(1):6–10.
28. Voinnet O: **Origin, biogenesis, and activity of plant microRNAs.** *Cell* 2009, **136**(4):669–687.
29. Wheeler DL, Barrett T, Benson DA, Bryant SH, Canese K, Chetverin V, Church DM, DiCuccio M, Edgar R, Federhen S, et al: **Database resources of the national center for biotechnology information.** *Nucleic Acids Res* 2007, **35**(Database issue):D5–D12.
30. Burnham JF: **Scopus database: a review.** *Biomed Digit Libr* 2006, **3**:1.
31. Henderson J: **Google scholar: a source for clinicians?** *CMAJ* 2005, **172**(12):1549–1550.
32. Fitzpatrick RB: **Decision support with ovid.** *Med Ref Serv Q* 2001, **20**(4):47–53.
33. Kozomara A, Griffiths-Jones S: **miRBase: integrating microRNA annotation and deep-sequencing data.** *Nucleic Acids Res* 2011, **39**(Database issue): D152–D157.
34. Magrane M, Consortium U: **UniProt knowledgebase: a hub of integrated protein data.** *Database (Oxford)* 2011, **2011**:bar009.
35. Zhou L, Liu Y, Liu Z, Kong D, Duan M, Luo L: **Genome-wide identification and analysis of drought-responsive microRNAs in *Oryza sativa*.** *J Exp Bot* 2010, **61**(15):4157–4168.
36. Griffiths-Jones S: **miRBase: microRNA sequences and annotation.** *Curr Protoc Bioinformatics* 2010, **Chapter 12**:Unit 12.19.11–10.
37. Kawashima CG, Matthewman CA, Huang S, Lee BR, Yoshimoto N, Koprivova A, Rubio-Somoza I, Todesco M, Rathjen T, Saito K, et al: **Interplay of SLIM1 and miR395 in the regulation of sulfate assimilation in *Arabidopsis*.** *Plant J* 2011, **66**(5):863–876.
38. Ding Y, Chen Z, Zhu C: **Microarray-based analysis of cadmium-responsive microRNAs in rice (*Oryza sativa*).** *J Exp Bot* 2011, **62**(10):3563–3573.
39. Wang T, Chen L, Zhao M, Tian Q, Zhang WH: **Identification of drought-responsive microRNAs in *Medicago truncatula* by genome-wide high-throughput sequencing.** *BMC Genomics* 2011, **12**:367.
40. Zhu C, Ding Y, Liu H: **MIR398 and plant stress responses.** *Physiol Plant* 2011, **143**(1):1–9.
41. Liu Z, Kumari S, Zhang L, Zheng Y, Ware D: **Characterization of miRNAs in response to short-term waterlogging in three inbred lines of *Zea mays*.** *PLoS One* 2012, **7**(6):e39786.
42. Wan P, Wu J, Zhou Y, Xiao J, Feng J, Zhao W, Xiang S, Jiang G, Chen JY: **Computational analysis of drought stress-associated miRNAs and miRNA co-regulation network in *Physcomitrella patens*.** *Genomics Proteomics Bioinformatics* 2011, **9**(1–2):37–44.
43. Gollmack D, Luking I, Yang O: **Plant tolerance to drought and salinity: stress regulating transcription factors and their functional significance in the cellular transcriptional network.** *Plant Cell Rep* 2011, **30**(8):1383–1391.

doi:10.1186/1471-2229-13-33

Cite this article as: Zhang et al.: *PASmiR*: a literature-curated database for miRNA molecular regulation in plant response to abiotic stress. *BMC Plant Biology* 2013 **13**:33.

Submit your next manuscript to BioMed Central and take full advantage of:

- **Convenient online submission**
- **Thorough peer review**
- **No space constraints or color figure charges**
- **Immediate publication on acceptance**
- **Inclusion in PubMed, CAS, Scopus and Google Scholar**
- **Research which is freely available for redistribution**

Submit your manuscript at
www.biomedcentral.com/submit

