



ORIGINAL RESEARCH

Open Access

Pain and quality of life in patients undergoing radiotherapy for spinal metastatic disease treatment

Edgar S Valesin Filho¹, Luiz Carlos de Abreu², Guilherme HV Lima¹, Daniel IG de Cubero¹, Fabrício H Ueno¹, Gustavo SL Figueiredo¹, Vitor E Valenti^{2,3}, Carlos Bandeira de Mello Monteiro^{2,4}, Rubens Wajnsztejn², Edison N Fujiki^{1,2}, Modesto Rolim Neto² and Luciano M Rodrigues^{1,2*}

Abstract

Background: Radiotherapy is an important tool in the control of pain in patients with spinal metastatic disease. We aimed to evaluate pain and of quality of life of patients with spinal metastatic disease undergoing radiotherapy with supportive treatment.

Methods: The study enrolled 30 patients. From January 2008 to January 2010, patients selection included those treated with a 20 Gy tumour dose in five fractions. Patients completed the visual analogue scale for pain assessment and the SF-36 questionnaire for quality of life assessment.

Results: The most frequent primary sites were breast, multiple myeloma, prostate and lymphoma. It was found that 14 spinal metastatic disease patients (46.66%) had restricted involvement of three or fewer vertebrae, while 16 patients (53.33%) had cases involving more than three vertebrae. The data from the visual analogue scale evaluation of pain showed that the average initial score was 5.7 points, the value 30 days after the end of radiotherapy was 4.60 points and the average value 6 months after treatment was 4.25 points. Notably, this final value was 25.43% lower than the value from the initial analysis. With regard to the quality of life evaluation, only the values for the functional capability and social aspects categories of the questionnaire showed significant improvement.

Conclusion: Radiotherapy with supportive treatment appears to be an important tool for the treatment of pain in patients with spinal metastatic disease.

Keywords: Radiotherapy, Neoplasm metastasis, Quality of life

Background

Spinal metastatic disease (SMD) is a relatively common disease and has received increased attention recently with the development of diagnostic and therapeutic tools that increase the life expectancy of these oncology patients [1,2]. Radiotherapy is an important tool for the control of pain and local neoplastic progression. Several studies have demonstrated its efficacy in stabilizing the progression of pain and in maintaining the ability of the

patient to walk, which are considered import goals of treatment [3-6].

The majority of tumors found in the spine are metastatic lesions, and approximately 18,000 new cases are diagnosed annually in the United States [7]. According to the literature, more than 10% of patients with cancer will develop symptomatic secondary spinal disease, and more than 40 to 70% of those cases will involve several vertebral levels [8-10].

Historically, radiotherapy has been used in the treatment of SMD and is considered the standard treatment by many authors [11-13]. Several studies have shown only negligible benefits from decompressive surgery (i.e., isolated laminectomy), associated or not with

* Correspondence: luciano.miller@uol.com.br

¹Hospital Estadual Mário Covas, Santo André, SP, Brazil

²Laboratório de Delineamento de Estudos e Escrita Científica, Departamento de Morfologia e Fisiologia, Faculdade de Medicina do ABC, Av. Príncipe de Gales, 821, Santo André, SP 09060-650, Brazil
Full list of author information is available at the end of the article

radiotherapy, when compared to radiotherapy alone for the treatment of pain and neurological dysfunction in patients [14]. The current dogma that dictates surgery for spinal metastatic disease as the procedure of choice to save the life of a patient has been established for some time.

However, previous studies have considered laminectomy inefficient in decompressing and resecting a tumor mass that is found in the vertebral body in most cases. Moreover, this procedure can lead to vertebral instability and deformities as well as the possible neurological and clinical deterioration of the patient. With recent advances in surgical techniques and the development of less invasive procedures, as well as earlier and more precise diagnosis, surgical approaches have shown improved outcomes in treating oncology patients, thereby making surgery a first-line treatment in selected cases [15].

Radiotherapy is still a fundamental therapeutic option for neoplasms sensitive to this treatment, such as lymphomas and multiple myelomas. This option is most commonly used with patients who have lesions at multiple levels or where surgery would be contraindicated by the clinical condition of the patient. There is still controversy with regard to the benefits of surgery, radiotherapy, chemotherapy or any combination of these treatments for different groups of patients [16].

Although previous studies have already investigated the radiation therapy effects on metastasis [17,18], its effects on immunological responses are hypothesized to impair the quality of life. Therefore, we aimed to analyze the progression of pain and of quality of life in patients undergoing radiotherapy with supportive treatment to treat SMD-related pain.

Methods

All patients considered for this study were diagnosed with SMD between January 2008 and January 2010. A total of 30 patients undergoing radiotherapy for pain management and to prevent the local progression of the tumor were included in the study. Patients that had previous decompressive or stabilization surgeries in addition to receiving radiotherapy were excluded.

Clinical outpatient evaluation was performed with the help of the Clinical Oncology team of the same service. Questionnaires were administered by the treating orthopedic physician specialized in spinal surgery. Patients enrolled in the study provided their informed and written consent. All procedures were approved by the Ethics Committee in Research of our University.


Details of other supportive treatments other than surgery and radiation therapy included steroid therapy (dexamethasone, 4 mg every 6 hours), spinal orthotics and physiotherapy.


The criteria used for radiotherapy indication were the following: patients selection included those treated with 20 Gy tumour dose, provided in five fractions, control of local pain that was not treatable by medication, patients with lesions in multiple levels, patients with pain and complete neurological deficits for more than 48 hours after neoplastic spinal cord compression and tumors sensitive to radiotherapy in patients without progressive neurological alteration during treatment. All patients were sent to the same radiotherapy clinic and underwent treatment with fractionated doses according to a specific protocol (20 Gy tumour dose in five fractions).

The questionnaire evaluated the pain in SMD subjects and its impact on quality of life at three different times: before radiotherapy, 30 days after the end of the treatment and 6 months after the end of the treatment. The visual analogue scale (VAS) [19] was used to measure pain, and the SF-36 questionnaire [20,21] was used to study possible quality of life variations. The SF-36 consists of 36 items that are grouped into the following eight areas: functional capacity, physical limitations, pain, overall health, vitality, social aspects, emotional limitations and mental health. Results were statistically evaluated using SPSS software (Statistical Package for Social Sciences, version 17.0) Statistical Package for Social Sciences software (SPSS, version 17.0) and were considered significant for $p < 0.05$.

Results

Thirty patients were chosen for the study, including 18 women (70%) and 12 men (30%). All patients participated in the evaluation performed 30 days after the end of radiotherapy. However, by the 6-month evaluation, there were 10 deaths, thereby resulting in a survival rate of 66.66%. The distribution of the primary sites of neoplasm afflicting these patients is presented in Figure 1. The patients were aged between 40 and 90 years old with an average age of 62.2 years old.





SMD involving three or fewer vertebrae was found in 14 cases (46.66%), whereas SMD that had disseminated (i.e., lesions in more than three vertebrae) was found in 16 patients (53.33%). Topography characteristic of restricted lesions was more frequently found in the thoracic spine (i.e., 8 cases, or 57.14% of the restricted lesions), although in 5 cases (35.71%) the lesions occurred in the lumbar spine. In 1 case (7.14%), the lesion occurred in the cervical spine.

The evaluation of pain according to the visual analogue scale (VAS) yielded an initial average value of 5.7 points. Thirty days after the end of radiotherapy, this was found to be 4.6 points on average. Six months after treatment, the average VAS score was 4.25 points (Figure 2), which is 25.43% lower than the initial value.

To describe and compare the observational periods regarding the variable VAS, we applied the *Friedman test*. The goal of this analysis was to verify possible differences among the three time points of observation (Table 1). Patients that died during the study were not included in this analysis.

Because we found significant difference between the three testing time points, we applied the *Wilcoxon test*, which was adjusted by a *Bonferroni correction*, in order to identify the observational periods that were significantly different (Table 2).

We observed the main changes in the VAS values when the pre-radiotherapy period was paired with the consecutive follow-up analyses, individually. Nevertheless, no statistically significant alteration in the VAS scores was

found when the 30-day post radiotherapy and 6-month post radiotherapy time points were compared.

To complete the quality of life evaluation using the SF-36 questionnaire, we compared the eight distinct subject areas across the observation time points. We applied the *Friedman test* to verify possible differences among the three observation time points.

Only in the areas of functional capacity (Figure 3) and social aspects (Figure 4) we found significant higher values at pre and 6 months post radiotherapy compared to 1 month post radiotherapy.

We found significant differences in these areas and we compared the three periods concomitantly. We applied the *Wilcoxon test*, which was adjusted by a *Bonferroni correction* in order to identify which observation points were significantly different (Tables 3 and 4).

The average score for functional capacity 6 months after radiotherapy was 38.75 points. This value is significantly higher than the average value obtained for functional capacity 30 days after the end of radiotherapy, which was 25 points. Analogously, the values for social aspects observed at the second and third time points were significantly different. Specifically, the average value for this parameter was 74.38 points 6 months post radiotherapy and was 50.63 points 30 days after the end of treatment.

We also performed statistical analyses to identify possible correlations between changes in pain levels and the SMD being restricted or disseminated, but we did not observe significant differences in the evolution of pain levels when comparing these two groups by either the VAS or SF-36 questionnaires.

Discussion

In this study we conducted a prospective analysis of pain and quality of life evolution in subjects undergoing radiotherapy with supportive treatment to treat pain caused by spinal metastatic disease. We found that radiotherapy and the related adjuvant treatments significantly improved the progression of pain at 30 days and 6 months after its application in the treatment of SMD.

Indication

The literature shows that a definitive cure is not an expected outcome for patients with SMD. The main goal of treatment is palliative, and life expectancy is generally

Table 1 Values of VAS for the analyzed time points

Variables	n	Average	Standard deviation	Min	Max	Median
VAS pre radiotherapy	20	5.7	3.05	0.00	10.00	6.5
VAS 1 month post radiotherapy	20	4.6*	2.52	0.00	10.00	5.00
VAS 6 months post radiotherapy	20	4.25*	3.16	0.00	10.00	4.00

*p=0.002: Different from VAS pre radiotherapy.

Table 2 Differences between the VAS values when comparing the time points analyzed in pairs

<i>Pair of variables</i>	<i>p</i>
VAS 1 month post radiotherapy vs. VAS pre radiotherapy	0.010
VAS 6 months post radiotherapy vs. VAS pre radiotherapy	0.009
VAS 6 months post radiotherapy vs. VAS 1 month post radiotherapy	0.546

(alpha for Bonferroni = 0.016952).

short, typically ranging from 4 to 15 months [20,22]. The main symptom of SMD is pain. It is presented as progressive night pains that are generally localized in segments of the thoracic spine with concomitant clinical deterioration. The pathophysiologic mechanisms of osseous pain secondary to neoplasms are multiple and become even more complex in the spine. These underlying factors include chemical mediators, increases in intraosseous pressure, microfractures, periosteum enlargement, pathological fractures and deformities [16,22].

Most patients with SMD and spinal cord compression syndrome (90%) present pain, while 47% present with symptomatic neurological alterations [23]. Moreover, only 11% to 34% of patients with spinal cord compression syndrome are able to walk at the time of diagnosis [24].

Our investigation indicates radiotherapy with supportive treatment as an efficient therapy. According to the literature, radiotherapy is indicated as the first-line therapy for SMD patients in the following situations: when complete neurological deficits are sustained for a period longer than 24 hours, when highly radiosensitive tumors (lymphoma, myeloma) are diagnosed, when there is involvement of multiple levels, when there is short life expectancy (less than 3 months) or when serious comorbidities contraindicate surgery [15]. Surgery to treat SMD is indicated in specific cases, such as neurologically

progressive deficits, pathologic deformities, resistance to radiotherapy and patients with an oncologic prognosis that is favorable.

Etiology and topography

Currently, there are three theories regarding the origin of extradural metastatic vertebral lesions: (1) progression of local disease, (2) retrograde dissemination via the spinal aivalvular venous plexus of Batson and (3) arterial embolisms [25-28].

All the patients' lesions were evaluated by magnetic resonance in our study and were all demonstrated to be extradural tumors. Classic studies involving autopsy have shown that the distribution of extradural metastases is more common in larger vertebrae. Therefore, the presence of SMD has been documented in the lumbar, thoracic and cervical regions with decreasing frequency [27,29]. Nevertheless, Papadopoulos et al. [28] showed that symptomatic lesions are most frequent in the thoracic region (70%), whereas the lumbar region represents approximately 20% cases and the cervical region about 10% [30]. Notably, our results are in agreement with the literature, whereby patients that presented restricted lesions (i.e., up to three vertebrae involved) were most frequently affected in the thoracic region. Overall, this group comprised 8 cases, or 57.14% of patients. We hypothesize that the small diameter of the vertebral canal in relation to the thoracic spinal cord caused this increase in symptoms [8].

The majority of published studies demonstrated that the major primary sites for extradural vertebral metastasis are the breast, lung and prostate, as well as lymphomas, multiple myelomas and renal cell tumors [27,29-32]. In our study, we observed a predominance of breast lesions (30%), followed by lymphomas and multiple myelomas (40%). Byrne et al. [31] observed that




Figure 3 Variation in the area of functional capacity using the SF-36.




Figure 4 Variation in the area of social aspects using the SF-36.

approximately 50% of vertebral metastases originated in one of the three following primary sites: breast, lung or prostate [33]. In a recent review, Klimo et al. [32] suggested that neoplasms originate from the following sites in order of decreasing frequency: renal tumors, gastrointestinal tissue, thyroid tissue, sarcomas and malignancies of lymphoreticular tissue (lymphomas and multiple myelomas) [34].

It is important to mention that the distribution of primary sites related to SMD may vary from one institution to another due to regional epidemiological differences and the specialty of the services involved. Our institution is a regional clinic for hematology, which justifies the greater portion of lymphoma and myeloma cases we encountered when compared to studies from other institutions.

Evaluation of SMD pain and quality of life with radiotherapy with supportive treatment

The literature suggests that the response to radiotherapy should be characterized in terms of pain relief and functional status [12,13]. In lesions sensitive only to radiotherapy, positive results occur in more than 80% of patients [23,24]. Lower rates of response are observed with lesions that are less sensitive to irradiation. As

reported by Jacobs and Perrin [19], more than 30% of patients showed improvement in neurological function due to epidural decompression and more than 60% achieved pain relief after radiotherapy [19,24]. Zaikova et al. [33] conducted a study involving an analysis of pain response among 355 patients that underwent radiotherapy for SMD. After 2 months of treatment, a favorable response (partial or total pain relief) was noted in 37% of patients, which is a lower percentage than is observed with osseous metastasis patients in general. Taken together, these results suggest that the presence of pathophysiological pain is more complex in nature than the pain associated with vertebral metastatic disease.

Chow et al. [34] reported strong positive response rates to radiotherapy in the treatment of osseous metastasis, which ranged from 60 to 90%. Nonetheless, the lowest rates of successful pain relief were observed in patients with vertebral lesions in other skeletal topographies [35,36]. Maranzano et al. administered radiotherapy to treat SMD in 209 patients. After an average of 49 months, 71% of the patients with pain (98% of the studies participants) showed relief from this symptom [13,21,35].

Our investigation recorded the regression of pain with an outpatient evaluation that was quantified by the VAS

Table 3 Wilcoxon test to evaluate the area of functional capacity at different time points

Pairs of variables	p
SF-36 functional capacity 1 month post radiotherapy vs. SF-36 functional capacity pre radiotherapy	0.067
SF-36 functional capacity 6 months post radiotherapy vs. SF-36 functional capacity pre radiotherapy	0.690
SF-36 functional capacity 6 months post radiotherapy vs. SF-36 functional capacity 1 month post radiotherapy	0.011

(alpha value for Bonferroni = 0.016952).

Table 4 Wilcoxon test to evaluate the area of social aspects at different time points

Pairs of variables	p
SF-36 social aspects 1 month post radiotherapy vs. SF-36 social aspects pre radiotherapy	0.196
SF-36 social aspects 6 months post radiotherapy vs. SF-36 social aspects pre radiotherapy	0.095
SF-36 social aspects 6 months post radiotherapy vs. SF-36 social aspects 1 month post radiotherapy	0.001

(alpha value for Bonferroni = 0.016952).

questionnaire. At 6 months after the radiotherapy, the average score was 25.43% lower than the initial value. This level of pain control is lower than the responses found in the literature. It is possible that the reduced pain control observed in our study was due to the fact that a higher portion of the patients we treated were afflicted with the advanced stages of the disease. The fact that 33% patients died within the 6 months following treatment illustrates this point.

No significant variations in paired evaluations of the data from the 30-day and 6-month post radiotherapy time points were found. On the other hand, there was a significant improvement in pain relief during the 30 days following radiotherapy that evolved over the course of the study.

The analysis of quality of life through the SF-36 questionnaire did not show significant results at the 6-month evaluation period. The only exceptions were in the areas of functional capacity and social aspects, which showed slight improvements when the 6-month time point was compared with the 30-day time point.

Analysis of the pain section of the SF-36 questionnaire did not indicate a significant improvement in pain levels, as was reported by the VAS questionnaire. This difference may have been because the latter consisted of an evaluation guided by the symptoms related to the spine region that was irradiated, while the evaluation of pain according to the SF-36 questionnaire was influenced by symptoms related to the overall clinical deterioration of the patient.

Analgesic medication was also used as an important resource to help control pain. Indeed, several studies have avoided the topic of analgesic usage for the further relief of pain in studies of radiotherapy due to the complexity of the prospective analysis required [2,36,37]. Nevertheless, we did not remove medication from the patients to avoid health's impairment of the patient.

The standard dose for efficacious irradiation in SMD cases consists of 20 to 30 Gy given over 5 to 10 sessions at the area of vertebral involvement. However, several variations of this regimen have been utilized [38,39]. Although we administered the radiotherapy in fractional doses (20 Gy in five fraction), the majority of studies in the literature do not show differences in pain relief from vertebral metastatic disease when radiotherapy is performed as a single dose or if it is fractionated [37,39,40].

We did not report significant toxicity in patients that underwent radiotherapy in the present study. Complications from the irradiation of the spine are generally mild. Esophagitis induced by radiation may occur in lesions of the superior thoracic region, while irradiation of the inferior thoracic region and lumbar superior can cause nausea and vomiting. These adverse reactions are more common at the

beginning of treatment and are regularly controlled by medication. In contrast, radiation-induced myelopathy is a complication that occurs more commonly in the latter during treatment, and although it is rare, it is extremely serious [37,39,40].

Some points are important to be raised. The supportive treatment other than surgery and radiation therapy included spinal orthotics, dexamethasone and physiotherapy. Unfortunately, we were not able to exclude these variables, since the withdrawal of those treatments would negatively affect the patient's health. Therefore, we indicate radiation therapy accompanied with those supportive treatment as effective. Although there are specific questionnaire for cancer's patient, we applied only the SF-36 questionnaire. However, this questionnaire is well accepted in the literature and it was applied in this disease [41-43]. Patients selection included those treated with a 20 Gy tumour dose, provided in five fractions according to a previous study [44]. This method was decided in order to avoid a possible influence of different radiotherapy regimens, i.e. 20 Gy in 5 fractions and 30 Gy in 10 fractions.

Although the use of radiation therapy for bony metastases with or without surgery was presented in previous studies, some studies still consider it as malefic regarding the immunological responses [45,46]. Therefore, our study provided important information to support this type of treatment. Our findings are relevant to the literature, since quality of life is a very important issue to be discussed in order to improve clinical practice [47-51].

Conclusion

Radiotherapy with supportive treatment resulted in a significant improvement in the progression of pain levels at 30 days and 6 months after its application in the treatment of SMD. The quality of life evaluation showed only significant improvement in functional capacity and social aspects. Thus, radiotherapy with supportive treatment is indicated a useful tool for the control of pain in patients with SMD.

Abbreviations

SMD: Spinal metastatic disease; VAS: Visual analogue scale.

Competing interest

The authors declare that they have no competing interests.

Authors' contribution

ESVF, LCA, GHV, DIGC, FHU, GSLF, VEV, CBMM, RW, ENF, MRN, LMR participated in the acquisition of data and revision of the manuscript. ESVF, GHV, DIGC, FHU, GSLF and LMRR conceived the study, determined the design, performed the statistical analysis, interpreted the data and drafted the manuscript. VEV, ENF and LCA determined the design, interpreted the data and drafted the manuscript. All authors read and gave final approval for the version submitted for publication.

Acknowledgements

This study received financial support from Faculdade de Medicina do ABC.

Author details

¹Hospital Estadual Mário Covas, Santo André, SP, Brazil. ²Laboratório de Delineamento de Estudos e Escrita Científica. Departamento de Morfologia e Fisiologia, Faculdade de Medicina do ABC, Av. Príncipe de Gales, 821, Santo André, SP 09060-650, Brazil. ³Faculdade de Filosofia e Ciências, Universidade Estadual Paulista, UNESP, Av. Higino Muzzi Filho, 737, Marília, SP 17.525-900, Brazil. ⁴Escola de Artes, Ciências e Humanidades da Universidade de São Paulo, São Paulo, Brazil.

Received: 23 April 2012 Accepted: 8 February 2013

Published: 18 February 2013

References

1. Loblaw DA, Lapierre NJ, Mackillop WJ: A population-based study of malignant spinal cord compression in Ontario. *Clin Oncol (R Coll Radiol)* 2003, 15:211–217.
2. Conway R, Graham J, Kidd J, Levack P, Scottish Cord Compression Group: What happens to people after malignant cord compression? Survival, function, quality of life, emotional well-being and place of care 1 month after diagnosis. *Int Oncol (R Coll Radiol)* 2007, 19:56–62.
3. Abbas H, Elyamany A, Salem M, Salem A, Biniaz S, Gamal B: The optimal sequence of radiotherapy and chemotherapy in adjuvant treatment of breast cancer. *Int Arch Med* 2011, 4:35.
4. Rades D, Stalpers LJ, Veninga T, Schulte R, Hoskin PJ, Obral N, Bajrovic A, Rudat V, Schwarz R, Hulshof MC, Poortmans P, Schild SE: Evaluation of five radiation schedules and prognostic factors for metastatic spinal cord compression. *J Clin Oncol* 2005, 23:3366–3375.
5. Maximous DW, Abdel-Wanis ME, El-Sayed MI, Abd-Elsayed AA: Preoperative gemcitabine based chemo-radiotherapy in locally advanced non metastatic pancreatic adenocarcinoma. *Int Arch Med* 2009, 2:7.
6. El-Sayed MI, Ali AM, Sayed HA, Zaky EM: Treatment results and prognostic factors of pediatric neuroblastoma: a retrospective study. *Int Arch Med* 2010, 3:37.
7. Gokslan ZL, York JE, Walsh GL, McCutcheon IE, Lang FF, Putnam JB Jr, Wildrick DM, Swisher SG, Abi-Said D, Sawaya R: Transthoracic vertebrectomy for metastatic spinal tumors. *J Neurosurg* 1998, 89:599–609.
8. Grant R, Papadopoulos SM, Greenberg HS: Metastatic epidural spinal cord compression. *Neurol Clin* 1991, 9:825–841.
9. Heldmann U, Myschitzky PS, Thomsen HS: Frequency of unexpected multifocal metastasis in patients with acute spinal cord compression. Evaluation by low-field MR imaging in cancerpatients. *Acta Radiol* 1997, 38:372–375.
10. Sundaresan N, DiGiacinto GV, Hughes JE, Cafferty M, Vallejo A: Treatment of neoplastic spinal cord compression: results of a prospective study. *Neurosurgery* 1991, 29:645–650.
11. Ippolito V, Micheletti E, Saccalani M, Barbera F, Tonoli S, Motta C: Radiotherapy and spinal brace: still first-choice treatment for vertebral metastases from breast cancer. *Chir Organi Mov* 1998, 83:177–183.
12. Katagiri H, Takahashi M, Inagaki J, Kobayashi H, Suguri H, Yamamura S, Iwata H: Clinical results of nonsurgical treatment for spinal metastases. *Int J Radiat Oncol Biol Phys* 1998, 42:1127–1132.
13. Maranzano E, Latini P, Checcaglini F, Ricci S, Panizza BM, Aristei C, Perrucci E, Beneventi S, Corgna E, Tonato M: Radiation therapy in metastatic spinal cord compression. A prospective analysis of 105 consecutive patients. *Cancer* 1991, 67:1311–1317.
14. Black P: Spinal metastasis: current status and recommended guidelines for management. *Neurosurgery* 1979, 5:726–746.
15. Steinmetz MP, Mekhail A, Benzel EC: Management of metastatic tumors of the spine: strategies and operative indications. *Neurosurg Focus* 2001, 11:e2.
16. Wise JJ, Fischgrund JS, Herkowitz HN, Montgomery D, Kurz LT: Complication, survival rates, and risk factors of surgery for metastatic disease of the spine. *Spine* 1999, 24:1943–1951.
17. Cheng J, Kong L, Huang W, Li B, Li H, Wang Z, Zhang J, Zhou T, Sun H: Explore the radiotherapeutic clinical target volume delineation for thoracic esophageal squamous cell carcinoma from the pattern of lymphatic metastases. *J Thorac Oncol* 2012, Dec 20. [Epub ahead of print].
18. Spraker MB, Bair E, Bair R, Connell PP, Mahmood U, Koshy M: An analysis of patient characteristics and clinical outcomes in primary pulmonary sarcoma. *J Thorac Oncol* 2012, Dec 20. [Epub ahead of print].
19. Jacobs WB, Perrin RG: Evaluation and treatment of spinal metastases: an overview. *Neurosurg Focus* 2001, 11:e10.
20. Ware JE Jr, Sherbourne CD: The MOS 36-item short-form health-survey (SF-36). I. Conceptual framework and item selection. *Med Care* 1992, 30:473–83.
21. Latini P, Maranzano E, Ricci S: Role of radiotherapy in metastatic spinal cord compression: preliminary results from a prospective trial. *Radiother Oncol* 1989, 15:227–33.
22. Weigel B, Maghsudi M, Neumann C, Kretschmer R, Müller FJ, Nerlich M: Surgical management of symptomatic spinal metastases. Postoperative outcome and quality of life. *Spine* 1999, 24:2240–6.
23. Janjan NA: Radiotherapeutic management of spinal metastases. *J Pain Symptom Manage* 1996, 11:47–56.
24. Janjan NA: Radiation for bone metastases: conventional techniques and the role of systemic radiopharmaceuticals. *Cancer* 1997, 80:1628–45.
25. Arguello F, Baggs RB, Duerst RE, Johnstone L, McQueen K, Frantz CN: Pathogenesis of vertebral metastasis and epidural spinal cord compression. *Cancer* 1990, 65:98–106.
26. Batson OV: The function of the vertebral veins and their role in the spread of metastases. *Ann Surg* 1940, 112:138–49.
27. Abdu WA, Provencier M: Primary bone and metastatic tumors of the cervical spine. *Spine* 1998, 23:2767–77.
28. Onimus M, Papin P, Gangloff S: Results of surgical treatment of spinal thoracic and lumbar metastases. *Eur Spine J* 1996, 5:407–11.
29. Hammerberg KW: Surgical treatment of metastatic spine disease. *Spine* 1992, 17:1148–53.
30. Sioutos PJ, Arbit E, Meshulam CF, Galichich JH: Spinal metastases from solid tumors. Analysis of factors affecting survival. *Cancer* 1995, 76:1453–9.
31. Byrne TN: Spinal cord compression from epidural metastases. *N Engl J Med* 1992, 327:614–9.
32. Klimo P Jr, Kestle JR, Schmidt MH: Treatment of metastatic spinal epidural disease: a review of the literature. *Neurosurg Focus* 2003, 15:E1.
33. Zaikova O, Fossa SD, Kongsgaard U, Kvaloy S, Giercksky KE, Skjeldal S: Pain after palliative radiotherapy for spine metastasis. *Clin Oncol (R Coll Radiol)* 2010, 22:828–36.
34. Chow E, Harris K, Fan G, Tsao M, Sze WM: Palliative radiotherapy trials for bone metastases: a systematic review. *J Clin Oncol* 2007, 25:1423–36.
35. Maranzano E, Latini P: Effectiveness of radiation therapy without surgery in metastatic spinal cord compression: final results from a prospective trial. *Int J Radiat Oncol Biol Phys* 1995, 32:959–67.
36. Maranzano E, Bellavita R, Rossi R, De Angelis V, Frattegiani A, Bagnoli R, Mignogna M, Beneventi S, Lupattelli M, Ponticelli P, Biti GP, Latini P: Short-course versus split-course radiotherapy in metastatic spinal cord compression: results of a phase III, randomized, multicenter trial. *J Clin Oncol* 2005, 23:3358–65.
37. van der Linden YM, Lok JJ, Steenland E, Martijn H, van Houwelingen H, Marijen CA, Leer JW, Dutch Bone Metastasis Study Group: Single fraction radiotherapy is efficacious: a further analysis of the Dutch Bone Metastasis Study controlling for the influence of retreatment. *Int J Radiat Oncol Biol Phys* 2004, 59:528–537.
38. Maranzano E, Latini P, Perrucci E, Beneventi S, Lupattelli M, Corgna E: Short-course radiotherapy (8 Gy X 2) in metastatic spinal cord compression: an effective and feasible treatment. *Int J Radiat Oncol Biol Phys* 1997, 38:1037–44.
39. Hartsell WF, Scott CB, Bruner DW, Scarantino CW, Ivker RA, Roach M 3rd, Suh JH, Demas WF, Movsas B, Petersen IA, Konski AA, Cleeland CS, Janjan NA, DeSilvio M: Randomized trial of short- versus long-course radiotherapy for palliation of painful bone metastases. *J Natl Cancer Inst* 2005, 97:798–804.
40. Kaasa S, Brenne E, Lund JA, Fayers P, Falkmer U, Holmberg M, Lagerlund M, Bruland O: Prospective randomized multicenter trial on single fraction radiotherapy (8 Gy x 1) versus multiple fractions (3 Gy x 10) in the treatment of painful bone metastases. *Radiat Oncol* 2006, 79:278–84.
41. Le Borgne G, Mercier M, Woronoff AS, Guizard AV, Abeillard E, Caravati-Jouvenceaux A, Klein D, Velten M, Joly F: Quality of life in long-term cervical cancer survivors: A population-based study. *Gynecol Oncol* 2012, Dec 29. [Epub ahead of print].

42. Tian J, Hong JS: Validation of the Chinese version of the Resilience Scale and its cutoff score for detecting low resilience in Chinese cancer patients. *Support Care Cancer* 2012, Dec 30. [Epub ahead of print].
43. Musarezaie A, Ghasemi TM, Esfahani HN: Investigation the quality of life and its relation with clinical and demographic characteristics in women with breast cancer under chemotherapy. *Int J Prev Med* 2012, 3:853–9.
44. Yarnold JR: 8 Gy single fraction radiotherapy for the treatment of metastatic skeletal pain: randomised comparison with a multifraction schedule over 12 months of patient follow-up. Bone Pain Trial Working Party. *Radiother Oncol* 1999, 52:111–21.
45. Seung SK, Curti BD, Crittenden M, Walker E, Coffey T, Siebert JC, Miller W, Payne R, Glenn L, Bageac A, Urba WJ: Phase 1 study of stereotactic body radiotherapy and interleukin-2-tumor and immunological responses. *Sci Transl Med* 2012, 4:137ra74.
46. Ivanov VN, Partridge MA, Huang SX, Hei TK: Suppression of the proinflammatory response of metastatic melanoma cells increases TRAIL-induced apoptosis. *J Cell Biochem* 2011, 112:463–75.
47. Centofanti G, Fujii EY, Cavalcante RN, Bortolini E, de Abreu LC, Valenti VE, Pires AC, Macedo H Jr, Yamazaki YR, Audi SG, Cisternas JR, Breda JR, Pereira VX, Fujiki EN, Correa JA: An experience of vascular access for hemodialysis in Brazil. *Int Arch Med* 2011, 4:16.
48. Vertamatti MAF, Abreu LC, Drezett J, Valenti VE, Barbosa CP: Time lapsed between sexual aggression and arrival at the Brazilian health service. *J Hum Growth Develop* 2013, 23:46–51.
49. Atrash HK, Carpentier R: The evolving role of public health in the delivery of health care. *J Hum Growth Develop* 2013, 22:396–399.
50. Carrasco JM, Pérez-Gómez B, García-Mendizábal MJ, Lope V, Aragónés N, Forjaz MJ, Guallar-Castillón P, López-Abente G, Rodríguez-Artalejo F, Pollán M: Health-related quality of life and mental health in the medium-term aftermath of the Prestige oil spill in Galiza (Spain): a cross-sectional study. *BMC Publ Health* 2007, 7:245.
51. Oliveira DKS, do Nascimento DDG, Marcolino FF: Perceptions of family caregivers and professionals in the family health strategy related to the care and neuropsychomotor development of children. *J Hum Growth Develop* 2013, 22:142–150.

doi:10.1186/1755-7682-6-6

Cite this article as: Valesin Filho et al.: Pain and quality of life in patients undergoing radiotherapy for spinal metastatic disease treatment. *International Archives of Medicine* 2013 6:6.

Submit your next manuscript to BioMed Central and take full advantage of:

- Convenient online submission
- Thorough peer review
- No space constraints or color figure charges
- Immediate publication on acceptance
- Inclusion in PubMed, CAS, Scopus and Google Scholar
- Research which is freely available for redistribution

Submit your manuscript at
www.biomedcentral.com/submit

