



Published in final edited form as:

J Nat Prod. 2011 September 23; 74(9): 1842–1850. doi:10.1021/np2000667.

Stereochemical Analysis of Leubethanol, an Anti-TB Active Serrulatane, from *Leucophyllum frutescens*

Gloria M. Molina-Salinas^{†,‡,⊕}, Verónica M. Rivas-Galindo^{†,⊕}, Salvador Said-Fernández^{‡,Δ}, David C. Lankin[§], Marcelo A. Muñoz^{||}, Pedro Joseph-Nathan[∇], Guido F. Pauli^{§,*}, and Noemí Waksman^{†,*}

[†]Department of Analytical Chemistry, Faculty of Medicine, Universidad Autónoma de Nuevo Leon, P.O. Box 2316, Sucursal Tecnológico, Monterrey, N.L., 64841 México

[‡]División de Biología Celular y Molecular, Centro de Investigación Biomédica del Noreste, Instituto Mexicano del Seguro Social, Monterrey, N.L., 64720 México

[§]Department of Medicinal Chemistry & Pharmacognosy, PCRPS, and Institute for Tuberculosis Research, College of Pharmacy, University of Illinois at Chicago, Chicago, IL 60612

^{||}Instituto de Química, Facultad de Ciencias, Universidad Austral de Chile, Valdivia, Chile

[∇]Departamento de Química, Centro de Investigación y de Estudios Avanzados del Instituto Politécnico Nacional, Apartado 14-740, México, D. F., 07000 México

Abstract

Bioactivity-guided fractionation of the methanolic root bark extract of *Leucophyllum frutescens* (Berl.) I.M. Johnst. led to the identification of leubethanol (**1**), a new serrulatane-type diterpene with activity against both multi drug-resistant and drug-sensitive strains of virulent *Mycobacterium tuberculosis*. Leubethanol (**1**) was identified by 1D/2D NMR data, as a serrulatane closely related to erogorgiane (**2**), and exhibited anti-TB activity with minimum inhibitory concentrations in the range 6.25–12.50 µg/mL. Stereochemical evidence for **1** was gleaned from 1D and 2D NOE experiments, ¹H-NMR full spin analysis, as well as by comparison of the experimental vibrational circular dichroism (VCD) spectrum to density functional theory calculated VCD spectra of two diastereomers.

Keywords

Leucophyllum frutescens; Scrophulariaceae; diterpenes; structure elucidation; anti-tuberculosis activity; drug-resistant; full spin analysis; circular dichroism

Introduction

Infection with *Mycobacterium tuberculosis* affects about one third of the human population.¹ In 2008, 11.1 million new cases and 1.8 million deaths from tuberculosis (TB) involving HIV-positive patients were reported.¹ One of the major problems is the increase of

^{*}To whom correspondence should be addressed. Tel & Fax +11 (5281) 8329 4185 (NW), Tel (312) 355-1949. Fax (312)-355-2693 (GFP). nwaksman@gmail.com (NW) and gfp@uic.edu (GFP).

^ΔActual address: Biochemistry Department, Facultad de Medicina, UANL

[⊕]Equally contributing authors.

Supporting Information

NMR spectra of **1** are available free of charge via the Internet at <http://pubs.acs.org>.

multidrug and extensive resistant strains of *M. tuberculosis*,² which leads to an urgent need of the synthesis of new anti-tuberculosis drugs^{3,4} or search from natural sources.^{5,6} Medicinal plants have attracted considerable attention as sources of novel natural products with intriguing novel structures and useful biological activities, including anti-TB activity.⁷

From the evaluation of the in vitro anti-TB activity of a panel of Mexican plant extracts, the MeOH extract obtained from the root bark of *Leucophyllum frutescens* (Berl.) I.M. Johnston. (Scrophulariaceae) was previously found to be the most active.⁸ Known as cenizo, *L. frutescens* is a medicinal plant of Mexican traditional medicine, which has long been used to treat lung complaints and specifically tuberculosis. A review of the literature revealed few phytochemical reports for the plant genus, with only one report each on the isolation of phytotoxic lignans from the species⁹ and on calmodulin inhibitors from the closely related *L. ambiguum*.¹⁰ Our work was aimed at the bioassay-guided isolation of the anti-TB active principles of *L. frutescens* and their detailed structure elucidation, including ¹³C INADEQUATE and ¹H NMR full spin analysis, as well as by density functional theory (DFT) calculated vibrational circular dichroism (VCD) spectra of two diastereomers, which were compared *in silico* to the experimental VCD spectrum of **1**.

Results and Discussion

The anti-TB active MeOH extract obtained from the root bark of *L. frutescens* was subjected to a bioassay-guided fractionation. After solvent partitioning, which concentrated the activity mainly in the *n*-hexane phase, this fraction was further purified by means of VLC and LPLC, to afford the active compound **1**.

Compound **1** was obtained as colorless oil. HR-EIMS established a molecular formula of C₂₀H₃₀O. The UV maximum at 280 nm indicated the presence of an aromatic ring. The ¹H NMR spectrum showed two *meta*-coupled protons at δ 6.448 and 6.615 indicating the presence of a 1,2,3,5-tetrasubstituted benzene ring. Other ¹H NMR features (Table 1) suggested that this compound was of terpenoid origin due to the presence of two methyl doublets at δ 0.984 and 1.224; a sharp three-proton singlet at δ 2.270 characteristic of an aromatic methyl; two broad methyl singlets at δ 1.582 and 1.683, characteristic of olefinic methyls, and two complex multiplets at δ 2.590 and 3.078, which suggested two benzylic hydrogens. The H,H-COSY and HMQC spectra indicated the presence of a single aliphatic spin system CH₃-CH-(CH₂)₂-CH-CH-(CH₃)-(CH₂)₂-CH=C(CH₃)₂. An olefinic proton at δ 5.018 exhibited cross peaks with both the olefinic methyls as well as one methylene group (δ 2.010 and 1.834, respectively). The latter methylene protons exhibited cross-peaks with a second methylene (δ 1.314 and 1.107) group, which in turn showed cross-peaks with a methine proton at δ 1.902; this methine proton showed correlations with the benzylic proton at δ 2.590 and a Me group at δ 0.984. The proton at δ 2.590 exhibited cross-peaks with the methylene protons at δ 1.870 and 1.740, which in turn showed cross-peaks with another methylene group at δ 1.967 and 1.520; the last methylene group showed cross-peaks with a methine proton at δ 3.078. Finally, coupling between this methine proton and a methyl group at δ 1.224 was observed. The ¹³C NMR spectrum exhibited 20 signals of a diterpenoid, including five methyl, four methylene, six methine and five quaternary carbons, as determined by a DEPT experiment. Their chemical shift values and multiplicity confirmed the presence of the tetrasubstituted aromatic ring. HMQC and HMBC experiments (Figure 1) were used to determine unambiguously the structure of the planar moiety and its attached aliphatic residues. Spectroscopic evidence was in agreement with compound **1** being a diterpene with a serrulatane-type skeleton, similar to erogorgiaene,^{11,12} bearing an OH substitution at C-7. A ¹³C-¹³C INADEQUATE experiment performed with **1** confirmed most of the C-C connectivity (Figure 2).

In order to establish the absolute configuration, a two-prong approach was taken: (a) NOE measurements (Figure 3) as well as a full spin analysis of the fingerprint characteristics of ^1H NMR spectra resulting in full J coupling information provided evidence for the relative configuration (Figure 4); (b) VCD experiments led to the assignment of absolute configuration. For the former (a), taking into consideration the severe overlap in the upper field spectral range, high-resolution spectra were acquired at 900 MHz, and a full spin analysis was carried out by iteration of the ^1H NMR spectra against the full δ/J parameter set using the PERCH tool which ultimately led to the determination of all spin-spin coupling constants (J). The methodology of full ^1H NMR spin analysis has been pioneered by Laatikainen, Niemitz, and co-workers, and previously been documented for a few natural products including alkaloids and mono-, sesqui-, and diterpenes.^{13–17} The method involves iterated quantum-mechanical simulation of the entire spin system of the molecule until full congruence is achieved between the experimental and simulated spectrum.

The results of the ^1H full spin analysis (900 MHz) of **1** are summarized in Table 1 and Figure 3. The coupling constants for proton pairs H-1/H-2_{ax} and H-1/H-2_{eq} were 6.16 and 2.58 Hz, respectively, and are consistent with the cyclohexene ring adopting a distorted chair conformation with H-1 being pseudo-equatorial. In an analogous way, as the couplings of H-4/H-3_{ax} and H-4/H-3_{eq} were 6.23 and 2.95 Hz respectively, H-4 must be also pseudo-equatorial. Therefore, H-1 and H-4 are located on opposite faces of the molecule, consistent with 1*S*,4*R* or 1*R*,4*S* relative configuration. This was further supported by the NOE correlations of **1** (Figure 3): Irradiation of Me-20 (δ 1.224, d) reduced the intensity of signals at δ 1.520 (H-2_{eq}) and 1.870 (H-3_{ax}), confirming that they are co-facial.

Considering the negative specific rotation of **1**, the configuration at C-1 and C-4 was assigned as 1*R*,4*S*, consistent with the rotation reported for (–)-erogorgiaene (*syn. ent*-erogorgiaene) isolated from the brown alga, *Dictyota dichotoma*.¹⁸ Due to the conformational flexibility of the aliphatic chain, it was problematic to determine the configuration at C-11 by means of NMR and molecular modeling. However, it was observed that the natural serrulatanes previously isolated from sea organisms exhibit an Me-18 signal in the ^1H NMR spectrum at relative high fields (δ 0.64–0.79),^{11,12,18–20} while in serrulatanes isolated from terrestrial plants the analogous signals appear at lower field (δ 0.95–1.12 ppm).^{21–24} Harmat and co-workers²⁵ assigned the stereochemistry at C-11 in different synthetic diastereomers of erogorgiaene, taking into consideration the position of the signal from the C-18 methyl group. They concluded that the position of this signal was dependent of the spatial relationship between H-4 and Me-18 (*syn* vs. *anti*), which can be visualized when drawing the serrulatanes as presently shown for **1** and **2**. In the *syn* compounds, Me-18 appears at higher fields, compared to lower field signals for the *anti*-series compounds. Therefore, considering that in **1** the Me-18 group appears at δ 0.984, H-4 and CH₃-18 must be *anti* which suggests the *S* configuration at C-4. In order to test this hypothesis and confirm the NMR interpretation, a VCD analysis was performed, which eventually confirmed the 1*R*,4*S*,11*S* absolute configuration for compound **1**.

Compound **1** is an interesting molecule for VCD studies as it contains the C₈ chain frequently found in terpenoids, ranging from bisabolene sesquiterpenes like perezone,²⁶ to sterols like desmosterol, which is the last biogenetic intermediate in the biosynthesis of cholesterol.²⁷ Due to the high conformational mobility of the side chain, the C-11 configuration is difficult to determine, and therefore both C-11 diastereomers of **1** were considered. This, in turn, provides evidence for the sensitivity of the VCD method, since it rarely has been used to study epimers.²⁸

Since rotational strength modes of VCD spectra depend on molecular conformation, initial searches for **1** and its 11*R* diastereomer were performed using a computational Monte Carlo

MMFF94 molecular mechanics method, in a 10 kcal/mol gap using Spartan'04 (Wavefunction, Irvine CA, USA), which provided 199 and 193 conformers, respectively. All conformers were then submitted to DFT for single point calculations using the B3LYP functional and the DGDZVP basis set implemented in Gaussian 03 (Gaussian Inc., Pittsburg, PA, USA) which provided 54 and 40 conformers for **1** and its 11*R* epimer, respectively, in a 3 kcal/mol gap. These conformers were further optimized at 25 °C and 1 atmosphere by DFT//B3LYP/DGDZVP calculations to provide 13 and 23 conformers for **1** and its 11*R* epimer, respectively, in a 2 kcal/mol range. The number of conformers was further reduced by eliminating those contributing less than 2% to the overall distribution. Finally, frequency calculations were performed for 10 and 13 conformers whose free energy ($\Delta G = RT \ln K$) at 25 °C and population, for **1** and its C-11 epimer, are listed in Tables 2 and 3, while conformational freedom is shown in Figures 5 and 6, respectively.

Inspection of Tables 2 and 3 reveals that **1** has significantly reduced conformational mobility than its 11*R* counterpart, not only from the fact that the former shows 10 conformers compared to 13 for the latter, but also from the relative conformational freedom. In **1** three conformers account for two thirds of the conformational distribution, while for the 11*R* epimer five conformers are required for an equivalent conformational abundance. Inspection of Figures 5 and 6 reveals the steric interaction derived from the *peri* arrangement of the C-8 hydroxy group and the C-1 secondary methyl group is responsible for the pseudo-axial nature of the secondary methyl group on a distorted chair conformation, in agreement with the ¹H NMR observations.

The IR and VCD calculated spectra were obtained from the dipole and rotational strengths using Lorentzian shapes, with a bandwidth of $\nu = 4 \text{ cm}^{-1}$. The average data for **1** and its 11*R* diastereomer, calculated according to the abundances reported in Tables 2 and 3, were used for the spectral comparison shown in Figure 6, from which the 11*S* absolute configuration seems evident. However, a reliable band-to-band comparison of observed and calculated VCD spectra is crucial for configuration assignment since measured IR and VCD frequencies derive from an anharmonic force field while calculated frequencies derive from a harmonic force field. Thus, the latter data are normally scaled using an anharmonicity factor which was obtained using the commercial available confidence level algorithm software (BioTools Co., Jupiter, FL, 33458 USA) based on neighbourhood similarity²⁹ that we recently applied for the study of 1*R*(-)-myrtenal at several levels of theory.³⁰ This algorithm uses a correlation function that describes the integrated overlap of the experimental and theoretical data as a function of a relative vibrational frequency shift, being the optimal shift known as the anharmonicity (*anH*) factor. Taking advantage of this methodology, the calculated spectra of both diastereomers were compared *in silico* to the experimental IR and VCD spectra of **1**, the pertinent results being summarized in Table 4. Although the *anH* factor, a measure of band alignment of experimental and theoretical absorption signals, is similar, the IR spectral similarity (S_{IR}) indicates the 11*S* configuration since for **1** it provides a value of 89.9 while for the 11*R* isomer no similarity is found. Further comparison data (Table 4) reveal that the VCD spectral similarity (S_E) for 11*S*-**1** is 70.6 and that of its enantiomer (S_{-E}) is 13.6, while for the 11*R* alternative those values are 62.9 and 18.5, respectively, confirming the 11*S* absolute configuration. Finally, the calculated confidence level for **1** is 100%, in contrast to 42%, for the 11*R* epimer, a reasonable value for a comparison where the absolute configuration of two of three stereogenic centers are correct.

Elisabethanol with the same planar structure as compound **1**, is reported³¹ to have 1*S*,4*R*, 11*S* configuration and, therefore, must be a different compound. Moreover, no information is available regarding the purification methodology or spectroscopic data obtained of elisabethanol, which essentially precludes structure dereplication. Taking these facts into

consideration as well as the differences in stereochemistry, **1** must be considered a new natural product. Similar stereochemical considerations have been reported for synthetic congeners.^{36,37} In order to reflect the structural analogy with elisabethanol, the trivial name leubethanol is proposed for **1**.

Of interest from a biological perspective, leubethanol exhibits significant activity against MDR resistant strains of *M. tuberculosis* such as CIBIN/UMF15:99, a clinical isolate that is resistant to all five first-line anti-TB drugs, streptomycin, isoniazid, rifampin, ethambutol, and pyrazinamide. Table 3 shows the MIC values of **1** against the clinical isolates of *M. tuberculosis*, compared with those of first-line anti-TB drugs. Another noteworthy feature of leubethanol is its increased activity against MDR strains compared to the drug-sensitive strain, H37Rv. Anti-TB activities with MIC values of around 10 µg/mL are in line with literature reports on serrulatanes: the structurally related 7-hydroxyerogorgiaene from the gorgonian octocoral, *Pseudoerogorgia elisabethae*, showed 77 % growth inhibition of *M. tuberculosis* H37Rv at 6.25 µg/mL.¹¹ While MIC values in the 10 µg/mL range are considered moderate activities in TB drug discovery projects, the potency of leubethanol is still noteworthy as it is 16 times more potent against MDR *M. tuberculosis* than the first-line anti-TB drugs streptomycin and rifampin. Therefore, the serrulatane diterpenes such as **1** represent potential leads for the development of anti-TB treatment of drug-resistant TB infections.

Experimental Section

General Experimental Procedures

UV Vis spectra were measured on a Beckman DU 7500 spectrophotometer. Optical rotation and IR were determined with a Perkin Elmer 341 polarimeter and a Perkin Elmer FT-IR Spectrum-One spectrometer, respectively. ¹H NMR spectra were measured at 400 and 900 MHz on a DPX 400 and Avance 900 MHz NMR spectrometers (Bruker, Zurich, Switzerland) in CDCl₃ as the solvent and ¹³C NMR spectra were measured at 100 MHz on a DPX 400 NMR spectrometer (Bruker, Zurich, Switzerland) in CDCl₃ as the solvent. Chemical shifts are expressed in ppm and were referenced to TMS using the signals from the solvent (δ_{H} 7.260, δ_{C} 77.000 relative to TMS) as internal standard. NMR data were processed with NUTS (Pro version for Microsoft Windows, Acorn NMR Co., USA). The ultra high-resolution 900 MHz ¹H NMR spectra were acquired with 64 K data points and zero-filled to 256 K data points prior to applying window functions (LG, EM) and Fourier transformation. The resulting digital resolution (real data points) exceeded 0.0001 ppm. Considering the higher order spin systems, the magnitude of small *J* couplings, and the total RMS of the spectral iteration (0.21 Hz), chemical shifts are reported to 0.001 ppm precision. Carbon multiplicities were determined by means of DEPT-135 and DEPT-90 (Bruker Avance DPX 400, at 100.62 MHz). NOE experiments were carried out under the following conditions: 5 mg of dry compound **1**, 0.5 mL of CDCl₃ 100 % *d* as solvent, degassing of the sample with Argon gas, relaxation time D1 = 1.5 s, decoupler power level for presaturation pL14 = 70 db, on a Bruker Avance DPX 400 spectrometer at 400.13 MHz. PERCH NMR software from Perch Solutions, Kuopio (Finland) was employed for the ¹H NMR spin system analysis. The HR ESMS was obtained on a Waters-Micromass Q-TOF micro. Column chromatography was performed on Sigel for TLC or LiChroprep RP-18 Lobar columns (Merck, 240 × 10 and 310 × 25). Analytical HPLC was carried out on a Waters 600 (DAD detector) with a C-18 symmetry column (150 × 3.9 mm, 5 µm); elution was accomplished with a gradient of MeOH-H₂O. All the solvents used were analytical grade.

Plant Material

L. frutescens was collected in Monterrey N.L. in October 2003. A voucher number specimen has been deposited in the herbarium of the Faculty of Biology, UANL (024165). It was authenticated by Biol. Marco Antonio Guzmán and M.C. María del Consuelo González de la Rosa, Faculty of Biology. UANL.

Extraction, Fractionation, and Isolation of Compounds

Root bark (200 g) of *L. frutescens* was air-dried, pulverized and extracted with 3×600 mL of MeOH at room temperature. The pooled methanolic extracts were evaporated to dryness *in vacuo* to yield 15.8 g of viscous mass. Liquid-liquid partition was carried out with *n*-hexane, EtOAc and *n*-BuOH. The active *n*-hexane fraction (5.7 g) was evaporated under reduced pressure and subjected to silica gel VLC using *n*-hexane-EtOAc (100:0 to 1:1 v/v) to give nine fractions (FVLC-1 to FVLC-9 in order of increasing polarity). The more active fraction FVLC-2 (812 mg) was further fractionated by RP-LPC (cartridge 310 \times 25) with mixtures of 300 mL MeOH:H₂O of increasing polarity. Six fractions were obtained, with FLP-3 (65 mg) and FLP-4 (250 mg) being the most active. Purification of these fractions were done with RP-LPC (cartridge of 240 X 10 and flow of 2 mL/min) using 60 mL of a gradient of MeOH:H₂O. One active compound **1** (216 mg) was isolated.

Compound 1: colorless oil; $[\alpha]_D -32$ (c 0.15, CHCl₃); UV (MeOH) 283.5 nm and 276.0 nm; IR (ATR) ν max 3463, 2922, 1618, 1578, 1451, 838 cm⁻¹; ¹H NMR (CDCl₃, 400 MHz) see Table 1; ¹³C NMR (CDCl₃, 100 MHz): δ 17.49 (C-17, CH₃), 18.64 (C-18, CH₃), 19.43 (C-3, CH₂), 20.96 (C-19, CH₃), 21.13 (C-20, CH₃), 25.59 (C-16, CH₃), 26.16 (C-13, CH₂), 26.56 (C-1, CH), 27.47 (C-2, CH₂), 33.33 (C-12, CH₂), 37.74 (C-11, CH), 42.37 (C-4, CH), 113.25 (C-7, CH), 122.30 (C-5, CH), 124.91 (C-14, CH), 126.38 (C-9, C), 130.95 (C-15, C), 134.83 (C-6, C), 140.81 (C-10, C), 153.01 (C-8, C); HR-ES-MS: *m/z* 287.2182 (2.56% BPI // 0.48 % TIC), 175 (100.00% BPI // 18.83 % TIC) *m/z* 287.2182 (M + H) (calcd. for C₂₀H₃₁O, 287.2376).

Anti-TB Bioassay

The anti-*M. tuberculosis* activity was assessed against *M. tuberculosis* H37Rv ATTC 27294 susceptible to all five first-line anti-TB drugs, streptomycin, isoniazid, rifampin, ethambutol, and pyrazinamide and CIBIN/UMF15:99, a clinical isolate resistant to the same drugs in a modified Microplate Assay Blue Alamar.³² The resistant *M. tuberculosis* strain (MDR) was isolated, identified, and characterized in the Mycobacteriology Laboratory of the Centro de Investigación Biomédica del Noreste, Instituto Mexicano del Seguro Social in Monterrey México, from a patient with advanced pulmonary TB. The organic extracts, fractions, and compounds for *M. tuberculosis* bioassays were prepared at a concentration of 2 mg/mL in 20% DMSO in Middlebrook 7H9 (Becton Dickinson and Co., Sparks MD, USA) broth. All solutions were sterilized by filtration using 13 mm diameter PTFE acrodiscs (0.22 μ m pore size, Millipore Co., Bedford, MA, USA). The concentrations for organic extracts, fractions, and compounds used ranged from 100 μ g/mL to 0.78 μ g/mL. Results are reported as minimal inhibition concentration (MIC). All biological assays were developed at least in triplicate.

VCD Analysis and Calculations

A sample of 9.8 mg of **1** was dissolved in 150 μ L of 100% CDCl₃, placed in a BaF₂ cell with a pathlength of 100 μ m and data were acquired on a BioTools BOMEM Chiral/IR FT-VCD spectrophotometer, equipped with dual photoelastic modulation, operated at a resolution of 4 cm⁻¹ during 4 h. The sample stability was verified by ¹H NMR measurement at 300 MHz on a Varian Mercury spectrometer using a 99.8% atom-D CDCl₃ solution

containing TMS as the internal standard immediately before and after VCD measurements. Geometry optimizations for both C-11 diastereomers were done using the MMFF94 force-field calculations as implemented in the Spartan'04 program. A Monte Carlo search protocol was carried out considering an energy cut-off of 10 kcal/mol providing 199 conformers for **1** and 193 for the 11*R* diastereoisomer. All conformers were optimized by single point DFT calculations at the B3LYP/DGDZVP level of theory, and those found in the 3 kcal/mol gap (54 for **1**, and 40 for its 11*R* epimer) were optimized using DFT//B3LYP/DGDZVP methodology^{33–35} implemented in the Gaussian 03W program. The 13 and 23 minimized structures found in the 2 kcal/mol window were further reduced by eliminating those conformers contributing less than 2% to the overall distribution to provide 10 and 13 conformers that were used to calculate the IR and VCD frequencies. The geometry optimization and vibrational calculations required between 12 and 20 h of computational time per conformer when using a desktop personal computer with 2 Gb RAM and operated with a 3 GHz processor.

Supplementary Material

Refer to Web version on PubMed Central for supplementary material.

Acknowledgments

The authors acknowledge financial support from CONACYT-México (grants 36544-N and 61122, and a doctoral fellowships for GMMS), IMSS (FOFOI: 2005/1/I/021), UANL (PAICYT), and CYTED (Project PIBATUB X.11). Ivonne Carrera is acknowledged for her technical assistance in the extraction procedures. Furthermore, GFP is indebted to Matthias Niemitz, Perch Solutions Ltd, and Reino Laatikainen, University of Kuopio and, both in Kuopio, Finland, for helpful discussions and PERCH support. The purchase of the 900 MHz NMR spectrometer and construction of the UIC Center for Structural Biology were funded by NIH Grant GM068944.

References and Notes

1. http://www.who.int/tb/publications/global_report/2009/en/index.html.
2. http://www.who.int/tb/publications/global_report/2010/en/index.htm.
3. Palomino JC, Ramos DF, da Silva PA. *Curr. Med. Chem.* 2009; 16:1898–1904. [PubMed: 19442153]
4. Rivers EC, Mancera RL. *Drug. Discov. Today.* 2008; 13:1090–1098. [PubMed: 18840542]
5. Negi AS, Kumar JK, Lugman S, Saikia D, Khanuja SP. *Med. Res. Rev.* 2010; 30:603–645. [PubMed: 19626592]
6. de Souza MV. *Fitoterapia.* 2009; 80:453–460. [PubMed: 19698768]
7. Copp BR, Pearce AN. *Nat. Prod. Rep.* 2007; 24:278–297. [PubMed: 17389998]
8. Molina-Salinas GM, Pérez-López A, Becerril-Montes P, Salazar-Aranda R, Said-Fernández S, Waksman de Torres N. *J. Ethnopharmacol.* 2007; 109:435–441. [PubMed: 17000069]
9. Rimando AM, Dayan FE, Mikell JR, Moraes RM. *Natural Toxins.* 1999; 7:39–43. [PubMed: 10441036]
10. Rojas S, Acevedo L, Macías M, Toscano RA, Bye R, Timmermann B, Mata R. *J. Nat. Prod.* 2003; 66:221–224. [PubMed: 12608853]
11. Rodriguez AD, Ramirez C. *J. Nat. Prod.* 2001; 64:100–102. [PubMed: 11170678]
12. Ata A, Kerr RG, Moya CE, Jacobs RS. *Tetrahedron.* 2003; 59:4215–4222.
13. Niemitz M, Laatikainen R, Chen SN, Kleps R, Kozikowski AP, Pauli GF. *Magn. Reson. Chem.* 2007; 45:878–882. [PubMed: 17729231]
14. Scher J, Schinkovitz A, Zapp J, Wang Y, Franzblau SG, Becker H, Lankin DC, Pauli GF. *J. Nat. Prod.* 2010; 73:656–663. [PubMed: 20353194]
15. Kolehmainen E, Laihia K, Laatikainen R, Vepsäläinen J, Niemitz M, Suontamo R. *Magn. Reson. Chem.* 1997; 35:463–467.

16. Inui T, Wang Y, Nikolic D, Smith D, Franzblau S, Pauli GF. *J. Nat. Prod.* 2010; 73:563–567. [PubMed: 20218656]
17. Kolesnikova S, Kaliniovsk A, Fedorov S, Shubina L, Stonik V. *Phytochemistry.* 2006; 67:2115–2119. [PubMed: 16842831]
18. Rodriguez II, Shi YP, García OJ, Rodriguez AD, Mayer AMS, Sánchez JA, Ortega-Barria E, González J. *J. Nat. Prod.* 2004; 67:1672–1680. [PubMed: 15497938]
19. Yan-Ping S, Wei X, Rodríguez II, Rodríguez AD, Mayer AMS. *Eur. J. Org. Chem.* 2009; 4:493–502.
20. Scher J, Schinkovitz A, Zapp J, Wang Y, Franzblau SG, Becker H, Lankin DC, Pauli GF. *J. Nat. Prod.* 2010; 73:656–663. [PubMed: 20353194]
21. Liu Q, Harrington D, Kohen JL, Vermulpad S, Jamie JF. *Phytochemistry.* 2006; 67:1256–1261. [PubMed: 16759677]
22. Smith JE, Tucker D, Watson K, Jones GL. *J. Ethnopharmacol.* 2007; 112:386–393. [PubMed: 17485184]
23. Ndi CP, Semple SJ, Griesser HJ, Pyke SM, Barton MD. *Phytochemistry.* 2007; 68:2684–2690. [PubMed: 17631925]
24. Ndi CP, Semple SJ, Griesser HJ, Pyke SM, Barton MD. *J. Nat. Prod.* 2007; 70:1439–1443. [PubMed: 17844993]
25. Harmata M, Hong X, Schreiner R. *J. Org. Chem.* 2008; 73:1290–1296. [PubMed: 18215060]
26. Burgueño-Tapia E, Castillo L, González-Coloma A, Joseph-Nathan P. *J. Chem. Ecol.* 2008; 34:766–771. [PubMed: 18528728]
27. Joseph-Nathan P, Mejía G, Abramo-Bruno D. *J. Am. Chem. Soc.* 1979; 101:1289–1291.
28. Muñoz MA, Chamy C, Carrasco A, J. Roviroso JA, San Martín A, Joseph-Nathan P. *Chirality.* 2009; 21:E208–E214. [PubMed: 19899155]
29. Kuppens T, Vandyck K, Van der Eycken J, Herrebout W, Van der Vecken BJ, Bultinck P. *J. Org. Chem.* 2005; 70:9103–9114. [PubMed: 16268579]
30. Burgueño-Tapia E, Zepeda LG, Joseph-Nathan P. *Phytochemistry.* 2010; 71:1158–1161. [PubMed: 20457457]
31. Jacobs RS, Kerr RG. Anti-inflammatory Compounds Derived from *Pseudopterogorgia elisabethae*. *PCT Int. Appl.* 2002:44. CODEN: PIXXD2 WO 2002044191 A2 20020606 CAN 137:15779 AN 2002:428919 CAPLUS.
32. Molina-Salinas GM, Ramos-Guerra MC, Vargas-Villarreal J, Mata-Cárdenas BD, Becerril-Montes P, Said-Fernandez S. *Arch. Med. Res.* 2006; 37:45–49. [PubMed: 16314185]
33. Parr, RG.; Yang, W. *Density-Functional Theory of Atoms and Molecules.* Oxford: Oxford Univ. Press; 1989.
34. Godbout N, Salahub DR, Andzelm J, Wimmer E. *Can. J. Chem.* 1992; 70:560–571.
35. Sosa C, Andzelm BC, Elkin E, Wimmer E, Dobbs KD, Dixon DA. *J. Phys. Chem.* 1992; 96:6630–6636.
36. Davies HML, Dai Z. *Tetrahedron.* 2006; 62:10477–10484.
37. Dai X, Wan Z, Kerr RG, Davies HM. *J. Org. Chem.* 2007; 72:1895–1900. [PubMed: 17309304]




Figure 1.
HMBC correlations found for **1**.




Figure 2.
 ^{13}C - ^{13}C INADEQUATE experiment performed with **1** confirming most of the C-C connectivity.




Figure 3.
Observed NOE correlations for **1**.




Figure 4. Full ^1H NMR spin analysis by PERCH iteration of 900 MHz data of compound **1** (simulated spectra in red; experimental spectra in blue).




Figure 5. Superposition of the three low energy conformers of **1**, calculated by DFT//B3LYP/DGDZVP, accounting for 68.6% of the conformational distribution according to Table 2.




Figure 6. Superposition of the five low energy conformers of the 11*R* diastereomer of **1**, calculated by DFT//B3LYP/DGDZVP, accounting for 65.2% of the conformational distribution according to Table 3.





Figure 7. Comparison of the experimental (center) VCD spectrum of **1** and the theoretical VCD spectra of **1** (top) and its C-11 diastereomer (bottom) calculated by DFT/B3LYP/DGDZVP.

**1**

Chemical Chart.


**2**

Table 1

Results of the ^1H NMR Full Spin Analysis of **1**, Sorted by Chemical Shifts in Support of Figures 2 and 3.

Hydrogen	δ in ppm	Multiplicity	J in Hz
CH ₃ -18	0.984	d	$J_{\text{H-11}} = 6.84$
H-12 _b	1.107	dddd (ddt-like)	$J_{\text{gem}} = -13.19$, $J_{\text{H-11}} = 10.02$, $J_{\text{H-13b}} = 9.28$, $J_{\text{H-13a}} = 4.86$
CH ₃ -20	1.224	d	$J_{\text{H-1}} = 6.97$
H-12 _a	1.314	dddd	$J_{\text{gem}} = -13.19$, $J_{\text{H-13a}} = 9.63$, $J_{\text{H-13b}} = 6.94$, $J_{\text{H-11}} = 2.96$
H-2 _{eq}	1.520	dddd (ddt-like)	$J_{\text{gem}} = -13.35$, $J_{\text{H-3ec}} = 5.20$, $J_{\text{H-3ax}} = 3.52$, $J_{\text{H-1}} = 2.58$
CH ₃ -17	1.582	dd/brs [<i>syn</i>]	$J_{\text{H-14}} = 1.36$, $J_{\text{H-13a}} = 0.89$, $J_{\text{H-13b}} = 0$
CH ₃ -16	1.683	ddd (=pseudo q) [<i>anti</i>]	$J_{\text{H-14}} = 1.38$, $J_{\text{H-13a}} = 1.18$, $J_{\text{H-13b}} = 1.14$
H-3 _{eq}	1.740	dddd	$J_{\text{gem}} = -13.84$, $J_{\text{H-2ec}} = 5.00$, $J_{\text{H-2ax}} = 3.37$, $J_{\text{H-4}} = 2.95$
H-13 _b	1.834	dddq [broad]	$J_{\text{gem}} = -14.37$, $J_{\text{H-12b}} = 9.28$, $J_{\text{H-14}} = 7.60$, $J_{\text{H-12a}} = 6.94$, $J_{\text{CH3-16}} = 1.14$, $J_{\text{CH3-17}} = 0$
H-3 _{ax}	1.870	dddd (tdd-like)	$J_{\text{gem}} = -13.84$, $J_{\text{H-2ax}} = 13.30$, $J_{\text{H-4}} = 6.23$, $J_{\text{H-2ec}} = 3.52$
H-11	1.902	dddq [broad]	$J_{\text{H-12b}} = 10.02$, $J_{\text{Me-18}} = 6.84$, $J_{\text{H-4}} = 5.21$, $J_{\text{H-12a}} = 2.96$
H-2 _{ax}	1.967	dddd (tdd-like)	$J_{\text{gem}} = -13.25$, $J_{\text{H-3ax}} = 13.25$, $J_{\text{H-1}} = 6.16$, $J_{\text{H-3ec}} = 3.37$
H-13 _a	2.010	dddq [broad]	$J_{\text{gem}} = -14.37$, $J_{\text{H-12a}} = 9.63$, $J_{\text{H-14}} = 6.76$, $J_{\text{H-12b}} = 4.86$, $^5J_{\text{CH3-16}} = 1.17$, $^5J_{\text{CH3-17}} = 0.89$
CH ₃ -19	2.270	s	
H-4	2.590	ddd	$J_{\text{H-3ax}} = 6.23$, $J_{\text{H-11}} = 5.21$, $J_{\text{H-3ec}} = 2.95$
H-1	3.078	ddq	$J_{\text{CH3-20}} = 6.97$, $J_{\text{H-2ax}} = 6.16$, $J_{\text{H-2ec}} = 2.58$
H-14	5.018	ddq=ddquint	$J_{\text{H-13b}} = 7.60$, $J_{\text{H-13a}} = 6.76$, $J_{\text{CH3-16}} = 1.38$, $J_{\text{CH3-17}} = 1.36$
H-7	6.448	brd	$J_{\text{H-5}} = 1.64$, $J_{\text{CH3-19}} = <0.20$
H-5	6.615	dq [brd]	$J_{\text{H-7}} = 1.64$, $J_{\text{H-4}} = 0.82$, $J_{\text{CH3-19}} = <0.20$

Table 2

Molecular Mechanics Relative Energy, Molecular Mechanics Population, DFT Thermochemical Parameters, and DFT Population for the 10 Minimum Energy Conformers of **1**.

conformer	ΔE_{MMFF}^a	p_{MMFF}^b	$\Delta E_0^{c,d}$	$\Delta H_{298}^{c,d,e}$	$\Delta G_{298}^{c,d}$	p_{DFT}^f
1a	0.38	22.03	0.00	0.00	0.00	37.93
1b	0.40	21.56	0.10	0.10	0.35	21.07
1c	1.30	4.69	0.78	0.79	0.82	9.57
1d	1.81	1.99	1.23	1.25	1.04	6.51
1e	1.27	4.94	0.90	0.90	1.17	5.28
1f	4.71	0.02	1.04	0.96	1.19	5.13
1g	0.00	41.99	0.57	0.45	1.20	5.02
1h	2.71	0.43	1.59	1.52	1.35	3.85
1i	1.76	2.16	1.40	1.40	1.41	3.53
1j	3.21	0.19	1.80	1.74	1.71	2.11

^aMolecular mechanics energy of conformers obtained from a systematic search, in kcal/mol relative to **1g** with $E_{\text{MMFF}} = 43.842$ kcal/mol.

^bMolecular mechanics population in %.

^cSum of electronic and zero-point energies (ΔE_0), thermal enthalpies (ΔH_{298}) and thermal free energies (ΔG_{298}) in kcal/mol relative to conformer **1a** and calculated at 298 K and 1 atm.

^dFor conformer **1a** the absolute values are $E_0 = -855.044971$ au, $H_{298} = -855.020784$ and $G_{298} = -855.099078$ au.

^e ΔE_{298} values equal to ΔH_{298} in all conformations.

^fDFT population in % calculated from ΔG_{298} values.

Table 3

Molecular Mechanics Relative Energy, Molecular Mechanics Population, DFT Thermochemical Parameters, and DFT Population for the 13 Minimum Energy Conformers of the C-11 Diastereomer of **1**.

conformer	ΔE_{MMFF}^a	p_{MMFF}^b	ΔE_0^c	ΔH_{298}^e	ΔG_{298}^d	p_{DFT}^f
a	1.20	7.35	0.00	0.00	0.00	18.23
b	1.13	8.26	0.19	0.18	0.04	16.92
c	1.19	7.47	0.10	0.08	0.26	11.79
d	2.17	1.41	0.75	0.80	0.36	9.97
e	1.06	9.33	0.48	0.41	0.46	8.32
f	2.53	0.78	1.41	1.47	0.69	5.67
g	2.63	0.67	1.11	1.18	0.80	4.69
h	2.62	0.66	1.11	1.17	0.85	4.31
i	2.24	1.27	1.33	1.38	0.86	4.29
j	3.32	0.20	1.30	1.29	0.88	4.16
k	0.00	55.45	0.63	0.52	0.89	4.07
l	2.05	1.74	1.18	1.18	0.91	3.90
m	1.38	5.41	1.04	0.96	0.95	3.68

^aMolecular mechanics energy of conformers obtained from a systematic search, in kcal/mol relative to **2k** with $E_{\text{MMFF}} = 43.215$ kcal/mol.

^bMolecular mechanics population in %.

^cSum of electronic and zero-point energies (ΔE_0), thermal enthalpies (ΔH_{298}) and thermal free energies (ΔG_{298}) in kcal/mol relative to conformer **2a** and calculated at 298 K and 1 atm.

^dFor conformer **2a** the absolute values are $E_0 = -855.044838$ au, $H_{298} = -855.020732$ and $G_{298} = -855.098357$ au.

^e ΔE_{298} values equal to ΔH_{298} in all conformations.

^fDFT population in % calculated from ΔG_{298} values.

Table 4

Confidence Level Data for the IR and VCD Spectra of **1** and its 11*R*-Diastereoisomer.

Epimer	<i>a</i> <i>anH</i>	<i>b</i> <i>S_{IR}</i>	<i>c</i> <i>S_E</i>	<i>d</i> <i>S_{-E}</i>	<i>e</i> <i>ESI</i>	<i>f</i> <i>C</i>
1	0.968	89.9	70.6	13.6	57.0	100
11 <i>epi</i> - 1	0.974	-	62.0	18.5	43.5	76

^aAnharmonicity factor.

^bIR spectral similarity.

^cVCD spectral similarity for the correct enantiomer.

^dVCD spectral similarity for the incorrect enantiomer.

^eEnantiomer similarity index, calculated as the *S_E*-*S_{-E}* difference.

^fConfidence level for the stereochemical assignment.

Table 5Anti-*M. tuberculosis* Activity of **1** in Comparison with First Line Anti-TB Drugs.

	MIC ($\mu\text{g/mL}$) vs <i>M. tuberculosis</i>	
	H37Rv strain	MDR strain
1	12.5	6.3
streptomycin	0.50	>100
isoniazide	0.06	3.1
rifampin	0.06	>100
ethambutol	2.0	8.0

MIC: minimal inhibition concentration.

^a *M. tuberculosis* H37Rv strain is sensitive to all five first-line anti-TB drugs and^b CIBIN/UMF15:99 a MDR clinical isolated resistant to the above drugs.