

RESEARCH ARTICLE

Open Access

Influence of high-density lipoprotein cholesterol on coronary collateral formation in a population with significant coronary artery disease

Po-Chao Hsu^{1,2}, Ho-Ming Su^{1,6,7}, Suh-Hang Juo^{3,4,5}, Hsueh-Wei Yen^{1,6}, Wen-Chol Voon^{1,6}, Wen-Ter Lai^{1,6}, Sheng-Hsiung Sheu^{1,6} and Tsung-Hsien Lin^{1,6*}

Abstract

Background: Coronary collateral circulation plays an important role in protecting myocardium from ischemia and reducing cardiovascular events. Low High-density lipoprotein cholesterol (HDL-C) level is a strong risk factor for coronary artery disease (CAD) and is associated with poor cardiovascular outcome. It was recently reported to be associated with poor coronary collateral development in Turkish population. Hence, we investigated the influence of HDL-C on coronary collateral formation in Chinese population.

Methods: We evaluated 970 consecutive patients undergoing coronary angiography, and 501 patients with significant coronary artery disease (SCAD) were finally analyzed. The collateral scoring system developed by Rentrop was used to classify patient groups as those with poor or good collaterals.

Results: The patients with poor collaterals had fewer diseased vessels (1.97 ± 0.84 vs 2.47 ± 0.68 , $p < 0.001$) and lower diffuse score (2.65 ± 1.63 vs 3.76 ± 1.78 , $p < 0.001$). There was no significant difference in HDL-C and other variables between good and poor collaterals. Multivariate analysis showed only number of diseased vessels (odds ratio 0.411, $p < 0.001$) was a significant predictor of poor collateral development.

Conclusions: The extent of CAD severity but not HDL-C level was the most powerful predictor of coronary collateral formation in our Chinese population with SCAD.

Keywords: Coronary artery disease, Coronary collateral circulation, High-density lipoprotein cholesterol

Background

Cardiovascular disease is one of the leading causes of morbidity and mortality in the world. Low high-density lipoprotein cholesterol (HDL-C) is an independent risk factor for coronary artery disease (CAD) and is associated with increased cardiovascular event [1-3].

The development of coronary collaterals is an adaptive response to chronic myoischemia and serves as a conduit bridging the significantly stenotic coronary vessels [4-6]. Collateral circulation can hence protect and preserve myocardium from episodes of ischemia, enhance residual myocardial contractility, and reduce angina

symptoms and cardiovascular events [7-9]. However, there is inter-individual difference of coronary collateral formation and the mechanisms for different individual ability to develop collateral circulation are still not well understood.

Furthermore, HDL-C was also recently reported to be associated with poor coronary collateral development in a Turkish population [10]. Hence, we wanted to investigate the influence of HDL-C on coronary collateral formation in a Chinese population with significant CAD (SCAD).

Methods

Study subjects

From February 2002 to March 2008, we evaluated 970 consecutive patients who had been scheduled to undergo diagnostic coronary angiography at Kaohsiung

* Correspondence: lth@kmu.edu.tw

¹Department of Internal Medicine, Division of Cardiology, Kaohsiung Medical University, 100 Tzyou 1st Road, Kaohsiung 80708, Taiwan, ROC

⁶Department of Internal Medicine, Faculty of Medicine, Kaohsiung, Taiwan

Full list of author information is available at the end of the article

Medical University Hospital in Taiwan. We excluded patients with coronary artery lumen diameter stenosis < 70%, history of coronary artery bypass surgery (CABG), history of percutaneous coronary intervention (PCI), or those who had inadequate angiograms for collateral evaluation. Finally, 501 patients with significant CAD were enrolled in this study. Other analyzed demographic and baseline data included gender, age, and any history of the following: diabetes mellitus, hypertension, hypercholesterolemia, body mass index, cigarette smoking, and medications. The research protocol was approved and registered by the ethics committee (Kaohsiung Medical University Hospital- Institutional Review Board) at our institution, and informed consent was obtained from all patients.

Coronary angiography and collateral scoring

The coronary artery angiography films were reviewed by 2 experienced cardiologists blinded to the clinical and demographic data for all patients. Any differences in interpretation were resolved by a third reviewer who was blinded to the readings of the first two reviewers. Coronary artery stenosis was determined by quantitative coronary angiography. The recorded data also included the number of diseased vessels, the vessel to which the collaterals were connected, and the grade of coronary collateral circulation. Vessels exhibiting a 70% or greater reduction in lumen diameter were classified as SCAD. In subjects with more than one SCAD vessel, the vessel with the highest collateral grade was chosen for analysis. The collateral scoring system developed by Rentrop and Cohen was used [11]. Grades of collateral filling from the contralateral vessel were: 0 = none; 1 = filling of side branches of the artery to be dilated via collateral channels without visualization of the epicardial segment; 2 = partial filling of the epicardial segment via collateral channels; 3 = complete filling of the epicardial segment of the artery being dilated via collateral channels. In subjects with more than one collateral vessel supplying the distal aspect of the diseased artery, the highest collateral grade was recorded. Patients were then classified according to their collateral grades as either poor (grade 0 or grade 1 collateral circulation) or good (grade 2 or grade 3 collateral circulation). The coronary artery angiography films were reviewed by two experienced cardiologists blinded to the clinical and demographic data for all patients. Any differences in interpretation were resolved by a third reviewer who was blinded to the readings of the first two reviewers. The two readers obtained a 96% agreement in the collateral classifications.

CAD severity scoring by coronary angiography

We evaluated the extent of coronary artery atherosclerosis by a "diffuse score" (DS) developed by Negri et al.

and modified by Birnie et al. [12,13]. In brief, the coronary circulation is divided into 15 segments, and eight of them are classified as first-order segments: proximal and middle right coronary arteries, left main coronary artery, proximal, middle, and distal left anterior descending, and proximal and distal circumflex. There are seven second-order segments: distal right coronary artery, posterior descending branch, obtuse marginal branch, postero-lateral branch of circumflex, and the first two diagonal branches of the left anterior descending artery. The first-order segments receive a score of 1 and the second order segments scored 0.5. The overall DS is the sum of the individual segment scores and the maximum score is 11.5.

Statistical analysis

All data were expressed as means \pm standard deviation. Student unpaired *t*-test was used to compare continuous variables between the two groups. Chi-square test was used to compare categorical data. Subsequently, age, sex, significantly correlated variables in the univariate analysis or relevant variables were further analyzed by logistic regression analysis. All *p* values were two-sided with a significance level of *p* < .05. The Statistical Package for the Social Sciences 11.0 for Windows (SPSS Inc., Chicago, IL) was used for statistical analysis.

Results

Clinical characteristics

Of the 970 subjects initially enrolled, 469 patients were excluded for the following reasons: coronary artery lumen diameter stenosis < 70%, history of CABG or PCI, or inadequate angiograms for collateral evaluation. The final study population was 501 subjects (396 male, 105 female; average age, 62.5 \pm 12.6 years). For SCAD, 134 patients (26.7%) were one-vessel disease, 153 patients (30.5%) were two-vessel disease, and 214 patients (42.7%) were three-vessel disease.

Coronary collaterals

The two readers obtained a 96% agreement in the collateral classifications. Of the 501 patients enrolled, 220 (43.9%) patients had no coronary collaterals. In subjects with collaterals, the coronary grade was distributed as follows: 91 (18.2%) with grade 1, 133 (26.5%) with grade 2 and 57 (11.4%) with grade 3. Table 1 compares the baseline characteristics between the poor group (*n* = 311) and the good group (*n* = 190). The patients in the poor group had a trend to have lower cholesterol level (199.7 \pm 45.1 vs 208.3 \pm 53.6, *p* = 0.06), lower low-density lipoprotein cholesterol (LDL-C) level (128.7 \pm 38.3 vs 136.2 \pm 43.5, *p* = 0.066), less angiotensin II receptor blocker (ARB) usage (15.9% vs 22.3%, *p* = 0.09), and less beta blocker usage (28% vs 36.7%, *p* = 0.054).

Table 1 Baseline characteristics in patients with poor and good collaterals

Variables	Poor collateral (n = 311)	Good collateral (n = 190)	P value
Age (years)	62.8 ± 12.9	62.1 ± 12.1	0.547
Gender (Female, %)	22.5	18.4	0.309
DM (%)	42.8	40.5	0.642
HTN (%)	64.3	65.3	0.848
Smoking (%)	60.5	58.2	0.639
BMI	24.3 ± 6.6	24.9 ± 5.7	0.277
eGFR (mL/min/1.73 m ²)	66.9 ± 26.9	68.0 ± 32.0	0.683
Laboratory data			
Glucose (mg/dl)	137.7 ± 62.2	135.4 ± 60.4	0.704
Cholesterol (mg/dl)	199.7 ± 45.1	208.3 ± 53.6	0.060
LDL-C (mg/dl)	128.7 ± 38.3	136.2 ± 43.5	0.066
HDL-C (mg/dl)	37.4 ± 9.9	36.0 ± 10.6	0.175
Triglyceride (mg/dl)	168.9 ± 124.9	163.4 ± 146.4	0.660
Uric acid (mg/dl)	6.54 ± 1.94	6.58 ± 1.81	0.819
Medication			
ACEI (%)	19.1	20.7	0.725
ARB (%)	15.9	22.3	0.09
Beta blocker (%)	28	36.7	0.054

ACEI: angiotensin converting enzyme inhibitor; ARB: angiotensin II receptor blocker; BMI: body mass index; DM: diabetes; HTN: hypertension; eGFR: estimated glomerular filtration rate; HDL-C: high-density lipoprotein cholesterol; LDL-C: low-density lipoprotein cholesterol.

Angiographic characteristics in patients with poor and good collaterals are shown in Table 2; the patients with poor collaterals had fewer diseased vessels numbers (1.97 ± 0.84 vs 2.47 ± 0.68, p < 0.001), and lower severity scores of CAD (2.65 ± 1.63 vs 3.76 ± 1.78, p < 0.001).

Binary logistic regression analysis

Age, sex, significant variable in the univariate analysis and relevant variables including cholesterol level, LDL-C level, HDL-C level, ARB usage, and beta blocker usage were included in the multivariate logistic regression analysis. We found the number of diseased vessels (OR =

Table 2 Angiographic characteristics in patients with poor and good collaterals

Mean±SD Number (%)	Poor collateral (n = 311)	Good collateral (n = 190)	P value
Number of diseased vessels	1.97 ± 0.84	2.47 ± 0.68	<0.001
Significant CAD, n (%)			
1 vessel disease (%)	114 (36.7)	20 (10.5)	
2 vessel disease (%)	93 (29.9)	60 (31.6)	<0.001
3 vessel disease (%)	104 (33.4)	110 (57.9)	
Diffuse score of CAD	2.65 ± 1.63	3.76 ± 1.78	<0.001

CAD: coronary artery disease.

0.410, 95% CI = 0.31-0.55, p < 0.001) was the only significantly independent predictor of poor coronary collaterals development (Table 3).

Discussion

In the current study, we investigated the influence of HDL-C on coronary collateral formation in Chinese population with SCAD and found two major findings. First, HDL-C was not significantly associated with coronary collateral formation in a Chinese population as a previous study in a Turkish population. Second, the number of diseased vessels was the only significantly independent predictor of poor coronary collaterals in the Chinese population. This finding was also different from previous studies in a Turkish population.

The association between HDL-C level and coronary collaterals

HDL-C has been reported to have anti-atherogenic, anti-inflammatory, and antioxidant properties in the literature [14-16]. Although Sumi M et al. also showed that reconstituted HDL-C directly stimulates endothelial progenitor cell differentiation via phosphatidylinositol 3-kinase/Akt pathway and enhances ischemia-induced angiogenesis, we did not find HDL-C was associated with coronary collateral development in this population study [17]. In addition, Low HDL-C level was a strong risk factor for coronary artery disease and was associated with poor cardiovascular outcome [1-3]. In a previous study, Turhan H et al. reported that the development of coronary collateral vessels is poorer in patients with metabolic syndrome with advanced ischemic heart disease than in control participants without metabolic syndrome in a Turkish population, which suggested that metabolic syndrome is one of the significant factors affecting the development of coronary collateral vessels adversely [18]. Although low HDL-C is one of the

Table 3 Multivariate logistic regression analysis of collateral circulation (good collateral group as reference group)

	B	OR	95% CI	p value
Gender (male vs female)	-0.268	0.765	0.42-1.41	0.390
Age	0.010	1.010	0.99-1.03	0.326
Cholesterol (mg/dl)	0.003	1.003	0.99-1.01	0.440
LDL-C (mg/dl)	-0.007	0.993	0.98-1.00	0.135
HDL-C (mg/dl)	0.018	1.018	0.99-1.04	0.160
ARB usage	-0.422	0.656	0.37-1.17	0.155
Beta blocker usage	-0.380	0.684	0.42-1.11	0.126
Number of diseased vessels	-0.892	0.410	0.31-0.55	<0.001

ARB: angiotensin II receptor blocker; HDL-C: high-density lipoprotein cholesterol; LDL-C: low-density lipoprotein cholesterol.

diagnostic criteria for metabolic syndrome, they did not report and we also did not find HDL-C was associated with coronary collateral formation. Recently, Kadi H et al. further investigated the relationship between HDL-C and coronary collateral formation in patients with SCAD and found that there was a positive relationship between HDL-C and coronary collateral formation and suggested that low HDL-C is an independent predictor of poor coronary collateral formation in a Turkish population [10]. However, Li CC et al. reported that severity of coronary stenosis was the major influencing factor in coronary collateral formation and function in a Chinese population. Serum HDL-C was not associated with coronary collateral formation, but related to coronary collateral function which was evaluated by the size of the collateral connection diameter [19]. In our study, we analyzed 501 Chinese patients with SCAD to evaluate the correlation between HDL-C and coronary collaterals. Our result showed there was no significant association between HDL-C and coronary collateral formation not only in univariate but also in multivariate analysis, which was different from a previous study in a Turkish population. The advantages of our study were that we had a larger sample size and selected more severe coronary artery lesions than the previous study [10]. In their study, the sample size was relatively small and vessels with 50-70% stenosis were selected for coronary collateral evaluation, which may not cause severe myoischemia to induce coronary collateral formation [6]. In addition, the influence of HDL-C for coronary collateral development might not be similar between different races because of genetic diversity. However, the detailed mechanism still needs further investigation.

The association between CAD severity and coronary collaterals

It is well known that coronary collateral formation is mainly dependent on CAD severity [20-22]. Coronary collateral formation is an adaptive response to chronic myoischemia and serves as a conduit bridging significantly stenotic coronary vessels. More severe and extensive myocardial ischemia such as multi-vessel diseases may further stimulate the development of coronary collateral circulation. Hence, patients with good coronary collaterals appear to have a more extensive CAD. In previous studies, number of diseased vessels was a significant predictor of good collateral formation [20,23]. In our study, number of diseased vessels was not only associated with good collateral formation in univariate analysis but in multivariate analysis as well. In addition, we also used DS to further evaluate the extent of CAD severity and found that there was also a significant difference of DS between patients with poor and good coronary collaterals. Hence, extent of coronary atherosclerosis significantly plays an important role in coronary collateral formation [24,25].

Limitations of the present study

First, the collateral formation was assessed by coronary angiography in this study. Measuring collateral flow index by intravascular Doppler guidewire may provide a more objective physiological measurement of collateral grade. However, the invasiveness of intravascular ultrasound limits its use in large-scale studies. Second, since this was only a clinical observational study, potential mechanisms were not fully elucidated. Third, we did not analyze the size of the collateral connection diameter to evaluate the coronary collateral function.

Conclusions

In conclusion, our data revealed that HDL-C was not associated with coronary collateral formation in a Chinese population with SCAD; however, CAD severity implicated by number of diseased vessels was the only significant predictor of coronary collateral development in our study.

Competing interests

The authors have no competing interests to declare.

Authors' contributions

PCH: study concept and design, analysis and interpretation of data and drafting the manuscript. HMS, SHJ, SHS, and WTL: study concept and design, acquisition of subjects. HWY, WCV: Providing intellectual content of critical importance to the work described. THL: study concept and design, preparation of manuscript, and final approval of the version. All authors read and approved the final manuscript.

Author details

¹Department of Internal Medicine, Division of Cardiology, Kaohsiung Medical University, 100 Tzyou 1st Road, Kaohsiung 80708, Taiwan, ROC. ²Graduate Institute of Medicine, Kaohsiung Medical University, Kaohsiung, Taiwan. ³Department of Medical Research, Kaohsiung Medical University Hospital, Kaohsiung, Taiwan. ⁴Medical Genetics, Kaohsiung, Taiwan. ⁵Center of Excellence for Environmental Medicine, Kaohsiung Medical University, Kaohsiung, Taiwan. ⁶Department of Internal Medicine, Faculty of Medicine, Kaohsiung, Taiwan. ⁷Department of Internal Medicine, Kaohsiung Municipal Hsiao-Kang Hospital, Kaohsiung, Taiwan.

Received: 24 October 2012 Accepted: 13 March 2013

Published: 20 March 2013

References

1. Gordon DJ, Rifkind BM: High-density lipoprotein - the clinical implications of recent studies. *N Engl J Med* 1989, **321**:1311-1316.
2. Rubins HB, Schectman G, Wilt TJ, Iwano MK: Distribution of lipid phenotypes in community-living men with coronary heart disease. High prevalence of isolated low levels of high-density lipoprotein cholesterol. *Arch Intern Med* 1992, **152**:2412-2416.
3. Abbott RD, Wilson PW, Kannel WB, Castelli WP: High-density lipoprotein in cholesterol, total cholesterol screening, and myocardial infarction. The Framingham Study. *Arteriosclerosis* 1998, **8**:207-211.
4. Fujita M, Sasayama S, Ohno A, Nakajima H, Asanoi H: Importance of angina for development of collateral circulation. *Br Heart J* 1987, **57**:139.
5. Tayebjee MH, Lip GY, MacFadyen RJ: Collateralization and the response to obstruction of epicardial coronary arteries. *QJM* 2004, **97**:259.
6. Levin DC: Pathways and functional significance of the coronary collateral circulation. *Circulation* 1974, **50**:831.
7. Cohen M, Rentrop KP: Limitation of myocardial ischemia by collateral circulation during sudden controlled coronary artery occlusion in human subjects: a prospective study. *Circulation* 1986, **74**(3):469-476.

8. Meier P, Gloekler S, Zbinden R, Beckh S, de Marchi SF, Zbinden S, Wustmann K, Billinger M, Vogel R, Cook S, Wenaweser P, Togni M, Windecker S, Meier B, Seiler C: **Beneficial effect of recruitable collaterals: a 10-year follow-up study in patients with stable coronary artery disease undergoing quantitative collateral measurements.** *Circulation* 2007, **116**(9):975–983.
9. Regieli JJ, Jukema JW, Nathoe HM, Zwinderman AH, Ng S, Grobbee DE, van der Graaf Y, Doevendans PA: **Coronary collaterals improve prognosis in patients with ischemic heart disease.** *Int J Cardiol* 2009, **132**(2):257–262.
10. Kadi H, Ozyurt H, Ceyhan K, Koc F, Celik A, Burucu T: **The relationship between high-density lipoprotein cholesterol and coronary collateral circulation in patients with coronary artery disease.** *J Investig Med* 2012, **60**(5):808–12.
11. Rentrop KP, Cohen M, Blanke H, Phillips RA: **Changes in collateral channel filling immediately after controlled coronary artery occlusion by an angioplasty balloon in human subjects.** *J Am Coll Cardiol* 1985, **5**(3):587–592.
12. Negri M, Sheiban I, Arigliano PL, Tonni S, Montresor G, Carlini S, Manzato F: **Interrelation between angiographic severity of coronary artery disease and plasma levels of insulin, C-peptide and plasminogen activator inhibitor-1.** *Am J Cardiol* 1993, **72**:397–401.
13. Birnie DH, Holme ER, McKay IC, Hood S, McColl KE, Hillis WS: **Association between antibodies to heat shock protein 65 and coronary atherosclerosis. Possible mechanism of action of Helicobacter pylori and other bacterial infections in increasing cardiovascular risk.** *Eur Heart J* 1998, **19**:387–94.
14. Oram JF, Vaughan AM: **ATP-binding cassette cholesterol transporters and cardiovascular disease.** *Circ Res* 2006, **99**:1031–1043.
15. Cockerill GW, Rye KA, Gamble JR, Vadas MA, Barter PJ: **High-density lipoproteins inhibit cytokine-induced expression of endothelial cell adhesion molecules.** *Arterioscler Thromb Vasc Biol* 1995, **15**:1987–1994.
16. Suc I, Escargueil-Blanc I, Trolly M, Salvayre R, Nègre-Salvayre A: **HDL and ApoA prevent cell death of endothelial cells induced by oxidized LDL.** *Arterioscler Thromb Vasc Biol* 1997, **17**:2158–2166.
17. Sumi M, Sata M, Miura S, Rye KA, Toya N, Kanaoka Y, Yanaga K, Ohki T, Saku K, Nagai R: **Reconstituted high-density lipoprotein stimulates differentiation of endothelial progenitor cells and enhances ischemia-induced angiogenesis.** *Arterioscler Thromb Vasc Biol* 2007, **27**(4):813–8.
18. Turhan H, Yasar AS, Erbay AR, Yetkin E, Sasmaz H, Sabah I: **Impaired coronary collateral vessel development in patients with metabolic syndrome.** *Coron Artery Dis* 2005, **16**(5):281–5.
19. Li CC, Yang TL, Pu XQ, Zheng ZF, Yu ZX, Chen XB, Chen F, Mo L, Hu DJ, Xie QY, He L, Deng JH, Meng SY: **Formation and function of coronary collateral circulation and their influencing factors.** *Zhong Nan Da Xue Xue Bao Yi Xue Ban* 2004, **29**(6):693–6.
20. Abaci A, Oğuzhan A, Kahraman S, Eryol NK, Unal S, Arinç H, Ergin A: **Effect of diabetes mellitus on formation of coronary collateral vessels.** *Circulation* 1999, **99**(17):2239–2242.
21. Fulton WFM: **The dynamic factor in enlargement of coronary arterial anastomoses, and paradoxical changes in the subendocardial plexus.** *Br Heart J* 1964, **26**:39–50.
22. Helfant RH, Kemp HG, Gorlin R: **Coronary atherosclerosis, coronary collaterals, and their relation to cardiac function.** *Ann Intern Med* 1970, **73**:189–193.
23. Sezer M, Ozcan M, Okcular I, Elitok A, Umman S, Umman B, Tayyareci Y, Olcay A, Nisanci Y, Bilge AK, Meric M: **A potential evidence to explain the reason behind the devastating prognosis of coronary artery disease in uraemic patients: renal insufficiency is associated with poor coronary collateral vessel development.** *Int J Cardiol* 2007, **115**(3):366–372.
24. Hsu PC, Lin TH, Su HM, Juo SH, Lai WT, Sheu SH: **Association between endostatin G4349A polymorphism and coronary collaterals in the Chinese population.** *Acta Cardiol Sin* 2012, **28**:10–16.
25. Hsu PC, Juo SH, Su HM, Chen SC, Tsai WC, Lai WT, Sheu SH, Lin TH: **Predictor of poor coronary collaterals in chronic kidney disease population with significant coronary artery disease.** *BMC Nephrol* 2012, **13**(1):98.

doi:10.1186/1756-0500-6-105

Cite this article as: Hsu et al.: Influence of high-density lipoprotein cholesterol on coronary collateral formation in a population with significant coronary artery disease. *BMC Research Notes* 2013 **6**:105.

Submit your next manuscript to BioMed Central and take full advantage of:

- Convenient online submission
- Thorough peer review
- No space constraints or color figure charges
- Immediate publication on acceptance
- Inclusion in PubMed, CAS, Scopus and Google Scholar
- Research which is freely available for redistribution

Submit your manuscript at
www.biomedcentral.com/submit

