



Published in final edited form as:

*Pediatr Res.* 2011 June ; 69(6): 538–543. doi:10.1203/PDR.0b013e3182139227.

## The 1p13.3 LDL (C)-Associated Locus Shows Large Effect Sizes in Young Populations

Joseph M. Devaney, Paul D. Thompson, Paul S. Visich, William A. Saltarelli, Paul M. Gordon, E. Funda Orkunoglu-Suer, Heather Gordish-Dressman, Brennan T. Harmon, Margaret K. Bradbury, Karuna Panchapakesan, Rahul Khianey, Monica J. Hubal, Priscilla M. Clarkson, Linda S. Pescatello, Robert F. Zoeller, Niall M. Moyna, Theodore J.

Angelopoulos, William E. Kraus, and Eric P. Hoffman

Department of Integrative Systems Biology [J.M.D., E.F.O-S., H.G-D., B.T.H., M.K.B., K.P., R.K., M.J.H., E.P.H.], Research Center for Genetic Medicine, Children's National Medical Center, Washington, DC 20010; Division of Cardiology [P.D.T.], Henry Low Heart Center, Hartford Hospital, Hartford, CT 06102; Human Performance Laboratory [P.S.V., W.A.S.], Central Michigan University, Mount Pleasant, MI 48859; Laboratory for Physical Activity and Exercise Intervention Research [P.M.G.], University of Michigan, Ann Arbor, MI 48108 ; Department of Kinesiology [P.M.C.], University of Massachusetts, Amherst, MA, 01003; Department of Kinesiology and Human Performance Laboratory [L.S.P.], University of Connecticut, Storrs, CT 06269; Department of Exercise Science and Health Promotion [R.F.Z.], Florida Atlantic University, Davie, FL 33314; Department of Sport Science and Health [N.M.M.], Dublin City University, Dublin 9, Ireland, UK; Center for Lifestyle Medicine and Department of Health Professions [T.J.A.], University of Central Florida, Orlando, FL 32816; Department of Medicine [W.E.K.], Duke University Medical Center, Durham, NC 27710

### Abstract

Genome-wide association studies (GWAS) have identified polymorphic loci associated with coronary artery disease (CAD) risk factors (i. e. serum lipids) in adult populations (42–69 yrs). We hypothesized that younger populations would show a greater relative genetic component due to fewer confounding variables. We examined the influence of 20 GWAS loci associated with serum lipids and insulin metabolism, in a university student cohort (n=548; mean age= 24 yrs), and replicated statistically associated results in a second study cohort of primary school students (n=810, mean age= 11.5 yrs). 19 loci showed no relationship with studied risk factors in young adults. However, the ancestral allele of the rs646776 (*SORT1*) locus was strongly associated with increased low density lipoprotein cholesterol {LDL (C)} in young adults (TT:  $97.6 \pm 1.0$  mg/dL {n=345}, vs. CT/CC:  $87.3 \pm 1.0$  mg/dL {n=203};  $p = 3 \times 10^{-6}$ ) and children (TT:  $94.0 \pm 1.3$  mg/dL {n=551}, vs. CT/CC:  $84.7 \pm 1.4$  mg/dL {n=259};  $p = 4 \times 10^{-6}$ ). This locus is responsible for 3.6% of population variance in young adults and 2.5% of population variance in children. The effect size of the *SORT1* locus is considerably higher in young populations (2.5%–4.1%) compared to older subjects (1%).

---

Copyright © 2011 International Pediatric Research Foundation, Inc. All rights reserved.

Corresponding author: Eric P. Hoffman, PhD, Department of Integrative Systems Biology, Research Center for Genetic Medicine, Children's National Medical Center, 111 Michigan Ave, NW, Washington DC 20010, USA, Phone: 202.476.6011, Fax: 202.476.6014, ehoffman@cnmcresearch.org.

**Publisher's Disclaimer:** This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final citable form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

## INTRODUCTION

Coronary artery disease is a major health problem of industrialized countries. Some risk factors for CAD such as insulin resistance and type II diabetes mellitus (T2DM) are also rising even among pediatric populations primarily because of changes in life-style including a high caloric intake and low physical activity.

Many GWA studies have been used to identify genetic risk factors for markers associated with selected CAD risk factors (1, 2). Typically, GWAS has used dichotomous traits (affected vs. non-affected controls) to identify new loci for such CAD risk factors as glucose levels (insulin resistance) (1, 2), T2DM (1), obesity (BMI) (3), blood pressure (4), and serum lipids (5).

A consistent finding of all of these GWAS studies of individual CAD risk factors and associated phenotypes has been the relatively weak contribution of any specific locus to the relative risk. Dichotomous studies of disease risk for T2DM, for example, have typically found only a 1.1-fold increase in relative risk for any specific genetic marker. This low effect size shows a marked difference from twin studies for each of the biochemical, physiological, and anthropomorphic measures associated with the development of such CAD risk factors as T2DM (6). Moreover, the poor predictive power of genetic loci, either singly or in isolation, have led some to conclude that clinical measures are in fact more prognostic than genetic predispositions (7). Likewise, quantitative (continuous) trait studies typically report that specific genetic components have an effect size of 1% or less. This has led to the hypothesis that traits associated with CAD risk factors are highly polygenic, with many loci each contributing a very small proportion of variation.

We hypothesized that study of younger populations could prove more valuable in identifying genetic contributions to CAD risk for two reasons: genetic markers associated with CAD risk factors could be highly valuable tools to identify individuals who might benefit most from early interventions and young adults lack many of the co-morbidities that complicate gene/phenotype associations. The co-variance of the co-morbidities in older subjects would serve to lessen the effect size of any specific genetic locus. Thus, young adults may show much larger effect sizes. Finally, the study of young adults should uncover the genetic loci associated with onset of the disease, rather than later disease progression. There are also disadvantages to studying young adults. The variance of values is considerably less in young adults, with fewer subjects showing abnormal values of the marker, and this might be expected to reduce statistical power. However, we hypothesized that the reduced statistical power would be offset by larger effect sizes of strong genetic components.

In this study, we analyzed 20 GWAS-identified single nucleotide polymorphisms (SNP) in younger populations. The loci studied have been replicated in numerous studies, with seven loci associated with serum lipids (5, 8, 9), 10 associated with T2DM (1), and three associated with development of CAD (8, 10). The initial cohort studied was college-aged individuals (Functional Polymorphism Associated with Human Muscle Size and Strength; FAMUSS), and gene associations queried for LDL (C), high-density lipoprotein cholesterol {HDL (C)}, triglycerides, glucose, homoeostasis model assessment (HOMA), and insulin. Validation of positive associations was performed in a cohort of 6<sup>TH</sup> grade volunteers (Cardiovascular Health Intervention Program; CHIP).

We establish that the 1p13.3 LDL (C) locus was strongly associated with a CAD risk factor in both of these young populations. LDL (C) is strong predictor of CAD (11), and this locus could be used to identify individuals who could benefit from early interventions to prevent CAD.

## Methods

Genotyping of the 20 GWAS-associated SNPs was performed in FAMUSS to identify loci associated with the following CAD risk factors: LDL (C), HDL (C), triglycerides (TG), glucose, HOMA, and insulin with and without adjustment for age and BMI. Significant loci were validated in the CHIP cohort.

### FAMUSS cohort

The FAMUSS study design is a population-based study of individuals (average age = 24 yrs old) designed to identify genetic factors associated with skeletal muscle changes as a result of resistance training (12). We collected selected baseline CAD risk factors such as serum lipids {LDL (C), HDL (C), and TG}, glucose, and insulin following an overnight fast (Quest Diagnostics, Chantilly, VA, 20153) in 548 White participants. The HOMA values were obtained using this calculation:  $(\text{insulin} \times \text{fasting glucose})/22.5$  (13). The study was approved by the Children's National Medical Center Institutional Review Board (protocol #2449) in compliance with the Helsinki Declaration. Informed consent was obtained from all subject participants.

### CHIP cohort

CHIP is a population-based study of 6<sup>th</sup> graders (11.5 yrs of age) in Michigan that includes a screening component and health education program (manuscript in preparation). The screenings have included the following CAD risk factor assessments: physical activity, cardiovascular fitness, body composition, blood pressure, family history, fasted blood lipids {LDL (C), HDL (C), and TG}, glucose, dietary assessments, grip strength, and a blood spot sample. Serum lipids and markers of glucose metabolism were available for 810 participants. The study was approved by the Central Michigan University in compliance with the Helsinki Declaration. Informed consent was obtained from all mothers and/or fathers of participants in the CHIP study. De-identified blood spots were sent to Children's National Medical Center for genotyping, and genotyping was approved by the CNMC IRB.

### The Study of Targeted Risk Reduction Intervention through Defined Exercise study (STRRIDE)

This study examines defined modes, doses and intensities of exercise in a structured exercise program, with the goal to define the relationship between mode, amount and intensities of exercise and improvements in risk parameters of cardiovascular health. The STRRIDE encompassed 555 middle-aged subjects with features of metabolic syndrome ( $25 < \text{BMI} < 35$ ; fasting hyperinsulinemia  $\{ > 10 \text{ IU/ml} \}$ , and dyslipidemia with LDL cholesterol between 130–190, and/or HDL  $< 45$  for women,  $< 35$  for men). Subjects were assigned to one of six groups addressing exercise mode, intensity and duration. STRRIDE participants signed informed consent associated with IRB approval by Duke University Medical Center.

### Genotyping Methods

Genomic DNA was isolated using Puregene kits (Gentra Systems, MN) from peripheral blood (FAMUSS), or from blood spots using Qiagen kits (CHIP). Genotyping was done using TaqMan allele discrimination assays (14). Both alleles were detected simultaneously and automatically called (see Table 1).

### Statistical Analyses

All genotype data sets were tested for Hardy-Weinberg equilibrium by using a  $\chi^2$  tests to compare the observed genotype frequencies to those expected under equilibrium. Normality

of each quantitative trait was tested using the Shapiro-Wilk normality test and any trait not meeting the criteria was appropriately transformed.

Bivariate correlation analyses of each quantitative measurement showed several significant correlations with age and baseline weight; therefore, associations between each SNP and quantitative phenotype were assessed using analysis of covariance methods using both dominant genetic models and quantitative trait models.

Linear regression analysis, including likelihood ratio tests between full (containing genotype and covariates) and constrained (containing covariates only) models, were performed to estimate the proportion of variance in volumetric measurements attributable to each SNP's genotype. All statistical analyses were performed using Stata V.8 (StataCorp, College Park, TX).

The CHIP cohort was used as a validation data set. Thus, only statistically significant associations surviving multiple testing thresholds seen in FAMUSS were subsequently tested in the CHIP cohort. Statistical significance threshold was established as  $p < 0.05$  as this was a validation cohort with no multiple testing.

## RESULTS

### Analysis of Selected CAD Risk Factor GWAS loci in young adults

The FAMUSS cohort was utilized as the test cohort. Traits used for association studies were fasting glucose (mmol/L), HOMA, insulin ( $\mu\text{U/ml}$ ), LDL (C) (mg/dl), HDL (C) (mg/dl), triglycerides (mg/dL), with and without adjustment for BMI ( $\text{kg/m}^2$ ). Average values for all subjects and stratified by sex are provided in Table 2.

The percentage of subjects showing abnormal levels of each CAD risk factor is provided in Table 3. BMI was  $>30$  in 12.0% of all subjects (9.5% of females; 16.5% of males). A relatively large proportion of volunteers showed abnormal HDL (C) levels (81% of males; 29% of females). Females showed an overall lower risk for mean blood pressure (mmHg) 100 (2.6% abnormal values in females; 14.5% in males), and for the diagnosis of metabolic syndrome (4.6% of females; 18.0% of males).

Loci previously reported to be associated with CAD, CAD risk factors, or glucose metabolism phenotypes were genotyped in FAMUSS (Table 1; Table 4). Resulting genotypes were tested for Hardy-Weinberg equilibrium; two SNPs showed divergence from expected allele frequencies (rs4846914 and rs1333049). Twenty loci were tested against six phenotypes {fasting glucose, triglycerides, HDL (C), LDL (C), fasting insulin, and HOMA} in FAMUSS. This required adjustment for multiple testing with a p-value threshold of  $< 0.00042$  for significance (adjusted for the testing of 20 SNPs and six phenotypes).

Nineteen of the 20 loci tested did not reach the adjusted p-value threshold of 0.00042 for any T2DM-related or lipid trait in the FAMUSS group. Given the lower average serum values in young adults compared to older adults, and lower variance, our data does not preclude the other 19 loci from having a phenotype effect in a larger population. We are underpowered to detect a  $<1\%$  effect size published in larger adult studies.

A single locus met our threshold for significance, and this was the rs646776 locus at 1p13.3, containing the *SORT1* gene with association for circulating LDL (C) levels (Table 5; Figure 1A). There was a dose effect of the polymorphism, suggesting a quantitative or additive effect of the alleles, however, the numbers of homozygotes for the rare allele were relatively low (Figure 1A). The association between rs646776 and LDL (C) was significant for the entire cohort ( $p = 3.0 \times 10^{-6}$ ), and significant in both male ( $p = 0.009$ ) and female ( $p =$

$8.8 \times 10^{-5}$ ) sub-cohorts. Inclusion of age and/or BMI as covariates in the statistical model showed little change in statistical support for the association of rs646776 with LDL (C). The amount of variation in serum LDL (C) explained by the rs646776 genotype was 3.6% for the entire cohort (4.1% in females; 3.0% in males; Table 5).

LDL (C) levels have been shown to have a weak correlation with BMI in children and adolescents (ages 3–18 yrs old) (15). Oxidized LDL (C) has a strong relationship to waist circumference and has been shown to have a strong relationship to LDL (C) levels (16–19). Our association data in young adults has shown that more of the variation of LDL (C) in 24 yr olds was due to rs646776 than to an increase BMI which is supported by the findings in the Lee et al. 2009 paper (15). To test this, regression models of LDL (C) were done with and without rs646766 genotype and BMI to determine the percent variation attributable to each factor. In both the FAMUSS and CHIP cohorts, more of the variation in LDL (C) can be attributed to rs646766 than BMI (3.6% versus 3.0% in FAMUSS; 2.5% versus 1.4% in CHIP). While there are significant associations between LDL (C) and BMI in both cohorts ( $p=0.03$  in FAMUSS;  $p=0.02$  in CHIP), the associations between LDL (C) and rs646766 are statistically much more significant ( $p=0.000003$  in FAMUSS;  $p=0.00008$  in CHIP).

### **The association between rs646776 and LDL (C) is replicated in the 11.5 yr old CHIP cohort**

The CHIP cohort is comprised of 810 predominantly White 6<sup>th</sup> grade students attending public school in Michigan. Average values for selected CAD risk factors in the 11.5 yr old CHIP volunteers were healthier compared to the FAMUSS, as expected given the well-accepted increase in risk factors with age (Table 2).

We then looked at the abnormal threshold levels for each phenotype, using age-adjusted criteria (Table 3) (20). In this population-based study, 44.6% of children met the criteria for an abnormally high BMI (41.8% for girls, 47.6% for boys). 40% showed abnormal total cholesterol, while about 25% of the children showed abnormal values for HDL (C), LDL (C), and TG as individual laboratory phenotypes.

As this was a replication cohort, we tested only the association of rs646776 with serum LDL (C). There was a strong statistical association between rs646776 and LDL (C), similar to that observed in FAMUSS (Figure 1B; Table 6). There was a similar dosage effect of the genotype, with homozygotes for the ancestral common allele showing the highest LDL (C) levels. The percentage of variation explained by genotype was 2.5% for the entire cohort (2.7% for females; 2.0% for males).

### **The rs646776 locus at 1p13.3 1 shows a greater effect in females than males**

In both populations, females showed a stronger percent variation attributable to *SORT1* genotype (see Tables 4 and 5). In the FAMUSS study, the *SORT1* locus contributed 4.1% to LDL (C) values in females while contributing 3.0% to males. We see a similar trend in the validation population, CHIP, where *SORT1* contributed 2.7% to LDL (C) values in females and 2.0% in males.

### **The rs646776 locus at 1p13.3 1 shows a greater effect in young populations, compared to older, sicker populations**

The association between rs646776 locus at 1p13.3 and LDL (C) has been studied in many older populations (5), some with overt T2DM (9). While each study has shown a similar effect of ancestral (common) allele associated with higher LDL (C), the effect size (proportion of variation explained by genotype) has been consistently quite small (~1%). We hypothesized that the larger effect size seen in the younger populations studied here (e.g. 4.1% in 24 yr old females; 2.7% in 11.5 yr old females) could be explained by the relative

absence of co-morbidities that serve to obscure underlying genetic associations. To test this, we compared the association data from FAMUSS and CHIP to STRRIDE. STRRIDE includes older (age = 54 yrs; Table 7) subjects with features of metabolic syndrome, but without a T2DM diagnosis (21). As expected, STRRIDE participants showed much higher average serum LDL (C) values (Figure 2) but not a significant association of LDL (C) with rs646776 genotype. This contrasts to the 11.5 yr old CHIP children, and 24 yr old FAMUSS young adults that showed a significant genotype correlation. However, the 24 yr olds showed a higher genotype effect than the 11.5 yr olds (3.6% vs. 2.5%, respectively).

## DISCUSSION

GWAS studies have enabled the identification of key polymorphisms linked to complex disease, including T2DM (1) and related phenotypes (see reviews from Rideerstrale and Groop 2009) (22). This approach is frequently cited as holding potential to transform the practice of medicine. Most frequently, the integration of genetic risk factors into clinical practice is envisioned through preventive medicine. For example, those at highest risk for T2DM and/or CAD can be identified prior to onset of clinical symptoms, with interventions targeted towards those at greatest risk, perhaps personalizing interventions for specific genotypes. There are some shortcomings to this rationale. First, clinical trials will be needed to provide the evidence-based data to bring such pre-symptomatic interventions into standard of care. Since T2DM and CAD take decades to develop, the progression to overt disease is unlikely to be a practical endpoint in such trials. Thus, alternative sensitive and reliable endpoints are needed that can be assessed over a reasonable amount of time (e.g. 1 yr). Second, many CAD risk factor loci such as those for T2DM have been identified with case/control cohorts, with advanced or end-stage disease. Thus, it is possible, if not likely, that existing GWAS CAD risk factor loci identified in adults are associated with advanced development of the risk factor. We hypothesized that the identification of T2DM- and dyslipidemia-related loci showing association with related biomarkers in young population-based cohorts would overcome many of these limitations.

We tested 20 CAD risk factors and T2DM GWAS loci (Table 4), and found that the large majority (19/20) showed no evidence of association with related phenotypes as quantitative traits in a young adult (24 yr old) population. The phenotypes tested were serum lipids (LDL (C), HDL (C), TG), glucose, HOMA, and insulin resistance with and without adjustment for age and BMI. One locus, rs646776 at 1p13.3, showed strong associations with serum LDL (C) in our 24 yr old FAMUSS cohort and this association (Table 5; Figure 1A) remained significant after adjustment for multiple testing in the entire cohort ( $p = 3.0 \times 10^{-6}$ ), males ( $p = 0.009$ ) and females ( $p = 8.8 \times 10^{-5}$ ). The association between rs646776 and LDL (C) was validated in a population-based 11.5 yr old replication cohort, and found to show similar significance (Table 6; Figure 1B).

We found a considerably larger effect size of the 1p13.3 LDL (C)-associated locus in young populations, compared to the previously published older cohorts. All previous studies have shown that rs646776 explains approximately 1% of the population variation in LDL (C) levels (5, 8). For comparison, a recent GWAS identified 30 loci for dyslipidemia (including the 1p13.3 locus) where the cumulative effect size of all 30 loci was approximately 5% of the population variance in lipoprotein levels (5). In contrast, our data in 24 yr olds showed the rs646776 SNP explained 3.6% of population variation – a considerably higher effect size than seen in most other loci (Table 5). This was despite lower average serum LDL (C) levels in our younger population (Figure 2), and lower population variance in LDL (C) values. Our data in 6<sup>th</sup> graders (Table 6) also showed larger effect sizes than previously published (2.5% vs. 1%); however, the effect size in our young population (11.5 yrs old) was not significantly higher than our 24 yr olds (FAMUSS; 3.6% effect size).

The specific functional variant at the 1p13.3 LDL (C)-associated locus has recently been shown to be a non-coding SNP in the gene promoter of the *SORT1* gene, a multi-ligand sorting receptor directly involved in lipid trafficking in the liver (23). This locus showed results consistent with our hypothesis that younger populations would show a stronger genetic component in population variation in serum markers. However, the other 19 loci tested did not show significance in the 24 yr olds. This could be due to poor statistical power in younger subjects due to lower average serum marker levels and lower variance in blood levels for the serum markers. Also, some genetic loci significant in older subjects may require the onset of age-related co-variables before showing the published genetic associations. For example, the well-studied *TCF7L2* T2DM locus has been associated with renal failure in diabetics and non-diabetics (24); a phenotype not expected to be expressed in younger subjects.

The rs646776 locus may provide the first opportunity to conduct clinical trials aimed at genotype-stratified prevention of dyslipidemia. Specifically, in our 11.5 yr old 6<sup>th</sup> grader cohort, 27.3% of children showed levels of LDL (C) above the threshold considered abnormal (Table 3). Increasing LDL (C) levels increase the risk of CAD (11). Interventions can be designed to test the efficacy of treatment strategies such as dietary interventions and low dose, lifelong lipid-lowering medication to decrease LDL (C) in children identified at risk by rs646776 genotype.

Our findings were consistent with previous reports showing that the ancestral common allele is associated with higher LDL (C) levels. Thus, the larger population is at high risk for elevated LDL (C), whereas the more recent derived allele imparts protection from high LDL (C) levels to the subset of rare allele carriers. The relatively low relative risk associated with rs646776 and T2DM (1.1 fold), and small effects sizes (1%) in older sicker populations is likely a consequence of the majority (2/3<sup>rd</sup>) of individuals carrying that high risk genotype for increased LDL (C) levels (homozygous for the common rs646776 genotype), and that multiple environmental and polygenic confounders obscure some of the stronger effects seen earlier in young adults (Figure 2).

## Acknowledgments

**Statement of financial support:** This work was supported by NIAMS—5R01AR055100-07 (FAMUSS), NIDDK—1R01DK081559-01A1 (STRIDE) and Owosso Memorial Health Care System (CHIP).

P.D.T is supported by a research grants from GlaxoSmithKline, Merck, Roche, Pfizer, AstraZeneca, B. Braun, Genomas, and Hoffman-LaRoche, served as a consultant for Astra Zenica, Merck, Schering-Plough, Takeda, Roche, Genomas, Abbott and Runners World; received speaker's fees from Merck, Pfizer, Abbott, Astra Zenica, Schering-Plough, and is a stock holder of Zoll, General Electric, JA Wiley Publishing, Zimmer, Medtronic, Merck, Isis and J&J.

## ABBREVIATIONS


<b>CAD</b>	Coronary Artery Disease
<b>CHIP</b>	Cardiovascular Health Intervention Program
<b>FAMUSS</b>	Functional Polymorphism Associated with Human Muscle Size and Strength
<b>GWAS</b>	Genome-Wide Association Study
<b>HOMA</b>	Homoeostasis Model Assessment
<b>T2DM</b>	Type II Diabetes Mellitus
<b>SNP</b>	Single Nucleotide Polymosphim

**STRIDE** Studies of a Targeted Risk Reduction Intervention through Defined Exercise**REFERENCES**

1. McCarthy MI, Zeggini E. Genome-wide association studies in type 2 diabetes. *Curr Diab Rep.* 2009; 9:164–171. [PubMed: 19323962]
2. Ioannidis JP. Prediction of cardiovascular disease outcomes and established cardiovascular risk factors by genome-wide association markers. *Circ Cardiovasc Genet.* 2009; 2:7–15. [PubMed: 20031562]
3. Walley AJ, Asher JE, Froguel P. The genetic contribution to non-syndromic human obesity. *Nat Rev Genet.* 2009; 10:431–442. [PubMed: 19506576]
4. Ehret GB. Genome-wide association studies: contribution of genomics to understanding blood pressure and essential hypertension. *Curr Hypertens Rep.* 2010; 12:17–25. [PubMed: 20425154]
5. Kathiresan S, Willer CJ, Peloso GM, Demissie S, Musunuru K, Schadt EE, Kaplan L, Bennett D, Li Y, Tanaka T, Voight BF, Bonnycastle LL, Jackson AU, Crawford G, Surti A, Guiducci C, Burt NP, Parish S, Clarke R, Zelenika D, Kubalanza KA, Morken MA, Scott LJ, Stringham HM, Galan P, Swift AJ, Kuusisto J, Bergman RN, Sundvall J, Laakso M, Ferrucci L, Scheet P, Sanna S, Uda M, Yang Q, Lunetta KL, Dupuis J, de Bakker PI, O'Donnell CJ, Chambers JC, Kooner JS, Hercberg S, Meneton P, Lakatta EG, Scuteri A, Schlessinger D, Tuomilehto J, Collins FS, Groop L, Altshuler D, Collins R, Lathrop GM, Melander O, Salomaa V, Peltonen L, Orho-Melander M, Ordovas JM, Boehnke M, Abecasis GR, Mohlke KL, Cupples LA. Common variants at 30 loci contribute to polygenic dyslipidemia. *Nat Genet.* 2009; 41:56–65. [PubMed: 19060906]
6. Beck-Nielsen H, Vaag A, Poulsen P, Gaster M. Metabolic and genetic influence on glucose metabolism in type 2 diabetic subjects—experiences from relatives and twin studies. *Best Pract Res Clin Endocrinol Metab.* 2003; 17:445–467. [PubMed: 12962696]
7. Jakobsdottir J, Gorin MB, Conley YP, Ferrell RE, Weeks DE. Interpretation of genetic association studies: markers with replicated highly significant odds ratios may be poor classifiers. *PLoS Genet.* 2009; 5:e1000337. [PubMed: 19197355]
8. Kathiresan S, Melander O, Anevski D, Guiducci C, Burt NP, Roos C, Hirschhorn JN, Berglund G, Hedblad B, Groop L, Altshuler DM, Newton-Cheh C, Orho-Melander M. Polymorphisms associated with cholesterol and risk of cardiovascular events. *N Engl J Med.* 2008; 358:1240–1249. [PubMed: 18354102]
9. Kathiresan S, Melander O, Guiducci C, Surti A, Burt NP, Rieder MJ, Cooper GM, Roos C, Voight BF, Havulinna AS, Wahlstrand B, Hedner T, Corella D, Tai ES, Ordovas JM, Berglund G, Vartiainen E, Jousilahti P, Hedblad B, Taskinen MR, Newton-Cheh C, Salomaa V, Peltonen L, Groop L, Altshuler DM, Orho-Melander M. Six new loci associated with blood low-density lipoprotein cholesterol, high-density lipoprotein cholesterol or triglycerides in humans. *Nat Genet.* 2008; 40:189–197. [PubMed: 18193044]
10. Erdmann J, Grosshennig A, Braund PS, König IR, Hengstenberg C, Hall AS, Linsel-Nitschke P, Kathiresan S, Wright B, Tregouet DA, Cambien F, Bruse P, Aherrahrou Z, Wagner AK, Stark K, Schwartz SM, Salomaa V, Elosua R, Melander O, Voight BF, O'Donnell CJ, Peltonen L, Siscovick DS, Altshuler D, Merlini PA, Peyvandi F, Bernardinelli L, Ardissino D, Schillert A, Blankenberg S, Zeller T, Wild P, Schwarz DF, Tiret L, Perret C, Schreiber S, El Mokhtari NE, Schafer A, Marz W, Renner W, Bugert P, Kluter H, Schrezenmeier J, Rubin D, Ball SG, Balmforth AJ, Wichmann HE, Meitinger T, Fischer M, Meisinger C, Baumert J, Peters A, Ouwehand WH, Deloukas P, Thompson JR, Ziegler A, Samani NJ, Schunkert H. New susceptibility locus for coronary artery disease on chromosome 3q22.3. *Nat Genet.* 2009; 41:280–282. [PubMed: 19198612]
11. Ingelsson E, Schaefer EJ, Contois JH, McNamara JR, Sullivan L, Keyes MJ, Pencina MJ, Schoonmaker C, Wilson PW, D'Agostino RB, Vasan RS. Clinical utility of different lipid measures for prediction of coronary heart disease in men and women. *JAMA.* 2007; 298:776–785. [PubMed: 17699011]
12. Thompson PD, Moyna N, Seip R, Price T, Clarkson P, Angelopoulos T, Gordon P, Pescatello L, Visich P, Zoeller R, Devaney JM, Gordish H, Bilbie S, Hoffman EP. Functional polymorphisms




- associated with human muscle size and strength. *Med Sci Sports Exerc.* 2004; 36:1132–1139. [PubMed: 15235316]
13. Keskin M, Kurtoglu S, Kendirci M, Atabek ME, Yazici C. Homeostasis model assessment is more reliable than the fasting glucose/insulin ratio and quantitative insulin sensitivity check index for assessing insulin resistance among obese children and adolescents. *Pediatrics.* 2005; 115:e500–e503. [PubMed: 15741351]
  14. Orkunoglu-Suer FE, Gordish-Dressman H, Clarkson PM, Thompson PD, Angelopoulos TJ, Gordon PM, Moyna NM, Pescatello LS, Visich PS, Zoeller RF, Harmon B, Seip RL, Hoffman EP, Devaney JM. INSIG2 gene polymorphism is associated with increased subcutaneous fat in women and poor response to resistance training in men. *BMC Med Genet.* 2008; 9:117. [PubMed: 19105843]
  15. Lee JM, Gebremariam A, Card-Higginson P, Shaw JL, Thompson JW, Davis MM. Poor performance of body mass index as a marker for hypercholesterolemia in children and adolescents. *Arch Pediatr Adolesc Med.* 2009; 163:716–723. [PubMed: 19652103]
  16. Holvoet P, Kritchevsky SB, Tracy RP, Mertens A, Rubin SM, Butler J, Goodpaster B, Harris TB. The metabolic syndrome, circulating oxidized LDL, and risk of myocardial infarction in well-functioning elderly people in the health, aging, and body composition cohort. *Diabetes.* 2004; 53:1068–1073. [PubMed: 15047623]
  17. Holvoet P, Lee DH, Steffes M, Gross M, Jacobs DR Jr. Association between circulating oxidized low-density lipoprotein and incidence of the metabolic syndrome. *JAMA.* 2008; 299:2287–2293. [PubMed: 18492970]
  18. Kopprasch S, Pietzsch J, Kuhlisch E, Fuecker K, Temelkova-Kurktschiev T, Hanefeld M, Kuhne H, Julius U, Graessler J. In vivo evidence for increased oxidation of circulating LDL in impaired glucose tolerance. *Diabetes.* 2002; 51:3102–3106. [PubMed: 12351454]
  19. Lapointe A, Couillard C, Piche ME, Weisnagel SJ, Bergeron J, Nadeau A, Lemieux S. Circulating oxidized LDL is associated with parameters of the metabolic syndrome in postmenopausal women. *Atherosclerosis.* 2007; 191:362–368. [PubMed: 16677652]
  20. Kavey RE, Daniels SR, Lauer RM, Atkins DL, Hayman LL, Taubert K. American Heart Association guidelines for primary prevention of atherosclerotic cardiovascular disease beginning in childhood. *Circulation.* 2003; 107:1562–1566. [PubMed: 12654618]
  21. Slentz CA, Aiken LB, Houmard JA, Bales CW, Johnson JL, Tanner CJ, Duscha BD, Kraus WE. Inactivity, exercise, and visceral fat. STRRIDE: a randomized, controlled study of exercise intensity and amount. *J Appl Physiol.* 2005; 99:1613–1618. [PubMed: 16002776]
  22. Ridderstråle M, Groop L. Genetic dissection of type 2 diabetes. *Mol Cell Endocrinol.* 2009; 297:10–17. [PubMed: 19000735]
  23. Musunuru K, Strong A, Frank-Kamenetsky M, Lee NE, Ahfeldt T, Sachs KV, Li X, Li H, Kuperwasser N, Ruda VM, Pirruccello JP, Muchmore B, Prokunina-Olsson L, Hall JL, Schadt EE, Morales CR, Lund-Katz S, Phillips MC, Wong J, Cantley W, Racie T, Ejebe KG, Orho-Melander M, Melander O, Kotliansky V, Fitzgerald K, Krauss RM, Cowan CA, Kathiresan S, Rader DJ. From noncoding variant to phenotype via SORT1 at the 1p13 cholesterol locus. *Nature.* 2010; 466:714–719. [PubMed: 20686566]
  24. Köttgen A, Hwang SJ, Rumpersaud E, Coresh J, North KE, Pankow JS, Meigs JB, Florez JC, Parsa A, Levy D, Boerwinkle E, Shuldiner AR, Fox CS, Kao WH. TCF7L2 variants associate with CKD progression and renal function in population-based cohorts. *J Am Soc Nephrol.* 2008; 19:1989–1999. [PubMed: 18650481]



**Figure 1. 1p13.3 variant (rs646776) shows an effect on LDL (C) in both FAMUSS and CHIP cohorts**

Both sexes are shown for the FAMUSS study (1A) and the CHIP study (1B). In both population-based studies, individuals with the TT genotype had significantly higher LDL (C) values than the CT/CC genotypes. Figure 1A: \* $p=0.000003$ ;  $N=345$  for TT,  $N=203$  for CT/CC. Figure 1B: \*\* $p=0.000008$ ;  $N=551$  for TT,  $N=259$  for CT/CC.



**Figure 2. LDL (C) cholesterol as a function of 1p genotype shows higher effect size in young population-based cohorts compared to older subjects with features of metabolic syndrome**  
For the CHIP cohort, LDL (C) was significantly different between genotypes (\* $p < 0.001$ ) with a mean  $\Delta$ LDL (C) = 0.103 and with genotype explaining 2.5% of the variability in LDL (C). For the FAMUSS cohort, LDL (C) was significantly different between genotypes (\* $p < 0.001$ ) with a mean  $\Delta$ LDL (C) = 0.116 and with genotype explaining 3.6% of the variability in LDL (C). For the STRRIDE cohort, LDL (C) was not significantly different between genotypes with a mean  $\Delta$ LDL (C) = 0.014 and with genotype explaining <0.1% of the variability in LDL (C).

**Table 1**

Assay ID#s for SNPs genotyped in FAMUSS Study.

Region	SNP	ABI ASSAY ID#
1p31.3	rs12130333	C__31145189_10
1p13.3	rs646776	C__3160062_10
1q42.13	rs4846914	C__27968379_10
3p25.2	rs1801282	C__1129864_10
3q22.3	rs9818870	C__26070086_10
3q27.2	rs4402960	C__2165199_10
6p22.3	rs7756992	C__2504058_20
6p22.3	rs10946398	C__31635230_10
7q11.23	rs17145738	C__2632544_10
8q24.11	rs13266634	C__357888_10
8q24.13	rs17321515	C__33068431_10
9p21.3	rs1333049	C__1754666_10
9p21.3	rs10811661	C__31288917_10
10q23.33	rs1111875	C__11214581_10
10q23.33	rs7923837	C__31982553_10
10q25.2	rs7901695	C__384583_10
11p15.1	rs5219	C__11654065_10
12q24.11	rs2338104	C__2877290_10
12q24.31	rs2259816	C__11753816_10
19p13.11	rs16996148	C__33703643_10

Table 2

Sample Characteristics for FAMUSS and CHIP Population-Based Cohorts.

Characteristic	FAMUSS (Test Cohort)				CHIP (Replication Cohort)				
	Mean ± SD				Mean ± SD				
	All	Females	Males	All	Females	Males	All	Females	Males
N	548	348	200	810	428	382	810	428	382
Mean Age (years)	23.86 ± 5.71	23.67 ± 5.71	24.19 ± 5.71	11.52 ± 0.81	11.44 ± 0.78	11.61 ± 0.83	11.52 ± 0.81	11.44 ± 0.78	11.61 ± 0.83
BMI (kg/m <sup>2</sup> )*	24.48 ± 4.84	23.89 ± 4.64	25.49 ± 5.03	21.56 ± 5.13	21.67 ± 5.45	21.44 ± 4.73	21.56 ± 5.13	21.67 ± 5.45	21.44 ± 4.73
Total cholesterol (mg/dl)	166.49 ± 31.35	166.62 ± 32.24	166.26 ± 29.81	165.39 ± 31.89	163.88 ± 30.58	167.08 ± 33.26	165.39 ± 31.89	163.88 ± 30.58	167.08 ± 33.26
LDL (C) (mg/dl)	97.54 ± 27.85	94.99 ± 28.02	101.97 ± 27.04	95.21 ± 28.95	93.17 ± 27.01	97.49 ± 30.85	95.21 ± 28.95	93.17 ± 27.01	97.49 ± 30.85
HDL (C) (mg/dl)	48.26 ± 12.82	52.32 ± 11.81	41.21 ± 11.40	51.26 ± 14.43	51.52 ± 14.53	50.97 ± 14.33	51.26 ± 14.43	51.52 ± 14.53	50.97 ± 14.33
Triglycerides (mg/dL)	103.48 ± 52.23	96.62 ± 45.76	115.41 ± 60.17	96.70 ± 51.92	99.17 ± 53.84	93.93 ± 49.61	96.70 ± 51.92	99.17 ± 53.84	93.93 ± 49.61
Insulin (µU/mL)	5.58 ± 4.65	5.26 ± 3.70	6.13 ± 5.92	N/A	N/A	N/A	N/A	N/A	N/A
Fasting Glucose (mmol/L)	86.84 ± 7.43	85.43 ± 7.11	89.30 ± 7.37	95.81 ± 10.25	94.67 ± 10.71	97.09 ± 9.56	95.81 ± 10.25	94.67 ± 10.71	97.09 ± 9.56
HOMA	1.21 ± 1.11	1.11 ± 0.82	1.38 ± 1.48	N/A	N/A	N/A	N/A	N/A	N/A
Mean Blood Pressure (mmHg)** , §	86.71 ± 9.19	85.07 ± 8.55	89.55 ± 9.58	82.40 ± 8.12	81.81 ± 8.51	83.07 ± 7.61	82.40 ± 8.12	81.81 ± 8.51	83.07 ± 7.61
Systolic blood pressure (mmHg)§	115.41 ± 12.07	112.59 ± 11.38	120.30 ± 11.69	108.81 ± 10.66	108.30 ± 10.49	109.37 ± 10.83	108.81 ± 10.66	108.30 ± 10.49	109.37 ± 10.83
Diastolic blood pressure (mmHg)§	72.36 ± 9.94	71.31 ± 9.21	74.18 ± 10.88	69.20 ± 8.90	68.56 ± 9.33	69.92 ± 8.34	69.20 ± 8.90	68.56 ± 9.33	69.92 ± 8.34

\* Mean Blood Pressure defined as (Systolic blood pressure + Diastolic blood pressure × 2)/3

\*\* FAMUSS Female N=347 for BMI due to missing weight

§ FAMUSS Female N=341 and Male N=197 due to missing BP measurement

**Table 3**

Incidence of abnormal values for FAMUSS and CHIP.

<b>FAMUSS (Ave. Age 24 yrs)</b>			
	<b>All</b>	<b>Females</b>	<b>Males</b>
<b>N</b>	548	348	200
<b>BMI (kg/m<sup>2</sup>) 25</b>	34.1%	28.4%	44.0%
<b>BMI (kg/m<sup>2</sup>) 30</b>	12.0%	9.5%	16.5%
<b>Total cholesterol (mg/dl) 240</b>	2.7%	3.5%	1.5%
<b>LDL (C) (mg/dl) 130</b>	12.2%	10.3%	15.5%
<b>HDL (C) (mg/dl) 40 (for males) 50 (for females)</b>	48.0%	29.0%	81.0%
<b>Triglycerides (mg/dL) 150</b>	15.1%	12.1%	20.5%
<b>Insulin resistance</b>	6.9%	5.5%	9.5%
<b>Mean BP (mmHg) 100</b>	6.9%	2.6%	14.5%
<b>Metabolic Syndrome</b>	9.5%	4.6%	18.0%
<b>CHIP (Ave. Age 11.5 yrs)</b>			
	<b>All</b>	<b>Females</b>	<b>Males</b>
<b>N</b>	810	428	382
<b>BMI (kg/m<sup>2</sup>)*</b>	44.6%	41.8%	47.6%
<b>Total cholesterol (mg/dl) 170</b>	41.2%	40.7%	42.0%
<b>LDL (C) (mg/dl)</b>	27.3%	25.9%	28.8%
<b>HDL (C) (mg/dl) 40</b>	23.5%	23.4%	23.6%
<b>Triglycerides (mg/dL) 110</b>	27.8%	29.4%	25.9%
<b>Insulin resistance</b>	ND**	ND**	ND**
<b>Mean BP (mmHg) &gt; 90<sup>th</sup> percentile, based on CDC growth charts</b>	11.4%	10.7%	12.0%
<b>Metabolic Syndrome</b>	7.6%	7.7%	7.6%

\* Thresholds for abnormal BMI were based on age and sex: Males 10 yrs 19; 11 yrs 20; 12 yrs 21. Females 10 yrs 20; 11 yrs 21; 12 yrs 22.

\*\* ND: Not Determined

BP: Blood Pressure

Table 4

SNPs genotyped in FAMUSS Study.

Region	SNP	Risk Allele	Risk allele Frequency	Nearest Gene (s)	Phenotype associated with SNP
1p31.3	rs12130333	C	0.77	<i>ANGPTL3, DOCK, ATG4C</i>	TG
1p13.3	rs646776	T	0.83	<i>CELSR2, PCSK1, SORT1</i>	LDL
1q42.13	rs4846914	G	0.43	<i>GALNT2</i>	HDL, TG
3p25.2	rs1801282	C	0.87	<i>PPARG2</i>	T2DM
3q22.3	rs9818870	T	0.16	<i>MRAS</i>	CAD
3q27.2	rs4402960	T	0.35	<i>IGFBP2</i>	T2DM
6p22.3	rs7756992	G	0.26	<i>CDKALI</i>	T2DM
6p22.3	rs10946398	C	0.31	<i>CDKALI</i>	T2DM
7q11.23	rs17145738	G	0.92	<i>BCL7B, TBL2, MLXIPL</i>	TG
8q24.11	rs13266634	C	0.71	<i>SCL30A8</i>	T2DM
8q24.13	rs17321515	A	0.55	<i>TRIB1</i>	TG
9p21.3	rs1333049	C	0.51	<i>ANRIL</i>	CAD
9p21.3	rs10811661	T	0.83	<i>CDKN2A/2B</i>	T2DM
10q23.33	rs1111875	G	0.39	<i>HHEX</i>	T2DM
10q23.33	rs7923837	G	0.61	<i>HHEX</i>	T2DM
10q25.2	rs7901695	T	0.72	<i>TCF7L2</i>	T2DM
11p15.1	rs5219	A	0.35	<i>KCNH1</i>	T2DM
12q24.11	rs2338104	G	0.46	<i>KCTD10</i>	HDL
12q24.31	rs2259816	T	0.35	<i>HNF1A</i>	CAD
19p13.11	rs16996148	G	0.07	<i>CILP2, PBX4</i>	LDL, TG

TG: Triglycerides; LDL (C): Low-density lipoprotein cholesterol; HDL (C): High-density lipoprotein; T2DM: Type 2 Diabetes Mellitus; CAD: Coronary Artery Disease

**Table 5**

1p13.3 variant (rs646776) is associated with LDL (C) in FAMUSS cohort (24 yr olds).

Phenotype	N: unadjusted mean $\pm$ SEM	<i>p</i> -value	% variation attributable to genotype
LDL (C) (log)	TT (N=345; 4.58 $\pm$ 0.0) CT/CC (N=203; 4.47 $\pm$ 0.02)	0.000003	3.6%
LDL (C) (log) in females	TT (N=212; 4.56 $\pm$ 0.02) CT/CC (N=136; 4.44 $\pm$ 0.02)	0.000088	4.1%
LDL (C) (log) in males	TT (N=133; 4.622 $\pm$ 0.022) CT/CC (N=67; 4.522 $\pm$ 0.031)	0.0094	3.0%



**Table 6**

Replication data for 1p13.3 variant (rs646776) shows association with LDL (C) in the CHIP cohort (11.5 yr olds).

Phenotype	N: unadjusted mean $\pm$ SEM	<i>p</i> -value	% variation attributable to genotype
LDL (C) (log)	TT (N=551; 4.54 $\pm$ 0.02) CT/CC (N=259; 4.44 $\pm$ 0.01)	0.0000076	2.5%
LDL (C) (log) in females	TT (N=280; 4.53 $\pm$ 0.03) CT/CC (N=148; 4.42 $\pm$ 0.02)	0.0006	2.7%
LDL (C) (log) in males	TT (N=271; 4.56 $\pm$ 0.03) CT/TT (N=; 111; 4.46 $\pm$ 0.02)	0.0061	2.0%

**Table 7**

Sample Characteristics for STRRIDE Study.

Phenotype	Females (N=114)	Males (N=117)
	Mean $\pm$ SD	Mean $\pm$ SD
Age (years)	53.65 $\pm$ 5.61	50.64 $\pm$ 6.64
Triglycerides (mg/dL)	125.02 $\pm$ 59.77	152.60 $\pm$ 78.59
Total cholesterol (mg/dL)	206.61 $\pm$ 22.05	184.39 $\pm$ 28.96
Fasting glucose (mg/dL) *	89.96 $\pm$ 11.32	94.77 $\pm$ 10.17
HDL (C) (mg/dL)	54.99 $\pm$ 14.15	39.22 $\pm$ 8.19
LDL (C) (mg/dL)	130.78 $\pm$ 24.00	121.17 $\pm$ 23.83
Height (m)	1.63 $\pm$ 0.07	1.78 $\pm$ 0.07
Weight (kg)	79.71 $\pm$ 10.56	95.37 $\pm$ 12.58
BMI	29.90 $\pm$ 3.20	30.12 $\pm$ 3.11

\* Glucose values available for only 111 females and 113 males