



CASK and CaMKII function in the mushroom body α'/β' neurons during *Drosophila* memory formation

Bilal R. Malik, John Michael Gillespie and James J. L. Hodge*

School of Physiology and Pharmacology, University of Bristol, Bristol, UK

Edited by:

Claude Desplan, NYU, USA

Reviewed by:

Makoto Sato, Kanazawa University, Japan

Gaia Tavasani, German Center for Neurodegenerative Diseases, Germany

*Correspondence:

James J. L. Hodge, School of Physiology and Pharmacology, University of Bristol, Medical Sciences Building, University Walk, Bristol BS8 1TD, UK.
e-mail: james.hodge@bristol.ac.uk


Ca^{2+} /CaM serine/threonine kinase II (CaMKII) is a central molecule in mechanisms of synaptic plasticity and memory. A vital feature of CaMKII in plasticity is its ability to switch to a calcium (Ca^{2+}) independent constitutively active state after autophosphorylation at threonine 287 (T287). A second pair of sites, T306 T307 in the calmodulin (CaM) binding region once autophosphorylated, prevent subsequent CaM binding and inactivates the kinase during synaptic plasticity and memory. Recently a synaptic molecule called Ca^{2+} /CaM-dependent serine protein kinase (CASK) has been shown to control both sets of CaMKII autophosphorylation events and hence is well poised to be a key regulator of memory. We show deletion of full length CASK or just its CaMK-like and L27 domains disrupts middle-term memory (MTM) and long-term memory (LTM), with CASK function in the α'/β' subset of mushroom body neurons being required for memory. Likewise directly changing the levels of CaMKII autophosphorylation in these neurons removed MTM and LTM. The requirement of CASK and CaMKII autophosphorylation was not developmental as their manipulation just in the adult α'/β' neurons was sufficient to remove memory. Overexpression of CASK or CaMKII in the α'/β' neurons also occluded MTM and LTM. Overexpression of either *Drosophila* or human CASK in the α'/β' neurons of the CASK mutant completely rescued memory, confirming that CASK signaling in α'/β' neurons is necessary and sufficient for *Drosophila* memory formation and that the neuronal function of CASK is conserved between *Drosophila* and human. At the cellular level CaMKII overexpression in the α'/β' neurons increased activity dependent Ca^{2+} responses while reduction of CaMKII decreased it. Likewise reducing CASK or directly expressing a phosphomimetic CaMKII T287D transgene in the α'/β' similarly decreased Ca^{2+} signaling. Our results are consistent with CASK regulating CaMKII autophosphorylation in a pathway required for memory formation that involves activity dependent changes in Ca^{2+} signaling in the α'/β' neurons.

Keywords: CASK, CaMKII, memory, *Drosophila*, mushroom body, calcium imaging, autophosphorylation, disease model

INTRODUCTION

Changes in neural activity and Ca^{2+} signaling in neural circuits of memory centers encode information during memory formation. One molecule critical for these processes is Ca^{2+} /CaM serine/threonine kinase II (CaMKII) whose activity is acutely sensitive to changes in Ca^{2+} during long-term potentiation (LTP) underlying hippocampal memory formation (Lisman et al., 2002). Further features that endow CaMKII with its central role in memory formation are its abundance in structures known to be required for memory. For instance, CaMKII is the main protein in the hippocampal post-synaptic density (PSD) (Kelly et al., 1984) and is similarly enriched in the mushroom body memory center of *Drosophila* (Takamatsu et al., 2003; Hodge et al., 2006). Finally CaMKII has also been dubbed “the molecular memory switch”; because after it associates with Ca^{2+} /CaM it undergoes a conformational change exposing a T286 on mammalian CaMKII and T287 on *Drosophila* CaMKII that can be autophosphorylated (Figure 1A), resulting in a Ca^{2+} independent constitutively

active kinase (Lisman and Zhabotinsky, 2001). Pharmacological blockade or knockout of CaMKII results in mice with deficits in LTP and memory (Silva et al., 1992a,b). Mice expressing Ca^{2+} dependent CaMKII-T286A have no LTP and memory and those expressing CaMKII-T286D also have abnormal LTP and memory (Mayford et al., 1996; Yasuda and Mayford, 2006). A second pair of autophosphorylation events within the CaM binding domain (TT305/6 equivalent to *Drosophila* TT306/7, Figure 1B) occur when Ca^{2+} /Calmodulin (CaM) dissociates from CaMKII and are inhibitory as autophosphorylation prevents subsequent CaM binding and hence inhibits CaMKII function. Mice with blocked inhibitory sites (CaMKII-TT305/6AA) show enhanced LTP while CaMKII-TT305/6DD expression also disrupts LTP and memory (Elgersma et al., 2002). In *Drosophila*, there is no CaMKII null, which would be expected to be lethal (Park et al., 2002; Mehren and Griffith, 2004), however peptide inhibition of CaMKII led to synaptic defects and memory deficits in the courtship-conditioning assay (Griffith et al., 1993, 1994). Therefore, the



control of CaMKII and its autophosphorylation is critical for synaptic plasticity and memory in *Drosophila* and mammals. But the mechanism of regulation of CaMKII autophosphorylation during memory formation is still unclear.

One molecule that in addition to CaM regulates CaMKII autophosphorylation is CASK (Ca^{2+} /CaM-dependent serine protein kinase, **Figure 1C**), a membrane-associated guanylate kinase (MAGUK) scaffolding protein that contains a CaMK-like and Lin-2/Lin-7 (L27) domain in addition to the canonical PDZ [Post-synaptic density protein (PSD95), *Drosophila* disc large tumor suppressor (Dlg1), and Zonula occludens-1 protein (Zo-1)], SH3 (SRC Homology 3), and GUK (guanylate kinase) domains with the CaMK and GUK domains likely kinase dead in *Drosophila* (Hata et al., 1996; Lu et al., 2003). The CaMK domain of CASK has low levels of Ca^{2+} /CaM independent activity against neurexin that unlike other kinases is magnesium independent (Mukherjee et al., 2008). Again the GUK domain of mammalian CASK encodes a pseudokinase. Two isoforms of CASK are present in flies, a long form, CASK- β and a short isoform, CASK- α (**Figure 1C**). The long form CASK- β contains the additional N-terminal CaMK-like and L27 domains, while the short form CASK- α contains just the canonical PDZ, SH3, and GUK domains which are common to both isoforms, and shows homology to the vertebrate MPP protein (Slawson et al., 2011). CASK- β associates with CaMKII at synapses and in the absence of Ca^{2+} /CaM promotes TT306/7 phosphorylation (**Figure 1A**), inactivating the kinase (Lu et al., 2003). Deletion of CASK in mice results in lethality, preventing their use in modeling CASK function in synaptic plasticity and memory (Atasoy et al., 2007). Flies completely lacking CASK are viable, have decreased levels of synaptic CaMKII-TT306/7 autophosphorylation and display abnormal habituation (Lu et al., 2003). Furthermore, CASK mutants increase T287 autophosphorylation thereby endowing CASK with the ability to regulate the CaMKII switch to Ca^{2+} independence (Hodge et al., 2006). CASK is expressed throughout the fly brain including the mushroom bodies (Martin and Ollo, 1996; Lu et al., 2003). In this study we determine the role of CASK and CaMKII autophosphorylation in memory and measure the accompanying changes in mushroom body Ca^{2+} signaling.

MATERIALS AND METHODS

DROSOPHILA STOCKS

Flies were grown on cornmeal molasses agar medium under standard conditions. *CASK- β null*, *uas-CASK (10,20MI)*, *uas-CaMKII*, *uas-CaMKII-T287D*, *uas-CaMKII-T287A*, *uas-CaMKII-TT306/7AA*, *Df(3R)x307*, and *Df(3R)x313* (Lu et al., 2003; Slawson et al., 2011) were kind gifts from Dr. Leslie Griffith (Brandeis University, US). *uas-CASK-RNAi* flies (stock #104793) were obtained from the Vienna *Drosophila* Stock Center (VDRC). *OK107-Gal4*, *c305a-Gal4*, *MB247-Gal4*, and wildtype flies [*CantonSw-*, (*CSw-*)] were from Dr. Scott Waddell (Oxford University, UK). All CASK and CaMKII mutants, Gal4, and UAS lines were outcrossed with the CSw- line for at least six generations prior to behavioral experiments. *GCaMP3.1* flies were a gift from Dr. Loren Looger (Janelia farm, VA, US). *MB247-Gal4; tubulin-Gal80^{ts}* and *OK107-Gal4; tubulin-Gal80^{ts}* were obtained from Dr. Yi Zhong (Cold Spring Harbor Laboratories, US).

uas-CaMKII-RNAi and *CaMKII-Gal4* flies were obtained from Dr. Sam Kunes (Harvard University, US).

CLONING

Human CASK cDNA isolated from human cerebellum was obtained from imaGenes (IMAGE full-length cDNA clone IRCMp5012G0614D, <http://www.imagenes-bio.de>) in a *pCR4-TOPO* vector. Forward (5'-CACC ATG GCC GAC GACGAC-3') and reverse (5'-CTA ATA GAC CCA GGA GAC AGG-3') primers (0.4 μ L at 0.5 μ M, Invitrogen), *dNTPs* (200 μ M), and CASK cDNA (1 μ l) was added to ddH₂O (13.4 μ l) before addition of High fidelity Phusion DNA polymerase (0.2 μ l, Finnzymes). The following reaction conditions were then used for PCR: 98°C for 30 s, 98°C for 10 s, 61°C for 20 s, 72°C for 60 s (25 cycles), 72°C for 5 min. The resultant PCR product was used for *pENTR*TM directional TOPO[®] cloning (Invitrogen) to create the plasmid *pEntr-CASK*. This was used to transfect α -Select Gold *E. Coli* (Bioline). This plasmid was then sequenced (Geneservice, London, <http://www.geneservice.co.uk>) and used in the Gateway LR cloning reaction (Invitrogen) with a *pTW* plasmid. The plasmid *pTW-CASK* was used for germline transformation (Bestgene, US) by microinjection into *Drosophila* embryos.

BEHAVIOR EXPERIMENTS

Behavior experiments were carried out at 25°C, 70% relative humidity and under dim red light. For *Gal80^{ts}* (TARGET) experiments the flies were grown at 18°C that allowed *Gal80^{ts}* inhibition of *Gal4*. Adult flies were collected everyday in the evening and maintained for another three days at 30°C. These flies were trained and tested at 30°C that relieved *Gal80^{ts}* inhibition allowing the expression of transgenes (McGuire et al., 2003). To measure learning (2 min memory) a mixed population of about one hundred 2–3 days (4 days for TARGET experiments) day old flies received one cycle of training during which they were exposed sequentially to one odor [conditioned stimulus, CS+; 3-octanol (1:100) or 4-methyl-cyclohexanol, 1:67] paired with electric shock (60V DC) (unconditioned stimulus, US) and then to a second odor (CS-odor) without electric shock. The flies were then allowed to choose between the two odors for 120 s in the T-maze (Tully and Quinn, 1985). To measure middle-term memory (MTM) flies were given one cycle of training and then stored in food containing vials for 3 h before they were tested as in learning experiments. A performance index (PI) was calculated as the number of flies avoiding the CS+ minus number of flies avoiding the CS-, divided by the total number of flies that participated in the test. A score of 1.0 would be equivalent to 100% learning, where all the flies avoided the CS+. In contrast a 50:50 distribution would give a PI of zero (no learning). For long-term memory a custom built maze was used which allowed simultaneous training of several batches of flies. The flies were administered five cycles of training either with an inter-cycle interval of 15 min (spaced) or without any inter-cycle interval (massed). They were then kept at 18°C until tested. Prior to testing, the flies were moved to 25°C and allowed to acclimatize for at least 1 h. For long-term memory (LTM), memory was assessed 24 h after training. All statistical analysis for behavioral data was performed and plotted with Graphpad Prism (Graphpad software, Inc) software.

CALCIUM IMAGING

Ca²⁺ imaging on dissected adult brains was performed as described previously (Ruta et al., 2010; Tessier and Broadie, 2011). Briefly, the fly brains were dissected in HL3.1, tethered to the bottom of a petri dish containing 5 ml of HL3.1. Images were collected using an Axio Examiner Z1 microscope (Zeiss) using a 10 \times water immersion objective and Axiovision software. The brains were stimulated by gently adding 500 μ l of 65 mM KCl in HL3.1 to the dish while the images were captured at 340 msec/frame.

IMAGE ANALYSIS

Image analysis was performed using the single channel ratio analysis of the physiology module of AxioVs40 V 4.8.0.0 (Zeiss). Regions of interest were selected by drawing around the mushroom bodies α'/β' neurons and the fluorescence values were obtained. An initial reference fluorescence (F_0) value of was calculated by averaging the fluorescence of first ten frames. Percent change in fluorescence, $\% \Delta F/F$, was calculated for each time point, which is given by $[(F-F_0)/F_0] \times 100$, where F is fluorescence at a given time. A ratio table was generated and the values were plotted as a function of number of time.

WESTERN BLOTTING AND RNAi VALIDATION

Extracts were prepared by freezing ten fly heads from either wildtype or *Elav-Gal4 > uas-RNAi* in liquid nitrogen followed by homogenization in 50 μ l of lysis buffer (50 mM Tris, pH 7.4, 150 mM NaCl, 1% Triton-x-100, 5 mM EDTA, 0.1% SDS, 1 mM Na₂VO₃, and complete mini protease inhibitor (Amersham Biosciences). The homogenate was incubated on ice for 10 min and then centrifuged at 14000 rpm. Supernatant was collected and mixed with 50 μ l sample buffer. 15 μ l of this sample were loaded per well. Following transfer to a nitrocellulose membrane, the membrane was probed with rabbit anti-CASK 1:800 antibodies. Bands were visualized using horseradish peroxidase-conjugated secondary antibodies (Amersham Biosciences) and enhanced chemiluminescence reagents (ECL, Amersham Biosciences). In order to validate the *RNAi* constructs, the CASK sequence from VDRC and the CaMKII sequence (Ashraf et al., 2006) were used in BLAST searches of the NCBI database and only the appropriate gene of interest came up as a significant hit suggesting no off-targets.

IMMUNOHISTOCHEMISTRY

Immunohistochemistry was performed essentially according to previously published protocols (Hodge et al., 2006). Briefly, the fly adult brains were dissected for 4–8 days old flies in HL3.1. The isolated brains were then fixed in 4% paraformaldehyde for 1 h followed by two washes with HL3.1-Tx (HL3.1 containing 0.1% Triton-X-100) for a total of 1 h. The brains were then blocked for 1 h with 0.1% bovine serum albumen (BSA) and 0.1% normal goat serum (NGS) in HL3-Tx. Brains were incubated overnight at 4°C with 1:40 dilution of a rabbit anti-CASK antibody (Lu et al., 2003) or 1:100 mouse anti-CaMKII (Takamatsu et al., 2003). Following an overnight washing HL3.1-Tx at 4°C the brains were then incubated with 1:400 anti-rabbit Alexa-648 or with 1:400 goat anti-mouse Alexa-488 (Invitrogen) secondary antibodies overnight. Following an overnight HL3.1-Tx wash the

brains were mounted in Vectashield (Vector laboratories) and stored at 4°C in the dark until they were imaged using a Leica TCS SP5 confocal microscope (Wolfson Bioimaging facility, University of Bristol). The images were then examined using Velocity imaging software (PerkinElmer) and projections were generated using the image processing software ImageJ (NIH).

SENSORIMOTOR CONTROLS

The odor acuity and shock reactivity were determined for all genotypes used in this study, as described previously (Tully et al., 1994). Briefly, for odor acuity ~80–100 flies were introduced into the T-maze. After 90 s the flies were taken to the choice point where they were allowed 2 min to make a choice between pure odors and air. The flies were then collected and counted. The percent avoidance was calculated by dividing the flies that chose odor by the total number of flies that participated in the test. For shock reactivity, flies were introduced into the shock chamber. After 90 s of rest they were given a 60 V DC electric shock from which time they were allowed to escape to a similar tube without electric shock on the other side. They were given 2 min to make a choice and then collected and counted. The percent shock avoidance was calculated by dividing the number of flies that avoided the shock by escaping the shock tube by the total number of flies in the experiment. The flies that remained in the central chamber were considered to have escaped the electric shock.

RESULTS

CASK- β ISOFORM CONTAINING CaMKII-LIKE AND L27 DOMAINS IS REQUIRED FOR MIDDLE-TERM MEMORY

In order to see if CASK plays a role in learning and memory flies were tested using the olfactory aversive conditioning assay (Tully and Quinn, 1985). We used deficiency lines: *Df(3R)x307* and *Df(3R)x313* which contain large chromosomal deficiencies both lacking CASK [called *camguk* (*cmg*) or *kaki*]. A cross between the two lines generates transheterozygote flies with only a short fragment of chromosome deleted that includes the whole of the CASK locus, therefore null for both CASK- α and CASK- β (Martin and O'Lo, 1996). All CASK mutant genotypes learned to avoid the shock-paired odor similar to controls when tested 2 min after training (Figure 2A). In order to investigate the role of the different CASK isoforms in learning (2 min memory) we used a mutant (CASK- β null) that completely removes the long isoform of CASK (CASK- β) but leaves the short (CASK- α) isoform intact (Figure 1C; Slawson et al., 2011). These mutant flies also did not show any defects in 2 min memory tested after 2 min of administering one training cycle (Figure 2B).

The majority of the CASK and CaMKII mutant genotypes tested showed normal shock reactivity and olfactory acuity demonstrating that any performance deficit was due to a defect in signal processing required for memory as opposed to a peripheral defect preventing the fly from being able to perform the behavioral task (Table 1). CASK- β null, *Df(3)x313/Df(3)x307* and the CASK heterozygous control deficiency line *Df(3)x313/+* reacted abnormally to electric shock. However, all the CASK and CaMKII mutant genotypes showed normal learning confirming that these flies are healthy and have mushroom bodies that are capable of detecting odor, respond to shock normally and able to support




FIGURE 2 | CASK is not required for learning. (A) Learning or initial (2 min) short-term memory (STM) was measured immediately after administering one cycle of shock-odor training. Flies lacking all forms of (Continued)

FIGURE 2 | Continued

CASK [*Df(3)x307/Df(3)x313*] learned equally well to heterozygote negative controls [*Df(3)x307/+* or *Df(3)x313/+*]. Data were analyzed using One-Way ANOVA followed by a Tukey's *post-hoc* test. In all figures the numbers denote n (typically ~100 flies used for each n), n.s. is not significant ($p > 0.05$). The brackets below the significance label denote the genotypes being compared. **(B)** Flies lacking CASK- β (CASK- β null), learned equally well to avoid the shock-paired odor as wildtype negative control. **(C)** Compared to wildtype, pan-neuronal (*elav-Gal4*) expression of *uas-CASK-RNAi* leads to reduced CASK- β immunofluorescence in whole mount adult brains. **(D)** Western blot of cell lysates from wildtype or *elav-Gal4, uas-CASK-RNAi* heads showed a similar reduction in CASK- β . Quantification of the intensity of the CASK- β band showed a ~50% reduction in CASK- β expression in *elav-Gal4, uas-CASK-RNAi*. **(E)** Flies with targeted reduction of CASK throughout their mushroom body (*OK107-Gal4 > uas-CASK-RNAi*) or just in the adult mushroom body (*OK107-Gal4; Gal80^{ts} > uas-CASK-RNAi*) displayed learning comparable to heterozygous wildtype negative controls. **(F)** Flies with targeted reduction of CASK in the mushroom body α'/β' neurons (*c305-Gal4 > uas-CASK-RNAi*) or just in the adult mushroom body α'/β' neurons (*c305a-Gal4; Gal80^{ts} > uas-CASK-RNAi*) displayed learning comparable to heterozygous wildtype negative controls. **(G)** Flies with targeted reduction of CASK in the mushroom body α/β and γ neurons (*MB247-Gal4 > uas-CASK-RNAi*) or just in the adult mushroom body α/β and γ neurons (*MB247-Gal4; Gal80^{ts} > uas-CASK-RNAi*) displayed learning comparable to heterozygous wildtype negative controls. **(H)** Flies with targeted reduction of CASK in CaMKII neurons (*CaMKII-Gal4 > uas-CASK-RNAi*) displayed learning comparable to heterozygous wildtype negative controls. All data were analyzed using One-Way ANOVA followed by a Tukey's *post-hoc* test.

initial learning, therefore the data shown in **Figures 2A,B** are negative controls for this issue. In addition none of the flies displayed any obvious developmental defect and neither displayed a wing phenotype or sluggishness (Park et al., 2002). This was reflected in the fact they were wildtype for peripheral controls (**Table 1**) and learning (**Figures 2, 4**), so therefore were able to choose to move away from the shock-paired odor in the T-maze the same as wildtype flies (**Figures 2, 4**).

Finally, we decided to investigate the effect of mushroom body specific reduction of CASK on learning. *Drosophila* mushroom bodies consist of three different classes of intrinsic neurons (α/β , α'/β' , and γ) that extend their axons into the five lobes of neuropil (Davis, 2011). We used a *CASK-RNAi* line which reduces the expression of CASK by ~50% (**Figures 2C,D**) to test if reduction of CASK in the mushroom body has an effect on learning in flies. Expression of *CASK-RNAi* transgene in either all mushroom body neurons [*OK107-Gal4* (Connolly et al., 1996)], mushroom body α'/β' neurons [*c305a-Gal4* (Krashes et al., 2007)], mushroom body α/β and γ neurons [*MB247-Gal4* (Zars et al., 2000)], or using a *CaMKII-Gal4* [that expresses in the mushroom body α/β , α'/β' , and dorsal anterior lateral (DAL) neurons, (Chen et al., 2012)] drivers did not lead to a significant decrease in 2 min memory (**Figures 2E–H**).

We then tested flies 3 h after one cycle of training (**Figure 3A**), CASK- β null reduced MTM to a similar extent as *Df(3)x307/Df(3)x313*. This showed that deletion of CASK- β alone was sufficient to cause the MTM defect, indicating an important role for the CaMK-like and L27 domains of CASK in MTM. Flies with CASK knockdown in either all mushroom body neurons (**Figure 3B**) or just α'/β' neurons (**Figure 3C**) similarly showed a drastic reduction in MTM, while restricting

Table 1 | The sensorimotor controls for CASK and CaMKII transgenic flies.

	Odor avoidance		Percent shock avoidance
	MCH	OCT	
	Mean \pm SEM	Mean \pm SEM	Mean \pm SEM
WT Control	0.77 \pm 0.06	0.69 \pm 0.03	62.9 \pm 3.8
MB247/+	0.6 \pm 0.08	0.59 \pm 0.03	62.7 \pm 3.5
c305a/+	0.59 \pm 0.03	0.54 \pm 0.04	93.9 \pm 2.1
OK107/+	0.65 \pm 0.05	0.58 \pm 0.07	87 \pm 2
CASK-RNAi/+	0.59 \pm 0.07	0.68 \pm 0.1	85.7 \pm 1.7
MB247 > CASK-RNAi	0.58 \pm 0.05	0.84 \pm 0.07	73.4 \pm 3.7
c305a > CASK-RNAi	0.61 \pm 0.03	0.83 \pm 0.11	79.4 \pm 3.2
OK107 > CASK-RNAi	0.63 \pm 0.06	0.87 \pm 0.04	79 \pm 2.4
MB247;G80 > CASK-RNAi	0.64 \pm 0.04	0.61 \pm 0.05	75.7 \pm 4.1
OK107;G80 > CASK-RNAi	0.71 \pm 0.02	0.65 \pm 0.05	78.5 \pm 3.5
Df(3)x307/+	0.65 \pm 0.03	0.53 \pm 0.1	72 \pm 5.6
Df(3)x313/+	0.6 \pm 0.05	0.52 \pm 0.07	41.5 \pm 1.4*
Df(3)x307/Df(3)x313	0.63 \pm 0.08	0.72 \pm 0.05	43.5 \pm 8.9*
T287D/+	0.61 \pm 0.07	0.61 \pm 0.05	70 \pm 6.1
MB247 > T287D	0.59 \pm 0.05	0.64 \pm 0.16	64.9 \pm 5.4
c305a > T287D	0.65 \pm 0.13	0.66 \pm 0.04	61.5 \pm 11.7
OK107 > T287D	0.87 \pm 0.02	0.73 \pm 0.03	69 \pm 2.6
MB247;G80 > T287D	0.7 \pm 0.03	0.61 \pm 0.11	75.5 \pm 7.6
OK107;G80 > T287D	0.7 \pm 0.07	0.68 \pm 0.08	81 \pm 3.1
CaMKII-Gal4 > T287D	0.48 \pm 0.04	0.51 \pm 0.12	93 \pm 3.5
CASK- β null	0.61 \pm 0.07	0.53 \pm 0.08	35.9 \pm 3.6*
CASK;CASK- β null	0.53 \pm 0.05	0.52 \pm 0.09	37 \pm 6.9*
c305a;CASK- β null	0.49 \pm 0.09	0.49 \pm 0.04	72.5 \pm 1.7
c305a > CASK;CASK- β null	0.63 \pm 0.06	0.65 \pm 0.07	60.3 \pm 4.4
CASK/+	0.64 \pm 0.1	0.56 \pm 0.05	73.1 \pm 3.6
MB247 > CASK	0.57 \pm 0.06	0.64 \pm 0.12	93.2 \pm 2.3
c305a > CASK	0.86 \pm 0.03	0.59 \pm 0.05	71.9 \pm 3.2
OK107 > CASK	0.86 \pm 0.03	0.59 \pm 0.05	82.2 \pm 2.2
T306A T307A/+	0.56 \pm 0.03	0.55 \pm 0.1	92.5 \pm 1
MB247 > T306A T307A	0.55 \pm 0.07	0.64 \pm 0.05	89.8 \pm 3.7
c305a > T306A T307A	0.56 \pm 0.06	0.6 \pm 0.02	91.8 \pm 1.7
OK107 > T306A T307A	0.62 \pm 0.03	0.57 \pm 0.03	88.8 \pm 1.4
CaMKII/+	0.64 \pm 0.05	0.6 \pm 0.11	79.6 \pm 2.4
MB247 > CaMKII	0.58 \pm 0.02	0.52 \pm 0.08	87 \pm 3.8
c305a > CaMKII	0.67 \pm 0.14	0.52 \pm 0.04	82.7 \pm 2.8
OK107 > CaMKII	0.74 \pm 0.08	0.5 \pm 0.03	92.9 \pm 1.2
CaMKII-Gal4 > CaMKII	0.48 \pm 0.03	0.51 \pm 0.11	93 \pm 3.5
T287A/+	0.43 \pm 0.04	0.61 \pm 0.08	85.5 \pm 2.2
MB247 > T287A	0.59 \pm 0.14	0.41 \pm 0.08	72.5 \pm 7.5
c305a > T287A	0.51 \pm 0.12	0.53 \pm 0.07	66.8 \pm 1.8
OK107 > T287A	0.59 \pm 0.14	0.6 \pm 0.05	69.5 \pm 7.6
CaMKII-RNAi/+	0.62 \pm 0.12	0.6 \pm 0.06	90.5 \pm 0.5
MB247 > CaMKII-RNAi	0.57 \pm 0.08	0.55 \pm 0.12	95.8 \pm 2.4
c305a > CaMKII-RNAi	0.62 \pm 0.07	0.55 \pm 0.12	92.3 \pm 0.3
OK107 > CaMKII-RNAi	0.63 \pm 0.12	0.54 \pm 0.12	89 \pm 4

CASK and CaMKII transgenic flies showed normal odor acuity for 3-Octanol (OCT) and methyl-cyclohexanol (MCH). However, some of the CASK genotypes have reduced shock reactivity compared to wildtype negative control (* $p < 0.05$).




FIGURE 3 | Continued

FIGURE 3 | CASK functions in the mushroom body α'/β' neurons during middle-term memory formation. (A) MTM measured 3 h post-training was completely removed in *CASK- β* null flies. Similarly transheterozygous [*Df(3)X313/Df(3)X307*] flies that lack both α -*CASK* and β -*CASK* have a similar reduction in MTM compared to wildtype or heterozygote negative controls [*Df(3)X313/+* or *Df(3)X307/+*]. (B) Flies with *uas-CASK-RNAi* expressed throughout their mushroom body using *OK107-Gal4* show a reduction in MTM compared to heterozygous wildtype negative controls. Adult specific reduction in mushroom body *CASK* using *OK107-Gal4; Gal80^{TS}* was sufficient to reduce MTM. (C) Flies expressing *uas-CASK-RNAi* in their α'/β' mushroom body neurons (*c305a-Gal4*) show a reduction in MTM. Reduction of *CASK* just in the adult α'/β' neurons using *Gal4-c305a; Gal80^{TS}* was sufficient to cause the reduction in MTM. (D) Reduction of *CASK* in α/β and γ neurons using *MB247-Gal4* did not affect MTM. Adult specific reduction of *CASK* in α/β and γ neurons using *MB247-Gal4; Gal80^{TS}* also did not affect MTM. (E) *CASK-RNAi* expression using *CaMKII-Gal4* also lead to a MTM defect compared to wildtype. (F) Flies that contained *Gal80^{TS}* in combination with either *Gal4-OK107* (G)

c305a-Gal4 (H) *MB247-Gal4* were reared and tested at 18°C, a temperature that prevented the expression of the *CASK-RNAi* (hence these are negative control experiments) displayed MTM similar to heterozygous wildtype controls. (I) MTM was completely removed in flies overexpressing full-length *CASK* throughout their mushroom body. (J) Overexpression of *CASK* in α'/β' neurons was sufficient to cause the decrease in MTM. (K) α/β and γ neuron overexpression of *CASK* did not affect MTM. (L) Expression of *CASK* in mushroom body α'/β' neurons in a *CASK- β* null background (*c305a-Gal4, uas-CASK; CASK- β* null) rescued the reduction in MTM seen in the *CASK- β* null mutants [*uas-CASK; CASK- β* null and *c305a-Gal4; CASK- β* null (the positive controls) compared to wildtype (the negative control)] to the same level as wildtype. (M) Overexpression of human *CASK* in the mushroom bodies α'/β' neurons in a fly otherwise completely lacking *CASK- β* rescued the reduction in MTM seen in the *CASK- β* null mutants (*uas-CASK; CASK- β* null and *Gal4-c305a; CASK- β* null compared to wildtype) to the same level as wildtype. All data were analyzed using One-Way ANOVA followed by a Tukey's *post-hoc* test. n.s. is not significant, ($p > 0.05$), * $p < 0.05$, ** $p < 0.01$, and *** $p < 0.001$.

expression to the remaining α/β and γ neurons had no effect (Figure 3D). This suggests *CASK* specifically controls memory formation via the α'/β' neurons. In order to distinguish the role of *CASK* in mushroom body development as opposed to an acute physiological role in signaling underlying memory we restricted the reduction of *CASK* to just the adult mushroom body using the TARGET system (McGuire et al., 2003). Reduction of *CASK* specifically in the adult mushroom body was sufficient to cause the reduction in MTM showing that the effects are post-developmental (Figure 3B). Again we confirmed that this deficit in MTM resulted from a function of *CASK* in the adult α'/β' neurons (Figure 3C) as opposed to the adult α/β and γ neurons (Figure 3D). The negative control flies reared and tested at 18°C, conditions where there was no transgene expression (Shuai et al., 2010) showed normal MTM (Figures 3F–H).

As previous work has showed that *CASK* influences plasticity and behavior via regulation of *CaMKII* autophosphorylation (Lu et al., 2003; Hodge et al., 2006) we used the *CaMKII*-specific promoter that appears to express in the mushroom body α/β , α'/β' , and DAL neurons and has been used to follow the changes in *CaMKII* transcription occurring during LTM (Chen et al., 2012). Knockdown of *CASK* in these *CaMKII* neurons was sufficient to completely remove MTM (Figure 3E). We believe it is the α'/β' neurons of the *CaMKII-Gal4* expression pattern that are most critical for mediating *CASK* and *CaMKII* effects on memory, as *CASK* and *CaMKII* memory phenotypes map to α'/β' (*c305a-Gal4*) neurons with α/β (*MB247-Gal4*) neurons having little effect and the DAL neurons thought to only affect certain aspects of LTM (Chen et al., 2012). The data suggests that *CASK* is needed in a subset of neurons that express *CaMKII* in order to get memory formation. In order to determine if increased levels of *CASK* also disrupted MTM we expressed *uas-CASK*, the cDNA corresponding to the long isoform called *CASK- β* (Figure 1C; Lu et al., 2003; Hodge et al., 2006; Slawson et al., 2011) throughout the mushroom body. This resulted in a dramatic reduction in MTM (Figure 3I), which again could be localized to the α'/β' neurons (Figure 3J) as opposed to the α/β and γ neurons where *CASK* overexpression had no effect (Figure 3K). Since the effects of *CASK* knockdown were also localized to the mushroom body α'/β' neurons, we tested whether expressing the

Drosophila CASK transgene in these neurons in a *CASK- β* null fly, would return their memory to normal (Figure 3L). Compared to *CASK- β* null mutant flies with *c305a-Gal4* alone or *uas-CASK* alone, mushroom body α'/β' expression of *CASK* in the *CASK- β* null background fully rescued the MTM defect to a level indistinguishable from wildtype, confirming that *CASK* signaling in mushroom body α'/β' is necessary and sufficient for *Drosophila* MTM formation.

HUMAN CASK OVEREXPRESSION IN MUSHROOM BODIES α'/β' NEURONS IS SUFFICIENT TO RESTORE THE MEMORY OF CASK NULL FLIES TO WILDTYPE

As human *CASK* and *CaMKII* display a high degree amino acid residue identity to *Drosophila CASK* (74% identical) and *CaMKII* (79% identical), it is likely that they might function in a similar way in both organisms (Cho et al., 1991; Hsueh, 2006). In order to test this hypothesis we overexpressed human *CASK* in mushroom body α'/β' neurons of flies that otherwise express no *CASK- β* . Whereas *CASK- β* null flies almost completely lack MTM, overexpression of human *CASK* just in mushroom body α'/β' neurons was sufficient to return memory to levels indistinguishable to wildtype (Figure 3M). This indicates that *Drosophila* and human *CASK* show conserved neuronal function in memory formation.

LEVELS OF CaMKII AUTOPHOSPHORYLATION REGULATE MIDDLE-TERM MEMORY FORMATION

In order to see if the *CaMKII* levels and autophosphorylation are important for aversive olfactory learning and memory we expressed a range of *CaMKII* transgenes in the mushroom body. These included a transgene that overexpresses *CaMKII* (Koh et al., 1999), a *CaMKII-hairpin* that allows targeted reduction of *CaMKII* (Ashraf et al., 2006; Akalal et al., 2010; Chen et al., 2012), a Ca^{2+} -independent constitutively active *CaMKII-T287D*, a Ca^{2+} dependent *CaMKII-T287A* (Park et al., 2002), and *CaMKII-TT306/7AA* containing phospho-blocking mutations of its inhibitory phosphorylation sites (Lu et al., 2003). We found that mushroom body expression of these transgenes did not affect learning with the avoidance of the shock-paired odor being similar between mutant and wildtype flies (Figures 4A–D) similar to what we found for all *CASK* genotypes. In order to




FIGURE 4 | Continued

FIGURE 4 | CaMKII autophosphorylation in the mushroom body is not required for initial memory formation. Initial memory or learning was measured immediately (2 min) after one cycle training. Flies expressing *CaMKII* transgenes either throughout the mushroom body (A), in the α/β' neurons (B), the α/β , and γ neurons (C) or CaMKII neurons (D) all learned similar to heterozygous wildtype negative control. Data were analyzed using One-Way ANOVA followed by a Tukey's *post-hoc* test.

see if the level of CaMKII in the mushroom body is important for MTM, we expressed *CaMKII-hairpin* in different parts of the mushroom body, however, none had a significant reduction in MTM compared to the heterozygote wildtype negative control (Figures 5A–C). However, when *CaMKII* is overexpressed throughout the mushroom body, there was a significant reduction in MTM compared to heterozygote wildtype negative control, an effect that localized to the mushroom body α'/β' neurons (Figure 6A).

In order to determine the contribution of the “molecular memory switch” (Figure 1A) to aversive olfactory memory we expressed either the Ca^{2+} -independent constitutively active form of *CaMKII-T287D* or Ca^{2+} dependent *CaMKII-T287A* (Park et al., 2002) in the mushroom body. Expression of *CaMKII-T287D* or *-T287A* either throughout the mushroom body (Figure 6B) or just in α'/β' neurons caused a dramatic reduction in MTM (Figure 6C), with expression in the remaining α/β and γ neurons having no effect (Figure 6D), suggesting that the state of T287 autophosphorylation in α'/β' is particularly important for memory formation. Restricted expression of *CaMKII-T287D* and *-T287A* transgenes to the adult mushroom body (Figure 6E) or just the adult α'/β' (Figure 6F) but not the adult α/β and γ neurons (Figure 6G) was sufficient to cause the reduction in MTM. Negative control flies reared and tested at 18°C, displayed wildtype MTM (Figures 6H–J). In order to see if CaMKII inhibitory autophosphorylation is also important for memory formation (Figure 1B), we overexpressed a transgene with these phosphorylation sites (T306A T307A) blocked (Lu et al., 2003). *CaMKII-T306A T307A* overexpression throughout the mushroom body (Figure 6K) or just the α'/β' neurons (Figure 6L) dramatically reduced MTM, while α/β and γ neuron expression had little effect (Figure 6M).

CASK AND CaMKII FUNCTIONALLY INTERACT TO REGULATE MIDDLE-TERM MEMORY FORMATION

The main effect of CASK is to increase inhibitory phosphorylation of T306 T307 on endogenous CaMKII resulting in a decrease in endogenous kinase activity (Figure 1A; Lu et al., 2003; Hodge et al., 2006) and we show that mushroom body overexpression of CASK removes MTM. Conversely flies overexpressing the *uas-CaMKII-T306A T307A* transgene would have an opposing effect with inhibitory phosphorylation being blocked resulting in increased transgenic kinase activity (Figure 1B), again with mushroom body overexpression of *CaMKII-T306A T307A* removing MTM. As expression of the two transgenes are predicted to have opposite effects on CaMKII activity we decided to co-express CASK and *CaMKII-T306A T307A* to see if they counteract each other's effect and return memory to normal. Indeed flies expressing both transgenes in their mushroom body showed




FIGURE 5 | Continued

FIGURE 5 | Levels of CaMKII in the mushroom body are not important for middle-term memory formation. MTM was equal in flies with reduced *CaMKII* either, throughout the mushroom body (A) the α'/β' neurons (B) or the α/β and γ neurons (C) compared to heterozygous wildtype negative controls. Data were analyzed using One-Way ANOVA followed by a Tukey's *post-hoc* test.

complete rescue of their memory deficit, confirming that CASK regulates CaMKII autophosphorylation during memory formation (Figure 6K). Expression of *CaMKII-T306A T307A* in just the α'/β' neurons was not sufficient to rescue the CASK overexpression memory defect (Figure 6L). Expression of any combination of the transgenes in the α/β and γ neurons had no effect on MTM (Figure 6M). This data suggests that as for CASK, changes in CaMKII autophosphorylation are required throughout the adult mushroom body during memory formation and that the effect of CASK on MTM formation is through CASK's regulation of CaMKII autophosphorylation.

CASK AND CaMKII ARE REQUIRED FOR LONG-TERM MEMORY FORMATION

In order to determine the role of CASK and CaMKII autophosphorylation in LTM, flies were subjected to five cycles of spaced training which is known to produce a form of consolidated memory that is protein synthesis and cyclic-AMP response element binding protein (CREB) dependent (Tully et al., 1994). CASK- β null flies were not able to form LTM (Figure 7A). Similarly mushroom body CASK knockdown or overexpression of *CaMKII-T287D, T287A, or TT306/7AA* throughout the mushroom body (Figure 7A), just the α'/β' (Figure 7B), but not the α/β and γ (Figure 7C) neurons reduced LTM compared to control. Previous studies have reported that *CaMKII* knockdown in α/β and γ mushroom body neurons or DAL neurons reduced LTM (Ashraf et al., 2006; Akalal et al., 2010; Chen et al., 2012). We therefore performed experiments using this *CaMKII-hairpin-RNAi* transgene but for the first time with a full complement of mushroom body neuron specific drivers (Figure 7D). Flies with a reduction of *CaMKII* in any of these sets of mushroom body neurons showed deficits in LTM indicating mushroom body CaMKII levels are crucial for normal LTM formation. This also demonstrates that the effect of changing the level of CaMKII as opposed to changing levels of autophosphorylated CaMKII can be qualitatively different in the α/β and γ neurons. Consistent with our MTM data we observed a similar reduction in LTM in flies overexpressing CASK or *CaMKII* throughout the mushroom body (Figure 7E), just in the α'/β' neurons (Figure 7F), but not in the α/β and γ neurons (Figure 7G). Again this data is consistent with CASK function and CaMKII autophosphorylation in the α'/β' neurons being critical for LTM memory formation.

CASK AND CaMKII LEVELS AND REDUCTION OF CaMKII AUTOPHOSPHORYLATION ARE REQUIRED FOR ANAESTHESIA RESISTANT MEMORY FORMATION

A second form of memory is generated by five cycles of training without rest intervals (massed training). This form of memory consists of anesthesia resistant memory (ARM) and is independent of CREB transcription (Tully et al., 1994). CASK- β nulls




FIGURE 6 | Continued

FIGURE 6 | CaMKII autophosphorylation in the mushroom body α'/β' neurons are critical for middle-term memory. (A) Overexpression of *CaMKII* throughout the mushroom body or just in the α'/β' neurons significantly decreased MTM, while expression in the α/β and γ neurons had little effect. (B) Overexpression of constitutively active *CaMKII-T287D* or Ca^{2+} dependent *CaMKII-T287A* throughout the mushroom body significantly decreased MTM. (C) Overexpression of *CaMKII-T287D* or *CaMKII-T287A* just in the α'/β' neurons significantly decreased MTM, (D) while expression in the α/β and γ neurons had little effect. (E) Adult specific mushroom body expression of *CaMKII-T287D* or *-T287A* with *OK107-Gal4*; *Gal80^{ts}* lead to a reduction in MTM compared to heterozygous wildtype negative controls. (F) Adult specific α'/β' neuron expression of *CaMKII-T287D* or *-T287A* with *c305a-Gal4*; *Gal80^{ts}* was sufficient to cause the reduction in MTM. (G) Expression of *CaMKII-T287D* or *-T287A* in the remaining adult α/β and γ neurons with *MB247-Gal4*; *Gal80^{ts}* did not affect MTM. Performance of flies that

contained *Gal80^{ts}* in combination with *OK107-Gal4* (H) *c305a-Gal4* (I) and *MB247-Gal4* (J) and that were reared and tested at 18°C; a temperature that prevented the expression of the *CaMKII* transgenes and hence is a negative controls had normal MTM. (K) Expression of *uas-CaMKII-T306A T307A* or *CASK* alone throughout the mushroom body reduced MTM (e.g., the positive controls) compared to heterozygous wildtype negative controls. Expression of *CaMKII-TT306/7AA* and *CASK* throughout the mushroom body (*OK107-Gal4*) rescued the MTM deficit seen with expression of either transgene alone to wildtype. (L) Expression of *uas-CaMKII-T306A T307A* in the α'/β' neurons was sufficient to reduce MTM compared to controls. However, simultaneous expression of *CASK* and *CaMKII-TT306/7AA* using *c305a-Gal4* was not sufficient to rescue this defect. (M) Expression of *uas-CaMKII-T306A T307A* in the α/β and γ neurons did not affect MTM. All data were analyzed using One-Way ANOVA followed by a Tukey's *post-hoc* test. n.s. is not significant, ($p > 0.05$), * $p < 0.05$, ** $p < 0.01$, and *** $p < 0.001$.

were not able to form ARM (Figure 8A), while mushroom body *CASK* knockdown or overexpression of *CaMKII-T287D*, *T287A*, or *TT306/7AA* throughout the mushroom body (Figure 8A), just the α'/β' neurons (Figure 8B) but not the α/β and γ neurons (Figure 8C) neurons removed ARM. Our results are consistent with *CASK* function and *CaMKII* autophosphorylation in the α'/β' neurons being critical for ARM formation.

CASK AND CaMKII REGULATE MUSHROOM BODY NEURAL ACTIVITY

Dynamic changes in neural activity and Ca^{2+} signaling in memory centers such as the mushroom body and hippocampus underlie memory formation (Lisman et al., 2002; Davis, 2011). Since *CASK* regulates MTM formation in adult mushroom body α'/β' neurons (as labeled by *c305a-Gal4*), we set out to determine the physiological basis of this defect by measuring dynamic changes in Ca^{2+} signaling as reported by changes in fluorescence of the genetically encoded Ca^{2+} reporter, *GCaMP3.1* in the relevant memory circuit (Tian et al., 2009). We imaged mushroom body Ca^{2+} induced fluorescence in response to acute application of high $[\text{K}^+]$ depolarizing solution that resulted in a robust increase in mushroom body intracellular Ca^{2+} levels (Figures 9A,B) and might reflect a proxy (although somewhat artificial) of the increase in synaptic activity occurring in α'/β' neurons during memory formation in the behavioral experiments. *CASK- β* null or *CASK* knockdown in the α'/β' neurons decreased maximum fluorescence (Figures 9B,C) indicating a disruption of neuronal signaling in the specific mushroom body neurons that cause the memory defect, consistent with this physiological change mediating the fly's inability to remember shown in Figures 3, 7, 8. *CaMKII* and *CaMKII-TT306/7AA* overexpression caused an increase in peak neural activity while reduced *CaMKII* caused a reduction in neural activity (Figures 9B,C), these bi-directional changes in neural activity provide an explanation for the disruption of memory seen with *CaMKII* misexpression in α'/β' neurons (Figures 6–8). In addition overexpression of *CaMKII-T287D* also reduced the peak Ca^{2+} response in a similar manner to reduced *CASK*, consistent with reductions in *CASK* increasing levels of *CaMKII* autophosphorylated at T287 (Figure 1B; Hodge et al., 2006) and suggesting a physiological mechanism for memory deficit resulting from α'/β' expression of *CaMKII-T287D* (Figures 6–8).

In order for relative changes in Ca^{2+} levels to encode information it would be expected that the baseline levels of Ca^{2+} would also be tightly regulated. Therefore, to see if plasticity molecules such as *CaMKII* and *CASK* are involved in setting basal Ca^{2+} , *GCaMP3* signals in α'/β' neurons were measured under baseline conditions. Compared to wildtype (Figure 9D) overexpression of *CaMKII*, *CaMKII-T287D*, or *CaMKII-TT306/7AA* increased basal Ca^{2+} levels. Reduced *CaMKII* or *CASK* caused a decrease in basal Ca^{2+} levels α'/β' neuron, while *CASK* overexpression also lowered baseline Ca^{2+} levels (Figure 9D), the later explaining the effect of α'/β' overexpression on *CASK* on memory (Figures 3, 6, 7). Since overexpression of *CaMKII-T287D* already drives neurons into a very high Ca^{2+} state under basal conditions (Figure 9D), stimulation of the neurons may not be able to increase Ca^{2+} concentrations any further, reducing the change in Ca^{2+} concentration measured for peak response (Figure 9C). These results suggest *CASK*, *CaMKII* levels and autophosphorylation regulate basal and activity-dependent changes in Ca^{2+} signaling in the mushroom body α'/β' neurons, revealing the likely neurophysiological basis for the disruption in memory found in these animals.

DISCUSSION

CASK REGULATES CaMKII AUTOPHOSPHORYLATION IN MUSHROOM BODY α'/β' NEURONS DURING MIDDLE-TERM MEMORY FORMATION

We found that *CASK- β* mutant flies that lack just the long isoform of *CASK* have reduced MTM, showing that the *CaMK*-like and L27 domains only present in this form of *CASK* (Figure 1C) are the key signaling domains required for regulating memory. Previous work has shown that *CASK- β* regulates *CaMKII* autophosphorylation by its *CaMK*-like domain (Figure 1A; Lu et al., 2003; Hodge et al., 2006), therefore based on this and the data presented here, it is likely that the way *CASK* functions in memory formation is via its control of *CaMKII* autophosphorylation mediated by its N-terminal *CaMK*-like domain. MTM formation was highly sensitive to the level and specific distribution of *CASK* in the mushroom body, with targeted reduction of *CASK* in the mushroom body α'/β' neurons impairing memory, but with no apparent contribution from the α/β and γ neurons. Decreased levels of *CASK* are known to increase *CaMKII-T287* autophosphorylation (Figure 1A; Lu




FIGURE 7 | CASK and CaMKII are required for long-term memory. (A) Flies null for *CASK-β* or overexpressing *CASK-RNAi*, *CaMKII-T287D*, *CaMKII-T287A*, or *CaMKII-T306A T307A* throughout their mushroom body completely lacked LTM induced by five cycles of spaced training compared to heterozygous wildtype negative controls. (B) α'/β' neuron *CASK* knockdown or overexpression of *CaMKII-T287D*, *CaMKII-T287A*, and *CaMKII-T306A T307A* was sufficient to cause this reduction in LTM. (C) Mushroom body α/β and γ neuron *CASK* knockdown or overexpression of *CaMKII-T287D*, *CaMKII-T287A*,

and *CaMKII-T306A T307A* did not affect LTM. (D) Reduction of *CaMKII* throughout the mushroom body, in the α'/β' or α/β and γ neurons decreased LTM. (E) Overexpression of *CaMKII* or *CASK* throughout the mushroom body decreased LTM. (F) Mushroom body α'/β' overexpression of *CaMKII* or *CASK* decreased spaced LTM. (G) Overexpression of *CaMKII* or *CASK* in the mushroom body α/β and γ neurons (*MB247-Gal4*) did not affect LTM. Data were analyzed using One-Way ANOVA followed by a Tukey's *post-hoc* test. n.s. is not significant, ($p > 0.05$), * $p < 0.05$, ** $p < 0.01$, and *** $p < 0.001$.

et al., 2003; Hodge et al., 2006). Consistent with this, we found that direct overexpression of the *CaMKII-T287D* transgene in the α'/β' neurons caused a similar reduction in MTM as knocking-down *CASK* in the same neurons. While expression of *CaMKII-T287D* in the α/β and γ neurons have no effect

on MTM. Expression of *CASK* just in the α'/β' neurons fully rescued the complete lack of memory in *CASK-β* null mutants to wildtype levels, showing *CASK* signaling in only the mushroom body α'/β' neurons is necessary and sufficient for MTM formation.

We have also determined for the first time the effect of CaMKII overexpression on memory, showing α'/β' neuron expression completely removed MTM. In addition reduction of CASK just in neurons that express CaMKII was sufficient to remove MTM.





FIGURE 9 | Continued

overtime in the α'/β' mushroom body lobes (*c305a-Gal4*) co-expressing the different CASK and CaMKII transgenes or *CASK* β null indicated compared to the negative control *c305a/+* expressing GCaMP3 (solid black line).

(C) Histogram showing that the % change in peak GCaMP3.1 fluorescence is reduced in *CASK* β null and when *CASK-RNAi*, *CaMKII-RNAi*, or *CaMKII-T287D* were expressed in the α'/β' neurons, while *CaMKII* overexpression increased the maximum response compared to negative control (*c305a-Gal4, uas-GCaMP3*) level (denoted by dotted line for comparison). (D) Histogram showing baseline Ca^{2+} levels were increased when *CaMKII*, *CaMKII-T287D*, or *CaMKII-T306A T307A* were overexpressed in α'/β' neurons compared to negative control. *CASK* β null or α'/β' neuron overexpression of *CASK-RNAi*, *CASK*, or *CaMKII-RNAi* led to a reduction in baseline Ca^{2+} signaling. Data were analyzed using One-Way ANOVA followed by a Tukey's *post-hoc* test. n.s. is not significant, ($p > 0.05$), * $p < 0.05$, ** $p < 0.01$, and *** $p < 0.001$.

Furthermore, increasing CASK in α'/β' neurons also greatly reduced MTM and decreased basal Ca^{2+} signaling. Such increases in CASK would be expected to block T287 autophosphorylation (Hodge et al., 2006), and indeed we found α'/β' neuron T287A overexpression gave a similar MTM phenotype. The role of CASK and CaMKII-T287 autophosphorylation in the memory neurons is an acute physiological one as opposed to a developmental one, as reducing CASK or changing CaMKII-T287 autophosphorylation just in the adult mushroom body α'/β' neurons was sufficient to remove memory. Recently a second pair of CaMKII autophosphorylation sites (TT306/7) has been shown to be important for the control of plasticity and memory in mammals (Figures 1A,B; Elgersma et al., 2002; Zhang et al., 2005). We found α'/β' neuron *CaMKII-TT306/7AA* overexpression removed MTM. Lastly we demonstrated that overexpression of CASK completely rescued the memory deficit due to mushroom body overexpression of *CaMKII-T306/7AA* (Figure 6K). However, *CaMKII T306A T307A* expression in α'/β' neurons was insufficient to rescue CASK overexpression (Figure 6L). The last result may suggest CASK does not regulate CaMKII T306 T307 in α'/β' neurons, or perhaps the *c305a-Gal4* promoter may not have adequate strength or the exact spatiotemporal pattern required for both CASK and *CaMKII-T306A T307A* expression to make the fly remember as wildtype. Overall our data suggests that CASK regulates CaMKII autophosphorylation in a common pathway required for memory formation in the mushroom body.

CASK REGULATES CaMKII AUTOPHOSPHORYLATION IN THE MUSHROOM BODY α'/β' NEURONS DURING LONG-TERM MEMORY FORMATION

Previous work has shown mushroom body expression of CaMKII-T287D enhanced training but did not affect memory in the courtship conditioning assay, while CaMKII-T287A expression changed habituation and neuronal excitability, but resulted in no change in courtship conditioning memory (Mehren and Griffith, 2004). However, mushroom body expression of the *CaMKII-hairpin* transgene has been shown to decrease LTM using the olfactory aversive conditioning assay (Ashraf et al., 2006) and was associated with decreased mushroom body Ca^{2+} signaling (Akalal et al., 2010). The differences in effects of CaMKII on courtship and olfactory learning phenotypes maybe due to

differences in the circuitry employed in the two memory tasks and also the timing of memory measured in the two assays. Recently CaMKII has been shown to undergo CREB-dependent gene transcription and translation in mushroom body and DAL neurons during LTM (Chen et al., 2012). Consistent with these studies we showed mushroom body expression of *CaMKII-hairpin* only affects LTM. In addition this is the only CASK or *CaMKII* transgene that gave a memory phenotype when expressed in the α/β or γ neuron, this suggests that LTM is particularly sensitive and requires a certain baseline level of CaMKII activity in every type of mushroom body neuron in order to form LTM. This is in contrast to transgenic manipulation of CaMKII autophosphorylation levels in the α/β or γ neuron that have no effect on LTM, possibly because the endogenous CaMKII in these neurons maybe adequate to support enough of the appropriate autophosphorylation activity to generate LTM. This is in contrast to the critical role of α'/β' neurons that require the correct level of CASK, CaMKII, and CaMKII autophosphorylation in order to form LTM. Therefore, our data is consistent with the other studies showing α/β or γ (they did not test α'/β') neuron expression of *CaMKII-RNAi* disrupts LTM, furthermore these studies showed that α/β or γ neuron *CaMKII-RNAi* expression decreased peak GCaMP3 Ca^{2+} response (Ashraf et al., 2006; Akalal et al., 2010).

We also measured a similar reduction in peak Ca^{2+} response in the α'/β' neurons with *CaMKII-hairpin*; however, this was never tested for in the previous studies. We also found that the reciprocal CaMKII overexpression caused a large increase in peak Ca^{2+} response. Previous electrophysiological studies have shown neuronal expression of *CASK-RNAi* or *CaMKII-T287D* both decreased neural excitability in response to stimulation (Chen and Featherstone, 2011). Likewise we find expression of these transgenes caused a reduction in α'/β' neuron peak Ca^{2+} signaling. Therefore, the GCaMP3 data is consistent with the current model of CASK regulation of CaMKII autophosphorylation (Figure 1A; Lu et al., 2003; Hodge et al., 2006).

Flies with the *CASK* β null mutation or reduced CASK in the α'/β' neurons reduced LTM. The LTM effects of CASK could be explained by its role in transcriptional activation of various plasticity molecules including NMDA receptors (Wang et al., 2002; Huang and Hsueh, 2009). NMDA receptors have recently been shown to be required for LTM in *Drosophila* (Wu et al., 2007). Furthermore, CaMKII itself is known to be a direct target of NMDA receptor activation (Thalhammer et al., 2006) leading to increased CaMKII-T286 autophosphorylation and subsequent phosphorylation and activation of molecules required for synaptic plasticity and LTM (Trinidad et al., 2006). At present there is no evidence that *Drosophila* CASK translocates to the nucleus; however, the effects of CASK on LTM maybe through changes in *CaMKII* expression that is known to occur during LTM (Ashraf et al., 2006; Akalal et al., 2010). We show that the CaMKII molecular memory switch (pT287) is required for mushroom body LTM formation with phospho-mimic or block removing both ARM and LTM. Again this seems to be an evolutionarily conserved memory mechanism with T286 mutant mice also not being able to form LTM after massed training (Irvine et al., 2011).

HUMAN CASK FUNCTION IN MUSHROOM BODY α'/β' NEURONS RESTORES MEMORY PERFORMANCE OF CASK NULL FLIES

Point mutations in human CASK have been associated with neurological and cognitive defects, including severe learning difficulties resulting from mutations in the CaMK-like and SH3 domains (Najm et al., 2008; Piluso et al., 2009; Tarpey et al., 2009). Recently CASK mutation has been shown to cause a number of cognitive defects in flies including disrupted sleep and place preference (Slawson et al., 2011; Donelson et al., 2012). In addition to these defects we show that CASK mutants with deletion of the CaMK-like and L27 domains have extreme impairment of MTM and LTM formation. Furthermore, we show that α'/β' neuron overexpression of human CASK can fully substitute for the lack of *Drosophila* CASK- β and rescue the CASK- β mutant memory defect to wildtype. This demonstrates that neuronal function of CASK is conserved between *Drosophila* to human, validating the use of this model to understand CASK function in both the healthy and diseased brain.

REFERENCES

- Akalal, D. B., Yu, D., and Davis, R. L. (2010). A late-phase, long-term memory trace forms in the gamma neurons of *Drosophila* mushroom bodies after olfactory classical conditioning. *J. Neurosci.* 30, 16699–16708.
- Ashraf, S. I., McLoon, A. L., Sclarsic, S. M., and Kunes, S. (2006). Synaptic protein synthesis associated with memory is regulated by the RISC pathway in *Drosophila*. *Cell* 124, 191–205.
- Atasoy, D., Schoch, S., Ho, A., Nadasy, K. A., Liu, X., Zhang, W., et al. (2007). Deletion of CASK in mice is lethal and impairs synaptic function. *Proc. Natl. Acad. Sci. U.S.A.* 104, 2525–2530.
- Chen, C. C., Wu, J. K., Lin, H. W., Pai, T. P., Fu, T. F., Wu, C. L., et al. (2012). Visualizing long-term memory formation in two neurons of the *Drosophila* brain. *Science* 335, 678–685.
- Chen, K., and Featherstone, D. E. (2011). Pre and postsynaptic roles for *Drosophila* CASK. *Mol. Cell. Neurosci.* 48, 171–182.
- Cho, K. O., Wall, J. B., Pugh, P. C., Ito, M., Mueller, S. A., and Kennedy, M. B. (1991). The alpha subunit of type II Ca^{2+} /calmodulin-dependent protein kinase is highly conserved in *Drosophila*. *Neuron* 7, 439–450.
- Connolly, J. B., Roberts, I. J., Armstrong, J. D., Kaiser, K., Forte, M., Tully, T., et al. (1996). Associative learning disrupted by impaired Gs signaling in *Drosophila* mushroom bodies. *Science* 274, 2104–2107.
- Davis, R. D. (2011). Traces of *Drosophila* memory. *Neuron* 70, 8–19.
- Donelson, N., Kim, E. Z., Slawson, J. B., Vecsey, C. G., Huber, R., and Griffith, L. C. (2012). High-resolution positional tracking for long-term analysis of *Drosophila* sleep and locomotion using the “tracker” program. *PLoS ONE* 7:e37250. doi: 10.1371/journal.pone.0037250
- Elgersma, Y., Fedorov, N. B., Ikonen, S., Choi, E. S., Elgersma, M., Carvalho, O. M., et al. (2002). Inhibitory autophosphorylation of CaMKII controls PSD association, plasticity, and learning. *Neuron* 36, 493–505.
- Griffith, L. C., Verselis, L. M., Aitken, K. M., Kyriacou, C. P., Danho, W., and Greenspan, R. J. (1993). Inhibition of calcium/calmodulin-dependent protein kinase in *Drosophila* disrupts behavioral plasticity. *Neuron* 10, 501–509.
- Griffith, L. C., Wang, J., Zhong, Y., Wu, C. F., and Greenspan, R. J. (1994). Calcium/calmodulin-dependent protein kinase II and potassium channel subunit eag similarly affect plasticity in *Drosophila*. *Proc. Natl. Acad. Sci. U.S.A.* 91, 10044–10048.
- Hata, Y., Butz, S., and Sudhof, T. C. (1996). CASK: a novel dlG/PSD95 homolog with an N-terminal calmodulin dependent protein kinase domain identified by interaction with neurexins. *J. Neurosci.* 16, 2488–2494.
- Hodge, J., Mullasseril, P., and Griffith, L. (2006). Activity-dependent gating of CaMKII autonomous activity by *Drosophila* CASK. *Neuron* 51, 327–337.
- Hsueh, Y. P. (2006). The role of the MAGUK protein CASK in neural development and synaptic function. *Curr. Med. Chem.* 13, 1915–1927.
- Huang, T.-N., and Hsueh, Y.-P. (2009). CASK point mutation regulates protein-protein interactions and NR2b promoter activity. *Biochem. Biophys. Res. Commun.* 382, 219–222.
- Irvine, E. E., Danhiez, A., Radwanska, K., Nassim, C., Lucchesi, W., Godaux, E., et al. (2011). Properties of contextual memory formed in the absence of alpha CaMKII autophosphorylation. *Mol. Brain* 4, 1–10.
- Kelly, P. T., McGuinness, T. L., and Greengard, P. (1984). Evidence that the major postsynaptic density protein is a component of a Ca^{2+} /calmodulin-dependent protein kinase. *Proc. Natl. Acad. Sci. U.S.A.* 81, 945–949.
- Koh, Y. H., Popova, E., Thomas, U., Griffith, L. C., and Budnik, V. (1999). Regulation of DLG localization at synapses by CaMKII-dependent phosphorylation. *Cell* 98, 353–363.
- Krashes, M., Keene, A., Leung, B., Armstrong, J., and Waddell, S. (2007). Sequential use of mushroom body neuron subsets during *Drosophila* odor memory processing. *Neuron* 53, 103–115.
- Lisman, J., Schulman, H., and Cline, H. (2002). The molecular basis of CaMKII function in synaptic and behavioural memory. *Nat. Rev. Neurosci.* 3, 175–190.
- Lisman, J. E., and Zhabotinsky, A. M. (2001). A model of synaptic memory: a CaMKII/PP1 switch that potentiates transmission by organizing an AMPA receptor anchoring assembly. *Neuron* 31, 191–201.
- Lu, C., Hodge, J., Mehren, J., Sun, X., and Griffith, L. (2003). Regulation of the Ca^{2+} /CaM-responsive pool of CaMKII by scaffold-dependent autophosphorylation. *Neuron* 40, 1185–1197.
- Martin, J. R., and Ollo, R. (1996). A new *Drosophila* Ca^{2+} /calmodulin-dependent protein kinase (Caki) is localized in the central nervous system and implicated in walking speed. *EMBO J.* 15, 1865–1876.
- Mayford, M., Bach, M. E., Huang, Y. Y., Wang, L., Hawkins, R. D., and Kandel, E. R. (1996). Control of memory formation through regulated expression of a CaMKII transgene. *Science* 274, 1678–1683.
- McGuire, S. E., Le, P. T., Osborn, A. J., Matsumoto, K., and Davis, R. L. (2003). Spatiotemporal rescue of memory dysfunction in *Drosophila*. *Science* 302, 1765–1768.
- Mehren, J. E., and Griffith, L. C. (2004). Calcium-independent calcium/calmodulin-dependent protein kinase II in the adult *Drosophila* CNS enhances the training of pheromonal cues. *J. Neurosci.* 24, 10584–10593.
- Mukherjee, K., Sharma, M., Urlaub, H., Bourenkov, G. P., Jahn, R., Sudhof, T. C., et al. (2008). CASK functions as a Mg^{2+} -independent neurexin kinase. *Cell* 133, 328–339.
- Najm, J., Horn, D., Wimplinger, I., Golden, J. A., Chizhikov, V. V., Sudi, J., et al. (2008). Mutations of CASK cause an X-linked brain malformation phenotype with microcephaly and hypoplasia of the brainstem

- and cerebellum. *Nat. Genet.* 40, 1065–1067.
- Park, D., Coleman, M., Hodge, J., Budnik, V., and Griffith, L. (2002). Regulation of neuronal excitability in *Drosophila* by constitutively active CaMKII. *J. Neurobiol.* 52, 24–42.
- Piluso, G., D'Amico, F., Saccone, V., Bismuto, E., Rotundo, I. L., Di Domenico, M., et al. (2009). A missense mutation in CASK causes FG syndrome in an Italian family. *Am. J. Hum. Genet.* 84, 162–177.
- Ruta, V., Datta, S. R., Vasconcelos, M. L., Freeland, J., Looger, L. L., and Axel, R. (2010). A dimorphic pheromone circuit in *Drosophila* from sensory input to descending output. *Nature* 468, 686–690.
- Shuai, Y., Lu, B., Hu, Y., Wang, L., Sun, K., and Zhong, Y. (2010). Forgetting is regulated through Rac activity in *Drosophila*. *Cell* 140, 579–589.
- Silva, A. J., Paylor, R., Wehner, J. M., and Tonegawa, S. (1992a). Impaired spatial learning in alpha-calcium-calmodulin kinase II mutant mice. *Science* 257, 206–211.
- Silva, A. J., Stevens, C. F., Tonegawa, S., and Wang, Y. (1992b). Deficient hippocampal long-term potentiation in alpha-calcium-calmodulin kinase II mutant mice. *Science* 257, 201–206.
- Slawson, J. B., Kuklin, E. A., Ejima, A., Mukherjee, K., Ostrovsky, L., and Griffith, L. C. (2011). Central regulation of locomotor behavior of *Drosophila melanogaster* depends on a CASK isoform containing CaMK-like and L27 domains. *Genetics* 187, 171–184.
- Takamatsu, Y., Kishimoto, Y., and Ohsako, S. (2003). Immunohistochemical study of Ca²⁺/calmodulin-dependent protein kinase II in the *Drosophila* brain using a specific monoclonal antibody. *Brain Res.* 974, 99–116.
- Tarpey, P. S., Smith, R., Pleasance, E., Whibley, A., Edkins, S., Hardy, C., et al. (2009). A systematic, large-scale resequencing screen of X-chromosome coding exons in mental retardation. *Nat. Genet.* 41, 535–543.
- Tessier, C. R., and Broadie, K. (2011). The Fragile X mental retardation protein developmentally regulates the strength and fidelity of calcium signaling in *Drosophila* mushroom body neurons. *Neurobiol. Dis.* 41, 147–159.
- Thalhammer, A., Rudhard, Y., Tigaret, C. M., Volynski, K. E., Rusakov, D. A., and Schoepfer, R. (2006). CaMKII translocation requires local NMDA receptor-mediated Ca²⁺ signaling. *EMBO J.* 25, 5873–5883.
- Tian, L., Hires, S. A., Mao, T., Huber, D., Chiappe, M. E., Chalasani, S. H., et al. (2009). Imaging neural activity in worms, flies and mice with improved GCaMP calcium indicators. *Nat. Methods* 6, 875–881.
- Trinidad, J. C., Specht, C. G., Thalhammer, A., Schoepfer, R., and Burlingame, A. L. (2006). Comprehensive identification of phosphorylation sites in postsynaptic density preparations. *Mol. Cell. Proteomics* 5, 914–922.
- Tully, T., Preat, T., Boynton, S. C., and Del Vecchio, M. (1994). Genetic dissection of consolidated memory in *Drosophila*. *Cell* 79, 35–47.
- Tully, T., and Quinn, W. G. (1985). Classical conditioning and retention in normal and mutant *Drosophila melanogaster*. *J. Comp. Physiol. A* 157, 263–277.
- Wang, Z., Wilson, G., and Griffith, L. (2002). Calcium/calmodulin-dependent protein kinase II phosphorylates and regulates the *Drosophila* eag potassium channel. *J. Biol. Chem.* 277, 24022–24029.
- Wu, C.-L., Xia, S., Fu, T.-F., Wang, H., Chen, Y.-H., Leong, D., et al. (2007). Specific requirement of NMDA receptors for long-term memory consolidation in *Drosophila* ellipsoid body. *Nat. Neurosci.* 10, 1578–1586.
- Yasuda, M., and Mayford, M. R. (2006). CaMKII activation in the entorhinal cortex disrupts previously encoded spatial memory. *Neuron* 50, 309–318.
- Zars, T., Fischer, M., Schulz, R., and Heisenberg, M. (2000). Localization of a short-term memory in *Drosophila*. *Science* 288, 672–675.
- Zhang, L., Kirschstein, T., Sommersberg, B., Merckens, M., Manahan-Vaughan, D., Elgersma, Y., et al. (2005). Hippocampal synaptic metaplasticity requires inhibitory autophosphorylation of Ca²⁺/calmodulin-dependent kinase II. *J. Neurosci.* 25, 7697–7707.

Conflict of Interest Statement: The authors declare that the research was conducted in the absence of any commercial or financial relationships that could be construed as a potential conflict of interest.

Received: 11 January 2013; accepted: 09 March 2013; published online: 27 March 2013.

Citation: Malik BR, Gillespie JM and Hodge JLL (2013) CASK and CaMKII function in the mushroom body α'/β' neurons during *Drosophila* memory formation. *Front. Neural Circuits* 7:52. doi: 10.3389/fncir.2013.00052

Copyright © 2013 Malik, Gillespie and Hodge. This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits use, distribution and reproduction in other forums, provided the original authors and source are credited and subject to any copyright notices concerning any third-party graphics etc.