




Contents lists available at ScienceDirect

Asian Pacific Journal of Tropical Biomedicine

journal homepage: www.elsevier.com/locate/apjtb



Document heading doi:10.1016/S2221-1691(12)60096-3 © 2012 by the Asian Pacific Journal of Tropical Biomedicine. All rights reserved.

Antifungal activity of four honeys of different types from Algeria against pathogenic yeast: *Candida albicans* and *Rhodotorula* sp.

Ahmed Moussa^{1*}, Djebli Noureddine², Aissat Saad¹, Meslem Abdelmelek¹, Benhalima Abdelkader³¹Institute of Veterinary Sciences University, Ibn-khaldoun Tiaret (14000), Algeria²Department of Biology, Faculty of Sciences, Mostaganem University, Algeria³Laboratory Science and Technology Environment and Development, Mostaganem University, Algeria

ARTICLE INFO

Article history:

Received 15 January 2012

Received in revised form 25 January 2012

Accepted 18 March 2012

Available online 28 July 2012

Keywords:

Honey

Antifungal activity

*Candida albicans**Rhodotorula* sp.

ABSTRACT

Objective: To evaluate the antifungal activity of four honeys of different types from Algeria against pathogenic yeast *i.e.* *Candida albicans* (*C. albicans*) and *Rhodotorula* sp. **Methods:** Four Algeria honeys of different botanical origin were analyzed to test antifungal effect against *C. albicans*, and *Rhodotorula* sp. Different concentrations (undiluted, 10%, 30%, 50% and 70% w/v) of honey were studied *in vitro* for their antifungal activity using *C. albicans* and *Rhodotorula* sp. as fungal strains. **Results:** The range of the diameter of zone of inhibition of various concentrations of tested honeys was (7–23 mm) for *Rhodotorula* sp., while *C. albicans* showed clearly resistance towards all concentrations used. The MICs of tested honey concentrations against *C. albicans* and *Rhodotorula* sp. were (70.09–93.48)% and (4.90–99.70)% v/v, respectively. **Conclusions:** This study demonstrates that, *in vitro*, these natural products have clearly an antifungal activity against *Rhodotorula* sp. and *C. albicans*.

1. Introduction

The increase in the resistance of antifungal drugs in use has attracted the attention of the scientific community. *Candida albicans* (*C. albicans*) is a dimorphic organism that commonly inhabits in oral and vaginal mucosa and gastro-intestinal tract of human beings as one of the commensal organisms[1–4]. In addition to *C. albicans*, *Rhodotorula* sp. has been implicated as the etiologic agent of central venous catheter infection and fungemia[5–9]. In recent years, there has been an increasing search for new antifungal compounds due to the lack of efficacy, side effects and or resistance associated with some of the existing drugs[10–12]. Recently, the potential antifungal effect of honey has attracted serious attention within the scientific community[13–16]. Most types of honey generate hydrogen peroxide when diluted because of the activation of the

enzyme glucose oxidase, which oxidizes glucose to gluconic acid and hydrogen peroxide[17,18]. Hydrogen peroxide is the major contributor to the antimicrobial activity of honey, and the different concentrations of this compound in different honeys result in their varying antimicrobial effects[19–22]. The *in vitro* antifungal activity of honey was reported by Maria *et al*[23], who observed that honey stops the growth of *C. albicans*, *Candida krusei* and *Cryptococcus neoformans*. Obaseik–Ebor and Afongo[24] compared the antifungal activity of honey distillate with some antimycotic preparations against *C. albicans* and found that all the strains resistant to conventional antimycotic agents are inhibited by the active fraction of honey distillate.

However, only limited data are available on the susceptibility of *Rhodotorula* sp. to antifungal and antiseptic agents[25,26]. This study was aimed to confirm the usage of Algeria honey as antifungal and antiseptic agents and evaluate this inhibitory action at different honey concentration against *C. albicans* and *Rhodotorula* sp.

2. Materials and methods

2.1. Honey samples

*Corresponding author: Ahmed Moussa, Institute of Veterinary Sciences University, Ibn-Khaldoun Tiaret (14000), Algeria.

Tel: +213 65234059

Fax: +213 46 425001

E-mail: moussa7014@yahoo.fr

Foundation Project: This work was financially supported by project CNEPRU, Institute of Veterinary Sciences (IVS), University Ibn-Khaldoun (TIARET), Algeria (grant No. F023 2009/0009).

During the 2011 flowering seasons, four honey samples were gathered and provided by various bee-keepers from two different areas of the western Algeria. These honey samples were aseptically collected in sterile screwed cups and kept in a cool and dry place (at room temperature) overnight before they were finally transported to the laboratory.

2.2. Preparation of honey solutions

Honey solutions were prepared immediately prior to testing by diluting honey to the required concentrations (undiluted, 10%, 30%, 50%, and 70%, w/v). All samples were then incubated for 30 min at 37 °C in a shaking water bath that allowed aeration of the solutions. Incubation was carried out in the dark because both hydrogen peroxide and glucose oxidase are light sensitive[27].

2.3. Yeast strains and susceptibility testing

Yeasts were maintained on Sabouraud dextrose agar (SDA; BioMérieux, Marcy l'Etoile, France) at 4 °C, and sub cultures were performed prior to each experiment in the same medium for 48 h at 35 °C. Turbidity standard and preparation of inocula: stock fungal inoculum suspensions were prepared in sterile saline from 48 h cultures on SDA at 35 °C. Each suspension was adjusted visually to 0.5 McFarland turbidity standard. Dilutions of these suspensions were subcultured on SDA to determine the number of CFU/mL. The adjusted inoculum was $1 \times 10^7 - 5 \times 10^7$ CFU/mL.

2.4. Antifungal assay

Three different methods were used to evaluate the antifungal activity of honey: disc diffusion, well and spectrophotometric methods[28].

Antifungal activity of honey was evaluated using agar disc diffusion method against test microorganisms. About 100 μ L of fresh culture suspension of the test microorganisms was spread on the respective media Sabouraud dextrose agar plates. The concentration of cultures was 1×10^7 CFU/mL. For screening, sterile filter paper discs (5 mm diameter) were impregnated

with 10 μ L of honey equivalent to 0.1 mg of honey after being placed on the surface of the inoculated media agar plates. The plates were stood at 4 °C for 2 h before being incubated under optimum conditions at 37 °C for 24 h. Clear inhibition zones around the discs indicated the presence of antimicrobial activity. The diameters of the inhibition zones were measured in millimeter, including the diameter of disc. The controls were set up with equivalent quantities of water.

The agar well diffusion method was employed. The honey samples were first inoculated separately on standard nutrient media with no test organisms so as to evaluate their possible contamination. Thereafter, solidified nutrient agar plates were separately flooded with the liquid inoculums of the different test organisms using the spread plate method. The plates were drained and allowed to dry at 37 °C for 30 min after which four equidistant wells of 5 mm in diameter were punched using a sterile cork borer at different sites on the plates. About 50 μ L of the different concentrations (undiluted, 30%, 50% and 70% w/v) of the honey samples were separately placed in the different punched wells with 1 mL sterile syringe. The plates were allowed to stay for 15 min for pre-diffusion to take place followed by an overnight incubation that lasted for 24 h at 37 °C. The diameter of inhibition zones, including the diameter of the well, was recorded. Each assay was carried out in triplicate.

2.5. Minimum inhibitory concentration (MIC) determination

Concentrations of honey suspensions (undiluted, 10%, 30%, 50% and 70%) were incorporated into media to test their efficiency against *C. albicans* and *Rhodotorula* sp. Each plate reaching final volume of 5 mL including both honey and media was inoculated and incubated at 37 °C for 48 h. The MIC was determined by finding the plates with the lowest concentration of honey on which the strain would not grow. All MIC values were expressed in % (v/v).

3. Results

Table 1 showed the inhibition zone sizes produced by various honeys at different dilutions. The diameters of zone

Table 1
Antifungal activity of honey at different concentrations against *C. albicans* and *Rhodotorula* sp.

Yeast strains	Honey dilution	Inhibition zone diameter (mm)							
		Disc diffusion method				Well diffusion method			
		Honey A	Honey B	Honey C	Honey D	Honey A	Honey B	Honey C	Honey D
<i>C. albicans</i>	Undiluted	ND	ND	ND	ND	ND	ND	ND	ND
	70%	ND	ND	ND	ND	ND	ND	ND	ND
	30%	ND	ND	ND	ND	ND	ND	ND	ND
	50%	ND	ND	ND	ND	ND	ND	ND	ND
	10%	ND	ND	ND	ND	ND	ND	ND	ND
<i>Rhodotorula</i> sp.	Undiluted	9	8	10	8	22	20	23	14
	70%	ND	ND	ND	ND	ND	ND	ND	ND
	30%	9	8	11	9	13	10	11	8
	50%	13	10	10	7	15	14	18	17
	10%	8	8	7	7	12	11	10	11

ND: No inhibition was detected.

Table 2
MIC of honey at different concentrations against *C. albicans* and *Rhodotorula* sp.

Honey concentrations	<i>C. albicans</i> MIC _% (v/v)				<i>Rhodotorula</i> sp. MIC _% (v/v)			
	Undiluted	50% (v/v)	25% (v/v)	12.5% (v/v)	Undiluted	50% (v/v)	25% (v/v)	12.5% (v/v)
Honey A	81.16	70.09	>100.00	>100.00	87.30	96.37	25.04	>100.00
Honey B	91.36	73.94	>100.00	>100.00	89.76	94.63	43.56	>100.00
Honey C	93.48	75.18	>100.00	>100.00	56.14	99.70	5.65	>100.00
Honey D	84.30	79.27	>100.00	>100.00	94.12	80.50	4.90	>100.00

of the inhibition of honey with various concentrations tested for *Rhodotorula* sp. ranged from 7 to 13 mm and 8 to 23 mm for disc and well diffusion method, respectively. While *C. albicans* showed resistance towards all honey concentrations used by both methods. The MIC ranges of the tested honey concentrations were (70.09–93.48) and (4.90–99.70)% v/v against *C. albicans* and *Rhodotorula* sp., respectively (Table 2).

4. Discussion

The conventional treatment of fungal disease is limited, and part of the reason is due to the limited spectrum of the currently antifungal drugs, and the expensive treatment, particularly due to the need of prolonged therapy. In recent years, several studies on the *in vitro* susceptibility of superficial mycoses to antifungal drugs have been done and the results have shown considerable variations[29,30]. Thus, nowadays many researches are focused on the therapeutical properties of natural compounds[31]. Honey is a natural product that is used for its antifungal activity[14]. Several factors may influence the antifungal activity of honey. For example, DeMera and Angert[32–38] reported that honeys from different phyto-geographic regions vary in their ability to inhibit the growth of yeasts, suggesting that botanical origin plays an important role in influencing the antifungal activity. In addition, there are a great variety of components, including phenolic acids, flavonoids and other biomolecules, in different honeys. Biological activity of honey is mainly attributed to the phenolic compounds reported by Estevinho *et al*[39]. In fact, the antimicrobial action of phenolics is well known and it is related to their ability to denature proteins, being generally classified as surfaceactive agents. Xesus and Maria[40] suggest that the honey mechanism for fungal growth inhibition is not related to the osmotic shock derived from the presence of sugar in the culture medium. Moreover, Wahdan[41] stated that high sugar concentration in honey leads to the high osmolarity that produces antimicrobial activity. Additionally, he found no inhibitory activity of the sugar solutions against *Trichophyton mentagrophytes* and *C. albicans* and noted that fungi are generally much more tolerant than bacteria to the high osmotic effect. Diekema *et al*[42] reported the *in vitro* activities of 8 antifungals against 64 *Rhodotorula* isolates. *Rhodotorula* strains were resistant *in vitro* to fluconazole (MIC₅₀, 1128 mg/ mL) and caspofungin (MIC₅₀, 18 mg/mL). In the present study, *Rhodotorula* sp. was susceptible to honey since growth inhibition was reached at the minor level. Honey samples inhibited completely the growth of *Rhodotorula* sp.

Our results showed that undiluted honey was able to

inhibit the growth of many species of *Rhodotorula* sp. but there was no effect on *C. albicans*. Al-Wailii[43] found that honey concentration ranging from 30% to 50% inhibits the growth of several pathogenic microorganisms, including *C. albicans*. Irish *et al*[6] reported antifungal efficacy of various honeys against clinical isolates of *C. albicans*, *Candida glabrata*, and *Candida dubliniensis*. Khosravi *et al*[44] reported that honey has antifungal activity against *Candida* species such as *C. albicans*, *Candida parapsilosis*, *Candida tropicalis*, *Candida kefyr*, *Candida glabrata*, and *Candida dubliniensis*. The results of this preliminary study demonstrated that Algeria honey is an effective inhibitor of *Rhodotorula* sp.

Conflict of interest statement

We declare that we have no conflict of interest.

Acknowledgements

Authors thank staff of Tiaret University for providing materials.

References

- [1] Arra BA, Zakaria Z, Sreenivasan S. A transmission electron microscopy study of the diversity of *Candida albicans* cells induced by *Euphorbia hirta* L. leaf extract *in vitro*. *Asian Pac J Trop Biomed* 2011; **1**: 20–22.
- [2] Cristina SM, José PM. Dimorphism in fungal pathogens: *Candida albicans* and *Ustilago maydis*—similar inputs, different outputs. *Curr Opin Microbiol* 2001; **4**(2): 214–221.
- [3] Gloria M, Rosalía DO, Federico NG, Lucía M, Jesús P, Concha G, et al. *Candida albicans*: genetics, dimorphism and pathogenicity. *Int Microbiol* 1998; **1**: 95–106.
- [4] Dean DA, Burchard KW. Surgical perspective on invasive *Candida* infections. *World J Surg* 1998; **22**: 127–134.
- [5] Kiehn E, Gorey E, Brown AE, Edwards FF, Armstrong D. Sepsis due to *Rhodotorula* related to use of indwelling central venous catheters. *Clin Infect Dis* 1992; **14**: 841–846.
- [6] Tuon FF, Costa SF. *Rhodotorula* infection. A systematic review of 128 cases from literature. *Rev Iberoam Micol* 2008; **25**: 135–140.
- [7] Samonis G, Anatoliotaki M, Apostolou H, Maraki S, Mavroudis D, Georgoulis V. Transient fungemia due to *Rhodotorula rubra* in a cancer patient: case report and review of the literature. *Infection* 2001; **29**: 173–176.
- [8] Tuon FF, de Almeida GM, Costa SF. Central venous catheter-

- associated fungemia due to *Rhodotorula* spp.—a systematic review. *Med Mycol* 2007; **45**: 441–447.
- [9] Rusthoven JJ, Feld R, Tuffeuell PJ. Systemic infection by *Rhodotorula* spp. in the immunocompromised host. *J Infect* 1984; **8**: 241–246.
- [10] Intzar A, Farrah GK, Krishan AS, Bishan DG, Naresh KS, Prabhu D, et al. *In vitro* antifungal activity of hydroxychavicol isolated from *Piper betle* L. *Ann Clin Microbiol Antimicrob* 2010; **9**: 7.
- [11] Klepser ME. *Candida* resistance its clinical relevance. *Pharmacotherapy* 2006; **26**: 68S–75S.
- [12] Barker KS, Rogers PD. Recent insights into the mechanisms of antifungal resistance. *Curr Infect Dis Rep* 2006; **8**: 449–456.
- [13] Ahmed M, Djebli N, Aissat S, Hammoudi SM, Bourabeh A, Hemida H. Additive potential of ginger starch on antifungal potency of honey against *Candida albicans*. *Asian Pac J Trop Biomed* 2012; **2**(1): 253–255.
- [14] Irish J, Carter DA, Shokohi T, Blair S. Honey has an antifungal effect against *Candida* species. *Med Mycol* 2006; **44**: 289–291.
- [15] Cancliracci M, Citterio B, Piatti E. Antifungal activity of the honey flavonoid extract against *Candida albicans*. *Food Chem* 2012; **131**(2): 493–499.
- [16] Khosravi AR, Shokri H, Katirae F, Ziglari T, Forzi M. Fungicidal potential of different Iranian honeys against some pathogenic *Candida* species. *J Apic Res* 2008; **47**(8): 256–260.
- [17] White JW, Subers MH, Schepartz AI. The identification of inhibine, the antibacterial factor in honey, as hydrogen peroxide and its origin in a honey glucose–oxidase system. *Biochim Biophys Acta* 1963; **73**: 57–70.
- [18] Bang LM, Bunting C, Molan PC. The effect of dilution on the rate of hydrogen peroxide production in honey and its implications for wound healing. *J Altern Complement Med* 2003; **9**(2): 267–273.
- [19] Molan PC. The antibacterial nature of honey: 1. The nature of the antibacterial activity. *Bee World* 1992; **73**(1): 5–28.
- [20] Cooper RA, Halas E, Molan PC. The efficacy of honey in inhibiting strains of *Pseudomonas aeruginosa* from infected burns. *J Burn Care Rehabil* 2002; **23**: 366–370.
- [21] Basualdo C, Sgroy V, Finola MS, Juam M. Comparison of the antibacterial activity of honey from different provenance against bacteria usually isolated from skin wounds. *Vet Microbiol* 2007; **124**: 375–381.
- [22] Adeleke OE, Olaitan JO, Okepekpe EI. Comparative antibacterial activity of honey and gentamicin against *Escherichia coli* and *Pseudomonas aeruginosa*. *Ann Burns Fire Disasters* 2006; **19**: 201–204.
- [23] Maria LE, Afonso SE, Xesús F. Antifungal effect of lavender honey against *Candida albicans*, *Candida krusei* and *Cryptococcus neoformans*. *J Food Sci Technol* 2011; **48**(5): 640–643.
- [24] Obaseiki–Ebor EE, Afonya TCA. *In vitro* evaluation of the anticandidiasis activity of honey distillate (HY–I) compared with that of some antimycotic agents. *J Pharm Pharmacol* 1984; **36**: 283–284.
- [25] Kiehn TE, Gorey E, Brown AE, Edwards FE, Armstrong D. Sepsis due to *Rhodotorula* related to use of indwelling central venous catheters. *Clin Infect Dis* 1992; **14**: 841–846.
- [26] Galan–Sanchez F, Garcia–Martos P, Rodriguez–Ramos C, Marin–Casoava P, Mira–Gutierrez J. Microbiological characteristics and susceptibility patterns of strains of *Rhodotorula* isolated from clinical samples. *Mycopathologia* 1999; **145**: 109–112.
- [27] White JW, Subers MH, Schepartz AI. The identification of inhibine, the antibacterial factor in honey, as hydrogen peroxide and its origin in a honey glucose–oxidase system. *Biochim Biophys Acta* 1963; **73**: 57–70.
- [28] Patton T, Barrett J, Brennan J, Moran N. Use of a spectrophotometric bioassay for determination of microbial sensitivity to manuka honey. *J Microbiol Methods* 2006; **64**: 84–95.
- [29] Jessup CJ, Warner J, Isham N, Hasan I, Ghannoum MA. Antifungal susceptibility testing of dermatophytes: establishing a medium for inducing conidial growth and evaluation of susceptibility of clinical isolates. *J Clin Microbiol* 2000; **38**: 341–344.
- [30] Grupta AK, Sauder DN, Shear NH. Antifungal agents: an overview Part II. *J Am Acad Dermatol* 1994; **30**: 911–933.
- [31] Ji HF, Li XJ, Zhang HY. Natural products and drug discovery. *EMBO Rep* 2009; **10**: 194–200.
- [32] DeMera JH, Angert ER. Comparison of the antimicrobial activity of honey produced by *Tetragonisca angustula* (Meliponinae) and *Apis mellifera* from different phylogeographic regions of Costa Rica. *Apidologie* 2004; **35**: 411–417.
- [33] Balakumar S, Rajan S, Thirunalasundari T, Jeeva S. Antifungal activity of *Ocimum sanctum* Linn. (Lamiaceae) on clinically isolated dermatophytic fungi. *Asian Pac J Trop Med* 2011; **4**: 654–657.
- [34] Madhumitha G, Saral AM. Preliminary phytochemical analysis, antibacterial, antifungal and anticandidal activities of successive extracts of *Crossandra infundibuliformis*. *Asian Pac J Trop Med* 2011; **4**: 192–195.
- [35] Shahbudin S, Muhammad T, Deny S, Haitham Q, Nurul Afifah Binti Abdul R. Antimicrobial activity of mangrove plant (*Lumnitzera littorea*). *Asian Pac J Trop Med* 2011; **4**: 523–525.
- [36] Bragadeeswaran S, Priyadharshini S, Prabhu K, Rani SRS. Antimicrobial and hemolytic activity of fish epidermal mucus *Cynoglossus arel* and *Arius caelatus*. *Asian Pac J Trop Med* 2011; **4**: 305–309.
- [37] Adwan G, Abu–Shanab B, Adwan K. Antibacterial activities of some plant extracts alone and in combination with different antimicrobials against multidrug–resistant *Pseudomonas aeruginosa* strains. *Asian Pac J Trop Med* 2011; **4**: 266–269.
- [38] Okoye TC, Akah PA, Okoli CO, Ezike AC, Mbaoji FN. Antimicrobial and antispasmodic activity of leaf extract and fractions of *Stachytarpheta cayennensis*. *Asian Pac J Trop Med* 2011; **4**: 189–192.
- [39] Estevinho L, Pereira AP, Moreira L, Dias LG, Pereira E. Antioxidant and antimicrobial effects of phenolic compound extracts of Northeast Portugal honey. *Food Chem Toxicol* 2008; **46**: 3774–3779.
- [40] Xesús F, Estevinho ML. A survey of the *in vitro* antifungal activity of *Heather* (*Erica* sp.) organic honey. *J Med Food* 2011; **14**: 1–5.
- [41] Wahdan HAL. Causes of the antimicrobial activity of honey. *Infection* 2008; **1**: 26–31.
- [42] Diekema DJ, Petroelje B, Messer SA, Hollis RJ, Pfaller MA. Activities of available and investigational antifungal agents against *Rhodotorula* species. *J Clin Microbiol* 2005; **43**: 476–478.
- [43] Al–Waili N. Therapeutic and prophylactic effects of crude honey on chronic seborrhic dermatitis and dandruff. *Eur J Med Res* 2001; **6**: 306–308.
- [44] Khosravi AR, Shokri H, Katirae F, Taherel F. Fungicidal potential of different Iranian honeys against some pathogenic *Candida* species. *J Apic Res* 2008; **47**: 256–260.