




Contents lists available at ScienceDirect

Asian Pacific Journal of Tropical Biomedicine

journal homepage: www.elsevier.com/locate/apjtb

Document heading doi:10.1016/S2221-1691(13)60015-5 © 2013 by the Asian Pacific Journal of Tropical Biomedicine. All rights reserved.

Anti-bacterial activity and brine shrimp lethality bioassay of methanolic extracts of fourteen different edible vegetables from Bangladesh

M. Obayed Ullah¹, Mahmuda Haque², Kaniz Fatima Urmi³, Abu Hasanat Md. Zulfiker², Elichea Synthi Anita², Momtaj Begum², Kaiser Hamid^{4*}

¹ School of Chemistry and Molecular Biosciences, University of Queensland, QLD, Australia

² Department of Pharmacy, Southeast University, Bangladesh

³ Department of Pharmacy, Jahangirnagar University, Bangladesh

⁴ Department of Pharmacy, East West University, Bangladesh

PEER REVIEW

Peer reviewer

Dr. Shaikh Jamal Uddin, Research Fellow, School of Pharmacy, Griffith University, Gold Coast campus, QLD, Australia

Tel: +61-07-55529236

Fax: +61-07-55528804

E-mail: s.uddin@griffith.edu.au

Comments

This is a good paper where authors evaluated the antibacterial activity against 12 different gram positive and gram negative bacteria and brine shrimp lethality bioassay of 14 edible vegetables from Bangladesh.

The results are interesting where it has been found that most of the vegetables are effective against gram negative bacteria rather than gram positive bacteria.

(Details on Page 5)

ABSTRACT

Objective: To investigate the antibacterial and cytotoxic activity of fourteen different edible vegetables methanolic extract from Bangladesh. **Methods:** The antibacterial activity was evaluated using disc diffusion assay method against 12 bacteria (both gram positive and gram negative). The plant extracts were also screened for cytotoxic activity using the brine shrimp lethality bioassay method and the lethal concentrations (LC₅₀) were determined at 95% confidence intervals by analyzing the data on a computer loaded with “Finney Programme”. **Results:** All the vegetable extracts showed low to elevated levels of antibacterial activity against most of the tested strains (zone of inhibition=5–28 mm). The most active extract against all bacterial strains was from *Xanthium indicum* which showed remarkable antibacterial activity having the diameter of growth inhibition zone ranging from 12 to 28 mm followed by *Alternanthera sessilis* (zone of inhibition=6–21 mm). All extracts exhibited considerable general toxicity towards brine shrimps. The LC₅₀ value of the tested extracts was within the range of 8.447 to 60.323 µg/mL with respect to the positive control (vincristine sulphate) which was 0.91 µg/mL. Among all studied extracts, *Xanthium indicum* displayed the highest cytotoxic effect with LC₅₀ value of 8.447 µg/mL. **Conclusions:** The results of the present investigation suggest that most of the studied plants are potentially good source of antibacterial and anticancer agents.

KEYWORDS

Bangladeshi vegetables, Antibacterial, Disc diffusion, Brine shrimp lethality

1. Introduction

Bangladesh is a developing and over populated country where the most of the people are living below the poverty level especially in rural area. Most of the people in Bangladesh are habituated with the local and indigenous food. Vegetables are very common in their everyday food menu and play an important role in the food chain. More than sixty, indigenous and exotic, vegetables are grown in

Bangladesh, most of which are very cheap, available and popular to people for daily food intake. The enriched food value as well as medicinal property are still unexplored and may play a pivotal role in the development of potential new drugs for chemotherapy which might help to overcome the growing problem of resistance and also the toxicity of the currently available commercial antibiotics. The traditional medicinal methods, especially the use of vegetable may play a vital role to cover the basic health needs in the

*Corresponding author: Kaiser Hamid, Lecturer, Department of Pharmacy, East West University, Dhaka, Bangladesh,
Tel: +8801926759309
E-mail: kaiserpharm_1134@yahoo.com

Article history:

Received 10 Oct 2012

Received in revised form 12 Nov 2012, 2nd revised form 21 Nov, 3rd revised form 28 Nov 2012

Accepted 20 Dec 2012

Available online 28 Jan 2013

developing countries like Bangladesh.

In the present study we screened fourteen vegetables for their antibacterial as well as cytotoxic properties. Most of them are reported for their individual as well as combined medicinal properties. *Alternanthera sessilis* was reported for antioxidant activity^[1], hepatoprotective activity^[2], antimicrobial and wound healing activities^[3], antiviral activity^[4]. *Amaranthus tricolor* was reported to have cyclooxygenase and human tumour cell growth inhibitory activities^[5]. *Benincasa hispida* was found to have antiulcerogenic activity, bronchospasm inhibition activity, antioxidant and angiotensin converting enzyme inhibition capacity, hypoglycemic and antihyperglycemic activity^[6–10], *Corchorus olitorius* was found to have antibacterial activity, antioxidant and antidiabetic effects, anti-angiogenic effects, anorectic activity, glucose and lipid lowering activity^[10–14], anticonvulsion and antisteroidogenic activity, cytotoxic and antimalarial activity^[15–18] for the source of histamine release inhibitors and phenolic antioxidants^[19,20]. *Diplazium esculentum* was found to have antioxidant and cytotoxic activity^[21,22].

Enhydra fluctuans was reported for antioxidant, analgesic, antimicrobial activity, and antibacterial activity^[23–26]. *Glinus oppositifolius* (L.) was found to contain immunomodulating peptic polymer, terpenoid saponine having antiprotozoal activity, and bioactives having antioxidant property^[27–29]. *Ipomoea aquatica* Forsk was reported for hypoglycemic activity, cytotoxic activity^[30–32]. *Lagenaria siceraria* was reported for its diuretic activity^[33], antihypertensive and cardioprotective effect^[34]. *Nymphaea nouchalli* was reported for hypoglycemic antitumor and antimicrobial activity^[35–37]. *Portulaca oleracea* Linn was reported for analgesic and anti-inflammatory effects, wound healing activity, neuropharmacological activity^[38–40]. *Spinacia oleracea* Linn has protective effect against radiation-induced oxidative stress, antimutagenic activity. *Xanthium indicum* has antitrypanosomal activity, antitumor and anti-inflammatory activity^[41–45].

The plant kingdom comprises many species of plants containing substances of medicinal value, which are yet to be explored. A large number of plants are constantly being screened for their possible medicinal value^[46]. In recent years, multiple drug resistance in human pathogenic microorganism has been developed due to indiscriminate use of commercial antimicrobial drugs commonly used in the treatment of various diseases^[47]. Therefore, it is of great interest to carry out a screening of these plants in order to unfold the dimension of use as antimicrobial and cytotoxic activity and to reveal the active principle by isolation and characterization of their constituents.

2. Materials and methods

2.1. Drugs and chemicals

Methanol and DMSO (dimethyl sulfoxide) were purchased from Merck, Germany. Kanamycin was collected from Square Pharmaceuticals Ltd., Bangladesh. Vinchristin sulphate was purchased from Cipla Ltd., Goa, India.

2.2. Collection of plant material

The fresh plants were collected during the month of December–January (2007–2008) from different areas of Dhaka city and their identity were authenticated from the Bangladesh National Herbarium, Dhaka where a voucher specimen was deposited having different accession number which is given here in parenthesis: *Alternanthera sessilis* Linn (DACB–32915), *Amaranthus tricolor* (DACB–34186), *Benincasa hispida* (DACB–34185), *Corchorus olitorius* (DACB–34187), *Diplazium esculentum* (DACB–34181), *Enhydra Fluctuans* (DACB–32910), *Glinus oppositifolius* (DACB–34183), *Ipomoea aquatica/Ipomoea alba* (DACB–32911) *Lagenaria siceraria* (DACB–34188), *Nymphaea nouchalli* (DACB–32912), *Portulaca oleracea* Linn (DACB–34182), *Spinacia oleracea* (DACB–32914), *Xanthium indicum* (DACB–34184). Leaves of *Corchorus olitorius* and whole herb for other plants were used for the experiments.

2.3 Extraction of plant materials

The whole plants were firstly sun dried for 5 d and then finally dried in an oven at 40–45 °C for 72 h. The dried plant parts were ground into coarse powders. About 100 grams of the powdered materials of the each plant part were extracted separately with Soxhlet Extractor for 2 d. Then each of the extracts were filtered using cotton and filter paper. The clear filtrates were then evaporated to dryness under vacuum in a rotary evaporator.

2.4. Determination of the antibacterial activity

2.4.1. Antibacterial assay

Twelve pathogenic bacterial strains were used as the test organisms for antibacterial screening of the dried extract, of which four gram were +ve and eight gram –ve. The antibacterial activity of the plant extracts against the test organisms was determined by disc diffusion method using Kanamycin disc 30 µg/disc as standard for comparison^[48,49]. Disc diffusion methods are extensively used to investigate the antibacterial activity of natural substances and plant extracts. These assays are based on the use of discs as reservoirs containing solutions of substances to be examined. In the case of solutions with a low activity, however, a large concentration or volume is needed. The limited capacity of discs means that holes or cylinders are preferably used^[50].

After placing the paper discs bearing the sample, the plates were incubated at (35±2) °C for 2 d and the zones of inhibition were determined. Clear inhibition zones around the discs indicated the presence of antibacterial activity.

2.4.2. Collection of microorganisms

The bacterial species used in the present study were *Bacillus megaterium*, *Bacillus subtilis*, *Staphylococcus aureus*, *Sarcina lutea*, *Salmonella paratyphi*, *Salmonella typhi*, *Vibrio parahemolyticus*, *Vibrio mimicus*, *Escherichia coli*, *Shigella dysenteriae*, *Shigella boydii* and *Pseudomonas aeruginosa*. These were collected as pure cultures from the Institute of Nutrition and Food Sciences, Dhaka University and International Center for Diarrheal Disease and Research, Bangladesh Dhaka, Bangladesh.

2.5. Determination of cytotoxicity

2.5.1. Brine shrimps

Brine shrimp, *Artemia salina* Leach, also known as sea monkeys, are marine invertebrates about 1 mm in size. Freeze-dried cysts were readily available at aquarium stores. The cysts lasted for several years and could be hatched without special equipment^[51].

2.5.2. Brine shrimp lethality bioassay

The cytotoxic activity of the plant was evaluated using Brine shrimp lethality bioassay method where 6 graded doses (*viz* 200 µg/mL, 100 µg/mL, 50 µg/mL, 20 µg/mL, 10 µg/mL, and 5 µg/mL) were used^[52]. Brine shrimps (*Artemia salina* Leach) nauplii (Ocean 90, USA) were used as test organisms. For hatching, eggs were kept in brine with a constant oxygen supply for 48 h. The mature nauplii were then used in the experiment. DMSO was used as a solvent and also as a negative control. Vincristine sulfate was used as a reference standard in this case. The numbers of survivors were counted after 24 h. Larvae were considered dead if they did not exhibit any internal or external movement during several seconds of observation. The larvae did not receive food. To ensure that the mortality observed in the bioassay could be attributed to bioactive compounds and not to starvation; we compared the dead larvae in each treatment to the dead larvae in the control.

2.6. Statistical analysis

The median lethal concentration (LC₅₀) and 95% confidence intervals of the test samples were calculated using the probit analysis method described by Finney^[53], as the measure of toxicity of the plant extract.

3. Results

In the present study, methanolic extracts of fourteen vegetables were screened for their antibacterial activity by disc diffusion method using dried extracts of 500 µg/disc against 12 pathogenic bacterial species of both gram

positive and gram negative organisms and the results were compared with the standard Kanamycin (30 µg/disc). The gram negative species were *Vibrio parahemolyticus*, *Pseudomonas sureus*, *Salmonella typhi*, *Salmonella paratyphi*, *Shigella dysenteriae*, *Shigella boydii*, *Vibrio mimicus* and *Escherichia coli*, gram positive species were *Sarcina lutea*, *Bacillus megaterium*, *Staphylococcus aureus* and *Bacillus subtilis*. Among all the extracts, *Xanthium indicum* and *Enhydra fluctuans* were active against all of the tested bacteria. On the other hand, *Ipomoea aquatica*, *Chenopodium album*, *Alternanthera sessilis*, *Portulaca grandifolia* and *Lagenaria siceraria* were active against most of the tested bacteria. The rest of the plants showed no or little antibacterial activities against the tested bacteria. Among the extracts, the largest spectrum of sensitivity was found in *Xanthium indicum* extract where higher potential of sensitivity were found against *Pseudomonas aeruginosa* and it was recorded as 28 mm (Table 1).

Xanthium indicum also showed potent inhibitory activity against other tested microorganisms with the zone of inhibition of 13–20 mm. On the other hand the weakest inhibitory activity was found against *Vibrio mimicus* by *Corchorus olitorius* (inhibition zone diameter 5 mm). Furthermore the growth of *Salmonella paratyphi* (21 mm), *Salmonella typhi* (20 mm), *Escherichia coli* (16 mm) and *Sarcina lutea* (13 mm) were strongly inhibited by *Alternanthera sessilis* (Table 1).

The LC₅₀ values of the brine shrimp lethality bioassay obtained for these extracts and that of the positive control, Vincristine sulphate, have been presented in Table 2 and Table 3 respectively. All extracts exhibited significant toxicity towards brine shrimps. The LC₅₀ values of the plant extracts were within the range of 8.447 to 60.323 µg/mL, whereas that of the positive control (vincristine sulphate) was 0.91 µg/mL, LC₉₀=6.523, 95% IC=0.46–1.79. So these extracts can be considered as a promising candidate for a plant derived anticancer compound. Among the extracts *Xanthium indicum* showed more cytotoxic effects with LC₅₀ value 8.447 µg/mL whereas *Spinacia oleracea* was the least toxic among the tested vegetables having LC₅₀ value 60.323 µg/mL.

Table 1
Antibacterial activity of fourteen different edible vegetables from Bangladesh.

Plants	Zone of inhibition (mm)											
	Gram negative bacteria						Gram positive bacteria					
	<i>E. coli</i>	<i>P. aeruginosa</i>	<i>S. typhi</i>	<i>S. paratyphi</i>	<i>S. dysenteriae</i>	<i>S. boydii</i>	<i>V. mimicus</i>	<i>V. parahemolyticus</i>	<i>B. subtilis</i>	<i>B. megaterium</i>	<i>S. aureus</i>	<i>S. lutea</i>
<i>Enhydra fluctuans</i>	6	6	6	8	6	6	6	6	6	6	6	6
<i>Ipomoea aquatica / Alba</i>	8	12	8	8	9	12	7	10	7	10	–	7
<i>Nymphaea nouchalli</i>	–	–	–	–	–	–	6	6	–	–	–	–
<i>Chenopodium album</i>	6	–	11	10	6	10	6	6	6	–	10	–
<i>Spinacia oleracea</i>	–	6	–	–	–	6	6	6	9	6	–	–
<i>Alternanthera sessilis</i>	16	6	20	21	–	6	–	6	7	6	6	13
<i>Diplazium esculcutum</i>	7	–	–	10	6	–	–	10	–	6	–	–
<i>Portulaca grandifolia</i>	11	9	–	9	–	–	6	–	–	9	8	9
<i>Gilinus oppositifolius</i>	6	8	6	7	–	–	–	–	–	–	6	6
<i>Xanthium indicum</i>	20	28	20	15	15	13	15	14	12	18	17	20
<i>Benincasa hispida</i>	–	6	–	–	–	–	–	6	–	–	–	–
<i>Amaranthus tricolor</i>	6	–	–	–	–	6	–	6	–	–	–	–
<i>Corchorus olitorius</i>	–	–	–	–	–	6	5	–	–	–	–	–
<i>Lagenaria siceraria</i>	6	–	–	7	6	6	6	10	6	8	–	–
<i>Kanamycin</i>	30	33	30	30	28	35	32	37	30	32	30	33

–: No inhibition zone (resistant). *E. coli*: *Escherichia coli*; *P. aeruginosa*: *Pseudomonas aeruginosa*; *S. typhi*: *Salmonella typhi*; *S. paratyphi*: *Salmonella paratyphi*; *S. dysenteriae*: *Shigella dysenteriae*; *S. boydii*: *Shigella boydii*; *V. mimicus*: *Vibrio mimicus*; *V. parahemolyticus*: *Vibrio parahemolyticus*; *B. subtilis*: *Bacillus subtilis*; *B. megaterium*: *Bacillus megaterium*; *S. aureus*: *Staphylococcus aureus*; *S. lutea*: *Sarcina lutea*.

Table 2

Cytotoxic activity of methanolic extract of fourteen different edible vegetables from Bangladesh on brine shrimp nauplii.

Tested material	Concentration tested ($\mu\text{g/mL}$)	Probit	LC ₅₀ ($\mu\text{g/mL}$)	LC ₉₀ ($\mu\text{g/mL}$)	95% confidence interval
<i>Alternanthera sessilis</i>	5, 10, 20	4.48, 4.75, 5.00	19.825	601.458	5.418–76.339
<i>Amaranthus tricolor</i>	5, 10, 20, 50, 100	4.16, 4.48, 5.00, 5.25, 5.52	28.319	467.834	12.003–66.811
<i>Benincasa hispida</i>	5, 10, 20, 50	4.16, 4.48, 4.75, 5.00	45.187	1522.153	13.563–150.541
<i>Chenopodium album</i>	5, 10, 20	4.48, 5.00, 5.52	10.000	55.080	5.096–19.623
<i>Corchorus olitorius</i>	5, 10, 20, 50	4.16, 4.48, 5.00, 5.25	26.254	352.338	10.797–63.839
<i>Diplazium esculentum</i>	5, 10, 20, 50, 100	4.48, 5.00, 5.00, 5.25, 5.52	18.561	1240.301	5.129–67.160
<i>Enhydra fluctuans</i>	5, 10, 20, 50	4.48, 5.00, 5.52, 5.84	10.522	89.646	5.055–21.900
<i>Glinus oppositifolius</i>	5, 10, 20, 50	4.16, 4.48, 4.48, 5.52	27.650	279.769	12.525–61.039
<i>Ipomoea aquatica / Alba</i>	5, 10, 20, 50	3.72, 4.16, 4.75, 5.25	32.668	215.645	15.498–68.857
<i>Lagenaria siceraria</i>	5, 10, 20, 50	4.16, 4.48, 5.00, 5.00	35.835	962.421	11.623–110.481
<i>Nymphaea nouchalli</i>	5, 10, 20, 50, 100	4.48, 5.00, 5.00, 5.00, 5.52	21.864	2550.281	5.095–93.818
<i>Portulaca grandifolia</i>	5, 10, 20, 50	4.48, 5.00, 5.25, 5.84	11.647	109.832	5.405–25.100
<i>Spinacia oleracea</i>	5, 10, 20, 50	4.16, 4.48, 4.48, 5.00	60.323	2693.980	16.441–221.325
<i>Xanthium indicum</i>	5, 10, 20, 50	4.75, 5.00, 5.52, 6.28	8.447	55.720	4.008–17.800

Table 3

Cytotoxicity of Vincristine sulphate on brine shrimp nauplii.

Concentration ($\mu\text{g/mL}$)	Probit
0.000	–
0.060	0
0.125	3.77
0.250	4.23
0.500	4.56
1.000	4.85
5.000	6.23
10.000	8.09

4. Discussion

Plants are important source of potentially useful structures for the development of new chemotherapeutic agents. The first step towards this goal is the *in vitro* antibacterial activity assay[54]. It seems very likely, therefore, that the antibacterial compounds extracted from *Xanthium indicum* may inhibit bacteria by a different mechanism than that of currently used antibiotics and may have therapeutic value as an antibacterial agent against multi-drug resistant bacterial strains[55]. Because infections caused by *Pseudomonas aeruginosa*, especially those with multi-drug resistance, are among the most difficult to treat with conventional antibiotics. From the results obtained, it appears that the antibacterial action of the extracts is more pronounced on gram-negative than on gram-positive bacteria in most cases or is even equal. These findings do not correlate with the observations of previous screenings of medicinal plants for antimicrobial activity, where most of the active plant extracts showed activity against Gram-positive strains only[56–59]. Because Gram-negative organisms were reported to be less susceptible to the action of antibacterials, since they possess an outer membrane surrounding the cell wall, which restricts diffusion of hydrophobic compounds through its lipopolysaccharide covering[60–67]. Besides the difference in sensitivity might be ascribed to the difference in morphological constitutions between Gram-

positive and Gram-negative organisms. Many plant species present inhibition zones of differing diameters; however, size difference of the inhibition zone depends primarily upon many factors for *e.g.* diffusion capacity of substances (present in the extracts) in the medium, antimicrobial activity of diffused substances, growth and metabolic activity of microorganisms in the medium[68]. However demonstration of antimicrobial activity against both Gram-positive and Gram-negative bacteria may be indicative of the presence of broad spectrum antibiotic compounds[69,70]. But the mechanisms behind the antibacterial activity are complex to understand and could be attributed to either inhibiting the cell division or to damaging the cell walls of bacteria; which however requires to be investigated in detail[71].

It was found that different parts of various plant extracts in our investigation were studied earlier and exhibited antimicrobial activity to some different extent. For instance, it was found that while *Alternanthera sessilis* whole plant was moderately active in *Escherichia coli* in our studies, but in an earlier report [3] the methanol extracts of leaves were more potential to *Escherichia coli* with no other activity to other organisms. Again *Corchorus olitorius* showed no activity except *Shigella boydi* and *Vibrio mimicus* while the methanol extracts of seeds were found to have potent antibacterial activity against many gram positive and gram negative bacteria. Likewise *Nymphaea nouchalli* showed no activity except *Vibrio mimicus* and *Vibrio parahemolyticus* whereas flowers and leaves of *Nymphea nouchalli* were found considerable antibacterial activity against *Pseudomonas aeruginosa*, *Bacillus cereus*, *Staphylococcus aureus* and *Escherichia coli*[37,72]. *Enhydra fluctuans* showed little activity in our studies which is in accordance with a previous study[25]. while significant antibacterial activities were found in another report[26]. Furthermore, *Portulaca oleracea* Linn showed little or no activity against most of the tested bacteria while it was found high antimicrobial activities against *Helicobacter pylori*, *Staphylococcus epidermidis*, *Staphylococcus aureus*, *Escherichia coli* and *Streptococcus mutans* in another studies[73].

The brine shrimp lethality bioassay (BSLB) has been used extensively in the primary screening of the crude extracts as well as the isolated compounds to evaluate

the toxicity towards brine shrimps, which could also provide an indication of possible cytotoxic properties of the test materials[52]. Because BLSB is the simple method useful for screening large number of extracts in the drug discovery process. The method allows the use of smaller quantity of the extracts and permits larger number of samples and dilutions within shorter time than using the original test vials[74]. Furthermore it has been established that the cytotoxic compounds generally exhibit significant activity in the BSLB, and this assay can be recommended as a guide for the detection of antitumour and pesticidal compounds because of its simplicity and low cost[75]. This bioassay has also a good correlation with the human solid tumour cell lines. The inhibitory effect of the extract might be due to the toxic compounds present in the active fraction that possess ovicidal and larvicidal properties. The metabolites either affected the embryonic development or slay the eggs. Therefore the cytotoxic effects of the plant extracts enunciate that it can be selected for further cell line assay because there is a correlation between cytotoxicity and activity against the brine shrimp nauplii using extracts[76]. Hence the present study supports BLSB as a reliable method for the assessment of bioactivity of Bangladeshi vegetables and lends support for their use in pharmacology. It was found that no investigation has been carried out for cytotoxic activities till now among our studied fourteen vegetables except *Diaplasium esculentum* and *Ipomoea aquatica*. In the present study, *Ipomoea aquatica* showed strong cytotoxic activity which is in accordance with the result obtained earlier[32]. On the other hand *Diaplasium esculentum* also displayed potent cytotoxic potential in our observations but in another report it was found to have no cytotoxic activity[77].

The *in vitro* cytotoxicity displayed by the plant extracts tested is an initial indicator of *in vivo* antitumour activity. However since a wide range of phytochemicals are capable of exhibiting nonspecific cytotoxicity, plant extracts with significant cytotoxic activity should be further assayed using animal models to confirm antitumour activity, and/or a battery of various cell lines to detect specific cytotoxicity. This step is necessary to eliminate cytotoxic compounds with little value for further investigation as anticancer agents[4].

The results obtained in the present study demonstrated that a number of Bangladeshi vegetables have promising antibacterial activity against tested microorganisms which can be used in the healing of various infectious diseases caused by resistant microorganisms. Especially there is a likelihood to obtain better therapeutic agent against microbial diseases from some of the studied plants such as *Xanthium indicum*, *Enhydra fluctuans*, *Ipomoea aquatica*, *Chenopodium album*, *Alternanthera sessilis*, *Portulaca grandifolia* and *Lagenaria siceraria*. Besides all the plant extracts showed moderate to potent cytotoxic activity as well which could serve for further ethnobotanical and phytochemical research to find the possible relationship between brine shrimp lethality and plant bioactivity. It is therefore recommended to explore and be studied in detail the nature and extent of active principles in each plant extract.

Conflict of interest statement

We declare that we have no conflict of interest.

Acknowledgements

The authors are thankful to the authority of Southeast University, Dhaka, Bangladesh for providing financial support in conducting this research work.

Comments

Background

In recent years, multiple drug resistance in human pathogenic microorganism has been developed due to indiscriminate use of commercial antimicrobial drugs commonly used in the treatment of various diseases. Therefore, it is of great interest to carry out a screening of these plants in order to unfold the dimension of use as antimicrobial and cytotoxic activity.

Research frontiers

This study is on the edible vegetables of Bangladesh to explore antimicrobial and cytotoxic effects. These types of studies on plants are available but studies on edible vegetables of a particular ethnicity are few.

Related reports

From the results obtained, it appears that the antibacterial action of the extracts is more pronounced on Gram-negative than on Gram-positive bacteria in most cases or is even equal. These findings do not correlate with the observations of previous screenings of medicinal plants for antimicrobial activity, where most of the active plant extracts showed activity against Gram-positive strains only [56–59].

Innovations and breakthroughs

Data regarding the antibacterial and cytotoxic activity of edible vegetables is few. This study revealed that edible vegetables are effective against bacteria particularly gram negative bacteria.

Applications

The results from this study can be used for isolation and characterization of new antibacterial and cytotoxic agents. As the vegetables are edible and consumed by an ethnic group for a long period of time without any noticeable unwanted effects, the isolated compounds may be safe as well.

Peer review

This is a good paper where authors evaluated the antibacterial activity against 12 different gram positive and gram negative bacteria and brine shrimp lethality bioassay of 14 edible vegetables from Bangladesh. The results are interesting where it has been found that most of the vegetables are effective against gram negative bacteria rather than gram positive bacteria.

References

- [1] Shyamala BN, Gupta S, Jyothi Lakshmi A, Prakash J. Leafy vegetable extracts–antioxidant activity and effect on storage stability of heated oils. *Innovative Food Sci Emerg Technol* 2005; **6**(2): 239–245.
- [2] Lin SC, Lin YH, SJo S, Lin CC. Hepatoprotective effects of Taiwan folk medicine: *Alternanthera sessilis* on liver damage induced by various hepatotoxins. *Phytother Res* 2006; **8**(7): 391–398.
- [3] Jalalpure SS, Agrawal N, Patil MB, Chimkode R, Ashish T. Antimicrobial and wound healing activities of leaves of *Alternanthera sessilis* Linn. *Int J Green Pharm* 2008; **2**(3): 141–144.
- [4] Ali AM, Mackeen MM, Saleh H, Ei-Sharkawy, Hamid JA, Ismail NH, et al. Antiviral and cytotoxic activities of some plants used in Malaysian indigenous medicine. *Pertanika J Trop Agric Sci* 1996; **19**(2/3): 129–136.
- [5] Jayaprakasam B, Zhang Y, Nair MG. Tumor cell proliferation and cyclooxygenase enzyme inhibitory compounds in *Amaranthus tricolor*. *J Agric Food Chem* 2004; **52** (23): 6939–6943.
- [6] Grover JK, Adiga G, Vats V, Rathi SS. Extracts of *Benincasa hispida* prevent development of experimental ulcers. *J Ethnopharm* 2001; **78**: 159–164.
- [7] Shetty BV, Arjuman A, Jorapur A, Samanth R, Yadav SK, Valliammai N, et al. Effect of extract of *Benincasa hispida* on oxidative stress. In rats with indomethacin induced gastric ulcers. *Indian J Physiol Pharmacol* 2008; **52**(2): 178–182.
- [8] Anil K, Ramu P. Effect of methanolic extract of *Benincasa hispida* against histamine and acetylcholine induced bronchospasm in guinea pigs. *Indian J Pharmacol* 2002; **34**(5): 365–366.
- [9] Huang HY, Huang JJ, Tso TK, Tsai YC, Chang CK. Antioxidant and angiotension–converting enzyme inhibition capacities of various parts of *Benincasa hispida* (wax gourd). *Mol Nutr Food Res* 2004; **48**(3): 230–233.
- [10] Battu GR, Mamidipalli SN, Parimi R, Viriyala RK, Patchula RP, Mood LR. Hypoglycemic and anti–hyperglycemic effect of alcoholic extract of *Benincasa hispida* in Normal and in alloxan induced diabetic rats. *Pharmacogn Mag* 2007; **3**(10): 101–105.
- [11] Lim SJ. Effects of fractions of *Benincasa hispida* on antioxidative status in streptozotocin induced diabetic rats. *Korean J Nutrition* 2007; **40**(4): 295–302.
- [12] Lee KH, Choi HR, Kim CH. Anti–angiogenic effect of the seed extract of *Benincasa hispida* Cogniaux. *J Ethnopharm* 2005; **97**(3): 509–513.
- [13] Kumar A, Vimalavathini R. Possible anorectic effect of methanol extract of *Benincasa hispida* (Thunb). Cogn, fruit. *Indian J Pharm* 2004; **36**(6): 348–350.
- [14] Lim SJ. Effects of fractions of *Benincasa hispida* on plasma levels of glucose and lipid in streptozotocin induced diabetic rats. *Korean J Nutr* 2005; **38**(10): 801–806.
- [15] Pal DK, Mandal M, Senthilkumar GP, Padhiari A. Antibacterial activity of *Cuscuta reflexa* stem and *Corchorus olitorius* seed. *Fitoterapia* 2006; **77**(7–8): 589–591.
- [16] Zakaria ZA, Somchit MN, Zaiton H, Mat Jais AM, Sulaiman MR, Farah WO, et al. The *in vitro* antibacterial activity of *Corchorus olitorius* extracts. *Int J Pharm* 2006; **2**(2): 213–215.
- [17] Gupta M, Mazumder UK, Pal DK, Bhattacharya S. Anti–steroidogenic activity of methanolic extract of *Cuscuta reflexa* roxb. stem and *Corchorus olitorius* Linn. seed in mouse ovary. *Indian J Exp Biol* 2003; **41**(6): 641–644.
- [18] Sathiyamoorthy P, Lugasi–Evgi H, Schlesinger P, Kedar I, Gopas J, Pollack Y, et al. Screening for cytotoxic and antimalarial activities in desert plants of the Negev and Bedouinmarket plant products. *Pharm Biol* 1999; **37**(3): 188–195.
- [19] Yoshikawa M, Shimada H, Saka M, Yoshizumi S, Yamahara J, Matsuda H. *Medicinal foodstuffs*. V. Moroheiya. (1): Absolute stereostructures of corchoionosides A, B, and C, histamine release inhibitors from the leaves of Vietnamese *Corchorus olitorius* L. (Tiliaceae). *Chem Pharm Bull* 1997; **45**(3): 464–469.
- [20] Azuma K, Nakayama M, Koshioka M, Ippoushi K, Yamaguchi Y, Kohata K, et al. Phenolic antioxidants from the leaves of *Corchorus olitorius* L. *J Agric Food Chem* 1999; **47**(10): 3963–3966.
- [21] Wong SP, Leong LP, Koh JHW. Antioxidant activities of aqueous extracts of selected plants. *Food Chem* 2006; **99**(4): 775–783.
- [22] Nanasombat S, Teckchuen N. Antimicrobial, antioxidant and anticancer activities of Thai local vegetables. *J Med Plants Res* 2009; **3**(5): 443–449.
- [23] Sannigrahi S, Mazuder UK, Pal DK, Parida S, Jain S. Antioxidant potential of crude extract and different fractions of *Enhydra fluctuans* Lour. *Iranian J Pharm Res* 2010; **9**(1): 75–82.
- [24] Rahman MT, Begum N, Alimuzzaman M, Khan MOF. Analgesic activity of *Enhydra fluctuans*. *Fitoterapia* 2002; **73**(7–8): 707–709.
- [25] Bhakta J, Majumdar P, Munekage Y. Antimicrobial efficacies of methanol extract of *Asteracantha longifolia*, *Ipomoea aquatica* and *Enhydra fluctuans* against *Escherichia coli*, *Pseudomonas aeruginosa*, *Staphylococcus aureus* and *Micrococcus luteus*. *Int J Alter Med* 2009; **7**(2).
- [26] Abu Ahmed AM, Rahman MS, Anwar MN. Antibacterial activities of extracts obtained from five important medicinal plants of Bangladesh. *Europ J Sci Res* 2005; **11**(3): 444–448.
- [27] Inngjerdingen KT, Debes SC, Inngjerdingen M, Hokputsa S, Harding SE, Rolstad B, et al. Bioactive pectic polysaccharides from *Glinus oppositifolius* (L.) Aug. DC., a Malian medicinal plant, isolation and partial characterization. *J Ethnopharmacol* 2005; **101**(1–3): 204–214.
- [28] Traore F, Faure R, Ollivier E, Gasquet M, Azas N, Debrauwer L, et al. Structure and antiprotozoal activity of triterpenoid saponins from *Glinus oppositifolius*. *Planta Med* 2000; **66**(4): 368–371.
- [29] AsokKumar K, UmaMaheswari M, Sivashanmugam AT, SubhadraDevi V, Subhashini N, Ravi TK. Free radical scavenging and antioxidant activities of *Glinus oppositifolius* (carpet weed) using different *in vitro* assay systems. *Pharm Biol* 2009; **47**(6): 474–482.
- [30] Sugandhika T, Malalavidhane, Wickramasinghe SMDN, Jansz ER. Oral hypoglycaemic activity of *Ipomoea aquatica*. *J Ethnopharmacol* 2000; **72**(1–2): 293–298.
- [31] Malalavidhane TS, Wickramasinghe SMDN, Perera MSA, Jansz ER. Oral hypoglycaemic activity of *Ipomoea aquatica* in streptozotocin–induced, diabetic wistar rats and type II diabetics. *Phytother Res* 2003; **17**(9): 1098–1100.
- [32] Prasad KN, Divakar S, Shivamurthy GR, Aradhya SM. Isolation of a free radical–scavenging antioxidant from water spinach (*Ipomoea aquatica* Forsk). *J Sci Food Agric* 2005; **85**(9): 1461–1468.
- [33] Ghule BV, Ghante MH, Yeole PG, Saoji AN. Diuretic activity of *Lagenaria siceraria* fruit extracts in rats. *Indian J Pharmaceutical Sci* 2007; **69**(6) 817–819.
- [34] Mali VR, Bodhankar SL. Cardioprotective effect of *Lagenaria siceraria* (LS) fruit powder in isoprenaline–induced cardiotoxicity in rats. *Europ J Integ Med* 2010; **2**(3): 143–149.
- [35] Rajagopal K, Sasikala K, Ragavan B. Hypoglycemic and antihyperglycemic activity of *Nymphaea stellata* flowers in normal and alloxan diabetic rats. *Pharm Biol* 2008; **46**: 654–659.
- [36] Murakami A, Ali AM, Mat–Salleh K, Koshimizu K, Ohigashi H. Screening for the *in vitro* anti–tumor–promoting activities of edible plants from Malaysia. *Biosci Biotech Biochem* 2000; **64**: 9–16.
- [37] Mohan Maruga Raja MK, Dhanabal SP, Patil MJ. Pharmacognostical investigation and antibacterial activity of *Nymphaea stellata* Willd. leaves. *Hamdard Med* 2008; **51**: 139–145.
- [38] Chan K, Islam MW, Kamil M, Radhakrishnan R, Zakaria MNM,

- Habibullah M, et al. The analgesic and anti-inflammatory effects of *Portulaca oleracea* L. subsp. *sativa* (Haw.) Celak. *J Ethnopharmacol* 2000; **73**(3): 445–451.
- [39] Rashed AN, Afifi FU, Disi AM. Simple evaluation of the wound healing activity of a crude extract of *Portulaca oleracea* L. (growing in Jordan) in *Mus musculus* JVI-1. *J Ethnopharmacol* 2003; **88**(2–3): 131–136.
- [40] Radhakrishnan R, Zakaria MNM, Islam MW, Chen HB, Kamil M, Chan K, et al. Neuropharmacological actions of *Portulaca oleracea* L. v. *sativa* (Hawk). *J Ethnopharmacol* 2001; **76**(2):171–176.
- [41] Bhatia AL, Jain M. *Spinacia oleracea* L. protects against gamma radiations: a study on glutathione and lipid peroxidation in mouse liver. *Phytomedicine* 2004; **11**(7–8): 607–615.
- [42] Rudolf E, Gernot K, Karl LP, Klaus KU. Isolation and characterization of structurally novel antimutagenic flavonoids from spinach (*Spinacia oleracea*). *J Agric Food Chem* 2001; **49** (6): 2767–2773.
- [43] Talakal TS, Dwivedi SK, Sharma SR. *In vitro* and *in vivo* antitrypanosomal activity of *Xanthium strumarium* leaves. *J Ethnopharm* 1995; **49**(3):141–145.
- [44] Haque N, Chowdhury SAR, Nutan MTH, Rahman GMS, Rahman KM, Rashid MA. Evaluation of antitumor activity of some medicinal plants of Bangladesh by potato disk bioassay. *Fitoterapia* 2000; **71**(5):547–552.
- [45] Namsaa ND, Hui T, Mandal M, Kalita P, Das AK. An ethnobotanical study of traditional anti-inflammatory plants used by the Lohit community of Arunachal Pradesh, India. *J Ethnopharmacol* 2009; **125**: 234–245.
- [46] Evans WC. *Trease and Evan's pharmacognosy*. 14th ed. UK: W.B. Saunders Company Limited; 1997, p. 3.
- [47] Mohanasundari C, Natarajan D, Srinivasan K, Umamaheswari S, Ramachandran A. Antibacterial properties of *Passiflora foetida* L. – a common exotic medicinal plant. *Afric J Biotech* 2007; **6**(23): 2650–2653.
- [48] Bauer AN, Kirby WMM, Sherris JC, Truck M. Antibiotic susceptibility testing by single disk method. *Amer J Clin Pathol* 1996; **45**: 493–496.
- [49] Kavanagh F, Grinnan E, Allanson E, Tunin D. *Dihydrostreptomycin produced by direct fermentation*. Antibiotics manufacturing and development division. Indianapolis, Indiana: Eli Lilly and Company; 1960, p. 160–162.
- [50] Bartner A, Pfeiffer KP, Batner, H. Applicability of disc diffusion methods required by the pharmacopoeias for testing antibacterial activity of natural compounds. *Pharmaz* 1994; **49**: 512–516.
- [51] Lieberman MA. Brine shrimp bioassay for measuring toxicity and remediation of chemicals. *J Chem Ed* 1999; **76**: 1689–1691.
- [52] Meyer BN, Ferrigni NR, Putnam JE, Jacobsen JB, Nicholsand DE, McLaughlin JL. Brine shrimp; a convenient general bioassay for active plant constituents. *Plant Med* 1982; **45**: 31–34.
- [53] Finney DJ. *Probit analysis*. 3rd ed. Cambridge: Cambridge University Press; 1971.
- [54] Tona L, Kambu K, Ngimbi N, Cimanga K, Vlietinck AJ. Antiamoebic and phytochemical screening of some Congolese medicinal plants. *J Ethnopharmacol* 1998; **61**: 57–65.
- [55] Abu-Shanab B, Adwan G, Abu-Safiya D, Jarrar N, Adwan K. Antibacterial activities of some plant extracts utilized in popular medicine in Palestine. *Turk J Biol* 2004; **28**: 99–102.
- [56] Kelmanson JE, Jager AK, Van Staden J. Zulu medicinal plants with antibacterial activity. *J Ethnopharm* 2000; **69**: 241–246.
- [57] Ali NAA, Julich WD, Kusnick C, Lindequist U. Screening of Yemeni medicinal plants for antibacterial and cytotoxic activities. *J Ethnopharm* 2001; **74**: 173–179.
- [58] Moshi MJ, Mbwambo ZH. Some pharmacological properties of extracts of *Terminalia sericea* roots. *J Ethnopharm* 2005; **97**:43–47.
- [59] Parekh J, Chanda S. *In vitro* antimicrobial activities of extract of *Launaea procumbens* Roxb. (Labiatae), *Vitis vinifera* (Vitaceae) and *Cyperus rotundus* (Cyperaceae). *Afr J Biomed Res* 2006; **9**: 89–93.
- [60] Skocibusic M, Bezic N, Dunkic V. Phytochemical composition and antimicrobial activities of the essential oils from *Satureja subspicata* Vis. growing in Croatia. *Food Chem* 2006; **96**: 20–28.
- [61] Anitha VT, Antonisamy JM, Jeeva S. Anti-bacterial studies on *Hemigraphis colorata* (Blume) H.G. Hallier and *Elephantopus scaber* L. *Asian Pac J Trop Med* 2012; **5**(9): 52–57.
- [62] SO Oyedemi, AJ Afolayan. Antibacterial and antioxidant activities of hydroalcoholic stem bark extract of *Schotia latifolia* Jacq. *Asian Pac J Trop Med* 2011; **4**(12): 952–958.
- [63] Sukumaran S, Kiruba S, Mahesh M, Nisha SR, Miller PZ, Ben CP, et al. Phytochemical constituents and antibacterial efficacy of the flowers of *Peltophorum pterocarpum* (DC.) Baker ex Heyne. *Asian Pac J Trop Med* 2011; **4**(9): 735–738.
- [64] Kannathasan K, Senthilkumar A, Venkatesalu V. *In vitro* antibacterial potential of some *Vitex* species against human pathogenic bacteria. *Asian Pac J Trop Med* 2011; **4**(8): 645–648.
- [65] Raja RDA, Jeeva S, Prakash JW, Antonisamy JM, Irudayaraj V. Antibacterial activity of selected ethnomedicinal plants from South India. *Asian Pac J Trop Med* 2011; **4**(5): 375–378.
- [66] Moussa A, Nouredine D, Abdelmelek M, Saad A. Antibacterial activity of various honey types of Algeria against pathogenic Gram-negative bacilli: *Escherichia coli* and *Pseudomonas aeruginosa*. *Asian Pac J Trop Dis* 2012; **2**(3): 211–214.
- [67] Gokulakrishnan R, Ravikumar S, Anandha Raj J. *In vitro* antibacterial potential of metal oxide nanoparticles against antibiotic resistant bacterial pathogens. *Asian Pac J Trop Dis* 2012; **2**(5): 411–413.
- [68] Bandeira MF, Teixeira MF, Abinader CD, Parente RC, Lima PS. Avaliação *in vitro* da sensibilidade da *Candida albicans* ao hidróxido de cálcio associado ao óleo de copaíba. *Rev Dentística* 2006; **6**:12–22.
- [69] Cichewicz RH, Thorpe PA. The antimicrobial properties of Chile peppers (*Capsicum species*) and their uses in Mayan medicine. *J Ethnopharmacol* 1996; **52**(2): 61–70.
- [70] Srinivasan D, Perumalsamy LP, Nathan S, Sures T. Antimicrobial activity of certain Indian medicinal plants used in folkloric medicine. *J Ethnopharmacol* 2001; **49**: 217–222.
- [71] Manikandan S, Sheela DR, Srikumar R, Thangaraj R, Ayyappan R, Jegadeesh R, et al. *In vitro* antibacterial activity of aqueous and ethanolic extracts of *Acorus calamus*. *Int J Applied Biol Pharm Tech* 2010; **1**(3): 1072–1075.
- [72] Vasu K, Singaracharya MA. Antimicrobial activity of certain aquatic angiosperms against some pathogenic bacteria. *Asian J Microbiol Biotech Envir Sci* 2008; **10**: 609–613.
- [73] Cho YJ, Ju IS, Kwon OJ, Chun SS, An BJ, Kim JH. Biological and antimicrobial activity of *Portulaca oleracea*. *J Korean Society App Biol Chem* 2008; **51**(1): 49–54.
- [74] Sam TW. Toxicity testing using the brine shrimp: *Artemia salina*. In: Colegate SM, Molyneux RJ (eds). *Bioactive natural products detection, isolation, and structural determination*. Boca Raton; CRC press; 1993, p. 442–456.
- [75] Mazid MA, Datta BK, Nahar L, Sarker SD. Assessment of antibacterial activity and brine shrimp toxicity of two *Polygonum* species. *Ars Pharm* 2008; **49**(2): 127–134.
- [76] Manilal A, Sujith S, Seghal KG, Selvin J, Shakir C. Cytotoxic potentials of red alga, *laurencia brandenii* collected from the Indian coast. *Global J Pharmacol* 2009; **3**(2): 90–94.
- [77] Rahmat A, Kumar V, Fong LM, Endrini S, Abdullah SH. Determination of total antioxidant activity in three types of local vegetables shoots and the cytotoxic effect of their ethanolic extracts against different cancer cell lines. *Asia Pacific J Clin Nutr* 2003; **12**(3): 292–295.