



Published in final edited form as:

*J Biomol NMR*. 2013 March ; 55(3): 249–256. doi:10.1007/s10858-013-9706-1.

## Fast hydrogen exchange affects $^{15}\text{N}$ relaxation measurements in intrinsically disordered proteins

Seho Kim<sup>1</sup>, Kuen-Phon Wu<sup>1</sup>, and Jean Baum<sup>1,2,\*</sup>

<sup>1</sup>Department of Chemistry and Chemical Biology, Rutgers University, Piscataway, New Jersey, 08854

<sup>2</sup>BioMaPS Institute for Quantitative Biology, Rutgers University, Piscataway, New Jersey, 08854

### Abstract

Unprotected amide protons can undergo fast hydrogen exchange (HX) with protons from the solvent. Generally, NMR experiments using the out-and-back coherence transfer with amide proton detection are affected by fast HX and result in reduced signal intensity. When one of these experiments,  $^1\text{H}$ - $^{15}\text{N}$  HSQC, is used to measure the  $^{15}\text{N}$  transverse relaxation rate ( $R_2$ ), the measured  $R_2$  rate is convoluted with the HX rate ( $k_{\text{HX}}$ ) and has higher apparent  $R_2$  values. Since the  $^{15}\text{N}$   $R_2$  measurement is important for analyzing protein backbone dynamics, the HX effect on the  $R_2$  measurement is investigated and described here by multi-exponential signal decay. We demonstrate these effects by performing  $^{15}\text{N}$   $R_2^{\text{CPMG}}$  experiments on  $\alpha$ -synuclein, an intrinsically disordered protein (IDP), in which the amide protons are exposed to solvent. We show that the HX effect on  $R_2^{\text{CPMG}}$  can be extracted by the derived equation. In conclusion, the HX effect may be pulse sequence specific and results from various sources including the J coupling evolution, the change of steady state water proton magnetization, and the  $\text{D}_2\text{O}$  content in the sample. To avoid the HX effect on the analysis of relaxation data of unprotected amides, it is suggested that NMR experimental conditions insensitive to the HX should be considered or that intrinsic  $R_2^{\text{CPMG}}$  values be obtained by methods described herein.

### Keywords

Hydrogen exchange; intrinsically disordered protein;  $^{15}\text{N}$   $R_2$  relaxation

### Introduction

$^{15}\text{N}$  backbone transverse relaxation experiments ( $R_2$ ) using the CPMG (Carr–Purcell–Meiboom–Gill) pulse train ( $^{15}\text{N}$   $R_2^{\text{CPMG}}$ ) have been used to investigate dynamics of backbone amides in globular proteins (Farrow et al. 1994; Kay et al. 1989; Mandel et al. 1995) and have also been applied to study NMR dynamics of intrinsically disordered proteins (IDPs) (Buevich and Baum 1999; Buevich et al. 2001; Eliezer 2009; Mittag and Forman-Kay 2007; Wright and Dyson 2009).  $^{15}\text{N}$   $R_2^{\text{CPMG}}$  rates have been important indicators for protein backbone dynamics and have provided information about the structural mobility on the ps to ns timescale and about conformational exchange processes on the  $\mu\text{s}$  to ms timescale (Mittermaier and Kay 2009; Palmer III 2004; Wang and Palmer 2003). When backbone amide protons in proteins are not protected from the solvent water, those protons undergo fast hydrogen exchange (HX). Although HX rates of amide protons and  $R_2^{\text{CPMG}}$  rates of amide nitrogens are measurements of different events in protein dynamics, fast HX

\*Corresponding author: 610 Taylor Road, Piscataway, NJ, 08854, jean.baum@rutgers.edu, Phone: 1-732-445-5666, Fax: 1-732-445-5312.

may affect the measurement of  $^{15}\text{N}$   $R_2^{\text{CPMG}}$  rates. Since  $^{15}\text{N}$   $R_2^{\text{CPMG}}$  rates are measured indirectly with  $^1\text{H}$ - $^{15}\text{N}$  HSQC using the out-and-back coherence transfer with a prolonged relaxation time in the pulse sequence, measured rates can be modulated due to the signal intensity of the HSQC vulnerable to fast HX. Some NMR experiments have been designed to measure fast HX rates indirectly during the  $^{15}\text{N}$  time evolution or relaxation. In the  $R_{1zz}$  relaxation experiment of two spin order ( $N_zH_z$ ) (Skrynnikov and Ernst 1999), the HX effect destroys the  $N_zH_z$  spin operator and this results in the combined HX and  $R_{1zz}$  relaxation rate. In the  $^{15}\text{N}$   $R_2$  relaxation experiment of the HCN-type coherence transfer (Kateb et al. 2007), the fast HX effect that dephases the  $^{15}\text{N}$  spin operator is correlated with the  $^1J_{\text{NH}}$  coupling evolution. The presence or absence of  $^1\text{H}$  decoupling during the relaxation time results in a significant difference in the HX effect on  $^{15}\text{N}$   $R_2$  relaxation rates. In the  $^{15}\text{N}^{\text{H/D}}$ -SOLEXSY experiment using the (HACACO)NH pulse sequence, H/D exchange rates were measured during the  $^{15}\text{N}$   $t_1$  evolution in  $\text{H}_2\text{O}/\text{D}_2\text{O}$  (50%/50%) (Chevelkov et al. 2010). Based on these experiments, it is expected that the  $^{15}\text{N}$   $R_2^{\text{CPMG}}$  relaxation rate measured by the HSQC-type NMR experiments may be modulated by fast HX although the mechanism of the HX effect is different from previously reported mechanisms and includes the spin state of  $^{15}\text{N}$ , the J-coupling evolution, or the solvent environment.

Here we show that  $R_2^{\text{CPMG}}$  rates are modulated by fast HX rates in  $\alpha$ -synuclein ( $\alpha$ Syn), a protein associated with Parkinson's disease (Chiti and Dobson 2006; Goedert 2001). If  $^{15}\text{N}$   $R_2^{\text{CPMG}}$  rates are modulated by both conformational exchange and HX, it will be complicated to interpret the relaxation data without knowing the magnitude of the HX effect on the  $^{15}\text{N}$   $R_2^{\text{CPMG}}$  rates. In particular, when residues with large conformational exchange are located at the solvent accessible surface areas,  $^{15}\text{N}$   $R_2^{\text{CPMG}}$  rates may be a combination of both conformational exchange and fast HX rates (Bertini et al. 2002; Tugarinov and Kay 2003).  $\alpha$ Syn is an intrinsically disordered protein (IDP) (Eliezer et al. 2001; Uversky et al. 2000; Wu et al. 2008) and forms structurally disordered ensembles in solution which possess enhanced HX rates at neutral pH and high temperature (Croke et al. 2008). We demonstrate the relationship between fast HX and the  $^{15}\text{N}$   $R_2^{\text{CPMG}}$  rates at different pH states and show that the  $^{15}\text{N}$  relaxation rates that are affected by fast HX can be corrected quantitatively. The methods discussed here may be applied to other IDPs or partially folded proteins that are studied at neutral pH to distinguish the inherent  $R_2$  values from those associated with fast hydrogen exchange.

## Materials and methods

### Preparation of $\alpha$ Syn protein

The expression and purification of  $^{15}\text{N}$ -labeled  $\alpha$ Syn was done by previously described protocols (Wu et al. 2009). Lyophilized  $\alpha$ Syn was dissolved in phosphate buffer at pH 7.4 and 6.2 (10 mM phosphate, 137 mM NaCl, 2 mM KCl). Prior to applying NMR experiments, samples were filtered by a 100 kDa filter (Millipore) to remove invisible small aggregates. Final sample concentrations were adjusted to 250  $\mu\text{M}$  with 10% (v/v)  $\text{D}_2\text{O}$  for all NMR experiments.

### NMR data acquisition

The  $^{15}\text{N}$   $R_2^{\text{CPMG}}$  experiment was performed using a modified pulse sequence of gNhsqc from the Varian BioPack library; the main modification is an optional condition for the proton steady state (SS) pulse in the beginning of the pulse sequence (Fig. 2). The NMR data were collected twice at a proton frequency of 800 MHz and 15 °C without and with the SS pulse. Relaxation time delays for  $R_2^{\text{CPMG}}$  experiments were 0.01, 0.03, 0.05, 0.07, 0.09, 0.13, 0.17, 0.21, and 0.25 (s) at pH 7.4 and pH 6.2. The CPMG inter-pulse delay ( $2\tau$ ) was 1.25 ms and the recycle delay ( $d_1$ ) for each scan was 2 seconds. Spectral complex points

were 1024 and 256 in the  $t_2$  and  $t_1$  dimensions, respectively. Relaxation data were acquired by interleaving relaxation time delays and  $t_1$  increments. Data were processed and analyzed by NMRPipe (Delaglio et al. 1995). The relaxation data were fit by two different methods to measure the apparent  $R_2^{\text{CPMG}}$  values; (1) single exponential fit as described in other previous studies (Bertoncini et al. 2007; Bussell and Eliezer 2001) and (2) multi-exponential fit introduced in this paper (discussed in results) to remove the HX effect.

To measure hydrogen exchange rates ( $k_{\text{HX}}$ ), CLEANEX (Hwang et al. 1998) experiments including a fast  $^1\text{H}$ - $^{15}\text{N}$  HSQC detection scheme were performed by using the pulse sequence of gCLNfhsqc from the Varian BioPack library. The following mixing times were used; 0.01, 0.02, 0.03, 0.04, 0.05, 0.06, 0.07 (s) at pH 7.4 and 0.01, 0.02, 0.04, 0.06, 0.08, 0.1, 0.12 (s) at pH 6.2. The reference fast  $^1\text{H}$ - $^{15}\text{N}$  HSQC spectra were acquired to serve as  $V_0$  in the recommended fitting equation (Hwang et al. 1998). The recycle delay for each scan was 2 seconds and spectral complex points were 1024 and 256 in the  $t_2$  and  $t_1$  dimensions, respectively. Hydrogen exchange data were acquired by interleaving mixing times and  $t_1$  increments. Data were processed and analyzed by NMRPipe (Delaglio et al. 1995). The intensity build-ups by mixing times were fitted to obtain two unknown  $k_{\text{HX}}$  and  $R_{1A, \text{app}}$  (apparent  $R_1$  of amide protons) with a fixed  $R_{1B, \text{app}}$  (apparent  $R_1$  of water protons, 0.6 Hz).

## Results and Discussion

The  $^{15}\text{N}$   $R_2^{\text{CPMG}}$  relaxation data of  $\alpha\text{Syn}$  acquired at pH 7.4 and pH 6.2 (Fig. 1) were acquired by using the pulse sequence (Fig. 2) without the proton steady state pulse and the data were analyzed with single exponential fits as described in other previous studies (Bertoncini et al. 2007; Bussell and Eliezer 2001). The resulting  $^{15}\text{N}$   $R_2^{\text{CPMG}}$  values showed that values at pH 7.4 were elevated relative to those at pH 6.2 and also showed some sequence-specific differences; specifically the N-terminal residues have slightly larger values overall than the C-terminal residues (Fig. 1). Individual  $R_2^{\text{CPMG}}$  values showed non-uniform deviations along residues and averages of reported  $^{15}\text{N}$   $R_2^{\text{CPMG}}$  values were  $4.21 \pm 0.55$  Hz at pH 7.4 and  $3.54 \pm 0.46$  Hz at pH 6.2. The discrepancy in  $^{15}\text{N}$   $R_2^{\text{CPMG}}$  values between two pH's may arise from changes in dynamics of  $\alpha\text{Syn}$  or by changes in conformational exchange or folding states of the protein.

However, pH-dependent water hydrogen exchange (HX) rates of amide protons may be a primary cause for large differences in  $^{15}\text{N}$   $R_2^{\text{CPMG}}$  values since  $\alpha\text{Syn}$  is known as an IDP (Eliezer et al. 2001; Uversky et al. 2000; Weinreb et al. 1996; Wu et al. 2008) and measurement of HX rates show that the protein undergoes very fast HX on the order of 5 to 20 Hz (Croke et al. 2008) and that its amide protons are not protected from H/D hydrogen exchange. In Supporting Information, HSQC spectra of  $\alpha\text{Syn}$  acquired at various experimental conditions were similar to each other except for the residue H50 due to the pH titration of the imidazole. This supports that the intrinsic properties (including  $R_2^{\text{CPMG}}$  values) of  $\alpha\text{Syn}$  are similar to each other at pH 7.4 and pH 6.2. In this paper, we show that the pH-dependent differences in  $^{15}\text{N}$   $R_2^{\text{CPMG}}$  can be derived from the pH-dependent hydrogen exchange (HX) rates. Improvement of the correlation between  $^{15}\text{N}$   $R_2^{\text{CPMG}}$  values between pH 7.4 and pH 6.2 is demonstrated and  $^{15}\text{N}$   $R_2^{\text{CPMG}}$  values without HX effect are obtained.

To describe the HX effect on the backbone  $^{15}\text{N}$   $R_2^{\text{CPMG}}$  NMR experiment, the conventional H/D exchange (exchange between amide hydrogen and solvent deuterium) is considered as one of HX mechanisms because of the mixed solvent environment of 90%  $\text{H}_2\text{O}$  and 10%  $\text{D}_2\text{O}$ . For H/D exchange, it is assumed temporarily that the protein  $^{15}\text{N}$  spin is 100% locked in the  $N_x$  spin state during the  $^{15}\text{N}$  relaxation measurement period ( $t$ ) and H/H exchange

(exchange between amide hydrogen and solvent hydrogen) has no HX effect. The pulse sequence of the  $^{15}\text{N}$   $R_2^{\text{CPMG}}$  NMR experiment using refocused HSQC is outlined in Fig. 2. The out-and-back nature of HSQC requires the NH chemical integrity, which selects the protein-NH by forward refocused INEPT and detects the same protein-NH by reverse refocused INEPT. By H/D exchange, the in-phase  $N_x$  spin state ( $H_E N_x$ ) after forward refocused INEPT can lose the attached H population ( $H_E$ ) and the deuterium populated  $D_E N_x$  spin state is generated. Then, the  $D_E N_x$  state fails to produce the NMR signal through reverse refocused INEPT.

In H/D exchange, the loss of proton occupancy on the amide is directly proportional to the  $D_2O$  content in the solvent. When  $H_2O/D_2O$  fractions in the mixed solvent are  $Q/(1-Q)$ , populations of NH/ND are  $Q/(1-Q)$  in equilibrium. However, the initial NH population selected by the forward refocused INEPT is repopulated into NH and ND by multiplying  $Q$  and  $(1-Q)$  if HX is completed during the relaxation period. Since the reverse refocused INEPT selects only the NH population, the HSQC signal is subject to signal loss by a maximum of  $(1-Q)$ . By introducing the HX rate constant ( $k_{\text{HX}}$ ), the NMR signal intensity decay ( $I$ ) can be a multi-exponential in equation 1 in combination with the  $^{15}\text{N}$   $R_2^{\text{CPMG}}$  during the CPMG relaxation time ( $t$ ).

$$I(t) = I_0 \{ Q + (1-Q)e^{-k_{\text{HX}}t} \} e^{-R_2^{\text{CPMG}}t} \quad (1)$$

Equation 1 is valid when the pseudo-first order kinetics can be applied for HX at the slow exchange regime ( $k_{\text{HX}} < R_2^{\text{CPMG}}$ ). At prolonged relaxation times under  $k_{\text{HX}} > R_2^{\text{CPMG}}$ , HX reaches equilibrium and the backward exchange (from ND to NH) cannot be ignored in the time scale of the  $R_2^{\text{CPMG}}$  relaxation. This requires the Bloch-McConnell equation for chemical exchange (Hansen and Led 2003; McConnell 1958) which includes two different  $R_2^{\text{CPMG}}$  rates ( $R_2^{\text{NH}}$  for the protonated nitrogen and  $R_2^{\text{ND}}$  for the deuterated nitrogen) instead of a single  $R_2^{\text{CPMG}}$  in equation 1. At the extreme fast exchange ( $k_{\text{HX}} \gg R_2^{\text{NH}}$ ), the  $R_2^{\text{CPMG}}$  (from equation 1) appears as a population-weighted average of  $R_2^{\text{NH}}$  and  $R_2^{\text{ND}}$  and the relative error of  $R_2^{\text{CPMG}}$  (from equation 1) to  $R_2^{\text{NH}}$  (from Bloch-McConnell equation) is a maximum of 5.5 % in our calculation (see Supporting Information). Especially under our experimental conditions ( $k_{\text{HX}} / R_2^{\text{CPMG}}$  about 10 and  $Q = 0.9$ ), the relative error of  $R_2^{\text{CPMG}}$  is estimated to be less than 4.5 % (see Supporting Information). Since it is difficult to obtain  $R_2^{\text{NH}}$  and  $R_2^{\text{ND}}$  separately from a single data set of a multi-exponential signal decay, equation 1 can be used as a good approximation to interpret the HX effect resulting in the large difference of  $R_2^{\text{CPMG}}$  values as shown in Figure 1.

In the case of H/H exchange happening during the  $^{15}\text{N}$  relaxation time period ( $t$ ), the  $H_E N_x$  state needs to be elaborated by using single element spin operators;  $H_E N_x = (0.5H^\alpha + 0.5H^\beta)N_x$  to understand the role of the exchanged protons. By H/H exchange, an exchanging proton spin replaces  $H^\alpha$  and  $H^\beta$  with  $c_1H^\alpha + c_2H^\beta$  where  $c_1$  and  $c_2$  are probability coefficients from the water proton and their ensemble averages are 0.5 each. As shown in equation 2, coefficients ( $c_1$  and  $c_2$ ) of the exchanged H are not correlated with coefficients of 0.5 but the ensemble average generates the original  $H_E N_x$  spin state back. If the water proton magnetization has  $c_1 = 0.5 + \delta$  and  $c_2 = 0.5 - \delta$  where  $\delta$  indicates a negligible population difference, the exchanged  $H_2 N_x$  term does not contribute to the signal through the reverse refocused INEPT. Thus the result of H/H exchange during the  $^{15}\text{N}$   $R_2^{\text{CPMG}}$  relaxation has no effect on the  $N_x$  spin operator.

$$\begin{aligned}
 H_E N_x &= (0.5H^\alpha + 0.5H^\beta) N_x \\
 \xrightarrow{HX} & \{0.5(c_1 H^\alpha + c_2 H^\beta) + 0.5(c_1 H^\alpha + c_2 H^\beta)\} N_x \\
 &= \{(0.5H^\alpha + 0.5H^\beta) + \delta(0.5H^\alpha - 0.5H^\beta)\} N_x \quad (2) \\
 &= (H_E + H_z) N_x \\
 &\approx H_E N_x
 \end{aligned}$$

The H/H exchange needs an additional consideration which is not necessary for H/D exchange since protons are the directly observed nucleus and variable pulses and durations are applied. In the pulse sequence (Fig. 2), 180° pulses are incremented by 2n-1 with the increase of the  $^{15}\text{N}$   $R_2^{\text{CPMG}}$  relaxation delay ( $t$ ). Even though the experiment starts with many dummy scans to reach a steady state, we need to be careful about the relationship between the maintenance of the steady state and the increase in the  $R_2^{\text{CPMG}}$  relaxation delay. The different number of proton pulses and delays with an increase in the  $^{15}\text{N}$   $R_2^{\text{CPMG}}$  relaxation delay may perturb the steady state of the proton magnetization, particularly the bulk water proton magnetization. The pulse sequence (Fig. 2) is designed to minimize the saturation of the water proton magnetization but additional pulses and delays may result in increased saturation of the water proton magnetization. With the increment in the number of 180° proton pulses, the water proton magnetization may be partially and progressively saturated. This causes a different amount of HX between amide and water protons during the fixed recycle delay ( $d_1$ ) when  $d_1$  is shorter than at least 5 times the water  $T_1$  relaxation time which is a typical experimental condition. It means that the steady state of the water proton magnetization can be altered depending on the  $^{15}\text{N}$   $R_2^{\text{CPMG}}$  relaxation delay and it modulates the steady state of the amide proton magnetization through HX.

It will be difficult to quantify how much water proton magnetization is affected by proton pulses and delays during the  $^{15}\text{N}$   $R_2^{\text{CPMG}}$  relaxation delay due to radiation damping but this effect can be removed by applying a steady state (SS) pulse (or a purge pulse) right after the acquisition time ( $t_2$ ) and before the recycle delay ( $d_1$ ). The proton SS pulse destroys the effect of the water flip-back pulse designed for minimizing water saturation (Fig. 2) and randomizes (saturates and dephases) the water proton magnetization right after acquisition. Although the SS pulse attenuates the  $^1\text{H}$ - $^{15}\text{N}$  HSQC signal intensity through HX when the amide hydrogen is under fast HX, it will ensure that the same amount of H/H exchange happens during the  $d_1$  delay regardless of the change of pulses and delays in the pulse sequence. For this reason,  $^{15}\text{N}$   $R_2^{\text{CPMG}}$  experiments were run with and without the proton SS pulse to test HX effect. The implemented SS pulse in the  $^{15}\text{N}$   $R_2^{\text{CPMG}}$  experiment is gradient -  $^1\text{H}$  90° pulse - gradient in Fig. 2 but the other type of SS pulses such as the x-axis trim pulse followed by the y-axis trim pulse will be effective, as well.

When the in-phase  $N_x$  spin state is not locked completely by the CPMG pulses during the relaxation period, a minor population of anti-phase  $^{15}\text{N}$  magnetization is present by the  $^1\text{J}_{\text{NH}}$  coupling evolution. The HX effect on the anti-phase  $^{15}\text{N}$  magnetization ( $2N_y H_z$ ) will be different from the HX effect on the in-phase magnetization  $N_x$ . The H/H exchange effect on  $2N_y H_z$  (equation 3) is considered in a similar way to the case of the H/H exchange of the in-phase  $N_x$  spin state (equation 2). The resulting coefficients of  $(c_1 - c_1)$  and  $(c_2 - c_2)$  by HX will become zero by ensemble averaging.

$$\begin{aligned}
 2H_z N_y &= (0.5H^\alpha - 0.5H^\beta) N_y \\
 \xrightarrow{HX} & \{0.5(c_1 H^\alpha + c_2 H^\beta) - 0.5(c_1 H^\alpha + c_2 H^\beta)\} N_y \quad (3) \\
 &= \{0.5(c_1 - c_1)H^\alpha + 0.5(c_2 - c_2)H^\beta\} N_y \\
 &\approx \text{zero}
 \end{aligned}$$

Since in-phase and anti-phase spin states undergo different HX effect, in-phase / anti-phase populations in the  $^{15}\text{N}$   $R_2^{\text{CPMG}}$  NMR experiment need to be known and have been formulated in equation 4 through the time-averaged  $R_2$  of the exchanging in-phase and anti-phase magnetizations via J-coupling

$$R_2^{\text{CPMG}} = 0.5\{1 + \text{sinc}(2\pi J\tau)\}R_2^{\text{in}} + 0.5\{1 - \text{sinc}(2\pi J\tau)\}R_2^{\text{anti}} \quad (4)$$

where  $R_2^{\text{in}}$  and  $R_2^{\text{anti}}$  are transverse relaxation rates of in-phase and anti-phase spin states,  $J$  is the  $^1\text{J}_{\text{NH}}$  coupling constant, and  $\tau$  is the CPMG refocusing delay (Griesinger and Ernst 1988; Palmer III et al. 1992). Coefficients of  $R_2^{\text{in}}$  and  $R_2^{\text{anti}}$  are effective fractions of in-phase and anti-phase states;  $f_{\text{in}} = 0.5\{1 + \text{sinc}(2\pi J\tau)\}$  and  $f_{\text{anti}} = 0.5\{1 - \text{sinc}(2\pi J\tau)\}$ . The fraction of the anti-phase state ( $f_{\text{anti}}$ ) is estimated as 0.011 with  $\tau = 0.625\text{ms}$  and  $^1\text{J}_{\text{NH}} = 93\text{Hz}$  under the experimental conditions. Since the HX effect on  $^{15}\text{N}$  relaxation is included only in the anti-phase spin state, the apparent  $R_2$  ( $R_2^{\text{app}}$ ) is approximated by  $R_2^{\text{CPMG}} + f_{\text{anti}}k_{\text{HX}}$  which includes the HX effect along with equation 4. By combining this and equation 1, the resulting multi-exponential function for HX effect is in equation 5.

$$I(t) = I_0\{Qe^{-f_{\text{anti}}k_{\text{HX}}t} + (1-Q)e^{-k_{\text{HX}}t}\}e^{-R_2^{\text{CPMG}}t} \quad (5)$$

Thus the total HX effect in the  $^{15}\text{N}$   $R_2^{\text{CPMG}}$  NMR experiment using refocused HSQC results from the combined effects of H/D exchange, H/H exchange, and in-phase/anti-phase J-coupling exchange. A relevant kinetic model is presented in scheme 1 and the derivation of equation 5 is in Supporting Information. In scheme 1, the total  $k_{\text{HX}}$  was divided into  $Qk_{\text{HX}}$  and  $(1-Q)k_{\text{HX}}$  where  $Q$  is the preserved hydrogen fraction through H/H exchange. Therefore,  $Qk_{\text{HX}}$  is the H/H exchange rate and  $(1-Q)k_{\text{HX}}$  is the H/D exchange rate in the mixed solvent of  $\text{H}_2\text{O}$  and  $\text{D}_2\text{O}$ . The multi-exponential fit of the  $R_2^{\text{CPMG}}$  relaxation data with five parameters ( $I_0$ ,  $Q$ ,  $f_{\text{anti}}$ ,  $k_{\text{HX}}$ , and  $R_2$ ) shown in equation 5 is not plausible but can be reduced to a fit with two unknown parameters ( $I_0$  and  $R_2$ ) and with three known  $Q$ ,  $f_{\text{anti}}$ , and  $k_{\text{HX}}$  parameters. Since  $Q (= 0.9)$  and  $f_{\text{anti}} (= 0.011)$  are estimated from the  $\text{H}_2\text{O}$  content and the anti-phase fraction under the controlled experimental condition,  $k_{\text{HX}}$  rates need to be measured separately.

To find  $k_{\text{HX}}$ , the CLEANEX (Hwang et al. 1998) experiments were performed at pH 7.4 and pH 6.2 and measured  $k_{\text{HX}}$  rates of  $\alpha\text{Syn}$  were plotted in Fig. 3. The  $k_{\text{HX}}$  values at pH 7.4 are significantly different from the values at pH 6.2 and also show variations across the protein sequence. Compared to the apparent  $R_2^{\text{CPMG}}$  values (Fig. 1) obtained from the single exponential decay, the difference in overall  $k_{\text{HX}}$  values between pH 7.4 and pH 6.2 were much greater than the difference in apparent  $R_2^{\text{CPMG}}$  values indicating that the  $R_2^{\text{CPMG}}$  experiment suppresses a significant amount of the HX effect despite the fact that the HX effect is still observable on the apparent  $R_2^{\text{CPMG}}$  values.

To ensure the steady state of amide and water proton magnetization, the  $R_2^{\text{CPMG}}$  pulse sequence with the proton SS pulse (Fig. 2) was used and  $R_2^{\text{CPMG}}$  relaxation data were acquired again at pH 7.4 and pH 6.2. With  $Q = 0.9$ ,  $f_{\text{anti}} = 0.011$ , and known  $k_{\text{HX}}$  values from Fig. 3, multi-exponential fits to the  $R_2^{\text{CPMG}}$  relaxation data were performed with equation 5. The optimized  $R_2^{\text{CPMG}}$  values at pH 7.4 and pH 6.2 were plotted in Fig. 4. The large discrepancies in the apparent  $R_2^{\text{CPMG}}$  values of  $\alpha\text{Syn}$  shown in Fig. 1 were eliminated by including the HX effect in the data analysis.  $R_2^{\text{CPMG}}$  values at pH 7.4 reported in Fig. 1 were corrected more than pH 6.2,  $R_2^{\text{CPMG}}$  values since  $k_{\text{HX}}$  values at pH 7.4 were generally much greater than these values at pH 6.2. Average  $^{15}\text{N}$   $R_2^{\text{CPMG}}$  values at pH 7.4 and pH 6.2 in Fig. 4 were  $3.27 \pm 0.46$  Hz and  $3.13 \pm 0.44$  Hz and the difference of average  $R_2^{\text{CPMG}}$

values between two pH's was reduced to 0.14 Hz comparing to the difference of 0.67 in Fig. 1.

To demonstrate how the HX impacts on the  $R_2^{\text{CPMG}}$  relaxation data, examples of the relaxation data with (filled squares) and without (filled circles) the proton SS pulse were plotted in Fig. 5 for residue G25 under fast HX ( $k_{\text{HX}} = 36.3$  Hz) at pH 7.4. Regardless of the SS pulse, multi-exponential fittings (solid lines) resulting from equation 5 improve the goodness of fit comparing to single exponential fittings (dotted line) in the presence of fast HX. Between multi-exponential fits of relaxation data with and without the proton SS pulse, it may be hard to judge whether the resulted  $R_2^{\text{CPMG}}$  values are correct or not by comparing the goodness of fits. However,  $R_2^{\text{CPMG}}$  values from the use of the SS pulse are smaller than those resulting from no SS pulse and represent the  $R_2^{\text{CPMG}}$  values without HX effect. The overall good agreement of the  $R_2^{\text{CPMG}}$  values between pH 7.4 and pH 6.2 (Fig. 4) can be achieved by acquiring the relaxation data with the proton SS pulse and analyzing the data with equation 5. Therefore, in the presence of observable HX, the  $R_2^{\text{CPMG}}$  relaxation data are affected by HX and these corrections are favorable.

In the presence of fast HX, it seems difficult to eliminate the HX effect in the  $R_2^{\text{CPMG}}$  experiment and also hard to identify the HX effect in the data analysis as shown in Fig. 5. Using a protein sample prepared in 100%  $\text{H}_2\text{O}$  (Iwahara et al. 2007) will suppress the HX effect on  $R_2^{\text{CPMG}}$  data but incomplete suppression is still expected due to no control of the water proton magnetization by the steady state pulse and the anti-phase populations. Without correcting the HX effect, the elevated apparent  $R_2^{\text{CPMG}}$  values could be mis-interpreted as a result of other contributions such as conformational exchange or protein aggregation. In this context, conformational exchange of  $\alpha\text{Syn}$  proposed at 37 °C may be due to HX effect on  $R_2^{\text{CPMG}}$  at these high temperatures (McNulty et al. 2006). When  $^1\text{H}$ - $^{15}\text{N}$  HSQC signal intensities are vulnerable by removing the water flip-back pulse and dephasing the water magnetization (Chen and Tjandra 2011), fast HX may play a role to attenuate the the signal intensity because of saturated water magnetization. In that situation, the HX effect needs to be under control in the acquisition scheme and in the data analysis as proposed here.

To minimize the HX effect on the  $R_2^{\text{CPMG}}$  experiment, relaxation rates need to be obtained by working under conditions for which fast HX does not exist such as low temperatures and low pH's. In Supporting Information, dramatic intensity changes of HSQC spectra by HX effect were demonstrated by acquiring spectra at (pH 7.4, 35 °C), (pH 7.4 15 °C), (pH 6.1, 35 °C), and (pH 6.1, 15 °C). When sample conditions for removing fast HX cannot be obtained, we have shown that the  $^{15}\text{N}$   $R_2^{\text{CPMG}}$  pulse sequence needs to contain SS pulses in order to remove water saturation effects and that quantitative corrections of the  $^{15}\text{N}$  relaxation rates for fast HX need to be performed. When the experiment is performed on a protein sample in mixed  $\text{H}_2\text{O}/\text{D}_2\text{O}$  solvent, use of the multi-exponential fit to the  $^{15}\text{N}$   $R_2^{\text{CPMG}}$  relaxation data by equation 5 is a necessary to extract  $^{15}\text{N}$   $R_2^{\text{CPMG}}$  values without HX effect. Other approaches that may simplify the data analysis include running the experiment in 100%  $\text{H}_2\text{O}$  as practiced by Clore's group (Iwahara et al. 2007). This removes the H/D exchange effect and the multi-exponential fit by equation 5 is changed to the single exponential fit with the rate constant of  $(f_{\text{anti}} \cdot k_{\text{HX}} + R_2^{\text{CPMG}})$ . The required HX rates in data analysis can be measured independently from  $R_{1\text{zz}}$  (Skrynnikov and Ernst 1999), MEXICO (Gemmecker et al. 1993), CLEANEX (Hwang et al. 1998), or  $^{15}\text{N}^{\text{H/D}}$ -SOLEXSY (Chevelkov et al. 2010) experiments. As an alternative to  $R_2^{\text{CPMG}}$ , the  $^{15}\text{N}$   $R_{1\rho}$  experiment has been also used for  $R_2$  measurement after correction of off-resonance effects (Korzhnev et al. 2002; Palmer III et al. 2001). If the anti-phase population is negligible ( $f_{\text{anti}} \cong 0$ ) in the  $^{15}\text{N}$   $R_{1\rho}$  experiment by applying a strong spin-lock on  $N_x$ , then equation 5 becomes the single exponential fit with the rate constant of  $R_{1\rho}$  with the use of 100%  $\text{H}_2\text{O}$ . A noticeable merit of this approach is no requirement of measuring  $k_{\text{HX}}$  values. However, it is important

that the SS pulses are used for any variation of experiments to assure the steady state HX equilibrium during the inter-scan recycle delay.

Although we demonstrated the HX effect on the backbone  $^{15}\text{N}$   $R_2^{\text{CPMG}}$  relaxation data applied to the IDP  $\alpha\text{Syn}$ , the HX effect may be present in other protein samples and various NMR experiments. In addition to IDPs, globular proteins with unstructured elements such as large loop regions, or partially folded proteins that have a large number of amides exposed to the solvent may have increased apparent  $^{15}\text{N}$   $R_2^{\text{CPMG}}$  values due to fast HX. Other NMR experiments that have long evolution times to measure relaxation and diffusion should be potentially considered for modulation by fast HX effects as well (Dempsey 2001; Zhang et al. 1995). Like  $^{15}\text{N}$   $R_2^{\text{CPMG}}$  experiments, NMR experiments that use the indirect detection scheme also need to be considered with caution although heteronuclear NMR properties other than  $^1\text{H}$  are observed.

## Supplementary Material

Refer to Web version on PubMed Central for supplementary material.

## Acknowledgments

K-PW was supported by an NIH Interdisciplinary Research Workforce Fellowship (5R90DK071502). This work has been supported by NIH grants (GM087012 and GM45302) and NSF grants (DBI-0403062 and DBI-0320746) to JB.


## References

- Bertini I, Carrano CJ, Luchinat C, Piccioli M, Poggi L. A  $^{15}\text{N}$  NMR mobility study on the dicalcium P43M calbindin D9k and its mono-La3+-substituted form. *Biochemistry*. 2002; 41:5104–5111. [PubMed: 11955058]
- Bertoncini CW, Rasia RM, Lamberto GR, Binolfi A, Zweckstetter M, Griesinger C, Fernandez CO. Structural characterization of the intrinsically unfolded protein beta-synuclein, a natural negative regulator of alpha-synuclein aggregation. *J Mol Biol*. 2007; 372:708–722. [PubMed: 17681539]
- Buevich AV, Baum J. Dynamics of unfolded proteins: incorporation of distributions of correlation times in the model free analysis of NMR relaxation data. *J Am Chem Soc*. 1999; 121:8671–8672.
- Buevich AV, Shinde UP, Inouye M, Baum J. Backbone dynamics of the natively unfolded pro-peptide of subtilisin by heteronuclear NMR relaxation studies. *J Biomol NMR*. 2001; 20:233–249. [PubMed: 11519747]
- Bussell R Jr, Eliezer D. Residual structure and dynamics in Parkinson's disease-associated mutants of alpha-synuclein. *J Biol Chem*. 2001; 276:45996–46003. [PubMed: 11590151]
- Chen J, Tjandra N. Water proton spin saturation affects measured protein backbone  $^{15}\text{N}$  spin relaxation rates. *J Magn Reson*. 2011; 213:151–157. [PubMed: 22015249]
- Chevelkov V, Xue Y, Rao DK, Forman-Kay JD, Skrynnikov NR.  $^{15}\text{N}^{\text{H/D}}$ -SOLESY experiment for accurate measurement of amide solvent exchange rates: application to denatured drkN SH3. *J Biomol NMR*. 2010; 46:227–244. [PubMed: 20195703]
- Chiti F, Dobson CM. Protein misfolding, functional amyloid, and human disease. *Annu Rev Biochem*. 2006; 75:333–366. [PubMed: 16756495]
- Croke RL, Sallum CO, Watson E, Watt ED, Alexandrescu AT. Hydrogen exchange of monomeric alpha-synuclein shows unfolded structure persists at physiological temperature and is independent of molecular crowding in *Escherichia coli*. *Protein Sci*. 2008; 17:1434–1445. [PubMed: 18493022]
- Delaglio F, Grzesiek S, Vuister GW, Zhu G, Pfeifer J, Bax A. NMRPipe: a multidimensional spectral processing system based on UNIX pipes. *J Biomol NMR*. 1995; 6:277–293. [PubMed: 8520220]
- Dempsey CE. Hydrogen exchange in peptides and proteins using NMR spectroscopy. *Prog Nucl Magn Reson Spectrosc*. 2001; 39:135–170.




- Eliezer D. Biophysical characterization of intrinsically disordered proteins. *Curr Opin Struct Biol.* 2009; 19:23–30. [PubMed: 19162471]
- Eliezer D, Kutluay E, Bussell R Jr, Browne G. Conformational properties of alpha-synuclein in its free and lipid-associated states. *J Mol Biol.* 2001; 307:1061–1073. [PubMed: 11286556]
- Farrow NA, Muhandham R, Singer AU, Pascal SM, Kay CM, Gish G, Shoelson SE, Pawson T, Forman-Kay JD, Kay LE. Backbone Dynamics of a Free and a Phosphopeptide-Complexed Src Homology 2 Domain Studied by  $^{15}\text{N}$  NMR Relaxation. *Biochemistry.* 1994; 33:5984–6003. [PubMed: 7514039]
- Gemmecker G, Jahnke W, Kessler H. Measurement of fast proton exchange rates in isotopically labeled compounds. *J Am Chem Soc.* 1993; 115:11620–11621.
- Goedert M. Alpha-synuclein and neurodegenerative diseases. *Nat Rev Neurosci.* 2001; 2:492–501. [PubMed: 11433374]
- Griesinger C, Ernst RR. Cross relaxation in time-dependent nuclear spin systems: Invariant trajectory approach. *Chem Phys Lett.* 1988; 152:239–247.
- Hansen DF, Led JJ. Implications of using approximate Bloch–McConnell equations in NMR analyses of chemically exchanging systems: application to the electron self-exchange of plastocyanin. *J Magn Reson.* 2003; 163:215–227. [PubMed: 12914837]
- Hwang TL, van Zijl PC, Mori S. Accurate quantitation of water-amide proton exchange rates using the phase-modulated CLEAN chemical EXchange (CLEANEX-PM) approach with a Fast-HSQC (FHSQC) detection scheme. *J Biomol NMR.* 1998; 11:221–226. [PubMed: 9679296]
- Iwahara J, Jung YS, Clore GM. Heteronuclear NMR spectroscopy for lysine  $\text{NH}_3$  groups in proteins: unique effect of water exchange on  $^{15}\text{N}$  transverse relaxation. *J Am Chem Soc.* 2007; 129:2971–2980. [PubMed: 17300195]
- Kateb F, Pelupessy P, Bodenhausen G. Measuring fast hydrogen exchange rates by NMR spectroscopy. *J Magn Reson.* 2007; 184:108–113. [PubMed: 17049891]
- Kay LE, Torchia DA, Bax A. Backbone dynamics of proteins as studied by  $^{15}\text{N}$  inverse detected heteronuclear NMR spectroscopy: application to staphylococcal nuclease. *Biochemistry.* 1989; 28:8972–8979. [PubMed: 2690953]
- Korzhnev DM, Skrynnikov NR, Millet O, Torchia DA, Kay LE. An NMR Experiment for the Accurate Measurement of Heteronuclear Spin-Lock Relaxation Rates. *J Am Chem Soc.* 2002; 124:10743–10753. [PubMed: 12207529]
- Mandel AM, Akke M, Palmer AG 3rd. Backbone dynamics of Escherichia coli ribonuclease HI: correlations with structure and function in an active enzyme. *J Mol Biol.* 1995; 246:144–163. [PubMed: 7531772]
- McConnell HM. Reaction rates by nuclear magnetic resonance. *J Chem Phys.* 1958; 28:430–431.
- McNulty BC, Tripathy A, Young GB, Charlton LM, Orans J, Pielak GJ. Temperature-induced reversible conformational change in the first 100 residues of alpha-synuclein. *Protein Sci.* 2006; 15:602–608. [PubMed: 16452621]
- Mittag T, Forman-Kay JD. Atomic-level characterization of disordered protein ensembles. *Curr Opin Struct Biol.* 2007; 17:3–14. [PubMed: 17250999]
- Mittermaier AK, Kay LK. Observing biological dynamics at atomic resolution using NMR. *Trends Biochem Sci.* 2009; 34:601–611. [PubMed: 19846313]
- Palmer AG III. NMR characterization of the dynamics of biomacromolecules. *Chem Rev.* 2004; 104:3623–3640. [PubMed: 15303831]
- Palmer AG III, Kroenke CD, Loria JP. Nuclear Magnetic Resonance Methods for Quantifying Microsecond-to-Millisecond Motions in Biological Macromolecules. *Method Enzymol.* 2001; 339:204–238.
- Palmer AG III, Skelton NJ, Chazin WJ, Wright PE, Rance M. Suppression of the effects of cross-correlation between dipolar and anisotropic chemical shift relaxation mechanisms in the measurement of spin-spin relaxation rates. *Mol Phys.* 1992; 75:699–711.
- Skrynnikov NR, Ernst RR. Detection of intermolecular chemical exchange through decorrelation of two-spin order. *J Magn Reson.* 1999; 137:276–280. [PubMed: 10053160]

- Tugarinov V, Kay LE. Quantitative NMR studies of high molecular weight proteins: application to domain orientation and ligand binding in the 723 residue enzyme malate synthase G. *J Mol Biol.* 2003; 327:1121–1133. [PubMed: 12662935]
- Uversky VN, Gillespie JR, Fink AL. Why are “natively unfolded” proteins unstructured under physiologic conditions? *Proteins.* 2000; 41:415–427. [PubMed: 11025552]
- Wang C, Palmer AG 3rd. Solution NMR methods for quantitative identification of chemical exchange in  $^{15}\text{N}$ -labeled proteins. *Magn Reson Chem.* 2003; 41:866–876.
- Weinreb PH, Zhen W, Poon AW, Conway KA, Lansbury PT Jr. NACP, a protein implicated in Alzheimer’s disease and learning, is natively unfolded. *Biochemistry.* 1996; 35:13709–13715. [PubMed: 8901511]
- Wright PE, Dyson HJ. Linking folding and binding. *Curr Opin Struct Biol.* 2009; 19:31–38. [PubMed: 19157855]
- Wu K-P, Kim S, Fela DA, Baum J. Characterization of conformational and dynamic properties of natively unfolded human and mouse alpha-synuclein ensembles by NMR: implication for aggregation. *J Mol Biol.* 2008; 378:1104–1115. [PubMed: 18423664]
- Wu K-P, Weinstock DS, Narayanan C, Levy RM, Baum J. Structural Reorganization of alpha-Synuclein at Low pH Observed by NMR and REMD Simulations. *J Mol Biol.* 2009; 391:784–796. [PubMed: 19576220]
- Zhang YZ, Paterson Y, Roder H. Rapid amide proton exchange rates in peptides and proteins measured by solvent quenching and two-dimensional NMR. *Protein Sci.* 1995; 4:804–814. [PubMed: 7613478]




**Fig. 1.**


(a) Apparent  $R_2^{\text{CPMG}}$  values of  $\alpha$ Syn at pH 7.4 (red circles) and pH 6.2 (blue rhombi) at 15°C.  $R_2^{\text{CPMG}}$  relaxation data acquired without the proton SS pulse (Fig. 2) were fit with a single exponential decay function. There are large and systematic differences between them in addition to small random deviations. (b) Correlation between  $R_2^{\text{CPMG}}$  values at pH 7.4 and pH 6.2.



**Fig. 2.** The  $^{15}\text{N}$   $R_2^{\text{CPMG}}$  pulse sequence. The gNhsqc (with  $T_2 = 'y'$ ) from the Varian BioPack library is outlined.  $90^\circ$  and  $180^\circ$  hard pulses are represented by narrow and wide rectangles. Pulse phases are written on the pulses but if not mentioned, phases are on the x-axis. The sinc shaped water flip-back pulse is used during the forward refocused INEPT to minimize the water saturation by bringing the water magnetization to the z-axis. Delay  $\tau_a$  is  $0.25/|J_{\text{NH}}|$ . Phase cycles are  $\phi_1 = x, -x$ ,  $\phi_2 = x, x, y, y, -x, -x, -y, -y$ , and  $\phi_R = x, -x, -x, x$ . To test the HX effect, an optional proton steady state (SS) pulse is introduced in the beginning of the pulse sequence. The implemented SS pulse is gradient – RF pulse – gradient where the RF pulse is a proton  $90^\circ$  hard pulse and the gradient is the strength of 10 gauss/cm and the duration of 0.5.  $2n-1$  ( $= 1, 3, 5$ , etc) cycles of  $^{15}\text{N}$  CPMG pulse trains are applied for the  $R_2^{\text{CPMG}}$  measurement where the single cycle takes the 10 ms relaxation delay and a  $180^\circ$  proton pulse. The inter-pulse delay between  $180^\circ$   $^{15}\text{N}$  pulses in CPMG is 1.25 ms. The gNhsqc adapts the sensitivity enhanced HSQC with PEP (Preservation and Equivalent Pathways) and the gradient coherence selection (by  $g_1$  and  $g_2$ ). Other experimental details are available from Varian BioPack.




**Fig. 3.** pH-dependent  $k_{HX}$  values.  $k_{HX}$  values of  $\alpha$ Syn were measured by CLEANEX method at pH 7.4 (red circles) and pH 6.2 (blue rhombi) at 15 °C. The large increase of  $k_{HX}$  values at pH 7.4 is prominent by increasing pH.




**Fig. 4.**

(a) Corrected  $R_2^{\text{CPMG}}$  values at pH 7.4 and 6.2.  $R_2^{\text{CPMG}}$  values at pH 7.4 (red circles) and pH 6.2 (blue rhombi) were obtained with the multi-exponential decay function (equation 5) using  $Q = 0.9$ ,  $f_{\text{anti}} = 0.011$ , and  $k_{\text{HX}}$  values from Fig. 3. The relaxation data were acquired with the SS pulse. The large differences seen in Fig. 1 were corrected. (b) Correlation between  $R_2^{\text{CPMG}}$  values at pH 7.4 and pH 6.2.



**Fig. 5.** Examples of data fitting for residue G25. Experimental  $R_2^{\text{CPMG}}$  relaxation data for residue G25 at pH 7.4 were obtained with and without the proton SS pulse. The data (filled circles) without the proton SS pulse were fit with a single exponential (dotted line,  $I_0 = 1.25 \times 10^6$ ,  $R_2^{\text{CPMG}} = 4.90$  Hz, and  $R^2 = 99.1$  %) and a multi-exponential (equation 5) (solid line,  $I_0 = 1.31 \times 10^6$ ,  $Q = 0.9$ ,  $f_{\text{anti}} = 0.011$ ,  $k_{\text{HX}} = 36.3$  Hz,  $R_2^{\text{CPMG}} = 4.13$  Hz, and  $R^2 = 99.8$  %), where  $R^2$  is the coefficient of determination for the goodness of fit. The data (filled squares) with the proton SS pulse were also fit with a single exponential (dotted line,  $I_0 = 8.85 \times 10^5$ ,  $R_2^{\text{CPMG}} = 4.11$  Hz, and  $R^2 = 99.2$  %) and a multi-exponential (equation 5) (solid line,  $I_0 = 9.35 \times 10^5$ ,  $Q = 0.9$ ,  $f_{\text{anti}} = 0.011$ ,  $k_{\text{HX}} = 36.3$  Hz,  $R_2^{\text{CPMG}} = 3.37$  Hz, and  $R^2 = 99.8$  %).

**Scheme 1.**

Hydrogen exchange effect during the  $^{15}\text{N}$   $R_2$  relaxation time period. The H/H exchange rate is  $Qk_{\text{HX}}$  and the H/D exchange rate is  $(1-Q)k_{\text{HX}}$  when the  $\text{H}_2\text{O}$  content in the sample solvent is  $Q$  and the  $\text{D}_2\text{O}$  content is  $(1-Q)$ . In H/H exchange, the product  $\text{NH}'$  is distinguished from the reactant  $\text{NH}$ .