



Published in final edited form as:

Fertil Steril. 2013 April ; 99(5): 1385–1391. doi:10.1016/j.fertnstert.2012.11.053.

Self-Reported Menopausal Symptoms, Coronary Artery Calcification and Carotid Intima-Media Thickness in Recently Menopausal Women Screened for the Kronos Early Estrogen Prevention Study (KEEPS)

Erin Foran Wolff, M.D.¹, Yunxiao He, Ph.D.², Dennis M. Black, Ph.D.³, Eliot A. Brinton, M.D., FAHA, FNLA⁴, Mathew J. Budoff, M.D.⁵, Marcelle I. Cedars, M.D.³, Howard N. Hodis, M.D.⁶, Rogerio A. Lobo⁷, JoAnn E. Manson, M.D., Dr.PH.⁸, George R. Merriam, M.D.⁹, Virginia M. Miller, Ph.D.¹⁰, Fredrick Naftolin, M.D., Ph.D.¹¹, Lubna Pal, MBBS, MRCOG, MS², Nanette Santoro, M.D.¹², Heping Zhang, Ph.D.², S. Mitchell Harman, M.D., Ph.D.¹³, and Hugh S. Taylor, M.D.²

¹Program in Reproductive and Adult Endocrinology, NICHD, NIH

²School of Medicine, Yale University

³School of Medicine, University of San Francisco

⁴School of Medicine, University of Utah

⁵Los Angeles biomedical Research institute at Harbor-UCLA

⁶University of Southern California, Atherosclerosis Research Unit Core Imaging and Reading Center

⁷College of Physicians and Surgeons, Columbia University

⁸Brigham and Women's Hospital, Harvard Medical School

⁹VA Puget Sound Health Care System and University of Washington School of Medicine

¹⁰College of Medicine, Mayo Clinic

¹¹School of Medicine, New York University

¹²School of Medicine, University of Colorado

¹³Kronos Longevity Research Institute

Abstract

Corresponding author: Erin Foran Wolff, MD, Reproductive Endocrinology and Infertility Fellow, Program in Reproductive and Adult Endocrinology, *Eunice Kennedy Shriver* National Institute of Child, Health and Human Development, NIH, Bldg 10, CRC; Rm 1E-3140, 10 Center Dr, Bethesda, MD 20892, 301-496-5800, Fax: 301-402-0884.

ClinicalTrials.gov number is NCT00154180.

Additional Contributions: We gratefully acknowledge the dedicated efforts of all the investigators and staff at the KEEPS clinical centers, the KEEPS Data Coordinating Center at KLRI, and the NIH Institutes supporting ancillary studies. The KEEPS investigators would like to thank the Aurora Foundation for study support and Bayer HealthCare Pharmaceuticals, Inc. and Abbott Laboratories for providing study medications. Above all, we recognize and thank the KEEPS participants for their dedication and commitment to the KEEPS research program.

Publisher's Disclaimer: This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final citable form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

Objective—To determine whether self-reported menopausal symptoms are associated with measures of subclinical atherosclerosis.

Setting—Multi-center, randomized controlled trial.

Patients—Recently menopausal women (n=868) screened for the Kronos Early Estrogen Prevention Study (KEEPS).

Design—Cross sectional analysis.

Interventions—None

Main Outcome Measures—Baseline menopausal symptoms (hot flashes, dyspareunia, vaginal dryness, night sweats, palpitations, mood swings, depression, insomnia, irritability), serum estradiol (E2) levels and measures of atherosclerosis were assessed. Atherosclerosis was quantified using Coronary Artery Calcium (CAC) Agatston scores (n=771) and Carotid Intima-Media Thickness (CIMT). Logistic regression model of menopausal symptoms and E2 was used to predict CAC. Linear regression model of menopausal symptoms and E2 was used to predict CIMT. Correlation between length of time in menopause with menopausal symptoms, estradiol (E2), CAC, and CIMT were assessed.

Results—In early menopausal women screened for KEEPS, neither E2 nor climacteric symptoms predicted the extent of subclinical atherosclerosis. Palpitations (p=0.09) and depression (p=0.07) approached significance as predictors of CAC. Other symptoms of insomnia, irritability, dyspareunia, hot flashes, mood swings, night sweats, and vaginal dryness were not associated with CAC. Women with significantly elevated CAC scores were excluded from further participation in KEEPS; in women meeting inclusion criteria, neither baseline menopausal symptoms nor E2 predicted CIMT. Years since menopause onset correlated with CIMT, dyspareunia, vaginal dryness and E2.

Conclusions—Self-reported symptoms in recently menopausal women are not strong predictors of subclinical atherosclerosis. Continued follow-up of this population will be performed to determine if baseline or persistent symptoms in the early menopause are associated with progression of cardiovascular disease.

Keywords

KEEPS; estrogen; cardiovascular; menopause; CAC; CIMT; palpitations; depression

Introduction

The efficacy of menopausal hormone therapy (MHT) in treating menopausal symptoms is well established and menopausal symptoms remain the primary indication for initiation of MHT in the perimenopausal and early postmenopausal years. Menopause also reflects a time when the risk for cardiovascular disease (CVD) increases, however, the precise effects of MHT on cardiovascular health remain controversial as they have been more difficult to delineate. Many large observational studies suggest that MHT has cardioprotective effects (1–8). However, this dogma was challenged by large, prospective, randomized controlled trials, which failed to show cardiovascular benefit of MHT when initiated remote (>10 yrs) from menopause (9) or in the setting of preexisting CVD (10).

Many preventive strategies have been established to reduce cardiovascular events in patients at risk. However, individuals at risk for CVD are not always easily identified. Baseline data from the Kronos Early Estrogen Prevention Study (KEEPS) were enlightening, revealing that 4% of the early postmenopausal population screened demonstrated an unexpected degree of asymptomatic coronary artery atherosclerosis (11). Additional screening tools to

better define cardiovascular risk may help to individualize the risk/benefit discussion of MHT.

As menopausal symptoms are the primary indication for MHT, it would be important to know if symptoms are related to cardiovascular disease risk. Conflicting evidence on this association exists. Menopausal complaints have been associated with cardiovascular disease risk factors such as hypertension, hypercholesterolemia, and body mass index (BMI) in one study (12). Increased cardiovascular events have been observed in women who experienced baseline vasomotor symptoms (VMS) compared to women who did not have baseline symptoms, but only in the first year after MHT was initiated (13). In the Study of Women Across the Nation (SWAN) study, perimenopausal women with VMS had significantly higher coronary artery calcium scores (CAC) when measured near menopause (14). In the Dutch Eindhoven Perimenopausal Osteoporosis Study (EPOS) study, night sweats (but not VMS) were associated with coronary heart disease events (16). Conversely, significantly lower CAC was observed at the end of the estrogen only arm of the Women's Health Initiative (WHI), suggesting that processes underlying atherosclerosis may be mitigated with postmenopausal estrogen therapy (15). Data from the WHI-observational study were particularly intriguing wherein a decreased hazard for cardiovascular events was observed in women experiencing VMS early in menopause, whereas, VMS occurring late in menopause were associated with increased hazard of cardiovascular events (17). These data suggest that a relationship between atherosclerosis and VMS, if any, may be modulated by the time since menopause. We have herein sought to determine if common menopausal symptoms relate to measures of atherosclerosis in an early menopausal population of overtly healthy women undergoing screening for KEEPS.

Given that symptoms in early menopause are the primary indication for initiation of MHT and the risks of initiating MHT in the setting of preexisting disease, we sought to characterize measures of atherosclerosis in this population to provide additional clarification on the association between menopausal symptoms with cardiovascular disease.

Materials and Methods

This multicenter trial was approved by an independent Institutional Review Board; in addition, the study was approved by the Institutional Review Board at each study site. Early menopausal women 42 to 58 years of age whose last menstrual period occurred at least 6 months and no more than 3 years previously were screened for eligibility for the KEEPS trial (NCT00154180) at 9 clinical study centers in the United States; study details have been reported elsewhere (18). As per study design, subclinical atherosclerosis was assessed by CAC and carotid intima-media thickness (CIMT). Coronary artery calcium was measured by computed tomography (CT) or electron beam tomography (EBT) and calculated two ways: 1) the mean Agatston score of two measurements and 2), and the mean of two CAC volume scores. Women with mean Agatston scores ≥ 50 U, taken to indicate coronary artery disease, were excluded from randomization. Women meeting inclusion criteria underwent baseline measurements of CIMT. CIMT was measured by computer image-processed B-mode ultrasound at two visits prior to randomization to study drug, and the average score was used for analysis. Emphasis of ultrasound imaging is on the distal centimeter of the common carotid artery (CCA) because least variability occurs in this area,(19) facilitating reproducible longitudinal imaging necessary for randomized controlled trials.(20–46) In addition, only longitudinal change in the distal centimeter of the (CCA) has been shown to be predictive of clinical coronary events(28); the CCA has been shown to be as good or better than other carotid artery segments in the prediction of cardiovascular events. The far wall is used for statistical purposes since measurement of near wall thickness is less accurate. (47) Per study design, study images were read at central facilities (CAC at the St.

Johns Cardiovascular Research Center, Harbor UCLA Medical Center and CIMT at the Atherosclerosis Research Unit, Core Imaging and Reading Center(25)) by one expert blinded to participant symptom burden and characteristics.

At screening, women were asked to rank the severity of their menopausal symptoms of depression, insomnia, irritability, dyspareunia, hot flashes, mood swings, night sweats, palpitations, and vaginal dryness on a scale of 0 (none), mild (1) to 4 (severe). Serum E2 concentrations were measured in pg/ml. These measures were correlated with self-reported menopausal symptoms and serum estradiol (E2).

Potential confounders for CVD were ascertained as previously reported (11): height (cm), weight (kg), body mass index (BMI, kg/m²), blood pressure (systolic, diastolic in mm Hg), waist circumference (cm), total cholesterol (mg/dL), low density cholesterol level (LDL, mg/dL), high density cholesterol level (HDL, mg/dL), triglyceride level (mg/dL), fasting glucose (mg/dL), Framingham Risk Score (FRS)(48), cigarette smoking (history of smoking and cigarettes per day), alcohol intake (number of alcoholic drinks consumed per month), and metabolic syndrome. Serum E2 levels (pg/ml) and serum follicle stimulating hormone (FSH, ng/ml) were ascertained at the local study site laboratories. Details of the study design for the randomized trial have been previously reported(11). We chose a more conservative exclusion cut off of CAC>50 to limit the risks to subjects associated with starting HT in the setting of significant heart disease, rather than an exclusion criteria of CAC>100. Women with high cardiovascular factors were excluded.

Statistical Analysis

Continuous data were presented as mean+standar deviation (range) and categorical data as number (%). Data distribution was assessed; since most (80.7%, 700/868) of the Agatston measurements and CAC volumes were scored as zero, this measure was recoded into a dichotomous variable as absent (0 score) or present (score>0). Logistic regression analysis was used to investigate the relationship between CAC (present/absent) and menopausal symptoms while controlling for the effects of the covariates (height, weight, BMI, blood pressure, waist circumference, total cholesterol, LDL, HCL, triglycerides, fasting glucose, FRS, smoking, alcohol use, metabolic syndrome, E2, and FSH). Linear regression analysis was performed to investigate the relationship between CIMT and the 9 measures of menopausal symptoms while controlling the effects of the same covariates as included in the logistic regression model. Spearman's rank correlation was used to test for correlation between years of menopause and each of the nine menopausal symptoms, Agatston score, and CAC volume since the former (years of menopause) is a continuous variable and the others are ordinal. Pearson's correlation (correlation coefficient, CC) test was used for testing correlations between years of menopause and CIMT and E2 given that all variables involved are continuous. Additional sensitivity analyses were conducted by atherosclerosis disease severity. Statistical analyses were performed using the statistical software R 2.11.1. A p-value of less than 0.05 is considered statistically significant.

RESULTS

Data from n=868 subjects screened for the KEEPS trial were included in the analysis. A summary of the important variables is shown in Table 1. The average age of women screened was 52.7±2.6 SD (42–58) years. As was the aim of the study design, all women were within 3 years since their last menstrual period. Participants had low levels of atherosclerotic burden, given the young age and early menopausal status of the study population(11). Due to the length of time since menopause, E2 values were skewed as expected.

Association between CAC and Menopausal Symptoms

For the logistic regression analysis, mean Agatston score and mean CAC volume were exactly the same after recoding into dichotomous variables as absent (0 score) or present (score>0). Therefore we only report the CAC result for the mean Agatston score. First, a logistic regression of CAC on all variables was performed with n= 771 fully observed records. The p-value from the unweighted sum-of-squares goodness-of-fit test was 0.283 (where the null hypothesis is that the model is fit to the data well) could not be rejected, suggesting that the model was fit to the data well. After excluding noncontributory covariates (specifically, E2 and alcohol use were excluded since they have P-values 0.826 and 0.738, respectively), a logistic regression model was refit by using the 9 measures of menopausal symptoms and the history of smoking whose p-value (0.073) was close to 0.05 in the original model. Under the re-fit model, n=822 records, including n=684 zero readings and n=138 nonzero readings for mean Agatston, were fully observed and used for analysis. Of the menopausal symptoms studied, palpitations and depression were the closest predictors of CAC. The p-value of 0.302 from the same goodness-of-fit test suggests that the null hypothesis that the model still fit the data well could not be rejected.

On univariate analyses summarized in Table 2, none of the nine menopausal symptoms were significantly associated with CAC, where the listed values indicate the log of the Odds Ratio and direction of the relationship in predicting CAC by a positive or negative coefficient. Two measures of menopausal symptoms, palpitations and depression, had p-values 0.091 and 0.070, respectively. Smoking history was shown to be strongly associated with CAC (p= 0.006).

For univariate logistic regression of mean Agatston on each of the nine symptoms, only depression approached significance (p=0.098). However, hot flash was significantly related to mean Agatston among women with evidence of calcification (univariate linear regression of mean Agatston (including only subjects with mean Agatston>0 and with log transformation of mean Agatston), p value = 0.047. However, the R-squared for the univariate model with hotflash is only 0.029. All other eight symptoms were not significant.

For univariate linear regression of mean Agatston (including only subjects with Mean Agatston>50) on each of the nine symptoms, hotflash is marginally significant (p=0.054) and sweats is significant (p= 0.043). The R-squared for the two univariate models are 0.104 and 0.115, respectively. All other seven symptoms are not significant. After controlling for the described confounders, an association between menopausal symptoms and CAC was also assessed in the subgroups with the highest disease burden (defined as CAC>0 and CAC>50) using linear regression. The R-squared measures are 0.08 and 0.18 respectively, suggesting that the menopausal symptoms have little value in predicting CAC.

Association between CIMT and Menopausal Symptoms

After excluding records with missing values, data from n=759 subjects were available for analysis. Intraobserver CIMT variability was estimated from 2 pretreatment scans on each patient, which were performed <6 (mean 3.9 ± 4.2) weeks apart. Coefficients of variation (SD/mean) were, on average, <1.0% (0.83 ± 0.98%). The results from the linear regression analysis are summarized in Table 3. The R square measure is a relatively low 0.131, which suggests that the current set of predictive variables is not adequate for modeling CIMT. In addition, the P-values for all the 9 menopausal symptoms and E2 are far larger than 0.05, showing that it is very unlikely that these variables are associated with CIMT.

Years Since Menopause Correlates with CIMT

The results of the correlation tests between years since menopause with atherosclerotic measures, menopausal symptoms and E2 levels are summarized in Table 4. Time since menopause (years) was significantly associated with E2 (CC=-0.084, P=0.015), CIMT (CC=0.075, P=0.029), dyspareunia (CC=0.084, P=0.015), and vaginal dryness (CC=0.151, P=1.0e-5), but not with CAC or other menopausal symptoms.

Discussion

As menopausal symptoms are the driving force for initiation of MHT, it is critical to understand the cardiovascular characteristics of the women who experience them. This will not only help with counseling women about the benefits and risks of MHT, but could also prove to be a valuable clinical tool to identify high risk patients for further evaluation and preventive strategies. However, menopausal symptoms are not statistically significantly correlated with CAC or CIMT, in this population of recently menopausal women screened for the KEEPS trial. The lack of linkage between menopausal symptoms and evidence of heart disease should provide some reassurance to clinicians about the cardiovascular safety of prescribing MHT to recently menopausal women.

Depression has been shown to be associated with clinical heart disease, but association with subclinical, asymptomatic disease is less studied. Other groups have shown that in asymptomatic middle aged women, coronary calcification is associated with recurrent major depression (49) as well as progression of disease (50), but single depressive episodes were not associated with CIMT plaque (51). However, the menopausal status and symptomatology of these women was not well characterized.

In this study, palpitations approached significance in predicting atherosclerosis measures. Although palpitations specifically have not previously been described to correlate with either clinical heart disease or subclinical markers of atherosclerosis, abnormal heart rate turbulence was found to be associated with cardiac mortality risk, and atrial fibrillation is well known to correlate with risk of stroke (52).

Interestingly, vasomotor symptoms of hot flashes and/or night sweats anytime up to enrollment in the estrogen only arm of the WHI were associated with reduced odds for CAC>0 at the conclusion of the study (15), contrasting findings were noted in a Dutch study (16). However, the women enrolled in the WHI had to be asymptomatic in order to be randomized, which represents a significant recruitment bias to study outcomes associated with menopausal symptoms. It will be interesting as KEEPS continues to assess how progression of CAC relates to vasomotor symptoms and if differences in early versus late symptoms will relate to changes in vascular anatomy (17, 53). Most studies have found some positive association of menopausal symptoms with cardiovascular disease (12–17, 54, 55), while only one study showed a reduced risk (15). Given that most of the participants in KEEPS had low CAC scores, we cannot know if the presence and severity of symptoms may relate to advanced CAC, as was likely present in the WHI subjects.

The strengths of this study include the narrow range of time since menopause of women enrolled in KEEPS. In addition, current and recent symptoms are ascertained, which are less subject to recall bias when recalled distantly. Further, the screened population in KEEPS is representative of the general population of recently menopausal women given that CAC measures were undertaken in all who were screened. Specifically, patients were not excluded for or against menopausal symptoms, which could significantly skew the population studied. The strength of these observations with regard to CIMT is that the relationship was defined in a population which was found to have a low CAC burden (<50)

and deemed at low risk for CVD (low risk low density lipoprotein cholesterol, triglycerides, systemic blood pressure and fasting blood glucose levels).

The cross-sectional study design, the relatively low CVD risk profile of the studied population, and the fact that this study was initiated post hoc following completion of the screening process and hence not designed to specifically assess a relations between atherosclerotic measures and self-reported symptoms are recognized as study limitations. Potential reasons why we failed to detect a significant association between common menopausal symptoms and atherosclerotic markers in this population could include that current and recent symptoms were reported by qualitatively assessment; a more specific, prospective, symptom journaling could potentially reveal an association. Since the evaluation of symptoms was retrospectively self-reported on a broad and subjective scale, it is possible that a prospective assessment of palpitations and depression would reach statistical significance as factors in predicting subclinical atherosclerotic disease. An alternative explanation for the negative study with regard to CIMT may be the study design wherein participants with CAC>50 were excluded from undergoing CIMT assessment; a relationship between menopausal symptoms and established atherosclerosis is thus still plausible.

Conclusion

In this cohort of relatively healthy early menopausal women screened for participation in KEEPS, we did not detect an association between self-reported common menopausal symptoms or E2 with CIMT or CAC. Results of this study support previous findings which suggest that symptoms do not predict atherosclerotic burden in early menopause.. The KEEPS cohort will provide an opportunity to further assess whether persistence and severity of menopausal symptoms relate to progression of atherosclerosis and if MHT modulates such a relationship.

Acknowledgments

Funding/Support:

The core KEEPS was funded by the Phoenix-based Kronos Longevity Research Institute, which is supported by the not-for-profit Aurora Foundation and carried out at nine U.S. academic medical centers. Other support included the National Institutes of Health (NIH) HL90639 to VMM, 1 UL1 RR024150, Mayo CTSA 1 UL1 RR024150, the Mayo Foundation, Brigham and Women's Hospital/Harvard Medical School CTSA, CTSA UL1 RR024139 and UCSF CTSA UL1 RR024131 from the National Center for Advancing Translational Sciences (NCATS), a component of the National Institutes of Health (NIH) and NIH Roadmap for Medical Research. The manuscript's contents are solely the responsibility of the authors and do not necessarily represent the official view of NCATS or NIH. Information on NCCR is available at <http://www.ncrr.nih.gov>. Bayer HealthCare Pharmaceuticals, Inc. supplied the CLIMARA[®] estradiol and placebo patches and Abbott Laboratories (formerly Solvay Pharmaceuticals) provided the micronized progesterone (PROMETRIUM[®]) and placebo capsules.

Heping Zhang and Yunxiao were partially supported by NIH grants R01DA016750 and U10HD55925.

Role of the Sponsors: The Aurora Foundation did not have input into the design or conduct of the study or the review or approval of this article.

KEEPs Investigators and Staff:

Albert Einstein College of Medicine: Genevieve Neal-Perry, Ruth Freeman, Hussein Amin, Barbara Isaac, Maureen Magnani, Rachel Wildman

Brigham and Women's Hospital/Harvard Medical School: JoAnn Manson, Maria Bueche, Marie Gerhard-Herman, Kate Kalan, Jan Lieson, Kathryn M. Rexrode, Barbara Richmond, Frank Rybicki, Brian Walsh

Columbia College of Physicians and Surgeons: Rogerio Lobo, Luz Sanabria, Maria Soto, Michelle P. Warren, Ralf C. Zimmerman

Kronos Longevity Research Institute: S. Mitchell Harman, Mary Dunn, Panayiotis D. Tsitouras, Viola Zepeda

Mayo Clinic: Virginia M. Miller, Philip A. Araoz, Rebecca Beck, Dalene Bott-Kitslaar, Sharon L. Mulvagh, Lynne T. Shuster, Teresa G. Zais

University of California, Los Angeles, CAC Reading Center: Matthew Budoff, Chris Dailing, Yanlin Gao, Angel Solano

University of California, San Francisco Medical Center: Marcelle I. Cedars, Nancy Jancar, Jean Perry, Rebecca S. Wong, Robyn Pearl, Judy Yee, Brett Elicker, Gretchen A. W. Gooding; UCSF Statistical Reading Center: Dennis Black, Lisa Palermo

University of Southern California, Atherosclerosis Research Unit/Core Imaging and Reading Center: Howard N. Hodis, Yanjie Li, Mingzhu Yan

University of Utah School of Medicine: Eliot Brinton, Paul N. Hopkins, M. Nazeem Nanjee, Kirtly Jones, Timothy Beals, Stacey Larrinaga-Shum

VA Puget Sound Health Care System and University of Washington School of Medicine: George Merriam, Pamela Asbery, SueAnn Brickle, Colleen Carney, Molly Carr, Monica Kletke, Lynna C. Smith

Yale University, School of Medicine: Hugh Taylor, Kathryn Czarkowski, Lubna Pal, Linda McDonald, Mary Jane Minkin, Diane Wall, Erin Wolff (now at NIH/NICHHD).

Others: Frederick Naftolin (New York University), Nanette Santoro (University of Colorado)

References

1. Bush T, Cowan L, Barrett-Connor E, Criqui M, Karon J, Wallace R, et al. Estrogen use and all-cause mortality. Preliminary results from the Lipid Research Clinics Program Follow-Up Study. *JAMA*. 1983; 249:903–6. [PubMed: 6823043]
2. Bush T, Barrett-Connor E, Cowan L, Criqui M, Wallace R, Suchindran C, et al. Cardiovascular mortality and noncontraceptive use of estrogen in women: results from the Lipid Research Clinics Program Follow-up Study. *Circulation*. 1987; 75:1102–9. [PubMed: 3568321]
3. Ettinger B, Friedman G, Bush T, Quesenberry CJ. Reduced mortality associated with long-term postmenopausal estrogen therapy. *Obstet Gynecol*. 1996; 87:6–12. [PubMed: 8532268]
4. Grodstein F, Stampfer M, Colditz G, Willett W, Manson J, Joffe M, et al. Postmenopausal hormone therapy and mortality. *N Engl J Med*. 1997; 336:1769–75. [PubMed: 9187066]
5. Grodstein F, Manson J, Colditz G, Willett W, Speizer F, Stampfer M. A prospective, observational study of postmenopausal hormone therapy and primary prevention of cardiovascular disease. *Ann Intern Med*. 2000; 133:933–41. [PubMed: 11119394]
6. Stampfer M, Colditz G. Estrogen replacement therapy and coronary heart disease: a quantitative assessment of the epidemiologic evidence. *Prev Med*. 1991; 20:47–63. [PubMed: 1826173]
7. Stampfer M, Colditz G, Willett W, Manson J, Rosner B, Speizer F, et al. Postmenopausal estrogen therapy and cardiovascular disease. Ten-year follow-up from the nurses' health study. *N Engl J Med*. 1991; 325:756–62. [PubMed: 1870648]
8. Wolf P, Madans J, Finucane F, Higgins M, Kleinman J. Reduction of cardiovascular disease-related mortality among postmenopausal women who use hormones: evidence from a national cohort. *Am J Obstet Gynecol*. 1991; 164:489–94. [PubMed: 1992690]
9. Manson J, Hsia J, Johnson K, Rossouw J, Assaf A, Lasser N, et al. Estrogen plus progestin and the risk of coronary heart disease. *N Engl J Med*. 2003; 349:523–34. [PubMed: 12904517]
10. Hodis H, Mack W, Azen S, Lobo R, Shoupe D, Mahrer P, et al. Hormone therapy and the progression of coronary-artery atherosclerosis in postmenopausal women. *N Engl J Med*. 2003; 349:535–45. [PubMed: 12904518]

11. Miller V, Black D, Brinton E, Budoff M, Cedars M, Hodis H, et al. Using basic science to design a clinical trial: baseline characteristics of women enrolled in the Kronos Early Estrogen Prevention Study (KEEPS). *J Cardiovasc Transl Res.* 2009; 2:228–39. [PubMed: 19668346]
12. Gast GC, Grobbee DE, Pop VJ, Keyzer JJ, Wijnands-van Gent CJ, Samsioe GN, et al. Menopausal complaints are associated with cardiovascular risk factors. *Hypertension.* 2008; 51:1492–8. [PubMed: 18391100]
13. Huang AJ, Sawaya GF, Vittinghoff E, Lin F, Grady D. Hot flushes, coronary heart disease, and hormone therapy in postmenopausal women. *Menopause.* 2009; 16:639–43. [PubMed: 19325499]
14. Thurston RC, Sutton-Tyrrell K, Everson-Rose SA, Hess R, Matthews KA. Hot flashes and subclinical cardiovascular disease: findings from the Study of Women's Health Across the Nation Heart Study. *Circulation.* 2008; 118:1234–40. [PubMed: 18765392]
15. Allison M, Manson J, Aragaki A, Langer R, Rossouw J, Curb D, et al. Vasomotor symptoms and coronary artery calcium in postmenopausal women. *Menopause.* 2010
16. Gast GC, Pop VJ, Samsioe GN, Grobbee DE, Nilsson PM, Keyzer JJ, et al. Vasomotor menopausal symptoms are associated with increased risk of coronary heart disease. *Menopause.* 2011; 18:146–51. [PubMed: 21127438]
17. Szmuiowicz ED, Manson JE, Rossouw JE, Howard BV, Margolis KL, Greep NC, et al. Vasomotor symptoms and cardiovascular events in postmenopausal women. *Menopause.* 2011
18. Harman S, Brinton E, Cedars M, Lobo R, Manson J, Merriam G, et al. KEEPS: The Kronos Early Estrogen Prevention Study. *Climacteric.* 2005; 8:3–12. [PubMed: 15804727]
19. O'Leary DH, Bryan FA, Goodison MW, Rifkin MD, Gramiak R, Ball M, et al. Measurement variability of carotid atherosclerosis: real-time (B-mode) ultrasonography and angiography. *Stroke.* 1987; 18:1011–7. [PubMed: 3317999]
20. Mack WJ, Selzer RH, Hodis HN, Erickson JK, Liu CR, Liu CH, et al. One-year reduction and longitudinal analysis of carotid intima-media thickness associated with colestipol/niacin therapy. *Stroke.* 1993; 24:1779–83. [PubMed: 8248954]
21. Hodis HN, Mack WJ, LaBree L, Selzer RH, Liu C, Alaupovic P, et al. Reduction in carotid arterial wall thickness using lovastatin and dietary therapy: a randomized controlled clinical trial. *Ann Intern Med.* 1996; 124:548–56. [PubMed: 8597317]
22. Hodis HN, Mack WJ, LaBree L, Mahrer PR, Sevanian A, Liu CR, et al. Alpha-tocopherol supplementation in healthy individuals reduces low-density lipoprotein oxidation but not atherosclerosis: the Vitamin E Atherosclerosis Prevention Study (VEAPS). *Circulation.* 2002; 106:1453–9. [PubMed: 12234947]
23. Hodis HN, Mack WJ, Dustin L, Mahrer PR, Azen SP, Detrano R, et al. High-dose B vitamin supplementation and progression of subclinical atherosclerosis: a randomized controlled trial. *Stroke.* 2009; 40:730–6. [PubMed: 19118243]
24. Hodis HN, Mack WJ, Kono N, Azen SP, Shoupe D, Hwang-Levine J, et al. Isoflavone Soy Protein Supplementation and Atherosclerosis Progression in Healthy Postmenopausal Women: A Randomized Controlled Trial. *Stroke.* 2011; 42:3168–75. [PubMed: 21903957]
25. Hodis HN, Mack WJ, Lobo RA, Shoupe D, Sevanian A, Mahrer PR, et al. Estrogen in the prevention of atherosclerosis. A randomized, double-blind, placebo-controlled trial. *Ann Intern Med.* 2001; 135:939–53. [PubMed: 11730394]
26. Hodis HN, Mack WJ, Zheng L, Li Y, Torres M, Sevilla D, et al. Effect of peroxisome proliferator-activated receptor gamma agonist treatment on subclinical atherosclerosis in patients with insulin-requiring type 2 diabetes. *Diabetes Care.* 2006; 29:1545–53. [PubMed: 16801577]
27. Xiang AH, Peters RK, Kjos SL, Ochoa C, Marroquin A, Goico J, et al. Effect of thiazolidinedione treatment on progression of subclinical atherosclerosis in premenopausal women at high risk for type 2 diabetes. *J Clin Endocrinol Metab.* 2005; 90:1986–91. [PubMed: 15623809]
28. Hodis HN, Mack WJ, LaBree L, Selzer RH, Liu CR, Liu CH, et al. The role of carotid arterial intima-media thickness in predicting clinical coronary events. *Ann Intern Med.* 1998; 128:262–9. [PubMed: 9471928]
29. Selzer RH, Hodis HN, Kwong-Fu H, Mack WJ, Lee PL, Liu CR, et al. Evaluation of computerized edge tracking for quantifying intima-media thickness of the common carotid artery from B-mode ultrasound images. *Atherosclerosis.* 1994; 111:1–11. [PubMed: 7840805]

30. Selzer RH, Mack WJ, Lee PL, Kwong-Fu H, Hodis HN. Improved common carotid elasticity and intima-media thickness measurements from computer analysis of sequential ultrasound frames. *Atherosclerosis*. 2001; 154:185–93. [PubMed: 11137099]
31. Mack WJ, LaBree L, Liu C, Selzer RH, Hodis HN. Correlations between measures of atherosclerosis change using carotid ultrasonography and coronary angiography. *Atherosclerosis*. 2000; 150:371–9. [PubMed: 10856529]
32. Mack WJ, Dhungana B, Dowsett SA, Keech CA, Feng M, Li Y, et al. Carotid artery intima-media thickness after raloxifene treatment. *J Womens Health (Larchmt)*. 2007; 16:370–8. [PubMed: 17439382]
33. Xiang AH, Hodis HN, Kawakubo M, Peters RK, Kjos SL, Marroquin A, et al. Effect of pioglitazone on progression of subclinical atherosclerosis in non-diabetic premenopausal Hispanic women with prior gestational diabetes. *Atherosclerosis*. 2008; 199:207–14. [PubMed: 18054942]
34. Baker JV, Henry WK, Patel P, Bush TJ, Conley LJ, Mack WJ, et al. Progression of carotid intima-media thickness in a contemporary human immunodeficiency virus cohort. *Clin Infect Dis*. 2011; 53:826–35. [PubMed: 21860012]
35. Currier JS, Kendall MA, Zackin R, Henry WK, Alston-Smith B, Torriani FJ, et al. Carotid artery intima-media thickness and HIV infection: traditional risk factors overshadow impact of protease inhibitor exposure. *AIDS*. 2005; 19:927–33. [PubMed: 15905673]
36. Kaplan RC, Kingsley LA, Gange SJ, Benning L, Jacobson LP, Lazar J, et al. Low CD4+ T-cell count as a major atherosclerosis risk factor in HIV-infected women and men. *AIDS*. 2008; 22:1615–24. [PubMed: 18670221]
37. Künzli N, Jerrett M, Mack WJ, Beckerman B, LaBree L, Gilliland F, et al. Ambient air pollution and atherosclerosis in Los Angeles. *Environ Health Perspect*. 2005; 113:201–6. [PubMed: 15687058]
38. Azen SP, Qian D, Mack WJ, Sevanian A, Selzer RH, Liu CR, et al. Effect of supplementary antioxidant vitamin intake on carotid arterial wall intima-media thickness in a controlled clinical trial of cholesterol lowering. *Circulation*. 1996; 94:2369–72. [PubMed: 8921775]
39. Vigen C, Hodis HN, Chandler WL, Lobo RA, Mack WJ. Postmenopausal oral estrogen therapy affects hemostatic factors, but does not account for reduction in the progression of subclinical atherosclerosis. *J Thromb Haemost*. 2007; 5:1201–8. [PubMed: 17389005]
40. Vigen C, Hodis HN, Selzer RH, Mahrer PR, Mack WJ. Relation of progression of coronary artery atherosclerosis to risk of cardiovascular events (from the Monitored Atherosclerosis Regression Study). *Am J Cardiol*. 2005; 95:1277–82. [PubMed: 15904629]
41. Karim R, Hodis H, Stanczyk F, Lobo R, Mack W. Relationship between serum levels of sex hormones and progression of subclinical atherosclerosis in postmenopausal women. *J Clin Endocrinol Metab*. 2008; 93:131–8. [PubMed: 17925335]
42. Chen YC, Guo X, Raffel LJ, Xiang AH, Fang B, Hsueh WA, et al. Carotid intima-media thickness (cIMT) cosegregates with blood pressure and renal function in hypertensive Hispanic families. *Atherosclerosis*. 2008; 198:160–5. [PubMed: 18028933]
43. Hodis HN, St John JA, Xiang M, Cushman M, Lobo RA, Mack WJ. Inflammatory markers and progression of subclinical atherosclerosis in healthy postmenopausal women (from the Estrogen in the Prevention of Atherosclerosis Trial). *Am J Cardiol*. 2008; 101:1131–3. [PubMed: 18394446]
44. Xiang AH, Azen SP, Buchanan TA, Raffel LJ, Tan S, Cheng LS, et al. Heritability of subclinical atherosclerosis in Latino families ascertained through a hypertensive parent. *Arterioscler Thromb Vasc Biol*. 2002; 22:843–8. [PubMed: 12006400]
45. Goodarzi MO, Taylor KD, Guo X, Quiñones MJ, Cui J, Li Y, et al. Association of the diabetes gene calpain-10 with subclinical atherosclerosis: the Mexican-American Coronary Artery Disease Study. *Diabetes*. 2005; 54:1228–32. [PubMed: 15793266]
46. Wang D, Yang H, Quiñones MJ, Bulnes-Enriquez I, Jimenez X, De La Rosa R, et al. A genome-wide scan for carotid artery intima-media thickness: the Mexican-American Coronary Artery Disease family study. *Stroke*. 2005; 36:540–5. [PubMed: 15692111]
47. Wendelhag I, Gustavsson T, Suurkula M, Berglund G, Wikstrand J. Ultrasound measurement of wall thickness in the carotid artery: fundamental principles and description of a computerized analysing system. *Clin Physiol*. 1991; 11:565–77. [PubMed: 1769190]

48. Wilson P, D'Agostino R, Levy D, Belanger A, Silbershatz H, Kannel W. Prediction of coronary heart disease using risk factor categories. *Circulation*. 1998; 97:1837–47. [PubMed: 9603539]
49. Agatista P, Matthews K, Bromberger J, Edmundowicz D, Chang Y, Sutton-Tyrrell K. Coronary and aortic calcification in women with a history of major depression. *Arch Intern Med*. 2005; 165:1229–36. [PubMed: 15956001]
50. Matthews K, Chang Y, Sutton-Tyrrell K, Edmundowicz D, Bromberger J. Recurrent Major Depression Predicts Progression of Coronary Calcification in Healthy Women: Study of Women's Health Across the Nation. *Psychosom Med*. 2010
51. Jones D, Bromberger J, Sutton-Tyrrell K, Matthews K. Lifetime history of depression and carotid atherosclerosis in middle-aged women. *Arch Gen Psychiatry*. 2003; 60:153–60. [PubMed: 12578432]
52. Stein PK, Barzilay JI. Relationship of Abnormal Heart Rate Turbulence and Elevated CRP to Cardiac Mortality in Low, Intermediate, and High-Risk Older Adults. *J Cardiovasc Electrophysiol*. 2011; 22:122–7. [PubMed: 21134026]
53. Szmuiłowicz ED, Manson JE. Menopausal vasomotor symptoms and cardiovascular disease. *Menopause*. 2011
54. Thurston RC, Kuller LH, Edmundowicz D, Matthews KA. History of hot flashes and aortic calcification among postmenopausal women. *Menopause*. 2010; 17:256–61. [PubMed: 20042895]
55. Thurston RC, Sutton-Tyrrell K, Everson-Rose SA, Hess R, Powell LH, Matthews KA. Hot flashes and carotid intima media thickness among midlife women. *Menopause*. 2011

Table 1

Baseline Characteristics of Subjects Screened for the KEEPS Trial

Variable (n=868 women)	Average	Range	SD*
Demographics			
Age of women screened (years)	52.7	42–58	2.6
Weight (kg)	68.8	42–104	11.7
Number of years in menopause	1.4	0.5–3.0	0.7
FSH (ng/ml)	87.6	6.3–194	33.4
E2 (pg/ml)	23.5	2–298	21.9
Drinks per month	11.9	0–122	17.9

	(N)
Marital Status	
Married	538
Never Married	153
Divorced	16
Separated	24
Partner-not married	24
Widowed	105
Race/Ethnicity	
Caucasian/White	671
Black/African American	65
Hispanic	57
Asian/Indian	13
Chinese	9
Philipino	4
Japanese	1
Korean	1
Other Asian	1
Decline to answer	38
Education	
Grade School	4
Some High School	3
High School Diploma or GED	67
Some College/Vocational	172
College Graduate	337
Some Graduate/Professional School	40
Graduate or Professional Degree	228
Declined to Answer	17
Employment Status	
Disabled	572
Employed full-time	130
Homemaker	66

	(N)
Employed part-time	35
Not working	28
Retired	21
Not answered	16
Hypertension	
Yes	75
No	784

Variable (n=868 women)	Average	Range	SD*
Baseline Atherosclerosis Measurements			
CIMT (mm)	0.721	0.532–1.168	0.1
Mean Agatston score	9.3	0–1091	54.3
Mean Agatston Vol	8.2	0–720.750	41.9
Baseline Menopausal Symptom Scores			
Depression score	0.483	0–3	
Insomnia score	1.096	0–3	
Irritability score	0.793	0–3	
Dyspareunia score	0.915	0–4	
Hot flashes score	1.362	0–3	
Mood swings score	0.764	0–3	
Night sweats score	1.096	0–3	
Palpitations score	0.349	0–3	
Vaginal dryness score	0.933	0–3	

Table 2

In a Logistic Regression Model, Menopausal Symptoms Do Not Predict CAC Scores

Variable (n=822)	Coefficient Estimate *	Std. Error	P
Hot flashes	-0.032	0.145	0.825
Dyspareunia	0.030	0.073	0.684
Vaginal dryness	0.008	0.108	0.944
Night sweats	-0.019	0.128	0.884
Palpitations	-0.298	0.176	0.091
Mood swings	0.126	0.189	0.505
Depression	0.280	0.155	0.070
Insomnia	0.044	0.113	0.696
Irritability	-0.244	0.188	0.194

* Log of Odd Ratio

Table 3

In a Linear Regression Model, Menopausal Symptoms Do Not Predict CIMT Measurements

Variable (n=759)	Coefficient Estimate *	Std. Error	P
HOTFLASH	0.004	0.005	0.419
DYSP	-0.002	0.003	0.392
VAGDRY	-0.003	0.004	0.393
SWEATS	0.002	0.005	0.691
PALPIT	-0.001	0.006	0.817
MOODSW	-1.2E-04	0.007	0.985
DEPR	-0.003	0.006	0.58
INSOM	-0.001	0.004	0.861
IRRIT	0.002	0.007	0.806
E2	-7.0E-06	1.7E-04	0.967

*
Log of Odd Ratio

Table 4

Correlation of Menopausal Symptoms, Estradiol and Atherosclerosis Measures with Time Since Menopause

Variable	Correlation coefficient (CC)	P
Hot flashes	0.016	0.651
Dyspareunia	0.084	0.015
Vaginal dryness	0.151	1.0e-5
Night sweats	-0.059	0.090
Palpitations	0.031	0.365
Mood swings	-0.032	0.357
Depression	0.051	0.142
Insomnia	0.019	0.595
Irritability	-0.031	0.373
Mean Agatston	-0.006	0.857
Mean Agatston Vol	-0.006	0.857
CIMT	0.075	0.029
E2	-0.084	0.015