

## FEATURE REVIEW

# A role for synaptic plasticity in the adolescent development of executive function

LD Selemon

Adolescent brain maturation is characterized by the emergence of executive function mediated by the prefrontal cortex, e.g., goal planning, inhibition of impulsive behavior and set shifting. Synaptic pruning of excitatory contacts is the signature morphologic event of late brain maturation during adolescence. Mounting evidence suggests that glutamate receptor-mediated synaptic plasticity, in particular long term depression (LTD), is important for elimination of synaptic contacts in brain development. This review examines the possibility (1) that LTD mechanisms are enhanced in the prefrontal cortex during adolescence due to ongoing synaptic pruning in this late developing cortex and (2) that enhanced synaptic plasticity in the prefrontal cortex represents a key molecular substrate underlying the critical period for maturation of executive function. Molecular sites of interaction between environmental factors, such as alcohol and stress, and glutamate receptor mediated plasticity are considered. The accentuated negative impact of these factors during adolescence may be due in part to interference with LTD mechanisms that refine prefrontal cortical circuitry and when disrupted derail normal maturation of executive function. Diminished prefrontal cortical control over risk-taking behavior could further exacerbate negative outcomes associated with these behaviors, as for example addiction and depression. Greater insight into the neurobiology of the adolescent brain is needed to fully understand the molecular basis for heightened vulnerability during adolescence to the injurious effects of substance abuse and stress.

*Translational Psychiatry* (2013) 3, e238; doi:10.1038/tp.2013.7; published online 5 March 2013

### Adolescent development of executive function

Adolescence is rather inexactly defined as the period beginning with the onset of puberty and ending with the shouldering of adult responsibilities.<sup>1</sup> It is a time of increased propensity to engage in risky behaviors that include experimentation with alcohol, tobacco, drugs and sexual behavior. Dahl<sup>1</sup> has called the adolescent brain a 'natural tinderbox' because gonadal hormones are actively stimulating affective and appetitive behaviors, such as sexual drive, increased emotional intensity, and risk taking, yet the brain systems that regulate and moderate these emotional and appetitive urges are not yet mature.

The prefrontal cortex (PFC) mediates executive functions, i.e., internally guided behavior, goal planning, and impulse control, that form the essence of rational thinking and serve to counter appetitive urges and check risk-taking behavior.<sup>2,3</sup> The PFC is the last brain region to mature,<sup>4-7</sup> and therefore not surprisingly the frontal lobe capacity for internally guided behavior, working memory, and organizational skills do not reach full adult functional capacity until mid to late adolescence.<sup>8-12</sup>

Crews *et al.*<sup>13</sup> have drawn parallels between adolescence and early sensory critical periods, which are dependent on plasticity of developing sensory connectivity and allow for environmental (sensory) modulation of maturing sensory connections. Specifically, they have suggested that in

adolescence PFC circuitry may be endowed with similar plasticity and responsiveness to environmental factors, and as a consequence with heightened vulnerability to the detrimental effects of substance abuse and stress.<sup>13</sup>

This review examines the literature on adolescent development across species and focuses on the role that glutamate-receptor mediated plasticity may play in maturation of PFC circuitry in adolescence. It is postulated that adolescence represents a phase of increased activity of long term depression (LTD) mechanisms that predispose to synaptic elimination and further that termination of this LTD-permissive phase marks the transition to adulthood. Finally, consideration is given to the possibility that greater vulnerability to substances of abuse and stress may represent an interaction between these environmental factors and the LTD mechanisms of plasticity that are accentuated during adolescence. The hypothesis put forward in this review, while speculative, is intended to spark further research into possible molecular mechanisms associated with adolescent development of the PFC. Certainly synaptic plasticity has been studied much less extensively in the PFC than in the hippocampus; nonetheless, mounting evidence suggests that both long term potentiation (LTP) and LTD play an important role in cognitive functioning mediated by the PFC and perhaps when disturbed in diseases related to malfunction of this cortex.<sup>14</sup>

Department of Neurobiology, Yale University School of Medicine, New Haven, CT, USA

Correspondence: Dr LD Selemon, Department of Neurobiology, PO Box 208001, New Haven, CT 06520-8001, USA. E-mail: ldselemon@aol.com

**Keywords:** alcohol; depression; dopamine; long term depression (LTD); prefrontal cortex; substance abuse

Received 4 September 2012; revised 15 December 2012; accepted 3 January 2013

## Preadolescent development and sensory critical periods

The specificity and topography of brain wiring are not entirely genetically preprogrammed but instead established via dynamic processes occurring in the developing brain. Adolescence represents the final epoch in a series of developmental stages that transform the immature brain into its adult form. In order to fully understand adolescent development, it is important to appreciate how it differs from earlier preadolescent maturation.

The developmental mechanisms that account for major remodeling of connectivity occur before the onset of adolescence, i.e., before postnatal day 28 (PD28) in rodents, 9 months in cats, and 3 years in non-human primates<sup>15–17</sup> and include prominent degeneration of neurons and axons.<sup>18,19</sup> Indeed, the immature mammalian brain is distinguished from its adult counterpart by the presence of connections between brain areas that are not interconnected in the mature brain and by overlap of terminal fields that are segregated in the adult brain. For example, in the newborn hamsters and rats, uncrossed retinocollicular projections, i.e. from the retina to the ipsilateral superior colliculus (SC), not only occupy a much expanded territory in the SC relative to that of the adult brain but also originate from nasal as well as temporal retinal ganglion cells.<sup>20–22</sup> Retraction of the terminal projections is associated with loss of these nasal, ipsilaterally-projecting ganglion cells.<sup>22</sup> More generally, in the central nervous system overproduction of neurons with ensuing neuronal death is a common mechanism employed by the developing brain to ensure that the appropriate balance of projection and receptive neurons is attained.<sup>19,23–25</sup>

A second, pervasive form of degeneration in the developing brain is degeneration limited to axonal connections leaving the neurons of origin intact. For instance, in the central nervous system, cortical callosal projections that are widespread in kittens and young rats are constricted to the adult patterning by retraction of callosal axons without cell loss.<sup>26–28</sup> Quantitative analysis of axon number in major tracts underscores the magnitude of this form of degeneration as the number of axons in the young non-human primate brain ranges from twice (optic tract) to 3.5 times (corpus callosum) the number in the adult brain.<sup>29–31</sup> Both forms of degeneration, involving loss of neurons or loss of axons, necessarily are associated with dissolution of established synapses.<sup>32</sup> However, these early developmental events are occurring at a time when, overall, synapses are increasing in density.<sup>33–38</sup> The classic example or early connective remodeling, that of reduction of polyneuronal input on a single muscle fiber to a single axon, illustrates how synaptic number may increase as the surviving single axon sprouts a much more elaborate terminal plexus.<sup>18,39</sup> Likewise, in the central nervous system regression of inappropriate synapses is more than compensated by growth and expansion of appropriate terminal fields.<sup>40</sup>

A wealth of evidence has established that reorganization of connections throughout the brain is activity dependent and therefore mediated by a Hebbian mechanism.<sup>41–45</sup> Although normal regression of connections in the visual system can proceed in the absence of visual input<sup>41</sup> there exists a period of plasticity during postnatal development that permits rewiring in response to altered sensory environments.<sup>43,46,47</sup>

It is noteworthy that critical periods for sensory plasticity occur in the same preadolescent period in which remodeling of connectivity occurs.<sup>34,48,49</sup>

## Adolescence: synaptic elimination and excitatory/inhibitory balance

The maturational event most consistently linked to the adolescent stage of development is reduction of synaptic density or 'synaptic pruning.' Quantitative analyses of synapses in the non-human primate uncovered a synchronous increase in synaptic density in multiple cortical areas that peaks during the postnatal 3rd month, declines slowly (10%) until ~2 years of age with a steeper decline (40%) occurring between 2.7 and 5 years (adulthood).<sup>35–38</sup> In the human cortex the timing of peak synaptic density is staggered in different regions, but the basic pattern of peak synaptic density in early childhood followed by robust synapse elimination throughout early (auditory cortex) or mid-adolescence (PFC) is in basic agreement with non-human primate studies.<sup>4,50</sup> More recent data have established that synapse elimination in humans does not end in adolescence but continues at a lower rate into early adulthood.<sup>51</sup> In addition, in human cortex the synaptic related proteins synaptophysin and postsynaptic density protein-95 (PSD-95) show similar patterns of peak in childhood and decline through adolescence,<sup>52</sup> though it should be noted that a recent study found increasing concentrations of synaptic-related molecules throughout the adolescent epoch.<sup>53</sup> Nonetheless, most evidence points to synaptic pruning as the signature late maturational process associated with adolescence. Other species have been studied less extensively but exhibit a comparable pattern. Peak synaptic density was observed by the 7th postnatal week in the cat.<sup>34</sup> In rat, recent data suggest that peak spine density in the PFC is present at PD31 with spine density decreasing thereafter until PD 57 or PD60, i.e., early adulthood.<sup>33</sup>

Synaptic elimination in adolescence is widely thought to account for the decline in gray matter volume detected via longitudinal magnetic resonance imaging (MRI) of human subjects. Although reduction of synaptic connectivity might be accompanied by retraction of glial and neuronal processes, elimination of neuronal cell bodies occurs much earlier in development.<sup>54</sup> One of the first longitudinal MRI studies of human subjects detected divergent developmental growth patterns in gray and white matter volumes: white matter volume increased linearly until approximately age 22 whereas cortical gray matter volume in the frontal and parietal lobes peaked just prior to adolescence (~10–12 years) and then declined to adult volumes.<sup>5</sup> Cross-sectional studies of children and adolescents, including one recent large multi-center study, also show opposing patterns for gray and white matter.<sup>55–57</sup> Interestingly, changing cortical volumes over this age range are most prominent in the frontal and parietal lobes.<sup>8,58,59</sup> Indeed, a recent study indicates that there is a progression in which higher cortical association areas like the PFC are last to show reduction of gray matter volume.<sup>7</sup>

The functional significance of synaptic elimination during adolescence, though still enigmatic, probably involves adjustment of the excitatory/inhibitory balance on individual neurons

and within networks. The main argument in support of this hypothesis stems from the specificity of the loss: excitatory synapses are selectively degenerated whereas inhibitory synapses are spared.<sup>35,37</sup> Even loss of chandelier axon boutons in the PFC, a finding that was originally interpreted as loss of inhibitory synapses,<sup>60</sup> now supports the elimination of excitatory input in light of new physiologic data.<sup>61</sup> Furthermore, recent evidence has established that D2 dopamine receptors on interneurons undergo a profound maturational change during adolescence.<sup>62–64</sup> Prior to adolescence, D2 stimulation elicits either no effect or only weak inhibition on interneurons. However, in adult animals stimulation of D2 receptors is strongly excitatory and therefore results in robust firing of interneurons and potent inhibition of their pyramidal cell targets. As a result, inhibition gains a position of ascendancy in adolescence via increased dopamine-mediated firing of interneurons as well as a relative gain in inhibitory/excitatory synapse ratio. In the PFC, neurophysiologic studies have established a critical role for inhibitory synapses in mediating information flow through local networks.<sup>65,66</sup> Moreover, fast-spiking interneurons mediate gamma oscillations which are essential to cortical computation in many areas of the cortex and to cognitive processing in the PFC.<sup>67,68</sup> Thus, the correct balance of inhibition and excitation seems to be critical for normative executive function, and conversely, disturbance of this balance is thought to be a fundamental component of psychiatric illness.<sup>69,70</sup>

### Molecular mechanisms associated with synaptic stabilization and synaptic pruning

Synapse stabilization and synapse elimination are primary players in the maturational processes associated with preadolescent and adolescent development. The transition of nascent synapses into mature synapses represents the first step in synapse stabilization. N-methyl-D-aspartate receptors (NMDAR) are localized very early to the postsynaptic membrane, but transition to a more mature synapse state is characterized by recruitment of alpha-amino-3-hydroxy-5-methyl-4-isoxazole propionic acid receptors (AMPA) to the synapse.<sup>71–74</sup> Expression of AMPARs on the postsynaptic membrane is induced by NMDAR-mediated long term potentiation (LTP), the same mechanism originally described in hippocampus for learning and memory.<sup>73–76</sup> A second NMDAR-mediated process, LTD, results when afferent stimulation fails to activate a target neuron.<sup>76</sup> In many respects LTP and LTD are opposite processes although they engage distinct intracellular signaling mechanisms.<sup>77–80</sup> Essentially stimulation of NMDARs can induce activity-dependent strengthening of synapses via LTP or weakening via LTD, and AMPAR insertion or removal from the postsynaptic membrane is the conduit for this change in synaptic strength.<sup>81,82</sup> Importantly, LTP and LTD do not just strengthen or weaken synaptic connections (short term plasticity) but actually trigger the addition or loss of synapses (long term plasticity) even in the adult brain.<sup>83–92</sup>

Well before the role of NMDAR-mediated LTP in establishing mature synaptic connections was recognized, Constantine-Paton *et al.*<sup>44</sup> postulated that activity-dependent remodeling of connectivity in the developing brain might be

mediated by NMDARs because these receptors are perfectly suited to detection of synchronized pre- and postsynaptic activation. Growing evidence now supports the idea that LTP and LTD are required for generation of whisker barrel field maps in the primary somatosensory cortex and ocular dominance columns in the primary visual cortex, both involving reorganization of thalamic inputs to layer 4.<sup>49,93–96</sup> In development, as in learning and memory, plasticity is bidirectional, i.e., synchronized activity of afferent inputs may trigger LTP and resulting synapse maturation and stabilization; conversely, asynchronous activity may diminish synaptic strength via LTD and predispose the synapse to elimination.<sup>97</sup>

Recently, changes in the NMDAR have been linked to critical periods of early developmental plasticity. NMDAR subunit composition shifts from NR2B predominant to NR2A prevalent forms in early development in the visual and somatosensory cortices.<sup>98–100</sup> Moreover, the shift in NR2B to NR2A shows a rough correspondence to critical periods for sensory plasticity: the beginning of the critical period is marked by an increase in NR2A expression, and the end of the critical period is associated with a decrease in NR2B expression.<sup>100,101</sup> Importantly, the switch is not locked to a specific age but in fact can be delayed by sensory deprivation, suggesting that it is controlled by activity.<sup>93,102–105</sup> In turn, the changeover from NR2B to NR2A receptor subtypes controls the sensitivity of these connections to stimulation by NMDARs. For example, in the primary visual cortex of the ferret, NR2B levels are already high at eye opening and decline in layer 4 at the end of the critical period for plasticity of ocular dominance columns but remain high in layer 2/3.<sup>106</sup> Correspondingly, physiologic studies in the cat visual cortex have shown that cortical layer 4 neurons, but not layer 2/3 cells, exhibit reduced sensitivity of visual and spontaneous activity to an NMDAR antagonist at the end of the critical period.<sup>107</sup> Together these findings suggest that the switch from NR2B to NR2A dominated receptors terminates the critical period of experience-dependent plasticity for establishment of ocular dominance columns in the visual cortex.

NMDAR-mediated LTP and LTD may also constitute the molecular underpinnings for adolescent synaptic pruning, albeit with greater emphasis on LTD and synaptic elimination. How could the same mechanism account for two very different developmental processes? Perhaps the period of adolescence corresponds to a widespread shift in the balance of LTP/LTD mechanisms and a corresponding prevalence of synaptic elimination over synaptic addition. In rat hippocampal slices, an increased NR2A/NR2B ratio has been linked to decreased spine motility and increased synaptic stabilization, suggesting a role in the NMDAR subunit composition in halting synaptogenesis.<sup>108</sup> Furthermore, the NR2A state is less conducive to LTP. This is because calcium/calmodulin-dependent protein kinase II (CaMKII), which has a well established role in LTP,<sup>109,110</sup> binds preferentially to the NR2B subunit.<sup>110–112</sup> Accordingly, NR2B expression on the postsynaptic membrane has been shown to be necessary for LTP induction, while a role for NR2A in LTP is not well established.<sup>113–115</sup> Moreover, NR2A expression is enhanced by ligand binding to NMDARs and therefore is activity-modulated whereas NR2B expression is not dependent on previous activity.<sup>116</sup> The NR2A subunit is therefore thought to

be responsible for metaplasticity of synapses, i.e., a change in the likelihood of subsequent synaptic plasticity.<sup>117,118</sup> With age and activity, NR2A subunits become incorporated into the postsynaptic membrane, replacing NR2B subunits.<sup>116</sup> The resulting increased NR2A/NR2B ratio translates into a higher threshold for induction of LTP and conversely a state that is more favorable for induction of LTD.<sup>118,119</sup>

The role of plasticity in the neocortex is not as well established as in the hippocampus. However, NMDAR-mediated LTP and LTD have been described in the visual neocortex<sup>120</sup> and at multiple synapses in the PFC.<sup>121–123</sup> Notably, LTD mediated through metabotropic glutamate receptors (mGluRs) has emerged as a major alternative to NMDAR-mediated LTD in widespread areas of the brain<sup>124–126</sup> and therefore deserves consideration as a possible molecular basis for synaptic pruning in the PFC. In this regard, mGluR plasticity has been described at the thalamocortical synapse in the somatosensory cortex,<sup>127</sup> perhaps indicating that this form of plasticity is also present at the mediodorsal thalamic synapses in the PFC. However, at the thalamocortical synapse, mGluR LTD acts presynaptically to decrease transmitter release and depress synaptic activity.<sup>127</sup> Such a mechanism would be unlikely to result in synapse loss and spine involution and therefore would not be a strong candidate for LTD-facilitated synaptic pruning during adolescence. Furthermore, mGluR LTD at postsynaptic sites in the hippocampus has been associated with large spines containing an abundance of AMPAR.<sup>128</sup> Unlike the hippocampus where large mushroom spines are in the majority, thin, filopodial spines predominate in the PFC.<sup>129</sup> Thus, strong evidence for mGluR in plasticity related to PFC synaptic pruning is presently lacking; nonetheless, possible involvement of mGluR-mediated LTD in prefrontal adolescent maturation cannot be discounted.

Many questions remain to be answered about the role of metaplasticity in the PFC as well. As the NR2A subtype promotes a LTD-receptive state in the synapse and LTD is associated with synaptic elimination, it would be interesting to know whether and when the NR2B to NR2A switch occurs in the PFC and how it relates to the synaptic pruning that refines connectivity associated with cognitive control of behavior. If the LTD-receptive state is a hallmark of adolescent development, a reasonable presumption is that there exists an additional molecular switch that greatly curtails the LTD-receptive state of adolescence into the much less receptive state of adulthood. This switch, although presently unidentified, would transform the synapse into a state that is less receptive to alterations in AMPAR expression on the postsynaptic membrane. Given that synaptic pruning continues into early adulthood albeit at a lower level than that of adolescence,<sup>33,51</sup> it seems likely that the transition phase is gradual rather than abrupt resulting in a much less plastic state by the end of the third decade in humans.

### Adolescence development of cognitive function and synaptic plasticity

Executive functions governed by the PFC exhibit a prolonged period of maturation reaching fulfillment only in late adolescence.<sup>11,130</sup> Volumetric changes occurring during adolescence have been correlated with improved cognitive performance, e.g., verbal and spatial memory performance is

positively correlated with gray matter thinning in the frontal lobes.<sup>6</sup> General intelligence has also been shown to bear a relationship to the trajectory of frontal cortical gray matter thinning, such that subjects with superior intelligence show a robust early adolescent increase in gray matter volume followed by equally robust thinning during later adolescence.<sup>131</sup> However, too much cortical thinning during adolescence has been associated with diseased states such as Attention Deficit Hyperactivity Disorder (ADHD).<sup>132</sup> Thus, there is an optimal level of synaptic pruning that is essential to normal development of adult cognitive function.

One recent study addressed the role of AMPAR expression and LTD in the development of PFC function in the mouse. Vazdarjanova *et al.*<sup>133</sup> utilized a transgenic mouse that over-expresses calcyon, a protein which mediates activity-dependent AMPAR internalization, and found that calcyon over-expression over the lifetime of the mouse resulted in marked impairment of contextual fear extinction (CFE) and working memory capacity, both dependent on normal PFC function. Most relevant to this discussion, adolescence was the critical period for production of these deficits. When over-expression was silenced specifically during the adolescent epoch, normal CFE function was rescued.<sup>133</sup> One possible explanation for these findings is that AMPAR internalization and associated functions like LTD are more sensitive to regulation during adolescence and this regulation is turned off, or at least greatly diminished, in the adult brain. Whether overactive LTD during adolescence translates into altered synaptic number in the PFC or elsewhere is currently not known. However, it is interesting that upregulated calcyon expression has been found in schizophrenia, a neurodevelopmental disease in which PFC gray matter deficits are prominent.<sup>134–136</sup>

In the PFC, synaptic plasticity is highly modulated by dopamine receptor, especially the D1 receptor.<sup>14,122,137</sup> This is not surprising since D1 receptor stimulation has been shown to trigger phosphorylation of AMPAR, which in turn promotes trafficking of these receptors to the external membrane.<sup>138,139</sup> The D1 receptor is therefore strategically positioned to effect changes in AMPAR synaptic expression and ultimately in synaptic strength and/or number. In the adult non-human primate, long term sensitizing regimens of amphetamine decrease spine density on pyramidal cells in the PFC and have detrimental effects on working memory performance.<sup>140</sup> Moreover, these effects seem to be due to changes at the D1 receptor because both cognitive and morphologic effects on PFC pyramidal neurons can be reversed by long term treatment with a D1 antagonist.<sup>141</sup> If AMPAR-mediated LTD expression is in a state of greater sensitivity to modulation in adolescence, then D1 receptor-stimulated interference with this mechanism could be magnified during adolescence resulting in exaggerated consequences at the synapse. Other known modulators of synaptic plasticity, e.g. D2,<sup>139</sup> muscarinic,<sup>142</sup> and cannabinoid<sup>143</sup> receptors, might have similarly increased potency during the adolescent period.

### Adolescence vulnerability to environmental factors

Adolescence has been described as a period of accentuated opportunity and of enhanced vulnerability.<sup>1</sup> It has long been recognized that early onset of substance abuse is associated

with greater propensity for problem drug use later in life.<sup>144–147</sup> In recent years, the period of adolescent plasticity has been shown to temporally correlate with the time of greatest vulnerability to addiction.<sup>148</sup> Some have postulated that addiction constricts the learning and memory pathways in a maladaptive fashion,<sup>149,150</sup> but the question of why addiction is more devastating in adolescence than in adulthood remains unanswered. Adolescence is also associated with onset of mental illness, as for example depression rates rise in adolescence especially for females,<sup>151</sup> and the prodromal phase of psychosis, including early onset schizophrenia, surfaces during the adolescent window.<sup>152</sup> Despite the fact that adolescents are bigger and stronger than younger children, mortality rates increase more than 200% from childhood mainly due to accidents, suicide, substance abuse, and eating disorders.<sup>1</sup>

One of the most studied environmental effects in adolescence is alcohol abuse. In adults, brain toxicity has been documented as a consequence of chronic alcohol abuse: cortical gray matter thinning is most prominent in the PFC<sup>153</sup> and associated with changes in neuronal and glia density in both the orbitofrontal<sup>154</sup> and superior frontal cortices.<sup>155</sup> Alarming, the detrimental effects of alcohol consumption seem to be magnified in adolescence. Studies in human subjects have shown that impairment of memory function is more pronounced following even acute exposure to alcohol in younger (ages 21–24) than in older (ages 25–29) subjects.<sup>156</sup> In adolescent rats, ethanol administration selectively impairs spatial memory whereas adult rats are unaffected by the same doses.<sup>157</sup> Moreover, ethanol consumption in rats that simulates binge drinking results in more widespread pathology in adolescent animals than in adults.<sup>158</sup>

The basis of the enhanced vulnerability to alcohol in adolescence is undoubtedly complex and involves interaction with multiple neurotransmitter systems.<sup>159</sup> With regard to neuroplasticity, there are well documented effects of alcohol on the glutamate system. Acutely, ethanol inhibits NMDAR neurotransmission whereas long term exposure results in homeostatic upregulation of NMDAR signaling.<sup>159,160</sup> There is also growing evidence to suggest that ethanol has a greater effect on glutamate neurotransmission during adolescence than in later life. Ethanol exposure at low doses in adolescent rats is associated with inhibition of NMDAR-mediated EPSCs in the CA1 region of the hippocampus while high doses are required to inhibit EPSCs in adults.<sup>161</sup> Ethanol also blocks LTP in CA1 neurons of the hippocampus in adolescent but not adult rats.<sup>162</sup> Thus, even acute alcohol consumption in adolescence could disrupt mechanisms of Hebbian plasticity, and more chronic alcohol consumption in adolescence may induce homeostatic upregulation of glutamate neurotransmission that could result in long term changes in synapse number and dendritic spine morphology.<sup>160</sup> Homeostatic regulation of synaptic activity, i.e., increases or decreases synaptic scaling across the whole population of synapses, is also thought to be mediated by increased or decreased expression of AMPAR receptors on the post-synaptic membrane.<sup>163</sup> This suggests a potential site of interaction between developmental plasticity and homeostatic plasticity since both involve trafficking of AMPARs. Furthermore, sites of homeostatic plasticity correlate with lamina that exhibit plasticity during critical periods in the visual and somatosensory cortices, suggesting a possible

mechanism for heightened vulnerability of selected circuitry during different phases of development.<sup>163</sup> If synaptic plasticity in adolescence is primarily occurring in the neural circuitry that mediates executive processing, then disruption of synaptic plasticity at this time might result in enduring deficits in control of emotion, logical thinking and inhibition of impulsivity. In turn this lack of executive control could exacerbate the addictive tendencies and result in more severe alcoholism.

The adolescent brain is also more responsive to stress than the adult brain<sup>164</sup> and as a consequence may be more vulnerable to depression.<sup>151</sup> Analogous to the manner in which alcohol has age-specific effects that may depend on which regions of the brain are most plastic, a recent study has shown that the effects of sexual abuse, presumably the stress associated with the abuse, produces different brain pathology at childhood and adolescent ages.<sup>165</sup> Notably, frontal gray matter volume deficits were most pronounced in adult subjects who experienced sexual abuse at ages 14–16.<sup>165</sup>

The neural pathways that mediate and modulate the stressful effects on cognitive function in the PFC involve monoamine signaling.<sup>164</sup> Given the prominence of dopamine neurotransmission in mediating stress, the development of dopamine innervation of the PFC during late maturation might provide insight into the enhanced sensitivity to stress at this age. In the non-human primate, dopamine innervation of the middle PFC layers peaks near the onset of puberty and then decreases rapidly to adult levels while innervation of other layers is stable throughout the postnatal period.<sup>166</sup> D1 receptor levels also peak and decline to adult levels around the beginning of puberty.<sup>167</sup> These findings which indicate that the adult D1 receptor pattern is reached early do not appear to support a role for dopamine in adolescent enhancement of plasticity. However, in rodent prefrontal cortex, cell specificity has been observed in the distribution of D1 receptors with pyramidal cell neurons, but not interneurons, expressing higher levels of D1 receptors in adolescence than in adulthood.<sup>168</sup> These rodent data suggest that changes in D1 receptor expression might accentuate dopamine signaling in adolescence and thereby account for greater plasticity during this critical period. However, a credible alternative explanation is that the LTD-receptive state of adolescence is more sensitive to modulators like dopamine and that critical differences might be found in the mechanisms of glutamate receptor-mediated synaptic plasticity in the adolescent brain compared its adult counterpart.

### Clinical considerations

Identifying the molecular basis for synaptic pruning in adolescence could have wide ranging clinical ramifications. If NMDA-mediated LTD were proved to underlie reduction of connectivity, then the intracellular pathways associated with LTD processes, including those that mediate AMPAR internalization, could be targeted to curtail excessive synaptic pruning in diseases such as schizophrenia and ADHD. Because the D1 receptor is a key modulator of synaptic plasticity in the PFC and can even determine polarity of plasticity, i.e. high dopamine levels can predispose prefrontal synapses to LTD over LTP,<sup>137</sup> treatment with dopaminergic

antagonists or drugs that target intracellular dopamine signaling might also be useful in decreasing overactive LTD mechanisms. Along these same lines, drugs that impact the D1 receptor or its signaling pathways could ameliorate the impact of stress on the adolescent brain of individuals at risk for depression. Likewise, the involvement of glutamate receptors, including mGluRs,<sup>126</sup> in drug and alcohol addiction raises the possibility that pharmacologic targeting of glutamate signaling might have the potential to diminish the long term consequences of substance abuse in adolescence. In the same manner that the discovery of aberrant mGluR5 mechanisms in Fragile X syndrome has spawned new therapeutic approaches for treating this disease,<sup>169–171</sup> greater insight into the molecular substrates of adolescent maturation of the prefrontal cortex might lead to similar novel drug development for disorders and environmental exposures linked to abnormal adolescent development.

## Conclusions

The adolescent epoch is a time when refinement of connectivity establishes the proper excitatory/inhibitory balance in the PFC, and it is a critical period for normal maturation of executive functioning. Adolescence is postulated to be a time when LTD-driven synaptic pruning is occurring at a high rate in regions that govern higher cognitive function like the PFC. Further, the transition to adulthood is hypothesized to be marked by changes in the synapse that make the mature neuron less sensitive to AMPAR internalization, less likely to undergo LTD and thus less likely to undergo retraction of synaptic contacts.

## Conflict of interest

The author declares no conflict of interest.

**Acknowledgements.** I thank Dr Keith Young for his pre-submission reading of this manuscript and helpful comments.

- Dahl RE. Adolescent brain development: A period of vulnerabilities and opportunities. *Ann NY Acad Sci* 2004; **1021**: 1–22.
- Goldman-Rakic PS. Circuitry of the prefrontal cortex and the regulation of representational knowledge. In Plum F, Mountcastle V (eds) *Handbook of Physiology* vol. 5 (American Physiological Society: Bethesda, MD 373–417, 1987.
- Fuster JM. Frontal lobe and cognitive development. *J Neurocytol* 2002; **31**: 373–385.
- Huttenlocher PR, Dabholkar AS. Regional differences in synaptogenesis in human cerebral cortex. *J Comp Neurol* 1997; **387**: 167–178.
- Giedd JN, Blumenthal J, Jeffries NE, Castellanos FX, Liu H, Zijdenbos A *et al.* Brain development during childhood and adolescence: A longitudinal MRI study. *Nature Neurosci* 1999; **2**: 861–863.
- Sowell ER, Delis D, Stiles J, Jernigan TL. Improved memory functioning and frontal lobe maturation between childhood and adolescence: A structural MRI study. *J Internat Neuropsychol Soc* 2001; **7**: 312–322.
- Gotay N, Giedd JN, Lusk L, Hayashi KM, Greenstein D, Vaituzis AC *et al.* Dynamic mapping of human cortical development during childhood through early adulthood. *Proc Natl Acad Sci USA* 2004; **101**: 8174–8179.
- Sowell ER, Trauner DA, Gamat A, Jernigan TL. Development of cortical and subcortical brain structures in childhood and adolescence: A structural MRI study. *Develop Med Child Neurol* 2002; **44**: 4–16.
- De Luca DR, Wood SJ, Anderson V, Buchanan JA, Proffitt TM, Mahony K *et al.* Normative data from the CANTAB. I: Development of function over the lifespan. *J Clin Exp Neuropsychol* 2003; **25**: 242–254.
- Luna B, Garver KE, Urban TA, Lazar NA, Sweeney JA. Maturation of cognitive processes from late childhood to adulthood. *Child Develop* 2004; **75**: 1357–1372.
- Luciana M, Conklin HM, Hooper CJ, Yarger RS. The development of nonverbal working memory and control processes in adolescents. *Child Develop* 2005; **76**: 697–712.
- Best JR, Miller PH. A developmental perspective on executive function. *Child Develop* 2010; **81**: 1641–1660.
- Crews F, He J, Hodge C. Adolescent cortical development: A critical period of vulnerability for addiction. *Pharmacol Biochem Behav* 2007; **86**: 189–199.
- Goto Y, Yang CR, Otani S. Functional and dysfunctional synaptic plasticity in prefrontal cortex: Roles in psychiatric disorders. *Biol Psychiatry* 2010; **67**: 199–207.
- Spear LP. Adolescent brain development and animal models. *Ann NY Acad Sci* 2004; **1021**: 23–26.
- Plant TM. A study of the role of the postnatal testes in determining the ontogeny of gonadotropin secretion in the male rhesus monkey (*Macaca mulatta*). *Endocrinol* 1985; **116**: 1341–1350.
- Butterwick RF, McConnell M, Markwell PJ, Watson TD. Influence of age and sex on plasma lipid and lipoprotein concentrations and associated enzyme activities in cats. *Am J Vet Res* 2001; **62**: 331–336.
- Purves D, Lichtman JW. Elimination of synapses in the developing nervous system. *Science* 1980; **210**: 153–157.
- Cowan WM, Fawcett JW, O'Leary DDM, Stanfield BB. Regressive events in neurogenesis. *Science* 1984; **225**: 1258–1265.
- Land PW, Lund RD. Development of the rat's uncrossed retinotectal pathway and its relation to plasticity studies. *Science* 1979; **205**: 698–700.
- Frost DO, So K-F, Schneider GE. Postnatal development of retinal projections in Syrian hamsters: A study using autoradiographic and anterograde degeneration techniques. *Neuroscience* 1979; **4**: 1649–1677.
- Insausti R, Blackmore C, Cowan WM. Ganglion cell death during development of ipsilateral retino-collicular projection in golden hamster. *Nature* 1984; **308**: 362–365.
- Williams RW, Herrup K. The control of neuron number. *Ann Rev Neurosci* 1988; **11**: 423–453.
- Williams RW, Rakic P. Elimination of neurons from the rhesus monkey's lateral geniculate nucleus during development. *J Comp Neurol* 1988; **272**: 424–436.
- Lotto RB, Asavartikra P, Vail L, Price DJ. Target-derived neurotrophic factors regulate the death of developing forebrain neurons after a change in their trophic requirements. *J Neurosci* 2001; **21**: 3904–3910.
- Innocenti GM. Growth and reshaping of axons in the establishment of visual callosal connections. *Science* 1981; **212**: 824–827.
- O'Leary DDM, Stanfield BB, Cowan WM. Evidence that the early postnatal restriction of the cells of origin of the callosal projection is due to the elimination of axonal collaterals rather than to the death of neurons. *Develop Brain Res* 1981; **1**: 607–617.
- Ivy GO, Killackey HP. Ontogenetic changes in the projections of neocortical neurons. *J Neurosci* 1982; **2**: 735–743.
- Rakic P, Riley KP. Overproduction and elimination of retinal axons in the fetal rhesus monkey. *Science* 1983; **219**: 1441–1444.
- LaMantia AS, Rakic P. Axon overproduction and elimination in the corpus callosum of the developing rhesus monkey. *J Neurosci* 1990; **10**: 2156–2175.
- LaMantia AS, Rakic P. Axon overproduction and elimination in the anterior commissure of the developing rhesus monkey. *J Comp Neurol* 1994; **340**: 328–336.
- Campbell G, Shatz CJ. Synapses formed by identified retinogeniculate axons during segregation of eye input. *J Neurosci* 1992; **12**: 1847–1858.
- Gourley SL, Olevska A, Sloan Warren M, Taylor JR, Koleski AJ. Arg kinase regulates prefrontal dendritic spine refinement and cocaine-induced plasticity. *J Neurosci* 2012; **32**: 2314–2323.
- Cragg BG. The development of synapses in the visual system of the cat. *J Comp Neurol* 1972; **160**: 147–166.
- Rakic P, Bourgeois J-P, Eckenhoff MF, Zecevic N, Goldman-Rakic PS. Concurrent overproduction of synapses in diverse regions of the primate cerebral cortex. *Science* 1986; **232**: 232–235.
- Zecevic N, Bourgeois J-P, Rakic P. Changes in synaptic density in motor cortex of rhesus monkey during fetal and postnatal life. *Develop Brain Res* 1989; **50**: 11–32.
- Bourgeois J-P, Rakic P. Changes of synaptic density in the primary visual cortex of the macaque monkey from fetal to adult stage. *J Neurosci* 1993; **13**: 2801–2820.
- Bourgeois J-P, Goldman-Rakic PS, Rakic P. Synaptogenesis in the prefrontal cortex of rhesus monkeys. *Cereb Cortex* 1994; **4**: 78–96.
- Redfern PA. Neuromuscular transmission in new-born rats. *J Physiol* 1970; **209**: 701–709.
- Sretavan D, Shatz CJ. Prenatal development of individual retinogeniculate axons during the period of segregation. *Nature* 1984; **308**: 845–848.
- Changeux J-P, Danchin A. Selective stabilisation of developing synapses as a mechanism for the specification of neuronal networks. *Nature* 1976; **264**: 705–712.
- Stryker MP, Harris WA. Binocular impulse blockade prevents the formation of ocular dominance columns in cat visual cortex. *J Neurosci* 1986; **6**: 2117–2133.
- Shatz CJ, Stryker MP. Ocular dominance in layer IV of the cat's visual cortex and the effects of monocular deprivation. *J Physiol* 1978; **281**: 267–283.
- Constantine-Paton M, Cline HT, Debski E. Patterned activity, synaptic convergence, and the NMDA receptor in developing visual pathways. *Ann Rev Neurosci* 1990; **13**: 129–154.
- Shatz CJ. Impulse activity and the patterning of connections during CNS development. *Neuron* 1990; **5**: 745–756.

46. Hubel H, Wiesel TN, LeVay S. Plasticity of ocular dominance columns in monkey striate cortex. *Philos Trans Roy Soc Lond. Ser B Biol Med* 1977; **278**: 377–409.
47. LeVay S, Wiesel TN, Hubel DH. The development of ocular dominance columns in normal and visually deprived monkeys. *J Comp Neurol* 1980; **191**: 1–51.
48. Hubel DH, Wiesel TN. The period of susceptibility to the physiological effects of unilateral eye closure in kittens. *J Physiol* 1970; **206**: 419–436.
49. Foeller E, Feldman DE. Synaptic basis for developmental plasticity in somatosensory cortex. *Curr Opin Neurobiol* 2004; **14**: 89–95.
50. Huttenlocher PR. Synaptic density in human frontal cortex – Developmental changes and effects of aging. *Brain Res* 1979; **163**: 195–205.
51. Petanjek A, Judas M, Simic G, Roko Rasin M, Uylings HBM, Rakic P *et al*. Extraordinary neoteny of synaptic spines in the human prefrontal cortex. *Proc Natl Acad Sci USA* 2011; **108**: 13281–13286.
52. Glantz LA, Gilmore JH, Hamer RM, Lieberman JA, Jarskog LF. Synaptophysin and postsynaptic density protein 95 in the human prefrontal cortex from mid-gestation into early adulthood. *Neuroscience* 2007; **149**: 582–591.
53. Webster MJ, Elashoff M, Weickert CS. Molecular evidence that cortical synaptic growth predominates in the first decade of life in humans. *Internat J Develop Neurosci* 2011; **29**: 225–236.
54. Finlay BL, Slattery M. Local differences in the amount of early cell death in the neocortex predict adult local specializations. *Science* 1983; **219**: 1349–1351.
55. Pfefferbaum A, Mathalon DH, Sullivan EV, Rawles JM, Zipursky RB, Lim KO. A quantitative magnetic resonance imaging study of changes in brain morphology from infancy to late adulthood. *Arch Neurol* 1994; **51**: 874–887.
56. Reiss AL, Abrams MT, Singer HS, Ross JL, Denckla MB. Brain development, gender and IQ in children: A volumetric imaging study. *Brain* 1996; **119**: 1763–1774.
57. Brain Development Cooperative Group. Total and regional brain volumes in a population-based normative sample from 4 to 18 years: The NIH MRI study of normal brain development. *Cereb Cortex* 2012; **22**: 1–12.
58. Jernigan TL, Trauner DA, Hesselink JR, Tallal PA. Maturation of human cerebrum observed *in vivo* during adolescence. *Brain* 1991; **114**: 2037–2049.
59. Sowell ER, Thompson PM, Holmes CJ, Bath R, Jernigan TL, Toga AW. Localizing age-related changes in brain structure between childhood and adolescence using statistical parametric mapping. *NeuroImage* 1999; **9**: 587–597.
60. Anderson SA, Classey JD, Conde F, Lund JS, Lewis DA. Synchronous development of pyramidal neuron dendritic spines and parvalbumin-immunoreactive chandelier neuron axon terminals in layer II of monkey prefrontal cortex. *Neuroscience* 1995; **67**: 7–22.
61. Woodruff AR, Anderson SA, Yuste R. The enigmatic function of chandelier cells. *Front Neurosci* 2010; **4**: 201.
62. Gorelova N, Seamans JK, Yang CR. Mechanisms of dopamine activation of fast-spiking interneurons that exert inhibition in rat prefrontal cortex. *J Neurophysiol* 2002; **88**: 3150–3166.
63. Tseng KY, O'Donnell P. D2 dopamine receptors recruit a GABA component for their attenuation of excitatory synaptic transmission in adult rat prefrontal cortex. *Synapse* 2007; **61**: 843–850.
64. Tseng KY, O'Donnell P. Dopamine modulation of prefrontal cortical interneurons changes during adolescence. *Cereb Cortex* 2007; **17**: 1235–1240.
65. Rao SG, Williams GV, Goldman-Rakic PS. Isodirectional turning of adjacent interneurons and pyramidal cells during working memory: Evidence for microcolumnar organization in PFC. *J Neurophysiol* 1999; **81**: 1903–1916.
66. Constantinidis C, Williams GV, Goldman-Rakic PS. A role for inhibition in shaping the temporal flow of information in the prefrontal cortex. *Nat Neurosci* 2002; **5**: 175–180.
67. Cho RY, Konecny RO, Carter CS. Impairments in frontal cortical  $\gamma$  synchrony and cognitive control in schizophrenia. *Proc Natl Acad Sci USA* 2006; **103**: 19878–19883.
68. Fries P. Neuronal gamma-band synchronization as a fundamental process in cortical computation. *Ann Rev Neurosci* 2009; **32**: 209–224.
69. Lewis DA, Hashimoto T, Volk DW. Cortical inhibitory neurons and schizophrenia. *Nat Rev Neurosci* 2005; **6**: 312–324.
70. O'Donnell P. Adolescent onset of cortical disinhibition in schizophrenia: Insights from animal models. *Schizophr Bull* 2011; **37**: 484–492.
71. Wu G, Malinow R, Cline HT. Maturation of a central glutamatergic synapse. *Science* 1996; **274**: 972–976.
72. Durand GM, Koyalchuk Y, Konnerth A. Long-term potentiation and functional synapse induction in developing hippocampus. *Nature* 1996; **381**: 71–75.
73. Isaac JT, Crair MC, Nicoll RA, Malenka RC. Silent synapses during development of thalamocortical inputs. *Neuron* 1997; **18**: 269–280.
74. Zhu JJ, Esteban JA, Hayashi Y, Malinow R. Postnatal synaptic potentiation: Delivery of GluR4-containing AMPA receptors by spontaneous activity. *Nat Neurosci* 2000; **3**: 1098–1106.
75. Liao D, Hessler NA, Malinow R. Activation of postsynaptically silent synapses during pairing-induced LTP in CA1 region of hippocampal slice. *Nature* 1995; **375**: 400–404.
76. Dudek SM, Bear MF. Long-term depression in area CA1 of hippocampus and effects of N-methyl-D-aspartate receptor blockade. *Proc Natl Acad Sci USA* 1992; **89**: 4363–4367.
77. Barria A, Muller D, Derkach V, Griffith LC, Soderling TR. Regulatory phosphorylation of AMPA-type glutamate receptors by CaM-KII during long term potentiation. *Science* 1997; **276**: 2042–2045.
78. Leonard AS, Lim IA, Hemsworth DE, Horne MC, Hell JW. Calcium/calmodulin-dependent protein kinase II is associated with the N-methyl-D-aspartate receptor. *Proc Natl Acad Sci USA* 1999; **96**: 3239–3244.
79. Lee HK, Kameyama K, Huganir RL, Bear MF. NMDA induces long-term synaptic depression and dephosphorylation of the GluR1 subunit of AMPA receptors in hippocampus. *Neuron* 1998; **21**: 1151–1162.
80. Lee HK, Barbarosie M, Kameyama K, Bear MF, Huganir RL. Regulation of distinct AMPA receptor phosphorylation sites during bidirectional synaptic plasticity. *Nature* 2000; **405**: 955–959.
81. Malenka RC. Synaptic plasticity in the hippocampus: LTP and LTD. *Cell* 1994; **79**: 535–538.
82. Malinow R, Malenka RC. AMPA receptor trafficking and synaptic plasticity. *Ann Rev Neurosci* 2002; **25**: 103–126.
83. Maletic-Savatic M, Malinow R, Svoboda K. Rapid dendritic morphogenesis in CA1 hippocampal dendrites induced by synaptic activity. *Science* 1999; **283**: 1923–1927.
84. Engert F, Bonhoeffer T. Dendritic spine changes associated with hippocampal synaptic plasticity. *Nature* 1999; **399**: 66–70.
85. Toni N, Buchs PA, Nikonenko I, Bron CR, Muller D. LTP promotes formation of multiple spine synapses between a single axon terminal and a dendrite. *Nature* 1999; **402**: 421–425.
86. Matsuzaki M, Honkura N, Ellis-Davies GC, Kasai H. Structural basis of long-term potentiation in single dendritic spines. *Nature* 2004; **429**: 761–766.
87. Nagerl UV, Eberhorn N, Cambridge SB, Bonhoeffer T. Bidirectional activity-dependent morphological plasticity in hippocampal neurons. *Neuron* 2004; **44**: 759–767.
88. Zhou Q, Homma KJ, Poo MM. Shrinkage of dendritic spines associated with long-term depression of hippocampal synapses. *Neuron* 2004; **44**: 749–757.
89. Tominaga-Yoshino K, Kondo S, Tamotsu S, Ogura A. Repetitive activation of protein kinase A induces slow and persistent potentiation associated with synaptogenesis in cultured hippocampus. *Neurosci Res* 2002; **44**: 357–367.
90. Shinoda Y, Kamikubo Y, Egashira Y, Tominaga-Yoshino K, Ogura A. Repetition of mGluR-dependent LTD causes slowly developing persistent reduction in synaptic strength accompanied by synaptic elimination. *Brain Res* 2005; **1042**: 99–107.
91. Kamikubo Y, Egashira Y, Tanaka T, Shinoda Y, Tominaga-Yoshino K, Ogura A. Long-lasting synaptic loss after repeated induction of LTD: independence to the means of LTD induction. *Eur J Neurosci* 2006; **24**: 1606–1616.
92. Bastrikova N, Gardner GA, Reece JM, Jeromin A, Dudek SM. Synapse elimination accompanies functional plasticity in hippocampal neurons. *Proc Natl Acad Sci USA* 2008; **105**: 3123–3127.
93. Crair MC, Malenka RC. A critical period for long-term potentiation at thalamocortical synapses. *Nature* 1995; **375**: 325–328.
94. Allen CB, Celikel T, Feldman DE. Long-term depression induced by sensory deprivation during cortical map plasticity *in vivo*. *Nat Neurosci* 2003; **6**: 291–299.
95. Lu H-C, She W-C, Plas DT, Neumann PE, Janz R, Crair MC. Adenylyl cyclase I regulates AMPA receptor trafficking during mouse cortical 'barrel' map development. *Nat Neurosci* 2003; **6**: 939–947.
96. Yoon B-J, Smith GB, Heynen AJ, Neve RL, Bear MF. Essential role for a long-term depression mechanism in ocular dominance plasticity. *Proc Natl Acad Sci USA* 2009; **106**: 9860–9865.
97. Bear MF. Synaptic plasticity: From theory to reality. *Philos Trans Biol Sci* 2003; **358**: 649–655.
98. Kato N, Artola A, Singer W. Developmental changes in the susceptibility to long-term potentiation of neurons in rat visual cortex slices. *Develop Brain Res* 1991; **60**: 43–50.
99. Flint AC, Maisch Us, Weishaupt JH, Kriegstein AR, Monver H. NR2A subunit expression shortens NMDA receptor synaptic currents in developing neocortex. *J Neurosci* 1997; **17**: 2469–2476.
100. Roberts EB, Romo AS. Enhanced NR2A subunit expression and decreased NMDA receptor decay time at the onset of ocular dominance plasticity in the ferret. *J Neurophysiol* 1999; **81**: 2587–2591.
101. Sheng M, Cummings J, Roldan LA, Jan YN, Jan LY. Changing subunit composition of heteromeric NMDA receptors during development of rat cortex. *Nature* 1994; **368**: 144–147.
102. Carnignoto G, Vicini S. Activity-dependent decrease in NMDA receptor responses during development of the visual cortex. *Science* 1992; **258**: 1007–1011.
103. Fox K, Daw N, Sato H, Czepita D. The effect of visual experience on development of NMDA receptor synaptic transmission in kitten visual cortex. *J Neurosci* 1992; **12**: 2672–2684.
104. Nase G, Weishaupt J, Stern P, Singer W, Monver H. Genetic and epigenetic regulation of NMDA receptor expression in the rat visual cortex. *Eur J Neurosci* 1999; **11**: 4320–4326.
105. Quinlan EM, Olstein DH, Bear MF. Bidirectional, experience-dependent regulation of N-methyl-D-aspartate receptor subunit composition in the rat visual cortex during postnatal development. *Proc Natl Acad Sci USA* 1999; **96**: 12876–12880.
106. Erisir A, Harris JL. Decline of the critical period of visual plasticity is concurrent with the reduction of NR2B subunit of the synaptic NMDA receptor in layer 4. *J Neurosci* 2003; **23**: 5208–5218.
107. Fox K, Sato H, Daw N. The location and function of NMDA receptors in cat and kitten visual cortex. *J Neurosci* 1989; **9**: 2443–2454.

108. Gambrell AC, Barria A. NMDA receptor subunit composition controls synaptogenesis and synapse stabilization. *Proc Natl Acad Sci USA* 2011; **108**: 5855–5860.
109. Barria A, Muller D, Derkach V, Griffith LC, Soderling TR. Regulatory phosphorylation of AMPA-type glutamate receptors by CaM-KII during long-term potentiation. *Science* 1997; **276**: 2042–2045.
110. Leonard AS, Lim IA, Hemworth DE, Horne MC, Hell JW. Calcium/calmodulin-dependent protein kinase II is associated with the N-methyl-D-aspartate receptor. *Proc Natl Acad Sci USA* 1999; **96**: 3239–3244.
111. Strack S, Colbran RJ. Autophosphorylation-dependent targeting of calcium/calmodulin-dependent protein kinase II by the NR2B subunit of the N-methyl-D-aspartate receptor. *J Biol Chem* 1998; **273**: 20689–20692.
112. Gardoni F, Schrama LH, van Dalen JJ, Gispén WH, Cattabeni F, Di Luca M. AlphaCaMKII binding to the C-terminal tail of NMDA receptor subunit NR2A and its modulation by autophosphorylation. *FEBS Lett* 1999; **456**: 394–398.
113. Barria A, Malinow R. NMDA receptor subunit composition controls synaptic plasticity by regulating binding to CaMKII. *Neuron* 2005; **48**: 289–301.
114. Zhao JP, Constantine-Paton M. NR2A<sup>-/-</sup> mice lack long-term potentiation but retain NMDA receptor and L-type Ca<sup>2+</sup> channel-dependent long-term depression in the juvenile superior colliculus. *J Neurosci* 2007; **27**: 13649–13654.
115. Foster KA, McLaughlin N, Edbauer D, Phillips M, Bolton A, Constantine-Paton M *et al.* Distinct roles of NR2A and NR2B cytoplasmic tails in long-term potentiation. *J Neurosci* 2010; **30**: 2676–2685.
116. Barria A, Malinow R. Subunit-specific NMDA receptor trafficking to synapses. *Neuron* 2002; **35**: 345–353.
117. Abraham WC, Bear MF. Metaplasticity: the plasticity of synaptic plasticity. *Trends Neurosci* 1996; **19**: 126–130.
118. Philpot BD, Cho KK, Bear MF. Obligatory role of NR2A for metaplasticity in visual cortex. *Neuron* 2007; **53**: 495–502.
119. Yashiro K, Philpot BD. Regulation of NMDA receptor subunit expression and its implications for LTD, LTP, and metaplasticity. *Neuropharmacol* 2008; **55**: 1081–1094.
120. Bear MF, Kirkwood A. Neocortical long-term potentiation. *Curr Opin Neurobiol* 1993; **3**: 197–202.
121. Herry C, Vouimba R-M, Garcia R. Plasticity in the mediodorsal thalamo-prefrontal cortical transmission in behaving mice. *J Neurophysiol* 1999; **82**: 2827–2832.
122. Gurden H, Tassin JP, Jay TM. Integrity of the mesocortical dopaminergic system is necessary for complete expression of in vivo hippocampal-prefrontal cortex long-term potentiation. *Neuroscience* 1999; **94**: 1019–1027.
123. Bueno-Junior LS, Lopes-Agular C, Ruggiero RN, Romcy-Pereira RN, Leite JP. Muscarinic and nicotinic modulation of thalamo-prefrontal cortex synaptic plasticity in vivo. *PLoS One* 2012; **7**: e47484.
124. Bellone C, Luscher C, Mameli M. Mechanisms of synaptic depression triggered by metabotropic glutamate receptors. *Cell Mol Life Sci* 2008; **65**: 2913–2923.
125. Gladding CM, Fitzjohn SM, Molnar E. Metabotropic glutamate receptor-mediated long-term depression: Molecular mechanisms. *Pharmacol Rev* 2009; **61**: 395–412.
126. Luscher C, Huber KM. Group 1 mGluR-dependent synaptic long-term depression: Mechanisms and implications for circuitry and disease. *Neuron* 2010; **65**: 445–459.
127. Mateo Z, Porter JT. Group II metabotropic glutamate receptors inhibit glutamate release at thalamocortical synapses in the developing somatosensory cortex. *Neuroscience* 2007; **146**: 1062–1072.
128. Holbro N, Grunditz A, Oertner TG. Differential distribution of endoplasmic reticulum controls metabotropic signaling and plasticity at hippocampal synapses. *Proc Natl Acad Sci USA* 2009; **106**: 15055–15060.
129. Dimitriu D, Hao J, Hara Y, Kaufman J, Janssen WGM, Lou W *et al.* Selective changes in thin spine density and morphology in monkey prefrontal cortex correlate with aging-related cognitive impairment. *J Neurosci* 2010; **30**: 7507–7515.
130. Andersen P. Assessment and development of executive function (EF) during childhood. *Child Neuropsychol* 2002; **8**: 71–82.
131. Shaw P, Greenstein D, Lerch J, Clasen L, Lenroot R, Gogtay N *et al.* Intellectual ability and cortical development in children and adolescents. *Nature* 2006; **440**: 676–679.
132. Shaw P, Sharp WS, Morrison M, Eckstrand K, Greenstein DK, Clasen LS *et al.* Psychostimulant treatment and the developing cortex in attention deficit hyperactivity disorder. *Am J Psychiatry* 2009; **166**: 58–63.
133. Vazdarjanova A, Bunting K, Muthusamy N, Bergson C. Calcyon upregulation in adolescence impairs response inhibition and working memory in adulthood. *Molec Psychiatry* 2011; **16**: 672–684.
134. Selemón LD. Regionally diverse cortical pathology in schizophrenia: Clues to the etiology of the disease. *Schizophr Bull* 2001; **27**: 349–377.
135. Koh P, Bergson C, Undie A, Goldman-Rakic PS, Lidow M. Up-regulation of the D1 dopamine receptor-interacting protein, calcyon, in patients with schizophrenia. *Arch Gen Psychiatry* 2003; **60**: 311–319.
136. Bai J, He F, Novikova S, Undie A, Dracheva S, Haroutunian V *et al.* Abnormalities in the dopamine system in schizophrenia may lie in altered levels of dopamine-interacting proteins. *Biol Psychiatry* 2004; **56**: 427–440.
137. Law-Tho D, Desce JM, Crepel F. Dopamine favours the emergence of long-term depression versus potentiation in slices of rat prefrontal cortex. *Neurosci Lett* 1995; **188**: 125–128.
138. Snyder GL, Allen PB, Fienberg AA, Valle CG, Hagan RL, Nairn AC *et al.* Regulation of phosphorylation of the GluR1 AMPA receptor in the neostriatum by dopamine and psychostimulants in vivo. *J Neurosci* 2000; **20**: 4480–4488.
139. Sun X, Zhao Y, Wolf ME. Dopamine receptor stimulation modulates AMPA receptor synaptic insertion in prefrontal cortex neurons. *J Neurosci* 2005; **25**: 7342–7351.
140. Selemón LD, Begovic' A, Goldman-Rakic PS, Castner SA. Amphetamine sensitization alters dendritic morphology in prefrontal cortical pyramidal neurons in the non-human primate. *Neuropsychopharmacology* 2007; **32**: 919–931.
141. Selemón LD, Begovic' A, Williams GV, Castner SA. Reversal of neuronal and cognitive consequences of amphetamine sensitization following chronic treatment with a D1 antagonist. *Pharmacol Biochem Behav* 2010; **96**: 325–332.
142. Caruana DA, Warburton EC, Bashir ZI. Induction of activity-dependent LTD requires muscarinic receptor activation in medial prefrontal cortex. *J Neurosci* 2011; **31**: 18464–18478.
143. Auclair N, Otani S, Soubrie P, Crepel F. Cannabinoids modulate synaptic strength and plasticity at glutamatergic synapses of rat prefrontal cortex pyramidal neurons. *J Neurophysiol* 2000; **83**: 3287–3293.
144. Anthony JC, Petronis KR. Early-onset drug use and risk of later drug problems. *Drug Alcohol Depend* 1995; **40**: 9–15.
145. Adriani W, Spiker S, Deroche-Gamonet V, Laviola G, Le Moal M, Smit AB *et al.* Evidence for enhanced neurobehavioral vulnerability to nicotine during periadolescence in rats. *J Neurosci* 2003; **23**: 4712–4716.
146. Kandel DB. Epidemiological and psychosocial perspectives on adolescent drug use. *J Am Acad Child Psychiatry* 1982; **21**: 4328–4347.
147. Taioli E, Wynder EL. Effect of the age at which smoking begins on frequency of smoking in adulthood. *New Engl J Med* 1991; **325**: 968–969.
148. Chambers RA, Taylor JR, Potenza MN. Developmental neurocircuitry of motivation in adolescence: A critical period of addiction vulnerability. *Am J Psychiatry* 2003; **160**: 1041–1052.
149. Kelley AE. Memory and addiction: Shared neural circuitry and molecular mechanisms. *Neuron* 2004; **44**: 161–179.
150. Hyman SE. Addiction: A disease of learning and memory. *Am J Psychiatry* 2005; **162**: 1414–1422.
151. Andersen SL, Stress Teicher MH. Sensitive periods and maturational events in adolescent depression. *Trends Neurosci* 2008; **31**: 183–191.
152. Hafner H, Maurer K, Löffler W, Riecher-Rössler A. The influence of age and sex on the onset and early course of schizophrenia. *Brit J Psychiatry* 1993; **162**: 80–86.
153. Pfefferbaum A, Sullivan EV, Rosenbloom MJ, Mathalon DH, Lim KO. A controlled study of cortical gray matter and ventricular changes in alcoholic men over a 5-year interval. *Arch Gen Psychiatry* 1998; **55**: 905–912.
154. Miguel-Hidalgo JJ, Overholser JC, Meltzer HY, Stockmeier CA, Rajkowska G. Reduced glial and neuronal packing density in the orbitofrontal cortex in alcohol dependence and its relationship with suicide and duration of alcohol dependence. *Alcohol Clin Exp Res* 2006; **30**: 1845–1855.
155. Harper C, Kril J. Patterns of neuronal loss in the cerebral cortex in chronic alcoholic patients. *J Neurol Sciences* 1989; **92**: 81–89.
156. Acheson SK, Stein RM, Swartzwelder HS. Impairment of semantic and figural memory by acute ethanol: Age-dependent effects. *Alcohol Clin Exp Res* 1998; **22**: 1437–1442.
157. Markwiese BJ, Acheson SK, Levin ED, Wilson WA, Swartzwelder HS. Differential effects of ethanol on memory in adolescent and adult rats. *Alcohol Clin Exp Res* 1998; **22**: 416–421.
158. Crews FT, Braun CJ, Hoplight B, Switzer RC 3rd, Knapp DJ. Binge ethanol consumption causes differential brain damage in young adolescent rats compared with adult rats. *Alcohol Clin Exp Res* 2000; **24**: 1712–1723.
159. Fadda F, Rossetti ZL. Chronic ethanol consumption: From neuroadaptation to neurodegeneration. *Progr Neurobiol* 1998; **56**: 385–431.
160. Carpenter-Hyland EP, Chandler LJ. Adaptive plasticity of NMDA receptors and dendritic spines: Implications for enhanced vulnerability of the adolescent brain to alcohol addiction. *Pharmacol Biochem Behav* 2007; **86**: 200–208.
161. Swartzwelder HS, Wilson WA, Tayyeb MI. Differential sensitivity of NMDA receptor-mediated synaptic potentials to ethanol in immature versus mature hippocampus. *Alcohol Clin Exp Res* 1995; **19**: 320–323.
162. Swartzwelder HS, Wilson WA, Tayyeb MI. Age-dependent inhibition of long-term potentiation by ethanol in immature versus mature hippocampus. *Alcohol Clin Exp Res* 1995; **19**: 1480–1484.
163. Turrigiano GG, Nelson SB. Homeostatic plasticity in the developing nervous system. *Nat Rev Neurosci* 2004; **5**: 97–107.
164. Arnsten AFT, Shansky RM. Adolescence: Vulnerable period for stress-induced prefrontal cortical function? *Ann NY Acad Sci* 2004; **1021**: 143–147.
165. Andersen LS, Tomada A, Vincow ES, Valente E, Polcari A, Teicher MH. Preliminary evidence for sensitive periods in the effect of childhood sexual abuse on regional brain development. *J Neuropsychiatry Clin Neurosci* 2008; **20**: 292–301.
166. Rosenberg DR, Lewis DA. Postnatal maturation of the dopaminergic innervation of monkey prefrontal and motor cortices: A tyrosine hydroxylase immunohistochemical analysis. *J Comp Neurol* 1995; **358**: 383–400.

167. Lidow MS, Rakic P. Scheduling of monoaminergic neurotransmitter receptor expression in the primate neocortex during postnatal development. *Cereb Cortex* 1992; **2**: 401–416.
168. Brenhouse HC, Sonntag KC, Andersen SL. Transient D1 dopamine receptor expression on prefrontal cortex projection neurons: Relationship to enhanced motivational salience of drug cues in adolescence. *J Neurosci* 2008; **28**: 2375–2382.
169. Bassell GJ, Warren ST. Fragile X syndrome: Loss of local mRNA regulation alters synaptic development and function. *Neuron* 2008; **60**: 201–214.
170. Berry-Kravis E, Sumis A, Hervey C, Nelson M, Porges SW, Weng N *et al*. Open-label treatment trial of lithium to target the underlying defect in fragile X syndrome. *J Dev Behav Pediatr* 2008; **29**: 293–302.
171. Berry-Kravis E, Hessel D, Coffey S, Hervey C, Schneider A, Yuhas J *et al*. A pilot open label, single dose trial of fenobam in adults with fragile X syndrome. *J Med Genet* 2009; **46**: 266–271.



**Translational Psychiatry is an open-access journal published by Nature Publishing Group. This work is licensed under the Creative Commons Attribution-NonCommercial-No Derivative Works 3.0 Unported License. To view a copy of this license, visit <http://creativecommons.org/licenses/by-nc-nd/3.0/>**