



Since January 2020 Elsevier has created a COVID-19 resource centre with free information in English and Mandarin on the novel coronavirus COVID-19. The COVID-19 resource centre is hosted on Elsevier Connect, the company's public news and information website.

Elsevier hereby grants permission to make all its COVID-19-related research that is available on the COVID-19 resource centre - including this research content - immediately available in PubMed Central and other publicly funded repositories, such as the WHO COVID database with rights for unrestricted research re-use and analyses in any form or by any means with acknowledgement of the original source. These permissions are granted for free by Elsevier for as long as the COVID-19 resource centre remains active.



ELSEVIER

What can we predict about viral evolution and emergence?

Edward C Holmes^{1,2}

Predicting the emergence of infectious diseases has been touted as one of the most important goals of biomedical science, with an array of funding schemes and research projects. However, evolutionary biology generally has a dim view of prediction, and there is a danger that erroneous predictions will mean a misuse of resources and undermine public confidence. Herein, I outline what can be realistically predicted about viral evolution and emergence, argue that any success in predicting what may emerge is likely to be limited, but that forecasting how viruses might evolve and spread following emergence is more tractable. I also emphasize that a properly grounded research program in disease prediction must involve a synthesis of ecological and genetic perspectives.

Addresses

¹Sydney Emerging Infections and Biosecurity Institute, School of Biological Sciences and Sydney Medical School, The University of Sydney, Sydney, NSW 2006, Australia

²Fogarty International Center, National Institutes of Health, Bethesda, MD 20892, USA

Corresponding author: Holmes, Edward C
(edward.holmes@sydney.edu.au)

Current Opinion in Virology 2013, 3:180–184

This review comes from a themed issue on **Emerging viruses**

Edited by **Ian Lipkin** and **Ab Osterhaus**

For a complete overview see the [Issue](#) and the [Editorial](#)

Available online 28th December 2013

1879-6257/\$ – see front matter, © 2013 Elsevier B.V. All rights reserved.

<http://dx.doi.org/10.1016/j.coviro.2012.12.003>

Introduction

The SARS epidemic of 2003, the global spread of swine-origin H1N1 influenza in 2009, the recent appearance of new hemorrhagic fevers [1,2^{*}] and pneumonias [3], and the continuing threat posed by H5N1 avian influenza highlight the pressing need to understand the mechanisms that underpin viral emergence. Not surprisingly, the potential health consequences of a new pandemic have stimulated attempts to predict what types of virus might emerge in the future, as well as where and when such an emergence event will occur [4,5^{*}]. As the process of emergence is synonymous with the cross-species transmission of viruses to new hosts [6–8], predicting what might emerge is essentially equivalent to predicting what viruses are better able to jump species boundaries and spread in new hosts.

The evolution of viral emergence

There is a general, and justifiable, nervousness about predictions in evolutionary biology. Mutations can be

fixed in populations by genetic drift or natural selection. Prediction in the case of genetic drift is necessarily hindered by a large sampling error. Although natural selection is a deterministic process, which ought to engender it with some inherent predictability, this will only be possible if the fitness of each relevant mutation is known in each relevant environment. For example, early attempts to predict the evolution of influenza virus as a guide for vaccine strain selection using the differential strength of positive selection among viral lineages gained little traction [9]. Although predictions are more robust when selection is extremely strong — it is trivial to predict that resistance (and often the causative mutations) will arise to commonly used antivirals — predictions of the evolution of more complex multifactorial traits such as host adaptation and emergence are a very different matter [10,11]. Even for traits where single mutations might confer a major fitness advantage, such as antiviral resistance, genome-scale interactions, including epistasis [12^{**},13^{*}] and permissive mutations [14^{**}], greatly complicate the exercise.

Fortunately, there are some evolutionary generalities that enable very broad-scale predictions about viral emergence, although none possess meaningful precision. One of the best established is that viruses are more likely to jump between closely related species [15,16^{**}]. Although informative, numerous exceptions arise because the probability of exposure does not match phylogenetic relatedness. For example, although humans are more closely related to other primates than to rodents, we probably have greater exposure to the latter. It is equally well known that RNA viruses jump species boundaries more often than DNA viruses and that this likely reflects their differing rates of evolutionary change [8,17]. RNA viruses have mutation rates of between 0.1 and 1.0 mutations per genome replication, several logs higher than those of double-strand DNA viruses, and which will generate abundant genetic variation when coupled with huge population sizes [17]. Although there is some variation in evolutionary rate among RNA viruses, this does not appear to be related to the propensity to jump hosts. Interestingly, single-stranded DNA viruses exhibit rates of evolutionary change similar to those of RNA viruses [17] and may frequently cross species boundaries [18]. Similarly, although rates of recombination (or reassortment) vary greatly among viruses, these appear to be of little predictive power: as a case in point, paramyxoviruses, such as measles and the emerging henipaviruses, experience extremely low rates of recombination (if they recombine at all) yet frequently jump species [19].

More useful predictors come from studies of the evolutionary correlates of past emergence. Both experimental and comparative analyses reveal that vector-borne viruses, which must simultaneously infect very different types of host, experience stronger selective constraints than those transmitted by other routes, and which likely hinders their adaptability [20,21]. Indeed, although vector-borne viruses frequently spill-over from one host to another, they are less likely to evolve sustained transmission in new hosts [8]. In contrast, viruses that utilize phylogenetically conserved cell receptors appear better able to jump host species boundaries than those that use more variable receptors [5]. However, receptor usage is not the only factor that mediates successful infection, and is difficult to assess until a virus has emerged.

The challenge of predicting viral evolution and emergence is reflected in the on-going debate over whether highly pathogenic avian H5N1 influenza virus will emerge in humans. Although only a small number of mutations may be necessary to make an avian influenza virus aerosol transmissible by mammals [22–24], how this relates to evolution in nature is less clear because some of the key mutations were generated artificially (i.e. via mutagenesis), and their fitness, both singularly and in combination, is unknown although essential to prediction. Indeed, it may be dangerous to use experimental studies, involving highly idealized conditions, to make strong predictions about evolution in nature.

Ecological aspects of viral emergence

As with evolution, there are a number of simple ecological ‘rules’ that in part explain the patterns and processes of viral emergence [7,25]. Indeed, all instances of viral emergence are likely to have their roots in an ecological perturbation, such as a change in land-use or human encroachment into a new locality [26]. Hence, it is easy to predict that human disturbance is likely to lead to future cross-species transmission events. In addition, excellent work has been done using epidemiological theory and quantitative models to predict future disease risks [27], including risk maps for the emergence of specific diseases [28], and which should undoubtedly play a central role in disease control.

At its most basic level ecology explains the relationship between host population size and density and the propensity for emergence. Trivially, the larger and more dense a host population, the greater the number of pathogens that population can carry and the greater their virulence can be [29]. This in part explains why bats are being increasingly recognized as major hosts for a diverse array of viruses, some of which may eventually make their way into humans [30,31,32*]. Indeed, bats have been found to be important reservoirs for coronaviruses [33,34], filoviruses [35], lyssaviruses [36], paramyxoviruses [37–40], and have recently been observed to

harbor both hantaviruses [41] and influenza viruses [42**]. Hence, future surveying will undoubtedly reveal even more novel bat viruses, although determining which of these might emerge in humans is more challenging.

One aspect of ecology with potentially far more predictive power centers on the existence of so-called ‘hot spots’ for emergence [43]. Although a useful advance, the delineation of any hot-spots will be adversely affected by ascertainment bias, with the large number of emerging pathogens documented in the developed world likely reflecting more intensive surveillance and reporting. For example, although the first AIDS cases were described in the United States, the first major African epidemic documented in the east of this continent (e.g. Uganda), the current burden of HIV/AIDS is highest in southern Africa, the virus clearly originated in central-western Africa where the chimpanzee reservoir resides [44*]. In short, the place where an emergent event was first documented is not necessarily the place where the initial cross-species transmission event actually took place. More generally, given the ever-changing impact of humans on the natural world it is not obvious that hot spots for past emergence have strong predictive power for where emergence will occur next. Indeed, documenting changing levels of animal biodiversity may be equally informative [26].

Predicting the virulence of emerging viruses

It is equally important to determine how a new virus will evolve after it has successfully emerged. Central to this is understanding the evolution of virulence (disease severity). The appearance of any new infection is often accompanied by speculations on how virulence will evolve, with the continual debate over whether HIV will evolve reduced virulence serving as a case in point [45]. Although conventional wisdom posits that evolution makes pathogens benign in the long run, theory tells us that there should be an evolutionary trade-off between virulence and transmissibility [46], such that the direction of virulence evolution following successful emergence can be predicted in principle.

Although it is tempting to think that next generation sequencing will solve the ‘virulence problem’, genetic studies of virulence evolution are only informative if it is possible to associate viral strains with specific virulence phenotypes. It is for this reason that the attenuation of myxoma virus (MYXV) following its introduction as a biological control into the European rabbit populations of Australia and Europe, within which virulence grade was carefully measured, has become the textbook example of virulence evolution [47]. Critically, MYXV involves a species jump, albeit one mediated by humans; the natural host for this virus is the South American tapeti (*Sylvilagus brasiliensis*) in which the virus causes only mild tumors, whereas severe myxomatosis appears when the virus

infects nonresistant European rabbits (*Oryctolagus cuniculus*), although intermediate levels of virulence (which maximized transmission rate) ultimately dominated in the field [47]. An analysis of the genome-scale evolution of MYXV from the original releases in the 1950s in both Australia and Europe to modern wild strains revealed that changes in virulence involved multiple genes, with no mutations common to specific virulence grades [48]. Hence, despite the similarity in selection pressures in both Australia and Europe, there are multiple genetic routes to attain either highly virulent or attenuated phenotypes in MYXV. Such flexibility sits in stark contrast to RNA viruses such as West Nile virus (WNV) in which a single amino acid substitution increases virulence and has evolved convergently multiple times during the history of WNV [49]. The difference between MYXV and WNV suggests a more profound relationship between virulence evolution and genome size, in which there is greater scope for virulence determining mutations in larger DNA-based organisms, from which evolutionary trade-offs between virulence and transmission rate can be optimized at the ecological scale, while these trade-offs appear in RNA viruses because of limited genomic flexibility such that the same mutations affect both virulence and transmissibility. Hence, it is possible that there will only ever be a limited number of virulence determining mutations in RNA viruses.

Conclusions

It is perhaps naïve to think that emergence prediction will be ever more than an inexact science, revealing broad-scale generalities at best [8]. Despite this, our understanding of future epidemics can be improved. It is obviously important to continue metagenomic surveys of the pathogens that circulate in potential reservoir species [50,51,52,53], although these will be costly if many animals need to be surveyed. As many metagenomic studies are opportunistic [54], they might be better focused by collating the global species range of likely reservoir species [55] and dissecting their pathogen load in those parts of their home range that most often overlap with humans or which are most prone to human disturbance. However, it is important to recall that identifying a virus through its genome sequence is not the same as isolating a virus, and that its exact biological properties cannot easily be determined from sequence data alone.

More generally, it is critical to recall that cross-species transmission and emergence represents an intricate balance between ‘genetics’ — defined here as the mechanisms and determinants by which a virus is able to productively infect the cells of a new host species and spread to multiple individuals within that species — and ‘ecology’, representing the likelihood that animals are exposed to a specific pathogen and that there are sufficient connections to enable the virus to maintain its

spread at the epidemiological scale. Only when all these conditions are satisfied will an epidemic occur.

Finally, more attention should be devoted to revealing the common evolutionary and epidemiological patterns exhibited by those viruses that have successfully jumped species boundaries. For example, a comprehensive survey of the phylodynamic patterns [56] exhibited by currently circulating viruses will do much to help us understand how a new virus will evolve and spread once it has emerged. Specifically, it should be possible to compile a cross virus data base of the parameters that correlate most with successful emergence, such as how rapidly each virus evolves, its mode of transmission, its major host or vector species, its cell receptors of choice, key aspects of phenotype such as virulence and antigenicity, its population growth rate, its phylogeography, and whether it has jumped species boundaries in the past. Although such data will not enable us to predict future emergence with any certainty, they may allow broad-scale conclusions as to which groups of viruses are most likely to emerge in humans, which animal species in which geographical locations need to be surveyed most intensively, and how evolution will proceed following a host jump.

Acknowledgements

The author acknowledges financial support from an NHMRC Australia Fellowship and NIH/NIGMS grant R01 GM080533-06.

References and recommended reading

Papers of particular interest, published within the period of review, have been highlighted as:

- of special interest
- of outstanding interest

1. Grard G, Fair JN, Lee D, Slikas E, Steffen I, Muyembe JJ, Sittler T, Veeraraghavan N, Ruby JG, Wang C *et al.*: **A novel rhabdovirus associated with acute hemorrhagic fever in central Africa.** *PLoS Pathog* 2012, **8**:e1002924.
2. Briese T, Paweska JT, McMullan LK, Hutchison SK, Street C, Palacios G, Khristova ML, Weyer J, Swanepoel R, Egholm M *et al.*: **Genetic detection and characterization of Lujo virus, a new hemorrhagic fever-associated arenavirus from southern Africa.** *PLoS Pathog* 2009, **5**:e1000455.
Informative example of how modern molecular methods are able to rapidly determine the microbial cause of an outbreak of emerging disease.
3. Zaki AM, van Boheemen S, Bestebroer TM, Osterhaus AD, Fouchier RA: **Isolation of a novel coronavirus from a man with pneumonia in Saudi Arabia.** *N Engl J Med* 2012, **367**:1814-1820.
4. Pike BL, Saylor KE, Fair JN, Lebreton M, Tamoufe U, Djoko CF, Rimoin AW, Wolfe ND: **The origin and prevention of pandemics.** *Clin Infect Dis* 2010, **50**:1636-1640.
5. Woolhouse M, Scott F, Hudson Z, Howey R: **Chase-topping M: human viruses: discovery and emergence.** *Philos Trans R Soc Lond B Biol Sci* 2012, **367**:2864-2871.
Overview of the key issues, both ecological and genetic, involved in the study of disease emergence and which provides a useful framework for future work in this area.
6. Parrish CR, Holmes EC, Morens DM, Park E-C, Burke DS, Calisher CH, Laughlin CA, Saif LJ, Dazsak P: **Cross-species viral transmission and the emergence of new epidemic diseases.** *Microbiol Mol Biol Rev* 2008, **72**:457-470.

7. Woolhouse ME, Gowtage-Sequeria S: **Host range and emerging and reemerging pathogens.** *Emerg Infect Dis* 2005, **11**:1842-1847.
8. Woolhouse ME, Haydon DT, Antia R: **Emerging pathogens: the epidemiology and evolution of species jumps.** *Trends Ecol Evol* 2005, **20**:238-244.
9. Bush RM, Bender CA, Subbarao K, Cox NJ, Fitch WM: **Predicting the evolution of human influenza A.** *Science* 1999, **286**:1921-1925.
10. Pepin KM, Lass S, Pulliam JR, Read AF, Lloyd-Smith JO: **Identifying genetic markers of adaptation for surveillance of viral host jumps.** *Nat Rev Microbiol* 2010, **8**:802-813.
11. Streicker DG, Altizer SM, Velasco-Villa A, Rupprecht CE: **Variable evolutionary routes to host establishment across repeated rabies virus host shifts among bats.** *Proc Natl Acad Sci USA* 2012, **109**:19715-19720.
12. Lalic J, Elena SF: **Epistasis between mutations is host-dependent for an RNA virus.** *Biol Lett* 2013, **9**:20120396.
Reveals the complex nature of epistasis in RNA viruses, and particularly a host-dependent effect. Genetic factors such as epistasis will greatly complicate any attempt to predict future viral evolution and emergence.
13. Tsetsarkin KA, Chen R, Leal G, Forrester N, Higgs S, Huang J, Weaver SC: **Chikungunya virus emergence is constrained in Asia by lineage-specific adaptive landscapes.** *Proc Natl Acad Sci USA* 2011, **108**:7872-7877.
Another important example of how epistatic interactions among mutations influence the probability of emergence.
14. Bloom JD, Gong LI, Baltimore D: **Permissive secondary mutations enable the evolution of influenza oseltamivir resistance.** *Science* 2010, **328**:1272-1275.
Although there is an understandable focus on the emergence of mutations of phenotypic importance, such as those that confer antiviral resistance, this paper shows that the evolution of major fitness enhancing mutations may be associated with those that occur elsewhere in the viral genome. Surveying these 'permissive' mutations is therefore a useful contributor to studies of disease emergence.
15. Davies TJ, Pedersen AB: **Phylogeny and geography predict pathogen community similarity in wild primates and humans.** *Proc R Soc Lond B* 2008, **275**:1695-1701.
16. Streicker DG, Turmelle AS, Vonhof MJ, Kuzmin IV, McCracken GF, Rupprecht CE: **Host phylogeny constrains cross-species emergence and establishment of rabies virus in bats.** *Science* 2010, **329**:676-679.
Comprehensive analysis of how the probability of emergence (i.e. successful cross-species transmission) is mediated by the phylogenetic relationships of the host species in question, in this case focusing on bat rabies viruses.
17. Holmes EC: **The evolutionary genetics of emerging viruses.** *Annu Rev Ecol Syst* 2009, **40**:353-372.
18. Sharp CP, LeBreton M, Kantola K, Nana A, Dippo J, Djoko CF, Tamoufe U, Kiyang JA, Babila TG, Ngole EM *et al.*: **Widespread infection with homologues of human parvoviruses B19, PARV4, and human bocavirus of chimpanzees and gorillas in the wild.** *J Virol* 2010, **84**:10289-10296.
19. Simon-Loriere E, Holmes EC: **Why do RNA viruses recombine?** *Nat Rev Microbiol* 2011, **9**:617-626.
20. Coffey LL, Vasilakis N, Brault AC, Powers AM, Tripet F, Weaver SC: **Arbovirus evolution in vivo is constrained by host alternation.** *Proc Natl Acad Sci USA* 2008, **105**:6970-6975.
21. Woelk CH, Holmes EC: **Reduced positive selection in vector-borne RNA viruses.** *Mol Biol Evol* 2002, **19**:2333-2336.
22. Herfst S, Schrauwen EJ, Linster M, Chutinimitkul S, de Wit E, Munster VJ, Sorrell EM, Bestebroer TM, Burke DF, Smith DJ *et al.*: **Airborne transmission of influenza A/H5N1 virus between ferrets.** *Science* 2012, **336**:1534-1541.
23. Imai M, Watanabe T, Hatta M, Das SC, Ozawa M, Shinya K, Zhong G, Hanson A, Katsura H, Watanabe S *et al.*: **Experimental adaptation of an influenza H5 HA confers respiratory droplet transmission to a reassortant H5 HA/H1N1 virus in ferrets.** *Nature* 2012, **486**:420-428.
24. Russell CA, Fonville JM, Brown AE, Burke DF, Smith DL, James SL, Herfst S, van Boheemen S, Linster M, Schrauwen EJ *et al.*: **The potential for respiratory droplet-transmissible A/H5N1 influenza virus to evolve in a mammalian host.** *Science* 2012, **336**:1541-1547.
25. Woolhouse M, Gaunt E: **Ecological origins of novel human pathogens.** *Crit Rev Microbiol* 2007, **33**:231-242.
26. Keesing F, Belden LK, Daszak P, Dobson A, Harvell CD, Holt RD, Hudson P, Jolles A, Jones KE, Mitchell CE *et al.*: **Impacts of biodiversity on the emergence and transmission of infectious diseases.** *Nature* 2010, **468**:647-652.
27. Woolhouse M: **How to make predictions about future infectious disease risks.** *Philos Trans R Soc Lond B Biol Sci* 2011, **366**:2045-2054.
28. Fuller T, Thomassen HA, Mulembakani PM, Johnston SC, Lloyd-Smith JO, Kisalu NK, Lutete TK, Blumberg S, Fair JN, Wolfe ND *et al.*: **Using remote sensing to map the risk of human monkeypox virus in the Congo Basin.** *Ecohealth* 2011, **8**:14-25.
29. Anderson RM, May RM: *Infectious Diseases of Humans: Dynamics and Control.* Oxford University Press; 1991.
30. Donaldson EF, Haskew AN, Gates JE, Huynh J, Moore CJ, Frieman MB: **Metagenomic analysis of the viromes of three North American bat species: viral diversity among different bat species that share a common habitat.** *J Virol* 2010, **84**:13004-13018.
31. Wang LF, Walker PJ, Poon LL: **Mass extinctions, biodiversity and mitochondrial function: are bats 'special' as reservoirs for emerging viruses?** *Curr Opin Virol* 2011, **1**:649-657.
32. Wood JL, Leach M, Waldman L, Macgregor H, Fooks AR, Jones KE, Restif O, Dechmann D, Hayman DT, Baker KS *et al.*: **A framework for the study of zoonotic disease emergence and its drivers: spillover of bat pathogens as a case study.** *Philos Trans R Soc Lond B Biol Sci* 2012, **367**:2881-2892.
Another useful overview of the mechanisms that drive disease emergence with a particular focus on the growing evidence that bats are a rich source of viral biodiversity, some of which could potentially emerge in humans.
33. Huynh J, Li S, Yount B, Smith A, Sturges L, Olsen JC, Nagel J, Johnson JB, Agnihothram S, Gates JE *et al.*: **Evidence supporting a zoonotic origin of human coronavirus strain NL63.** *J Virol* 2012, **86**:12816-12825.
34. Li W, Shi Z, Yu M, Ren W, Smith C, Epstein JH, Wang H, Cramer G, Hu Z, Zhang H *et al.*: **Bats are natural reservoirs of SARS-like coronaviruses.** *Science* 2005, **310**:676-679.
35. Leroy EM, Kumulungui B, Pourrut X, Rouquet P, Hassanin A, Yaba P, Délicat A, Paweska JT, Gonzalez JP, Swanepoel R: **Fruit bats as reservoirs of Ebola virus.** *Nature* 2005, **438**:575-576.
36. Rupprecht CE, Turmelle A, Kuzmin IV: **A perspective on lyssavirus emergence and perpetuation.** *Curr Opin Virol* 2011, **1**:662-670.
37. Baker KS, Todd S, Marsh G, Fernandez-Loras A, Suu-Ire R, Wood JL, Wang LF, Murcia PR, Cunningham AA: **Co-circulation of diverse paramyxoviruses in an urban African fruit bat population.** *J Gen Virol* 2012, **93**:850-856.
38. Drexler JF, Corman VM, Müller MA, Maganga GD, Vallo P, Binger T, Gloza-Rausch F, Rasche A, Yordanov S, Seebens A *et al.*: **Bats host major mammalian paramyxoviruses.** *Nat Commun* 2012, **3**:796 <http://dx.doi.org/10.1038/ncomms1796>.
39. Mahalingam S, Herrero LJ, Playford EG, Spann K, Herring B, Rolph MS, Middleton D, McCall B, Field H, Wang LF: **Hendra virus: an emerging paramyxovirus in Australia.** *Lancet Infect Dis* 2012, **12**:799-807.
40. Weiss S, Nowak K, Fahr J, Wibbelt G, Mombouli JV, Parra HJ, Wolfe ND, Schneider BS, Leendertz FH: **Henipavirus-related sequences in fruit bat bushmeat, Republic of Congo.** *Emerg Infect Dis* 2012, **18**:1536-1537.
41. Weiss S, Witkowski PT, Auste B, Nowak K, Weber N, Fahr J, Mombouli JV, Wolfe ND, Drexler JF, Drosten C *et al.*: **Hantavirus in bat, Sierra Leone.** *Emerg Infect Dis* 2012, **18**:159-161.

42. Tong S, Li Y, Rivaille P, Conrardy C, Castillo DA, Chen LM, Recuenco S, Ellison JA, Davis CT, York IA *et al.*: **A distinct lineage of influenza A virus from bats.** *Proc Natl Acad Sci USA* 2012, **109**:4269-4274.

Bats contain a diverse array of both RNA and DNA viruses. Here a highly divergent lineage of influenza virus is observed in South American bats, and which may ultimately change our understanding of the origins and evolution of this perennially important human pathogen.

43. Jones KE, Patel NG, Levy MA, Storeygard A, Balk D, Gittleman JL, Daszak P: **Global trends in emerging infectious diseases.** *Nature* 2008, **451**:990-993.
44. Sharp PM, Hahn BH: **The evolution of HIV-1 and the origin of AIDS.** *Philos Trans R Soc Lond B Biol Sci* 2010, **365**:2487-2494. Although there are many outstanding issues in HIV research, the key questions on viral origins have been resolved and illustrate the power of molecular and evolutionary methods to reveal key aspects of disease emergence.
45. Ndung'u T, Weiss RA: **On HIV diversity.** *AIDS* 2012, **26**:1255-1260.
46. Bolker BM, Nanda A, Shah D: **Transient virulence of emerging pathogens.** *J R Soc Interface* 2010, **7**:811-822.
47. Fenner F, Ratcliffe FN: *Myxomatosis.* Cambridge University Press; 1965.
48. Kerr PJ, Ghedin E, DePasse JV, Fitch A, Cattadori IM, Hudson PJ, Tscharke DC, Read AF, Holmes EC: **Evolutionary history and attenuation of myxoma virus on two continents.** *PLoS Pathog* 2012, **8**:e1002950.

This study provides insights into the genomic basis of virulence evolution in the textbook example of myxoma virus. The central observation is that despite a similarity in selection pressures through time and across continents, and there is only convergent evolution for viral phenotype and not genotype.

49. Brault AC, Huang CY-H, Langevin SA, Kinney RM, Bowen RA, Ramey WN, Panella NA, Holmes EC, Powers AM, Miller BR: **A**

single positively selected West Nile viral mutation confers increased avian virogenesis in American crows. *Nat Genet* 2007, **39**:1162-1166.

50. Daszak P, Lipkin WI: **The search for meaning in virus discovery.** *Curr Opin Virol* 2011, **1**:620-623.
51. Ge X, Li Y, Yang X, Zhang H, Zhou P, Zhang Y, Shi Z: **Metagenomic analysis of viruses from bat fecal samples reveals many novel viruses in insectivorous bats in China.** *J Virol* 2012, **86**:4620-4630.
52. Kapoor A, Simmonds P, Gerold G, Qaisar N, Jain K, Henriquez JA, Firth C, Hirschberg DL, Rice CM, Shields S *et al.*: **Characterization of a canine homolog of hepatitis C virus.** *Proc Natl Acad Sci USA* 2011, **108**:11608-11613.

The majority of human RNA viruses have zoonotic origins. Until recently a notable outlier was hepatitis C virus (HCV). This paper revealed that a very similar virus circulates in dogs (animals which have also been identified as hosts for influenza virus), although whether human HCV emerged from dogs is uncertain.

53. Phan TG, Kapusinszky B, Wang C, Rose RK, Lipton HL, Delwart EL: **The fecal viral flora of wild rodents.** *PLoS Pathog* 2011, **7**:e1002218.
54. Ng TF, Marine R, Wang C, Simmonds P, Kapusinszky B, Bodhidatta L, Oderinde BS, Wommack KE, Delwart E: **High variety of known and new RNA and DNA viruses of diverse origins in untreated sewage.** *J Virol* 2012, **86**:12161-12175.
55. Jetz W, McPherson JM, Guralnick RP: **Integrating biodiversity distribution knowledge: toward a global map of life.** *Trends Ecol Evol* 2012, **27**:151-159.

An important initiative by ecologists to determine the global distribution of animal species. By associating viruses with specific reservoir hosts, this database could also be used to help predict the global distribution of potentially emerging diseases.

56. Holmes EC, Grenfell BT: **Discovering the phylodynamics of RNA viruses.** *PLoS Comput Biol* 2009, **5**:e1000505.