

## Therapeutic potential of targeting the renin angiotensin system in portal hypertension

Chandana B Herath, Josephine A Grace, Peter W Angus

Chandana B Herath, Josephine A Grace, Department of Medicine, University of Melbourne, Austin Health, Heidelberg, Victoria 3084, Australia

Josephine A Grace, Peter W Angus, Department of Gastroenterology, Austin Health, Heidelberg, Victoria 3084, Australia

Author contributions: Herath CB wrote and reviewed the manuscript; Grace JA contributed to the work and reviewed the manuscript; Angus PW reviewed and assisted with editing the manuscript.

Supported by Grant from the National Health and Medical Research Council of Australia

Correspondence to: Chandana B Herath, PhD, Department of Medicine, University of Melbourne, Austin Health, 145 Studley Road, Heidelberg, Victoria 3084, Australia. [cherath@unimelb.edu.au](mailto:cherath@unimelb.edu.au)

Telephone: +61-3-94962549 Fax: +61-3-94575485

Received: September 24, 2012 Revised: November 15, 2012

Accepted: January 5, 2013

Published online: February 15, 2013

### Abstract

Portal hypertension is responsible for the bulk of the morbidity and mortality in patients with cirrhosis. Drug therapy to reduce portal pressure involves targeting two vascular beds. The first approach is to reduce intra hepatic vascular tone induced by the activity of powerful vasoconstrictors such as angiotensin II, endothelin-1 and the sympathetic system and mediated *via* contraction of perisinusoidal myofibroblasts and perversular smooth muscle cells. The second approach is to reduce mesenteric and portal blood flow. Non-selective  $\beta$ -blockers are widely used and have been shown to prolong patient survival and reduce oesophageal variceal bleeding in advanced cirrhosis. However many patients are unable to tolerate these drugs and they are ineffective in a significant proportion of patients. Unfortunately there are no other drug therapies that have proven efficacy in the treatment of portal hypertension and prevention of variceal bleeding. This review briefly outlines current therapeutic approaches to the

management of portal hypertension, and the evidence supporting the role of the renin angiotensin system (RAS) and the use of RAS blockers in this condition. It will also outline recent advances in RAS research that could lead to the development of new treatments focusing in particular on the recently discovered "alternate axis" of the RAS.

© 2013 Baishideng. All rights reserved.

**Key words:** Angiotensin-(1-7); Portal hypertension; Intrahepatic resistance; Mesenteric vasodilatation; Variceal bleeding; Non-selective  $\beta$ -blockers; Renin angiotensin system; Mas receptor; Angiotensin receptor; Cirrhosis

Herath CB, Grace JA, Angus PW. Therapeutic potential of targeting the renin angiotensin system in portal hypertension. *World J Gastrointest Pathophysiol* 2013; 4(1): 1-11 Available from: URL: <http://www.wjgnet.com/2150-5330/full/v4/i1/1.htm> DOI: <http://dx.doi.org/10.4291/wjgp.v4.i1.1>

### INTRODUCTION

Hepatic fibrosis and its end-stage sequelae of cirrhosis and liver cancer are major causes of morbidity and mortality throughout the world and their prevalence is rising, largely due to the increasing impact of chronic viral hepatitis and non alcoholic steatohepatitis. Much of the morbidity and mortality that occurs in cirrhosis is due to the development of portal hypertension. However, despite major advances in our understanding of the pathogenesis of portal hypertension, current treatment options are limited.

It is clear that the renin angiotensin system (RAS) contributes to organ dysfunction and chronic tissue injury in a range of conditions including diabetes, cardiovascular and renal disease, primarily through the vasoactive and profibrotic effects of its key effector peptide, angiotensin II<sup>[1]</sup>. More recently, the RAS has also been

implicated in the pathogenesis of both hepatic fibrosis and portal hypertension<sup>[2,3]</sup>. This is supported by studies which have shown that RAS blockers are able to reduce fibrosis in experimental models of chronic liver injury and that they can lower portal pressure in both animal models and in man, primarily by inhibiting angiotensin II mediated intrahepatic vasoconstriction<sup>[4-6]</sup>. This review will briefly outline current therapeutic approaches to the management of portal hypertension, and focus on the evidence supporting the role of the RAS and the use of RAS blockers in this condition, and recent advances in RAS research that could lead to the development of new treatments.

## CURRENT TREATMENT OF PORTAL HYPERTENSION

The initiating mechanism for the development of portal hypertension is thought to be the development of increased intrahepatic resistance to portal inflow. This is mostly caused by increased deposition of extracellular matrix and disruption of the normal hepatic vascular architecture<sup>[2]</sup>. However, a significant proportion of portal resistance is attributable to intrahepatic vasoconstriction which is caused by the contraction of activated perisinusoidal hepatic stellate cells and of vascular smooth muscle cells in portal venules<sup>[7]</sup>. It is this variable component of intrahepatic resistance, mediated by powerful intrahepatic vasoconstrictors such as angiotensin II and endothelin<sup>[8-11]</sup>, which is potentially amenable to pharmacological therapies.

The second and equally important contributor in the development of portal hypertension is splanchnic vasodilatation which increases portal blood flow<sup>[12]</sup>. The mechanisms responsible for splanchnic vasodilatation in portal hypertension are incompletely understood, however, there is considerable evidence from both animal and human studies to suggest that nitric oxide (NO) generated by endothelial NO synthase (NOS) plays a central role<sup>[13-16]</sup>. One of the key consequences of portal hypertension is the development of portosystemic collaterals, of which the most important clinically are oesophageal varices. These vessels divert much of the increased mesenteric inflow away from the liver. However, even when portal blood flow is entirely diverted through collaterals, portal hypertension persists because of concomitant increases in portal venous inflow caused by increasing splanchnic vasodilatation. It should be noted that although the formation of these collaterals has been assumed to be the result of dilatation of pre-existing vascular channels, recent studies have implicated a process of neoangiogenesis which has been shown to contribute to both the formation of portosystemic collaterals and increased splanchnic blood flow<sup>[17]</sup>.

Bleeding from oesophageal varices is responsible for much of the mortality and morbidity that occurs in patients with portal hypertension. Varices are present in about 50% of patients at diagnosis, and this increases to

about 90% on long-term follow up. The risk of variceal rupture is 10%-30% per year depending on their size and appearance and the severity of liver disease, and the risk of mortality from a single episode is around 15%-20%<sup>[18]</sup>. Thus prevention or control of variceal bleeding has been the primary aim of the drug treatments that have been used in an attempt to lower portal pressure.

In theory portal pressure should fall in response to drugs that reduce portal inflow or those that lower intra hepatic resistance to portal inflow. The drugs most widely used in the prevention of variceal bleeding are non-selective  $\beta$ -blockers (NSBB). They reduce portal pressure by reducing cardiac output *via*  $\beta$ -1 receptors and causing splanchnic vasoconstriction by blocking  $\beta$ -2 receptors, resulting in unopposed  $\alpha$ -1 activity. Randomized clinical trials showed NSBB reduce portal pressure and the risk of bleeding from oesophageal varices<sup>[18-26]</sup>. However around 15% of cirrhotic patients are intolerant of NSBB treatment, and up to 60% fail to achieve the treatment response required to prevent variceal bleeding defined as a fall in hepatic venous pressure gradient (HVPG) to less than 12 mmHg or a decrease of greater than 20% from baseline<sup>[27]</sup>. Although portal pressure is directly correlated with the presence of varices, lowering pressure with NSBB does not prevent the development of varices in patients with cirrhosis<sup>[28]</sup>.

Another approach is to reduce intrahepatic resistance with drugs that increase the delivery of NO to the intrahepatic circulation (*e.g.*, nitrates), or drugs that block  $\alpha$ -adrenergic activity (*e.g.*, prazosin, clonidine). Although modest reduction in HVPG can be achieved with these drugs, their use as monotherapy is not recommended as they not only act on the intrahepatic circulation but also exert a vasodilatory effect on the systemic circulation, leading to arterial hypotension<sup>[29]</sup>. A recent small placebo-controlled randomized study showed that simvastatin, a drug that originally developed for hypercholesterolemia and shown to act through the posttranslational modification of endothelial NOS (eNOS), significantly reduced HVPG in cirrhotic patients without altering the blood flow. This suggested that simvastatin improved HVPG by reducing intrahepatic vascular resistance<sup>[30]</sup>.

A number of vasoconstrictor drugs which increase splanchnic vascular tone have been shown to be effective in controlling acute variceal bleeding. The vasopressin analogue, terlipressin, acts on vascular V1 receptors in both the mesenteric and systemic arterial beds to mediate vasoconstriction, and as a result, the drug lowers mesenteric inflow and portal pressure. Terlipressin is generally well-tolerated, but there remains a small incidence of ischaemic events which respond to cessation of the drug<sup>[31]</sup>. This drug reduces the relative risk of mortality from acute variceal bleeding by approximately one third<sup>[32]</sup>. Moreover, terlipressin (plus albumin) is the only treatment shown to prolong short-term survival in type 1 hepatorenal syndrome (HRS)<sup>[33]</sup>. Somatostatin and its analogues octreotide and vapreotide are splanchnic

vasoconstrictors which act by inhibiting glucagon secretion and by a local mesenteric vasoconstrictive effect<sup>[34]</sup>. Although these drugs have a role in the treatment of acute variceal bleeding in combination with endoscopic therapy<sup>[35]</sup>, they do not reduce mortality in this setting compared to endoscopic therapy alone<sup>[35,36]</sup> and are ineffective in HRS<sup>[37]</sup>.

In summary, NSBBs are widely accepted as the main pharmacotherapy currently available for prevention of variceal bleeding. However, a significant proportion of patient fail to achieve an optimal response or do not tolerate treatment<sup>[27]</sup> and no other drugs have an established role in the long-term treatment of portal hypertension. Thus there remains a major need to develop more safe and effective treatment options for the treatment of portal hypertension.

## NEW CONCEPTS IN RAS PHYSIOLOGY

In recent years it has been shown that the RAS is a much more complex enzymatic pathway than previously thought. It has been long recognized that the RAS plays a central role in cardiovascular and fluid homeostasis *via* the formation of the potent vasoconstrictor angiotensin II<sup>[38]</sup>. However it is now clear that in addition to its vasoactive roles the “classical” axis of the RAS, comprising angiotensin II, angiotensin converting enzyme (ACE) and the angiotensin II type 1 receptor (AT1R), plays a role in the wound healing response to chronic tissue injury and contributes to inflammation, cell proliferation and fibrogenesis<sup>[39,42]</sup>. In addition an “alternate” axis of the RAS has been characterized comprising ACE2, a structural homologue of ACE, its peptide product angiotensin-(1-7) and the Mas receptor, which has effects that counterbalance those mediated by the classical axis (Figure 1).

Early studies showed that angiotensin-(1-7) can be generated from angiotensin I by the actions of endopeptidases such as prolyl oligopeptidase<sup>[43]</sup> and thimet oligopeptidase<sup>[44]</sup> in tissue, and in the circulation by neutral endopeptidase<sup>[45]</sup>. Whilst the various endopeptidases have been shown to produce angiotensin-(1-7) depending upon their tissue localization and access to substrates, emerging evidence suggests that ACE2 which has a distinct enzyme activity<sup>[46]</sup>, plays a key role in angiotensin-(1-7) production in several tissues. ACE2 is able to generate angiotensin-(1-7) from angiotensin I indirectly through an intermediary peptide angiotensin-(1-9); however, in comparison, ACE2 has an approximately 400-fold higher substrate preference for angiotensin II<sup>[47]</sup> which suggests that ACE2 is important not only for production of angiotensin-(1-7) but also in degrading angiotensin II. Recently, Westwood and Chappell described another pathway in which angiotensin-(1-7) is generated directly from angiotensin-(1-12) or *via* angiotensin I generation<sup>[48]</sup>.

Angiotensin-(1-7), an effector peptide of the alternate axis of the RAS, is a vasodilator in several vascular


beds and has been shown to act mainly *via* its receptor Mas<sup>[49,54]</sup>. However, the existence of a receptor population that is insensitive to blockade with Mas receptor blocker A779 has also been reported<sup>[54,55]</sup>. It appears that angiotensin-(1-7), upon binding to its receptor, activates diverse pathways of intracellular signalling, leading to vasodilatation. For example, vasodilatory prostacyclin and/or NO appear to be involved in the response to angiotensin-(1-7) in the regional vascular beds<sup>[50,53,54,56-58]</sup>. It therefore appears that angiotensin-(1-7)-stimulated intracellular signaling leading to vasodilatation is depending upon the vascular bed under study and under differing pathophysiological condition.

Most components of the RAS are expressed in the liver, which is the primary source of angiotensinogen synthesis. Recent findings from our laboratory and others suggest that this intrahepatic RAS plays an important role in liver fibrosis since it is markedly upregulated in liver injury<sup>[11,59-61]</sup>, and blockade of the RAS improves experimental hepatic fibrosis<sup>[5,62-64]</sup>. The alternative axis of the RAS is also expressed in the liver and upregulated in liver disease leading to the generation of angiotensin-(1-7)<sup>[11,61,63]</sup>. The major pathway responsible for the generation of angiotensin-(1-7) in the cirrhotic liver is degradation of angiotensin II by ACE2 (Figure 2)<sup>[60]</sup> confirming previous *in vitro* findings that ACE2 has the highest substrate preference towards angiotensin II<sup>[47]</sup>. However there is limited data regarding the possible role of the alternate RAS in liver fibrosis and in modulating intrahepatic blood flow.


## RAS AND INTRAHEPATIC RESISTANCE

As outlined above, in patients with cirrhosis, the development of portal hypertension results from both an increase in the intrahepatic resistance to portal flow and an associated vasodilatation of the mesenteric vascular bed which leads to an increase in mesenteric blood flow. Splanchnic and systemic vasodilatation leads to secondary activation of vasoconstrictor pathways such as the sympathetic nervous system and the RAS in an attempt to maintain systemic vascular filling and blood pressure<sup>[67]</sup>. However these changes fail to correct the underlying circulatory hemodynamics<sup>[68]</sup>. There is now increasing evidence that in addition to its well recognized role in the homeostatic response to vasodilatation in cirrhosis the RAS may also play a primary role in the pathogenesis and maintenance of portal hypertension.

Hepatic structural changes such as tissue remodeling and scarring play a central role in increasing hepatic resistance to portal flow in the cirrhotic liver. However, when activated, hepatic stellate cells adopt a contractile myofibroblast phenotype, express the AT1R and have been shown *in vitro* to contract in response to angiotensin II and other vasoconstrictors such as endothelin-1<sup>[10,69]</sup>. The vasoconstriction response to angiotensin II is markedly increased in the perfused cirrhotic liver compared to normal livers, presumably mediated *via*



**Figure 1 Overview of the renin-angiotensin system.** The effects of the renin-angiotensin system (RAS) are determined by the balance between the angiotensin II (Ang II)-mediated “classical” axis, depicted in blue, which is vasoconstrictive and the angiotensin-(1-7) [Ang-(1-7)]-mediated “alternate” axis, depicted in orange, which is vasodilatory. Both Ang II and Ang-(1-7) can stimulate the Ang II type 2 (AT2) receptor, depicted in green; the effects of which are often analogous to those mediated by the Ang-(1-7) receptor Mas. Recent evidence indicates that a new member, Ang-(1-12), which is cleaved from angiotensinogen, also contributes either indirectly via Ang I or directly to the pool of Ang-(1-7). NEP: Neutral endopeptidase; ACE: Angiotensin converting enzyme; ACE2: Angiotensin converting enzyme 2; AT1 receptor: Angiotensin II type 1 receptor.




**Figure 2 Intrahepatic enzymatic pathways of renin-angiotensin system peptide production.** Schematic representation of the pathways responsible for the generation of vasodilator peptide angiotensin-(1-7) [Ang-(1-7)] in rat liver. The colour of the arrows represents the relative contribution of each pathway for *ex vivo* formation of Ang-(1-7) from the substrates angiotensin I (Ang I) and angiotensin II (Ang II) in cirrhotic rat liver. The most efficient pathway is represented by blue line, followed by green line. The black line is the least efficient pathway whilst red represents the efficiency less than green but higher than black. The yellow colour represents an efficient pathway to generate Ang-(1-7) directly from Ang I by the action of neutral endopeptidase (NEP) but it appears that this pathway is masked by angiotensin converting enzyme 2 in cirrhotic rat liver. Ang-(1-9): Angiotensin-(1-9); Ang-(1-5): Angiotensin-(1-5); ACE: Angiotensin converting enzyme; ACE2: Angiotensin converting enzyme 2; NEP: Neutral endopeptidase. Figure was adapted from our previous publication.

upregulated AT1R and AT1R expressing perivascular myofibroblasts<sup>[66]</sup>. Furthermore intrahepatic angiotensin II generation is increased in the cirrhotic liver<sup>[66]</sup>. These findings provide a rationale for the use of RAS blockers in the management of portal hypertension.

There is considerable evidence that another important contributor to elevated vascular tone in the cirrhotic liver is endothelial dysfunction of the hepatic microcirculation which diminishes the response to vasodilators<sup>[70]</sup>. It has been proposed that the reduced activity of hepatic vascular eNOS with concomitant reduction in NO synthesis impairs intrahepatic vasodilatation and thus, shifts the balance towards vasoconstriction<sup>[71]</sup>. This reduction in eNOS activity is linked to an increase in the expression of caveolin, a protein which is highly expressed in endothelial cells of the hepatic vasculature with predominant expression found in venous and sinusoidal endothelial cells in cirrhotic livers<sup>[72,73]</sup>. Interestingly, the calcium binding protein calmodulin competitively binds eNOS and reduces caveolin binding, thus increasing eNOS activity<sup>[72]</sup>.

Recent findings from our laboratory demonstrated that in *in-situ* perfused cirrhotic rat liver elicited a marked endothelium-dependent vasodilatory effect of exogenous angiotensin-(1-7) on the vasoconstrictive response evoked by angiotensin II (Figure 3)<sup>[55]</sup>. This finding suggests that as in other vascular beds<sup>[50,51,53,54]</sup>, in the cirrhotic liver angiotensin-(1-7) may cause a vasodilatory response that antagonizes the increase in portal pressure mediated by angiotensin II and other local vasoconstrictors. Although eNOS activity was not measured in this



**Figure 3 Angiotensin-(1-7)-induced portal pressure reduction.** Portal pressure changes in response to angiotensin II (Ang II) in cirrhotic rat liver. Portal pressure responses were measured using a vertically positioned graduated fluid-filled column open to atmospheric pressure. Ang II bolus (60 pmole) was injected into the portal vein of *in-situ* perfused cirrhotic rat liver preparations in the presence of an angiotensin converting enzyme (ACE) inhibitor lisinopril (0.7  $\mu$ mol/L). A: Percentage increases in pressure in response to Ang II bolus in the absence (closed circles) or presence (open circles) of angiotensin-(1-7) [Ang-(1-7), 0.7  $\mu$ mol/L]. The highest pressure that was recorded 45 s after the Ang II bolus injection was designated as 100% response and all other responses were calculated relative to this maximal response. B: The total area under Ang II response curve (AUC) in the absence (filled bar) or presence (open bar) of Ang-(1-7). Portal pressure change at each time point in panel A and AUC in B represents the mean  $\pm$  SEM from 21 cirrhotic rat liver preparations. Pre-incubation with Ang-(1-7) significantly reduced the portal pressure response evoked by Ang II bolus injection. <sup>a</sup> $P < 0.05$ , <sup>b</sup> $P < 0.01$  vs those with Ang II alone.

study, the eNOS inhibitor nitro-L-arginine methyl ester, L-NAME, completely abolished eNOS phosphorylation at Ser1177 and the response to angiotensin-(1-7)<sup>[55]</sup>. Whilst bradykinin *via* its B2 receptor mediates vasodilatation in response to angiotensin-(1-7) in porcine and canine coronary arteries<sup>[56,58,74]</sup>, it increases intrahepatic resistance and portal pressure<sup>[75]</sup>, possibly through acting on B2 receptors on stellate cells. However, angiotensin-(1-7)-induced vasodilatation in the cirrhotic liver was not affected by bradykinin B2 receptor blockade<sup>[55]</sup>. Possible mechanisms for these effects of angiotensin-(1-7) in the cirrhotic rat liver include increased phosphorylation of eNOS Ser1177<sup>[55]</sup> with simultaneous dephosphorylation at Thr495 and/or effects on calmodulin binding<sup>[76]</sup>. These findings suggest that it may be possible to reduce intrahepatic resistance and portal pressure by targeting the alternate axis of the RAS in the liver.

## RAS AND SPLANCHNIC VASODILATION

In contrast to intrahepatic hypervascular tone, the systemic and splanchnic circulation in cirrhosis is characterized by vasodilatation and hyporesponsiveness to vasoconstrictors including angiotensin II<sup>[8,67,77]</sup>. Interestingly, recent studies have shown that systemic levels of the vasodilatory peptide angiotensin-(1-7) increase as liver fibrosis progresses, whereas angiotensin II levels do not generally rise until cirrhosis is established with concurrent portal hypertension<sup>[65,78]</sup>. Furthermore, regional levels of the hormone are different from systemic levels such that in cirrhotic patients at transplant, the angiotensin-(1-7)/angiotensin II ratio is elevated in the splanchnic compared to the peripheral circulation, and negatively cor-

relates with systemic vascular resistance<sup>[78]</sup>. The recent findings suggest that angiotensin-(1-7) might contribute to vasodilatation in cirrhosis. Recent work from our laboratory provides support for this hypothesis<sup>[79]</sup>. We have shown that ACE2 is upregulated in cirrhotic mesenteric vessels and although angiotensin-(1-7) has no effect in the normal mesenteric circulation, it significantly reduces mesenteric vascular contractility in cirrhotic mesenteric beds *via* activation of the Mas receptor and the release of NO<sup>[79]</sup>.

Studies using isolated vessel preparations from cirrhotic animal models or portal hypertensive rats have led to the concept that vasodilatation is also linked to an intrinsic vascular hyporesponsiveness to endogenous vasoconstrictors such as angiotensin II,  $\alpha$ -adrenergic agonists and endothelin-1<sup>[80-82]</sup>. This concept is supported by the findings that peripheral vessels are hyporeactive to angiotensin II,  $\alpha$ -adrenergic agonists and endothelin-1 from cirrhotic animals and cirrhotic patients<sup>[77,83-90]</sup>, despite the fact that expression of AT1R and  $\alpha$ -adrenergic receptor subtypes 1a and 1b in the peripheral vessels is either normal or upregulated in both cirrhotic animals and patients<sup>[79,87,91]</sup>. However, previous studies in patients with cirrhosis or in peripheral resistance vessels obtained from such patients reported variable results in this regard<sup>[84,90,92]</sup>, probably attributable to the differences between conditions in different studies. Indeed, small resistance omental vessels from cirrhotic patients had a larger vasoconstriction response to  $\alpha$ -adrenergic agonists norepinephrine and methoxamine than similar vessels from healthy controls<sup>[92]</sup>. The same vessels vasodilated in response to substance P but this was inhibited by blocking NO or prostacyclin synthesis, suggesting that intrinsic hyporeactivity that is present in the peripheral circulation in cirrhosis is related to increased levels of NO

and prostanoids<sup>[93-95]</sup>. Hyporeactivity to angiotensin II infusion is also improved after inhibiting systemic NO production using an NO-clamp in cirrhotic patients<sup>[85]</sup>. This is in keeping with a wealth of literature suggesting that vasodilatory molecules including angiotensin-(1-7) are produced in excess in cirrhosis and that the final *in vivo* pressor effect is governed by a balance between the pressor and depressor arms of the circulation<sup>[96]</sup>.

There are also data, from both *in vitro* and *in vivo* studies, suggesting that vascular hyporeactivity to a range of endogenous pressors is attributed to changes that are downstream of the G-protein coupled receptors<sup>[86-88]</sup>. Evidence supporting the existence of vascular endothelium/NO independent pathways comes from studies in which endothelium denudation and pharmacological blockade of NOS in isolated vessels from cirrhotic animals and patients failed to improve vascular hyporeactivity to a range of vasoconstrictors<sup>[83,84,86,87,97-99]</sup>. One of the important NO-independent pathways is an impaired signaling by RhoA and Rho kinase, leading to a decreased phosphorylation of Ca<sup>2+</sup>-sensitizing proteins and increased myosin light chain phosphatase activity<sup>[87]</sup>. Moreover, increased expression of receptor desensitizing proteins, G protein-coupled receptor kinase 2 and  $\beta$ -arrestin-2, have also been implicated in this hyporeactivity to angiotensin II in vessels isolated from cirrhotic patients and rats<sup>[87]</sup>. It was also shown that mesenteric arteries from portal hypertensive rats had a reduced level of membrane associated RhoA, probably reflecting a diminished activity of RhoA/Rho kinase pathway which in turn results in increased activity of myosin light chain phosphatase and vasodilatation<sup>[68,100]</sup>.

## THERAPIES TARGETING THE RAS IN PORTAL HYPERTENSION

### **Therapeutic potential of the classical RAS**

The evidence from studies in experimental cirrhosis showing the angiotensin II contributes to the variable component of intrahepatic resistance in portal hypertension have provided a rationale for a number of trials examining the effects of ACE inhibitors and angiotensin receptors blockers (ARBs)<sup>[5,23,87]</sup> on portal pressure. Unfortunately many of these studies are small or non-randomized and there is very little long-term data. However Tandon *et al*<sup>[23]</sup> in a recent meta-analysis of individual patient data from three and nine studies that used ACE inhibitors and ARBs, respectively, showed that patients with Child Pugh A cirrhosis receiving ARBs/ACE inhibitors had a similar reduction in HVPG (17%) compared to patients with Pugh A cirrhosis that received NSBB (21%). There was no improvement of HVPG in patients with Child Pugh B/C cirrhosis receiving ARBs. Furthermore, several studies reported that RAS blockade can result in significant hypotension and renal impairment in patients with decompensated (Child Pugh B/C) cirrhosis<sup>[23]</sup> in whom there is activation of the systemic RAS.

Thus, although ARBs/ACE inhibitors do lower por-


tal pressure in early cirrhotic patients where the activation of RAS may be a predominant pathway responsible for increased intrahepatic tone, they have less effect in late stages of cirrhosis. This probably reflects the fact that the hypotension induced by RAS blockers increases activation of other vasoconstrictive pathways such as the sympathetic nervous system that in turn increase intrahepatic vascular tone<sup>[28,101-103]</sup>. Further studies are needed to clarify whether this class of drugs could be useful in the prevention of variceal bleeding in patients with compensated cirrhosis as an alternative to or possibly even in combination with  $\beta$ -blockers.

### **Alternate RAS-a novel potential target for the treatment of portal hypertension**

Recent animal studies focusing on the alternate RAS have led to the suggestion that new generation antihypertensives developed to target this axis could serve as effective therapeutic agents to treat arterial, pulmonary and portal hypertension<sup>[1,65,104,105]</sup>. Recent work outlined in this review demonstrates the presence of all of the key components of the alternate RAS in the liver and mesenteric vasculature of both healthy and cirrhotic animals as well as in the liver of healthy and cirrhotic patients<sup>[11,61,66,79]</sup>. Furthermore circulating angiotensin-(1-7) levels are increased<sup>[11,65,78]</sup> and the system is upregulated in the liver and mesenteric circulation in cirrhosis suggesting that it plays an important role in the pathophysiology of hepatic fibrosis and portal hypertension (Figure 4). This evidence linking elevated angiotensin-(1-7) levels to mesenteric and systemic vasodilatation in cirrhosis suggests that blocking the alternate axis could reduce mesenteric flow and thus lower portal pressure. In line with this, Mas receptor blockade would provide an intervention option in portal hypertension<sup>[79]</sup> as this treatment regime does not appear to compromise the vasodilatory response of angiotensin-(1-7) within the hepatic vasculature in experimental cirrhosis<sup>[55]</sup>. Further studies are clearly needed examining the effects of Mas blockade and angiotensin-(1-7) on hepatic and mesenteric haemodynamics in experimental cirrhosis *in vivo*.

## CONCLUSION

Recent developments in our understanding of the complexities of the RAS and its role in the pathogenesis of chronic liver disease and portal hypertension have opened up new therapeutic possibilities. It is clear that the classical axis of the RAS and its key effector peptide angiotensin II play a central role in hepatic fibrogenesis and in regulating intrahepatic vascular tone in cirrhosis and that despite the mixed results achieved in previous trials, consideration should be given to further prospective studies examining the effects of RAS blockers in patients with compensated cirrhosis. There is also fascinating new evidence showing that there is increased regional production of angiotensin-(1-7) in the mesenteric vascular bed in cirrhosis, and that this vasodilatory



**Figure 4 Overview of the renin angiotensin system-mediated pathophysiological changes in portal hypertension.** In cirrhosis, the effects of classical axis of the renin angiotensin system (RAS), mediated by its potent vasoconstrictor peptide angiotensin II (Ang II), are predominant within the hepatic vasculature, resulting in increased hepatic resistance to portal inflow. In this, apart from fixed barrier due to increased deposition of extracellular matrix (ECM) proteins, vascular tone is exacerbated by Ang II action on myfibroblastic cells (activated hepatic stellate cells-HSCs) and vascular smooth muscle cells (VSMCs). In contrast, the effects of the alternate axis of the RAS, mediated by its vasodilator peptide angiotensin-(1-7) (Ang1-7), are predominant within the splanchnic vasculature, resulting in increased nitric oxide (NO) production by vascular endothelial cells (ECs). This consequently exacerbates portal hypertension as a result of increased inflow to the portal circulation. Ang1-5: Angiotensin-(1-5); ACE: Angiotensin converting enzyme; ACE2: Angiotensin converting enzyme 2; eNOS: Endothelial nitric oxide synthase; Bk: Bradykinin; BkB2R: Bradykinin B2 receptor; AT1R: Angiotensin II type 1 receptor; AT2R: Angiotensin II type 2 receptor; MasR: Mas receptor.

peptide of the alternate axis of the RAS, contributes to mesenteric vasodilatation and the hyperdynamic circulation in cirrhosis. These novel data suggest that ACE2-angiotensin-(1-7)-Mas receptor axis is a potential target for the management of portal hypertension.

**REFERENCES**

- 1 Ferrario CM, Trask AJ, Jessup JA. Advances in biochemical and functional roles of angiotensin-converting enzyme 2 and angiotensin-(1-7) in regulation of cardiovascular function. *Am J Physiol Heart Circ Physiol* 2005; **289**: H2281-H2290 [PMID: 16055515 DOI: 10.1152/ajpheart.00618.2005]
- 2 Bataller R, Brenner DA. Liver fibrosis. *J Clin Invest* 2005; **115**: 209-218 [PMID: 15690074 DOI: 10.1172/JCI200524282]
- 3 Töx U, Steffen HM. Impact of inhibitors of the Renin-Angiotensin-aldosterone system on liver fibrosis and portal hypertension. *Curr Med Chem* 2006; **13**: 3649-3661 [PMID: 17168728 DOI: 10.2174/092986706779026138]
- 4 Dickstein K, Kjekshus J. Effects of losartan and captopril on mortality and morbidity in high-risk patients after acute myocardial infarction: the OPTIMAAL randomised trial. *Lancet* 2002; **360**: 752-760 [PMID: 12241832 DOI: 10.1016/S0140-6736(02)09895-1]
- 5 Paizis G, Gilbert RE, Cooper ME, Murthi P, Schembri JM, Wu LL, Rumble JR, Kelly DJ, Tikellis C, Cox A, Smallwood RA, Angus PW. Effect of angiotensin II type 1 receptor blockade on experimental hepatic fibrogenesis. *J Hepatol* 2001; **35**: 376-385 [PMID: 11592599 DOI: 10.1016/S0168-8278(01)00146-5]
- 6 Parving HH, Persson F, Lewis JB, Lewis EJ, Hollenberg NK. Aliskiren combined with losartan in type 2 diabetes and nephropathy. *N Engl J Med* 2008; **358**: 2433-2446 [PMID: 18525041 DOI: 10.1056/NEJMoa0708379]

- 7 Pinzani M, Gentilini P. Biology of hepatic stellate cells and their possible relevance in the pathogenesis of portal hypertension in cirrhosis. *Semin Liver Dis* 1999; **19**: 397-410 [PMID: 10643625 DOI: 10.1055/s-2007-1007128]
- 8 Angus PW. Role of endothelin in systemic and portal resistance in cirrhosis. *Gut* 2006; **55**: 1230-1232 [PMID: 16905694 DOI: 10.1136/gut.2005.088633]
- 9 Bataller R, Gäbele E, Schoonhoven R, Morris T, Lehnert M, Yang L, Brenner DA, Rippe RA. Prolonged infusion of angiotensin II into normal rats induces stellate cell activation and proinflammatory events in liver. *Am J Physiol Gastrointest Liver Physiol* 2003; **285**: G642-G651 [PMID: 12773299]
- 10 Bataller R, Gines P, Nicolas JM, Gorbis MN, Garcia-Ramallo E, Gasull X, Bosch J, Arroyo V, Rodes J. Angiotensin II induces contraction and proliferation of human hepatic stellate cells. *Gastroenterology* 2000; **118**: 1149-1156 [PMID: 10833490 DOI: 10.1016/S0016-5085(00)70368-4]
- 11 Herath CB, Warner FJ, Lubel JS, Dean RG, Jia Z, Lew RA, Smith AI, Burrell LM, Angus PW. Upregulation of hepatic angiotensin-converting enzyme 2 (ACE2) and angiotensin-(1-7) levels in experimental biliary fibrosis. *J Hepatol* 2007; **47**: 387-395 [PMID: 17532087 DOI: 10.1016/j.jhep.2007.03.008]
- 12 Wiest R, Groszmann RJ. Nitric oxide and portal hypertension: its role in the regulation of intrahepatic and splanchnic vascular resistance. *Semin Liver Dis* 1999; **19**: 411-426 [PMID: 10643626 DOI: 10.1055/s-2007-1007129]
- 13 Albornoz L, Motta A, Alvarez D, Estevez A, Bandi JC, McCormack L, Matera J, Bonofiglio C, Ciardullo M, De Santibanes E, Gimeno M, Gadan A. Nitric oxide synthase activity in the splanchnic vasculature of patients with cirrhosis: relationship with hemodynamic disturbances. *J Hepatol* 2001; **35**: 452-456 [PMID: 11682028 DOI: 10.1016/S0168-8278(01)00168-4]
- 14 Sarela AI, Mihaimeed FM, Batten JJ, Davidson BR, Mathie

- RT. Hepatic and splanchnic nitric oxide activity in patients with cirrhosis. *Gut* 1999; **44**: 749-753 [PMID: 10205218 DOI: 10.1136/gut.44.5.749]
- 15 Sieber CC, Lopez-Talavera JC, Groszmann RJ. Role of nitric oxide in the in vitro splanchnic vascular hyporeactivity in ascitic cirrhotic rats. *Gastroenterology* 1993; **104**: 1750-1754 [PMID: 8388839]
  - 16 Wiest R, Shah V, Sessa WC, Groszmann RJ. NO overproduction by eNOS precedes hyperdynamic splanchnic circulation in portal hypertensive rats. *Am J Physiol* 1999; **276**: G1043-G1051 [PMID: 10198349]
  - 17 Fernandez M, Vizzutti F, Garcia-Pagan JC, Rodes J, Bosch J. Anti-VEGF receptor-2 monoclonal antibody prevents portal-systemic collateral vessel formation in portal hypertensive mice. *Gastroenterology* 2004; **126**: 886-894 [PMID: 14988842 DOI: 10.1053/j.gastro.2003.12.012]
  - 18 D'Amico G, Pagliaro L, Bosch J. The treatment of portal hypertension: a meta-analytic review. *Hepatology* 1995; **22**: 332-354 [PMID: 7601427 DOI: 10.1002/hep.1840220145]
  - 19 García-Pagán JC, Villanueva C, Albillos A, Bañares R, Morillas R, Abraldes JG, Bosch J. Nadolol plus isosorbide mononitrate alone or associated with band ligation in the prevention of recurrent bleeding: a multicentre randomised controlled trial. *Gut* 2009; **58**: 1144-1150 [PMID: 19218249]
  - 20 Groszmann RJ, Bosch J, Grace ND, Conn HO, Garcia-Tsao G, Navasa M, Alberts J, Rodes J, Fischer R, Bermann M. Hemodynamic events in a prospective randomized trial of propranolol versus placebo in the prevention of a first variceal hemorrhage. *Gastroenterology* 1990; **99**: 1401-1407 [PMID: 2210246]
  - 21 Pagliaro L, D'Amico G, Sørensen TI, Lebrec D, Burroughs AK, Morabito A, Tiné F, Politi F, Traina M. Prevention of first bleeding in cirrhosis. A meta-analysis of randomized trials of nonsurgical treatment. *Ann Intern Med* 1992; **117**: 59-70 [PMID: 1350716]
  - 22 Patch D, Sabin CA, Goulis J, Gerunda G, Greenslade L, Merkel C, Burroughs AK. A randomized, controlled trial of medical therapy versus endoscopic ligation for the prevention of variceal rebleeding in patients with cirrhosis. *Gastroenterology* 2002; **123**: 1013-1019 [PMID: 12360462 DOI: 10.1053/gast.2002.35955]
  - 23 Tandon P, Abraldes JG, Berzigotti A, Garcia-Pagan JC, Bosch J. Renin-angiotensin-aldosterone inhibitors in the reduction of portal pressure: a systematic review and meta-analysis. *J Hepatol* 2010; **53**: 273-282 [PMID: 20570385 DOI: 10.1016/j.jhep.2010.03.013]
  - 24 Villanueva C, Aracil C, Colomo A, Lopez-Balaguer JM, Piqueras M, Gonzalez B, Torras X, Guarner C, Balanzo J. Clinical trial: a randomized controlled study on prevention of variceal rebleeding comparing nadolol + ligation vs. hepatic venous pressure gradient-guided pharmacological therapy. *Aliment Pharmacol Ther* 2009; **29**: 397-408 [PMID: 19006538 DOI: 10.1111/j.1365-2036.2008.03880.x]
  - 25 Villanueva C, Balanzó J, Novella MT, Soriano G, Sáinz S, Torras X, Cussó X, Guarner C, Vilardell F. Nadolol plus isosorbide mononitrate compared with sclerotherapy for the prevention of variceal rebleeding. *N Engl J Med* 1996; **334**: 1624-1629 [PMID: 8628357 DOI: 10.1056/NEJM199606203342502]
  - 26 Villanueva C, Miñana J, Ortiz J, Gallego A, Soriano G, Torras X, Sáinz S, Boadas J, Cussó X, Guarner C, Balanzó J. Endoscopic ligation compared with combined treatment with nadolol and isosorbide mononitrate to prevent recurrent variceal bleeding. *N Engl J Med* 2001; **345**: 647-655 [PMID: 11547718 DOI: 10.1056/NEJMoa003223]
  - 27 Feu F, Garcia-Pagan JC, Bosch J, Luca A, Teres J, Escorsell A, Rodes J. Relation between portal pressure response to pharmacotherapy and risk of recurrent variceal haemorrhage in patients with cirrhosis. *Lancet* 1995; **346**: 1056-1059 [PMID: 7564785 DOI: 10.1016/S0140-6736(95)91740-3]
  - 28 Groszmann RJ, Garcia-Tsao G, Bosch J, Grace ND, Burroughs AK, Planas R, Escorsell A, Garcia-Pagan JC, Patch D, Matloff DS, Gao H, Makuch R. Beta-blockers to prevent gastroesophageal varices in patients with cirrhosis. *N Engl J Med* 2005; **353**: 2254-2261 [PMID: 16306522 DOI: 10.1056/NEJMoa044456]
  - 29 García-Tsao G. Current management of the complications of cirrhosis and portal hypertension: variceal hemorrhage, ascites, and spontaneous bacterial peritonitis. *Gastroenterology* 2001; **120**: 726-748 [PMID: 11179247 DOI: 10.1053/gast.2001.22580]
  - 30 Abraldes JG, Albillos A, Bañares R, Turnes J, González R, García-Pagán JC, Bosch J. Simvastatin lowers portal pressure in patients with cirrhosis and portal hypertension: a randomized controlled trial. *Gastroenterology* 2009; **136**: 1651-1658 [PMID: 19208350 DOI: 10.1053/j.gastro.2009.01.043]
  - 31 Testro AG, Wongseelashote S, Angus PW, Gow PJ. Long-term outcome of patients treated with terlipressin for types 1 and 2 hepatorenal syndrome. *J Gastroenterol Hepatol* 2008; **23**: 1535-1540 [PMID: 17784863 DOI: 10.1111/j.1440-1746.2007.05176.x]
  - 32 Ioannou G, Doust J, Rockey DC. Terlipressin for acute esophageal variceal hemorrhage. *Cochrane Database Syst Rev* 2003; (1): CD002147 [PMID: 12535432]
  - 33 Gluud LL, Christensen K, Christensen E, Krag A. Systematic review of randomized trials on vasoconstrictor drugs for hepatorenal syndrome. *Hepatology* 2010; **51**: 576-584 [PMID: 19885875 DOI: 10.1002/hep.23286]
  - 34 Wiest R, Tsai MH, Groszmann RJ. Octreotide potentiates PKC-dependent vasoconstrictors in portal-hypertensive and control rats. *Gastroenterology* 2001; **120**: 975-983 [PMID: 11231951 DOI: 10.1053/gast.2001.22529]
  - 35 Bañares R, Albillos A, Rincón D, Alonso S, González M, Ruiz-del-Arbol L, Salcedo M, Molinero LM. Endoscopic treatment versus endoscopic plus pharmacologic treatment for acute variceal bleeding: a meta-analysis. *Hepatology* 2002; **35**: 609-615 [PMID: 11870374 DOI: 10.1053/jhep.2002.31354]
  - 36 D'Amico G, Politi F, D'Antonio A, Giannuoli G, Pasta L, Vizziani G, Traina M, Morabito A, Pagliaro L. Second study shows that octreotide may prevent early rebleeding in cirrhosis. *BMJ* 1998; **316**: 1320 [PMID: 9554914 DOI: 10.1136/bmj.316.7140.1320]
  - 37 Pomier-Layrargues G, Paquin SC, Hassoun Z, Lafortune M, Tran A. Octreotide in hepatorenal syndrome: a randomized, double-blind, placebo-controlled, crossover study. *Hepatology* 2003; **38**: 238-243 [PMID: 12830007 DOI: 10.1053/jhep.2003.50276]
  - 38 Lubel JS, Herath CB, Burrell LM, Angus PW. Liver disease and the renin-angiotensin system: recent discoveries and clinical implications. *J Gastroenterol Hepatol* 2008; **23**: 1327-1338 [PMID: 18557800 DOI: 10.1111/j.1440-1746.2008.05461.x]
  - 39 Kagami S, Border WA, Miller DE, Noble NA. Angiotensin II stimulates extracellular matrix protein synthesis through induction of transforming growth factor-beta expression in rat glomerular mesangial cells. *J Clin Invest* 1994; **93**: 2431-2437 [PMID: 8200978 DOI: 10.1172/JCI117251]
  - 40 Marrero MB, Schieffer B, Paxton WG, Heerdt L, Berk BC, Delafontaine P, Bernstein KE. Direct stimulation of Jak/STAT pathway by the angiotensin II AT1 receptor. *Nature* 1995; **375**: 247-250 [PMID: 7746328 DOI: 10.1038/375247a0]
  - 41 Mezzano SA, Ruiz-Ortega M, Egido J. Angiotensin II and renal fibrosis. *Hypertension* 2001; **38**: 635-638 [PMID: 11566946 DOI: 10.1161/hy09t1.094234]
  - 42 Suzuki Y, Ruiz-Ortega M, Lorenzo O, Ruperez M, Esteban V, Egido J. Inflammation and angiotensin II. *Int J Biochem Cell Biol* 2003; **35**: 881-900 [PMID: 12676174 DOI: 10.1016/S1357-2725(02)00271-6]
  - 43 Santos RA, Brosnihan KB, Jacobsen DW, DiCorleto PE, Ferrario CM. Production of angiotensin-(1-7) by human vascular endothelium. *Hypertension* 1992; **19**: II56-II61 [PMID: 1310484 DOI: 10.1161/01.HYP.19.2\_Suppl.II56]



- 44 **Chappell MC**, Gomez MN, Pirro NT, Ferrario CM. Release of angiotensin-(1-7) from the rat hindlimb: influence of angiotensin-converting enzyme inhibition. *Hypertension* 2000; **35**: 348-352 [PMID: 10642323 DOI: 10.1161/01.HYP.35.1.348]
- 45 **Yamamoto K**, Chappell MC, Brosnihan KB, Ferrario CM. In vivo metabolism of angiotensin I by neutral endopeptidase (EC 3.4.24.11) in spontaneously hypertensive rats. *Hypertension* 1992; **19**: 692-696 [PMID: 1317352 DOI: 10.1161/01.HYP.19.6.692]
- 46 **Warner FJ**, Smith AI, Hooper NM, Turner AJ. Angiotensin-converting enzyme-2: a molecular and cellular perspective. *Cell Mol Life Sci* 2004; **61**: 2704-2713 [PMID: 15549171 DOI: 10.1007/s00018-004-4240-7]
- 47 **Vickers C**, Hales P, Kaushik V, Dick L, Gavin J, Tang J, Godbout K, Parsons T, Baronas E, Hsieh F, Acton S, Patane M, Nichols A, Tummino P. Hydrolysis of biological peptides by human angiotensin-converting enzyme-related carboxypeptidase. *J Biol Chem* 2002; **277**: 14838-14843 [PMID: 11815627 DOI: 10.1074/jbc.M200581200]
- 48 **Westwood BM**, Chappell MC. Divergent pathways for the angiotensin-(1-12) metabolism in the rat circulation and kidney. *Peptides* 2012; **35**: 190-195 [PMID: 22490446 DOI: 10.1016/j.peptides.2012.03.025]
- 49 **Benter IF**, Yousif MH, Anim JT, Cojocel C, Diz DI. Angiotensin-(1-7) prevents development of severe hypertension and end-organ damage in spontaneously hypertensive rats treated with L-NAME. *Am J Physiol Heart Circ Physiol* 2006; **290**: H684-H691 [PMID: 16403946 DOI: 10.1152/ajp-heart.00632.2005]
- 50 **Castro CH**, Santos RA, Ferreira AJ, Bader M, Alenina N, Almeida AP. Evidence for a functional interaction of the angiotensin-(1-7) receptor Mas with AT1 and AT2 receptors in the mouse heart. *Hypertension* 2005; **46**: 937-942 [PMID: 16157793 DOI: 10.1161/01.HYP.0000175813.04375.8a]
- 51 **Lemos VS**, Silva DM, Walther T, Alenina N, Bader M, Santos RA. The endothelium-dependent vasodilator effect of the nonpeptide Ang(1-7) mimic AVE 0991 is abolished in the aorta of mas-knockout mice. *J Cardiovasc Pharmacol* 2005; **46**: 274-279 [PMID: 16116331 DOI: 10.1097/01.fjc.0000175237.41573.63]
- 52 **Neves LA**, Williams AF, Averill DB, Ferrario CM, Walkup MP, Brosnihan KB. Pregnancy enhances the angiotensin (Ang)-(1-7) vasodilator response in mesenteric arteries and increases the renal concentration and urinary excretion of Ang-(1-7). *Endocrinology* 2003; **144**: 3338-3343 [PMID: 12865311 DOI: 10.1210/en.2003-0009]
- 53 **Santos RA**, Simoes e Silva AC, Maric C, Silva DM, Machado RP, de Buhr I, Heringer-Walther S, Pinheiro SV, Lopes MT, Bader M, Mendes EP, Lemos VS, Campagnole-Santos MJ, Schultheiss HP, Speth R, Walther T. Angiotensin-(1-7) is an endogenous ligand for the G protein-coupled receptor Mas. *Proc Natl Acad Sci USA* 2003; **100**: 8258-8263 [PMID: 12829792 DOI: 10.1073/pnas.1432869100]
- 54 **Silva DM**, Vianna HR, Cortes SF, Campagnole-Santos MJ, Santos RA, Lemos VS. Evidence for a new angiotensin-(1-7) receptor subtype in the aorta of Sprague-Dawley rats. *Peptides* 2007; **28**: 702-707 [PMID: 17129638 DOI: 10.1016/j.peptides.2006.10.007]
- 55 **Herath CB**, Mak K, Burrell LM, Angus PW. Angiotensin-(1-7) reduces the perfusion pressure response to angiotensin II and methoxamine via an endothelial nitric oxide-mediated pathway in cirrhotic rat liver. *Am J Physiol Gastrointest Liver Physiol* 2013; **304**: G99-G108 [PMID: 23086915 DOI: 10.1152/ajpgi.00163.2012]
- 56 **Brosnihan KB**, Li P, Ferrario CM. Angiotensin-(1-7) dilates canine coronary arteries through kinins and nitric oxide. *Hypertension* 1996; **27**: 523-528 [PMID: 8613197 DOI: 10.1161/01.HYP.27.3.523]
- 57 **Ferrario CM**, Chappell MC, Tallant EA, Brosnihan KB, Diz DI. Counterregulatory actions of angiotensin-(1-7). *Hypertension* 1997; **30**: 535-541 [PMID: 9322978 DOI: 10.1161/01.HYP.30.3.535]
- 58 **Pörsti I**, Bara AT, Busse R, Hecker M. Release of nitric oxide by angiotensin-(1-7) from porcine coronary endothelium: implications for a novel angiotensin receptor. *Br J Pharmacol* 1994; **111**: 652-654 [PMID: 8019744 DOI: 10.1111/j.1476-5381.1994.tb14787.x]
- 59 **Bataller R**, Gäbele E, Parsons CJ, Morris T, Yang L, Schoonhoven R, Brenner DA, Rippe RA. Systemic infusion of angiotensin II exacerbates liver fibrosis in bile duct-ligated rats. *Hepatology* 2005; **41**: 1046-1055 [PMID: 15841463 DOI: 10.1002/hep.20665]
- 60 **Paizis G**, Cooper ME, Schembri JM, Tikellis C, Burrell LM, Angus PW. Up-regulation of components of the renin-angiotensin system in the bile duct-ligated rat liver. *Gastroenterology* 2002; **123**: 1667-1676 [PMID: 12404241 DOI: 10.1053/gast.2002.36561]
- 61 **Paizis G**, Tikellis C, Cooper ME, Schembri JM, Lew RA, Smith AI, Shaw T, Warner FJ, Zuilli A, Burrell LM, Angus PW. Chronic liver injury in rats and humans upregulates the novel enzyme angiotensin converting enzyme 2. *Gut* 2005; **54**: 1790-1796 [PMID: 16166274 DOI: 10.1136/gut.2004.062398]
- 62 **Jonsson JR**, Clouston AD, Ando Y, Kelemen LI, Horn MJ, Adamson MD, Purdie DM, Powell EE. Angiotensin-converting enzyme inhibition attenuates the progression of rat hepatic fibrosis. *Gastroenterology* 2001; **121**: 148-155 [PMID: 11438504 DOI: 10.1053/gast.2001.25480]
- 63 **Yoshiji H**, Kuriyama S, Fukui H. Angiotensin-I-converting enzyme inhibitors may be an alternative anti-angiogenic strategy in the treatment of liver fibrosis and hepatocellular carcinoma. Possible role of vascular endothelial growth factor. *Tumour Biol* 2002; **23**: 348-356 [PMID: 12677092 DOI: 10.1159/000069792]
- 64 **Yoshiji H**, Kuriyama S, Noguchi R, Ikenaka Y, Yoshiji J, Yanase K, Namisaki T, Kitade M, Yamazaki M, Asada K, Akahane T, Tsujimoto T, Uemura M, Fukui H. Amelioration of liver fibrogenesis by dual inhibition of PDGF and TGF-beta with a combination of imatinib mesylate and ACE inhibitor in rats. *Int J Mol Med* 2006; **17**: 899-904 [PMID: 16596278]
- 65 **Lubel JS**, Herath CB, Tchongue J, Grace J, Jia Z, Spencer K, Casley D, Crowley P, Sievert W, Burrell LM, Angus PW. Angiotensin-(1-7), an alternative metabolite of the renin-angiotensin system, is up-regulated in human liver disease and has antifibrotic activity in the bile-duct-ligated rat. *Clin Sci (Lond)* 2009; **117**: 375-386 [PMID: 19371232 DOI: 10.1042/CS20080647]
- 66 **Herath CB**, Lubel JS, Jia Z, Velkoska E, Casley D, Brown L, Tikellis C, Burrell LM, Angus PW. Portal pressure responses and angiotensin peptide production in rat liver are determined by relative activity of ACE and ACE2. *Am J Physiol Gastrointest Liver Physiol* 2009; **297**: G98-G106 [PMID: 19389807 DOI: 10.1152/ajpgi.00045.2009]
- 67 **Macgillchrist AJ**, Howes LG, Hawksby C, Reid JL. Plasma noradrenaline in cirrhosis: a study of kinetics and temporal relationship to ascites formation. *Eur J Clin Invest* 1991; **21**: 238-243 [PMID: 1905638 DOI: 10.1111/j.1365-2362.1991.tb01816.x]
- 68 **Hennenberg M**, Trebicka J, Sauerbruch T, Heller J. Mechanisms of extrahepatic vasodilation in portal hypertension. *Gut* 2008; **57**: 1300-1314 [PMID: 18445644 DOI: 10.1136/gut.2007.144584]
- 69 **Pinzani M**, Milani S, De Franco R, Grappone C, Caligiuri A, Gentilini A, Tosti-Guerra C, Maggi M, Failli P, Ruocco C, Gentilini P. Endothelin 1 is overexpressed in human cirrhotic liver and exerts multiple effects on activated hepatic stellate cells. *Gastroenterology* 1996; **110**: 534-548 [PMID: 8566602 DOI: 10.1053/gast.1996.v110.pm8566602]
- 70 **Hernández-Guerra M**, García-Pagán JC, Bosch J. Increased hepatic resistance: a new target in the pharmacologic therapy of portal hypertension. *J Clin Gastroenterol* 2005; **39**:

- S131-S137 [PMID: 15758648]
- 71 **Rockey DC.** Vasoactive agents in intrahepatic portal hypertension and fibrogenesis: implications for therapy. *Gastroenterology* 2000; **118**: 1261-1265 [PMID: 10833501 DOI: 10.1016/S0016-5085(00)70379-9]
  - 72 **Shah V,** Cao S, Hendrickson H, Yao J, Katusic ZS. Regulation of hepatic eNOS by caveolin and calmodulin after bile duct ligation in rats. *Am J Physiol Gastrointest Liver Physiol* 2001; **280**: G1209-G1216 [PMID: 11352814]
  - 73 **Shah V,** Toruner M, Haddad F, Cadelina G, Papapetrooulos A, Choo K, Sessa WC, Groszmann RJ. Impaired endothelial nitric oxide synthase activity associated with enhanced caveolin binding in experimental cirrhosis in the rat. *Gastroenterology* 1999; **117**: 1222-1228 [PMID: 10535886 DOI: 10.1016/S0016-5085(99)70408-7]
  - 74 **Gorelik G,** Carhini LA, Scicli AG. Angiotensin 1-7 induces bradykinin-mediated relaxation in porcine coronary artery. *J Pharmacol Exp Ther* 1998; **286**: 403-410 [PMID: 9655885]
  - 75 **Gioli-Pereira L,** Nascimento EA, Santos EL, Bracht A, Juliano MA, Pesquero JB, Borges DR, Kouyoumdjian M. Fate of bradykinin on the rat liver when administered by the venous or arterial route. *J Gastroenterol Hepatol* 2005; **20**: 463-473 [PMID: 15740493 DOI: 10.1111/j.1440-1746.2005.03580.x]
  - 76 **Fleming I,** Fisslthaler B, Dimmeler S, Kemp BE, Busse R. Phosphorylation of Thr(495) regulates Ca(2+)/calmodulin-dependent endothelial nitric oxide synthase activity. *Circ Res* 2001; **88**: E68-E75 [PMID: 11397791 DOI: 10.1161/hh1101.092677]
  - 77 **Ferlitsch A,** Pleiner J, Mittermayer F, Schaller G, Homoncik M, Peck-Radosavljevic M, Wolzt M. Vasoconstrictor hyporeactivity can be reversed by antioxidants in patients with advanced alcoholic cirrhosis of the liver and ascites. *Crit Care Med* 2005; **33**: 2028-2033 [PMID: 16148476]
  - 78 **Vilas-Boas WW,** Ribeiro-Oliveira A, Pereira RM, Ribeiro Rda C, Almeida J, Nadu AP, Simões e Silva AC, dos Santos RA. Relationship between angiotensin-(1-7) and angiotensin II correlates with hemodynamic changes in human liver cirrhosis. *World J Gastroenterol* 2009; **15**: 2512-2519 [PMID: 19469002 DOI: 10.3748/wjg.15.2512]
  - 79 **Grace JA,** Jia Z, Herath CB, Burrell LM, Angus PW. Angiotensin-(1-7) is a mesenteric vasodilator in experimental cirrhosis. *Hepatology* 2010; **52**: 919A-1020A
  - 80 **Miñano C,** Garcia-Tsao G. Clinical pharmacology of portal hypertension. *Gastroenterol Clin North Am* 2010; **39**: 681-695 [PMID: 20951924 DOI: 10.1016/j.gtc.2010.08.015]
  - 81 **Schrier RW,** Arroyo V, Bernardi M, Epstein M, Henriksen JH, Rodés J. Peripheral arterial vasodilation hypothesis: a proposal for the initiation of renal sodium and water retention in cirrhosis. *Hepatology* 1988; **8**: 1151-1157 [PMID: 2971015 DOI: 10.1002/hep.1840080532]
  - 82 **Sieber CC,** Groszmann RJ. In vitro hyporeactivity to methoxamine in portal hypertensive rats: reversal by nitric oxide blockade. *Am J Physiol* 1992; **262**: G996-G1001 [PMID: 1616049]
  - 83 **Hartleb M,** Moreau R, Cailmail S, Gaudin C, Lebrec D. Vascular hyporesponsiveness to endothelin 1 in rats with cirrhosis. *Gastroenterology* 1994; **107**: 1085-1093 [PMID: 7523215]
  - 84 **Heller J,** Schepke M, Gehnen N, Molderings GJ, Muller A, Erhard J, Spengler U, Sauerbruch T. Altered adrenergic responsiveness of endothelium-denuded hepatic arteries and portal veins in patients with cirrhosis. *Gastroenterology* 1999; **116**: 387-393 [PMID: 9922320 DOI: 10.1016/S0016-5085(99)70136-8]
  - 85 **Helmy A,** Newby DE, Jalan R, Johnston NR, Hayes PC, Webb DJ. Nitric oxide mediates the reduced vasoconstrictor response to angiotensin II in patients with preascitic cirrhosis. *J Hepatol* 2003; **38**: 44-50 [PMID: 12480559 DOI: 10.1016/S0168-8278(02)00319-7]
  - 86 **Hennenberg M,** Biecker E, Trebicka J, Jochem K, Zhou Q, Schmidt M, Jakobs KH, Sauerbruch T, Heller J. Defective RhoA/Rho-kinase signaling contributes to vascular hypocontractility and vasodilation in cirrhotic rats. *Gastroenterology* 2006; **130**: 838-854 [PMID: 16530523 DOI: 10.1053/j.gastro.2005.11.029]
  - 87 **Hennenberg M,** Trebicka J, Biecker E, Schepke M, Sauerbruch T, Heller J. Vascular dysfunction in human and rat cirrhosis: role of receptor-desensitizing and calcium-sensitizing proteins. *Hepatology* 2007; **45**: 495-506 [PMID: 17256744 DOI: 10.1002/hep.21502]
  - 88 **Hennenberg M,** Trebicka J, Kohistani AZ, Heller J, Sauerbruch T. Vascular hyporesponsiveness to angiotensin II in rats with CCl(4)-induced liver cirrhosis. *Eur J Clin Invest* 2009; **39**: 906-913 [PMID: 19522833 DOI: 10.1111/j.1365-2362.2009.02181.x]
  - 89 **Kiel JW,** Pitts V, Benoit JN, Granger DN, Shepherd AP. Reduced vascular sensitivity to norepinephrine in portal-hypertensive rats. *Am J Physiol* 1985; **248**: G192-G195 [PMID: 3970200]
  - 90 **Newby DE,** Jalan R, Masumori S, Hayes PC, Boon NA, Webb DJ. Peripheral vascular tone in patients with cirrhosis: role of the renin-angiotensin and sympathetic nervous systems. *Cardiovasc Res* 1998; **38**: 221-228 [PMID: 9683925 DOI: 10.1016/S0008-6363(98)00008-X]
  - 91 **Neef M,** Biecker E, Heller J, Schepke M, Nischalke HD, Wolff M, Spengler U, Reichen J, Sauerbruch T. Portal hypertension is associated with increased mRNA levels of vasopressor G-protein-coupled receptors in human hepatic arteries. *Eur J Clin Invest* 2003; **33**: 249-255 [PMID: 12641544 DOI: 10.1046/j.1365-2362.2003.01131.x]
  - 92 **Vaughan RB,** Angus PW, Angus JA. Enhanced vascular responses to noradrenaline in isolated omental arteries from patients with advanced cirrhosis: evidence against intrinsic vascular hyporeactivity in cirrhosis. *Gut* 2004; **53**: 1544-1545 [PMID: 15361510]
  - 93 **Langer DA,** Shah VH. Nitric oxide and portal hypertension: interface of vasoreactivity and angiogenesis. *J Hepatol* 2006; **44**: 209-216 [PMID: 16297493 DOI: 10.1016/j.jhep.2005.10.004]
  - 94 **Shah V.** Cellular and molecular basis of portal hypertension. *Clin Liver Dis* 2001; **5**: 629-644 [PMID: 11565134 DOI: 10.1016/S1089-3261(05)70185-9]
  - 95 **Vaughan RB,** Angus JA, Angus PW. Vasoconstrictor responses are normal but prostanoid-mediated vasodilatation is enhanced in human cirrhotic mesenteric arteries. *J Gastroenterol Hepatol* 2005; **20**: 1158-1164 [PMID: 16048562 DOI: 10.1111/j.1440-1746.2005.03946.x]
  - 96 **Wiest R,** Groszmann RJ. The paradox of nitric oxide in cirrhosis and portal hypertension: too much, not enough. *Hepatology* 2002; **35**: 478-491 [PMID: 11826425 DOI: 10.1053/jhep.2002.31432]
  - 97 **Colle IO,** De Vriese AS, Van Vlierberghe HR, Lameire NH, De Vos MM. Vascular hyporesponsiveness in the mesenteric artery of anaesthetized rats with cirrhosis and portal hypertension: an in-vivo study. *Eur J Gastroenterol Hepatol* 2004; **16**: 139-145 [PMID: 15075986 DOI: 10.1097/00042737-200402000-00004]
  - 98 **Mathie RT,** Ralevic V, Burnstock G. Portal vascular responsiveness to sympathetic stimulation and nitric oxide in cirrhotic rats. *J Hepatol* 1996; **25**: 90-97 [PMID: 8836907 DOI: 10.1016/S0168-8278(96)80333-3]
  - 99 **Schepke M,** Heller J, Paschke S, Thomas J, Wolff M, Neef M, Malago M, Molderings GJ, Spengler U, Sauerbruch T. Contractile hyporesponsiveness of hepatic arteries in humans with cirrhosis: evidence for a receptor-specific mechanism. *Hepatology* 2001; **34**: 884-888 [PMID: 11679958 DOI: 10.1053/jhep.2001.28794]
  - 100 **Zhang HY,** Shirasawa Y, Chen X, Yu H, Benoit JN. Impaired agonist-dependent myosin phosphorylation and decreased RhoA in rat portal hypertensive mesenteric vasculature. *Am J Physiol Gastrointest Liver Physiol* 2005; **288**: G603-G608 [PMID: 15513955 DOI: 10.1152/ajpgi.00116.2004]
  - 101 **Alam I,** Bass NM, Bacchetti P, Gee L, Rockey DC. He-

- patic tissue endothelin-1 levels in chronic liver disease correlate with disease severity and ascites. *Am J Gastroenterol* 2000; **95**: 199-203 [PMID: 10638583 DOI: 10.1111/j.1572-0241.2000.01684.x]
- 102 **Gracia-Sancho J**, Laviña B, Rodríguez-Vilarrupla A, García-Calderó H, Bosch J, García-Pagán JC. Enhanced vasoconstrictor prostanoid production by sinusoidal endothelial cells increases portal perfusion pressure in cirrhotic rat livers. *J Hepatol* 2007; **47**: 220-227 [PMID: 17459512 DOI: 10.1016/j.jhep.2007.03.014]
- 103 **Guturu P**, Shah V. New insights into the pathobiology of portal hypertension. *Hepatol Res* 2009; **39**: 1016-1019 [PMID: 19796039 DOI: 10.1111/j.1872-034X.2009.00553.x]
- 104 **Ferreira AJ**, Raizada MK. Are we poised to target ACE2 for the next generation of antihypertensives? *J Mol Med (Berl)* 2008; **86**: 685-690 [PMID: 18449520 DOI: 10.1007/s00109-008-0339-x]
- 105 **Shenoy V**, Qi Y, Katovich MJ, Raizada MK. ACE2, a promising therapeutic target for pulmonary hypertension. *Curr Opin Pharmacol* 2011; **11**: 150-155 [PMID: 21215698 DOI: 10.1016/j.coph.2010.12.002]

**P- Reviewers** Gurvits GE, Zhao D, Saavedra J  
**S- Editor** Song XX **L- Editor** A **E- Editor** Lu YJ

