

Published in final edited form as:


*Angew Chem Int Ed Engl.* 2013 March 25; 52(13): 3672–3675. doi:10.1002/anie.201209457.

## Aerobic Oxidative Heck/Dehydrogenation Reactions of Cyclohexenones: Efficient Access to *meta*-Substituted Phenols\*\*

 Yusuke Izawa<sup>+</sup>, Changwu Zheng<sup>+</sup>, and Shannon S. Stahl<sup>\*</sup>

Department of Chemistry, University of Wisconsin-Madison, 1101 University Avenue, Madison, WI 53706 (USA)

### Abstract



A new dicationic Pd<sup>II</sup> catalyst, employing a 6,6'-dimethylbipyridine ligand, has been identified that is capable of promoting both aerobic oxidative Heck coupling and dehydrogenation reactions of cyclohexenones. These reactions may be combined in a one-pot sequence to enable the straightforward synthesis of *meta*-substituted phenols.

### Keywords

Palladium; Heck Coupling; Cyclohexenone; Phenol; Dehydrogenation; Aerobic Oxidation

Phenol derivatives are common and important structural motifs in bioactive natural products and pharmaceuticals,<sup>[1]</sup> and the selective synthesis of substituted phenols is facilitated by the strong *ortho/para*-directing effect of the hydroxyl group. The same directing effect, however, limits access to analogous *meta*-substituted derivatives. In recent years, considerable efforts have targeted C–H functionalization reactions that enable preparation of *meta*-substituted arenes via steric<sup>[2]</sup> or directing-group<sup>[3]</sup> control over the site selectivity. The overall efficiency of these methods is often limited by functional group interconversions or installation/removal of directing groups needed to access the final product.<sup>[4]</sup> Moreover, in molecules with more than one electronically or sterically active substituent, competition between the directing groups can lead to product mixtures. Following our recent development of Pd-catalyzed aerobic dehydrogenation reactions of ketones,<sup>[5,6,7]</sup> we envisioned that *meta*-substituted phenols could be accessed efficiently via an aerobic

\*\*We thank Dr. Doris Pun for assistance in product purification. Financial support was provided by the NIH (R01 GM100143), Mitsubishi Chemical Corporation, and Shanghai Institute of Organic Chemistry (postdoctoral fellowship for CWZ).

<sup>\*</sup>Fax: (+1) 608-262-6143, stahl@chem.wisc.edu.

<sup>+</sup>These authors contributed equally to this work.

Supporting information for this article is available on the WWW under <http://www.angewandte.org> or from the author.

oxidative Heck/dehydrogenation sequence with cyclohexenone (Scheme 1).<sup>[8]</sup> Cyclohexenone is a convenient and inexpensive phenol precursor, and the proposed strategy exploits the intrinsic regioselectivity of additions to electron-deficient alkenes to enable functionalization of the "meta" C–H bond. Here, we describe a new Pd catalyst and reaction conditions compatible with this sequence, and we showcase their utility in the synthesis of a pharmaceutically active phenol derivative.

The proposed sequence in Scheme 1 faces several challenges. The oxidative Heck reaction must be more facile than the dehydrogenation step in order to avoid direct conversion of the cyclohexenone starting material to unsubstituted phenol. Furthermore, while aerobic oxidative Heck reactions have extensive precedent with terminal alkenes,<sup>[9, 10]</sup> analogous reactions with cyclohexenone tend to be more difficult.<sup>[11,12]</sup> With this substrate, the Pd<sup>II</sup>-enolate must isomerize to place the Pd atom on the opposite side of the ring in order to undergo  $\beta$ -hydride elimination (Scheme 2).<sup>[13]</sup> Finally, the catalyst and conditions must be compatible with both reactions in the sequence. The only general method for dehydrogenation of cyclohexenones to phenols employs a strong-acid additive (*p*-TsOH; Scheme 3),<sup>[5a]</sup> which interferes with oxidative Heck reactions.<sup>[14]</sup>

Our initial studies targeted the identification of non-acidic reaction conditions for aerobic dehydrogenation of 3-methylcyclohexenone. Upon screening diverse PdX<sub>2</sub> sources, ligands, additives and solvents (see Supp. Info. for full screening data), we found that the dicationic Pd<sup>II</sup> complex [Pd(CH<sub>3</sub>CN)<sub>4</sub>](BF<sub>4</sub>)<sub>2</sub> was particularly effective as a catalyst (Table 1). Formation of Pd black and gradual loss of catalytic activity during the reaction prompted us to test ancillary ligands to stabilize the catalyst. Most of the ligands tested inhibited the reaction (cf. Table 1 and Supp. Info.); however, 4,5-diazafluorenone **L4**<sup>[15]</sup> and 6,6'-dimethyl-2,2'-bipyridine **L5** enabled good product yields to be obtained. While screening of numerous additives, including Brønsted bases, Cu<sup>II</sup> and Ag<sup>I</sup> salts, and quinones showed little beneficial effect, nearly quantitative yield of the phenol product (95%) was obtained when 9 mol % AMS (anthraquinone-2-sulfonic acid sodium salt) was included in the reaction with ligand **L5**.<sup>[16]</sup> The optimal result was obtained upon addition of water (20 vol %) to enhance the solubility of AMS.

The optimized conditions proved to be effective with a number other substituted cyclohexenones, including those with heteroatom substituents (Table 2). These neutral reaction conditions revealed some advantages over the previously reported conditions in Scheme 3. For example, 6-phenylcyclohexanone underwent dehydrogenation to *o*-phenyl phenol in only 33% yield under the previous conditions, but this product is obtained in excellent yields under the present conditions (entries 1 and 2). The successful reaction of 3-arylcyclohexenones, prepared via oxidative Heck reactions with cyclohexenone (entries 9–11), provided a useful starting point for the investigation of oxidative Heck and tandem oxidative Heck/dehydrogenation reactions.

Preliminary experiments showed that this catalyst was quite effective for the oxidative Heck coupling of 4-methoxyphenylboronic acid and cyclohexenone. Moreover, the reaction could take place at 50 °C, a temperature at which no conversion of cyclohexenone to phenol was observed. In DMSO, the oxidative Heck reaction proceeded in 65% yield. Upon heating of this reaction mixture to 80 °C, nearly complete in situ conversion to the 3-aryl phenol was observed (i.e., 64% yield of the phenol; Table 3, entry 1). Several other solvents, including DMF, *N*-methylpyrrolidone (NMP) and 1,4-dioxane, proved to be better for the oxidative Heck reaction (entries 2–7); however, they proved less effective for the tandem sequence (e.g., entry 2). Further studies revealed that an effective one-pot sequence could be achieved by performing the oxidative Heck reaction in NMP at 50 °C, followed by addition of DMSO

and heating to 80 °C for the dehydrogenation step. This protocol enabled a good yield of the phenol to be obtained (84%, entry 12).

$[\text{Pd}(\text{CH}_3\text{CN})_4](\text{BF}_4)_2/\mathbf{L5}$  proved to be very effective as a standalone catalyst for oxidative Heck reactions with cyclohexenone. Good yields of the 3-arylcyclohexenones were obtained with diverse arylboronic acids (Table 4). Reactions with the electron-rich arylboronic acids typically led to higher yields than electron-deficient substrates as the latter substrates were more susceptible to the formation of homocoupling products. Halogenated arylboronic acids (X = F, Cl, Br) were tolerated in the oxidative Heck reaction, with yields ranging from 68% to 86% (entries 4–6 and 18). The same arylboronic acids were then employed in the one-pot oxidative Heck/dehydrogenation to afford the 3-substituted phenol derivatives. In most cases, the phenol yields correlate closely with the yields of the 3-aryl cyclohexenones in the independent oxidative Heck reaction.

In order to demonstrate the potential utility of the aerobic oxidative Heck/dehydrogenation sequence and further test its functional group compatibility, we investigated the synthesis of URB597 from cyclohexenone and the commercially available benzamide-derived boronic acid **1** (Scheme 4). URB597 is a potent inhibitor of fatty acid amide hydrolase (FAAH) and an important focus of efforts to treat pain, anxiety and depression.<sup>[17,18]</sup> The phenol intermediate **3** was prepared via stepwise oxidative Heck coupling of **1** and cyclohexenone, followed by catalytic dehydrogenation of the isolated intermediate **2**, and in a direct, one-pot process. The  $[\text{Pd}(\text{CH}_3\text{CN})_4](\text{BF}_4)_2/\mathbf{L5}$  catalyst was employed for each of these steps, and both pathways led to the phenol product **3** in good yield (approx. 72%, in each case).

The results above highlight a new catalyst system that mediates both aerobic oxidative Heck reactions with cyclohexenone and aerobic dehydrogenation of cyclohexenones. The one-pot sequence developed for these reactions represents an efficient strategy for the preparation of *meta* substituted phenols, which should be advantageous or highly competitive with other approaches based on C–H functionalization of an aromatic ring.

## Supplementary Material


Refer to Web version on PubMed Central for supplementary material.

## References


1. For example, see, Tyman JHP. *Synthetic and Natural Phenols*. 1996 New York: Elsevier; Hoarau C, Pettus TRR. *Synlett*. 2003;127. [PubMed: 18563206] Dewick PM. *Medicinal Natural Products: A Biosynthetic*.
2. a) Cho J-Y, Iverson CN, Smith MR III. *J. Am. Chem. Soc.* 2000; 122:12868. b) Ishiyama T, Takagi J, Ishida K, Miyaura N, Anastasi NR, Hartwig JF. *J. Am. Chem. Soc.* 2002; 124:390. [PubMed: 11792205] c) Ishiyama T, Takagi J, Hartwig JF, Miyaura N. *Angew. Chem.* 2002; 114:3182. *Angew. Chem. Int. Ed.* 2002, 41, 3056; d) Cho J-Y, Tse MK, Holmes D, Maleczka RE, Smith MR III. *Science*. 2002; 295:305. [PubMed: 11719693] e) Maleczka RE Jr, Shi F, Holmes D, Smith MR III. *J. Am. Chem. Soc.* 2003; 125:7792. [PubMed: 12822984] f) Zhang YH, Shi BF, Yu JQ. *J. Am. Chem. Soc.* 2009; 131:5072. [PubMed: 19296661]
3. a) Phipps RJ, Gaunt MJ. *Science*. 2009; 323:1593. [PubMed: 19299616] b) Duong HA, Gilligan RE, Cooke ML, Phipps RJ, Gaunt MJ. *Angew. Chem.* 2011; 123:483. *Angew. Chem. Int. Ed.* 2011, 50, 463; c) Saidi O, Marafie J, Ledger AEW, Liu PM, Mahon MF, Kociok-Köhn G, Whittlesey MK, Frost CG. *J. Am. Chem. Soc.* 2011; 133:19298. [PubMed: 22047022] d) Leow D, Li G, Mei TS, Yu JQ. *Nature*. 2012; 486:518. [PubMed: 22739317]
4. For other recent advances that enable efficient preparation of substituted phenols, see: Anderson KW, Ikawa T, Tundel RE, Buchwald SL. *J. Am. Chem. Soc.* 2006; 128:10694. [PubMed: 16910660] Tlili A, Xia N, Monnier F, Taillefer M. *Angew. Chem.* 2009; 121:8881. *Angew. Chem.*

- Int. Ed.* **2009**, *48*, 8725; Zhang YH, Yu JQ. *J. Am. Chem. Soc.* 2009; 131:14654. [PubMed: 19788192] Maurer S, Jiang YW, Ma DW, Liu W, Zhang XJ. *Synlett.* 2010:976. Zou YQ, Chen JR, Liu XP, Lu LQ, Davis RL, Jørgensen KA, Xiao WJ. *Angew. Chem. Int. Ed.* **2012**, *51*, 784; Yu CW, Chen GS, Huang CW, Chern JW. *Org. Lett.* 2012; 14:3688.
5. a) Izawa Y, Pun D, Stahl SS. *Science.* 2011; 333:209. [PubMed: 21659567] b) Diao T, Stahl SS. *J. Am. Chem. Soc.* 2011; 133:14566. [PubMed: 21851123] c) Diao T, Wadzinski TJ, Stahl SS. *Chem. Sci.* 2012; 3:887. [PubMed: 22690316]
6. For precedents and leading references, see: Horning EC, Horning MG. *J. Am. Chem. Soc.* 1947; 69:1359. Hirao T, Mori M, Ohshiro Y. *J. Org. Chem.* 1990; 55:358. Shvo Y, Arisha AH. *J. Org. Chem.* 1998; 63:5640. Moriuchi T, Kikushima K, Kajikawa T, Hirao T. *Tetrahedron Lett.* 2009; 50:7385. For a review, see: Muzart J. *Eur. J. Org. Chem.* 2010:3779.
7. For related, recently reported dehydrogenation methods, see: Hajra A, Wei Y, Yoshikai N. *Org. Lett.* 2012; 14:5488. [PubMed: 23072451] Kandukuri SR, Oestreich M. *J. Org. Chem.* 2012; 77:8750. [PubMed: 22950832] Imahori T, Tokuda T, Taguchi T, Takahata H. *Org. Lett.* 2012; 14:1172. [PubMed: 22296212] Girard SA, Hu X, Knauber T, Zhou F, Simon M, Deng GJ, Li CJ. *Org. Lett.* 2012; 14:5606. [PubMed: 23067013] Simon M, Girard SA, Li CJ. *Angew. Chem.* 2012; 124:7655. *Angew. Chem. Int. Ed.* **2012**, *51*, 7537; Xie Y, Liu S, Liu Y, Wen Y, Deng GJ. *Org. Lett.* 2012; 14:1692. [PubMed: 22409600]
8. For examples of sequential reactions involving dehydrogenation steps, see: Ohmura T, Kijima A, Suginome M. *J. Am. Chem. Soc.* 2009; 131:6070. [PubMed: 19354214] Xia X-F, Shu X-Z, Ji K-G, Yang Y-F, Shaikat A, Liu X-Y, Liang Y-M. *J. Org. Chem.* 2010; 75:2893. [PubMed: 20345090] Yip KT, Nimje RY, Leskinen MV, Pihko PM. *Chem. Eur. J.* 2012; 18:12590. [PubMed: 22915444] Moon Y, Kwon D, Hong S. *Angew. Chem.* 2012; 124:11495. *Angew. Chem. Int. Ed.* **2012**, *51*, 11333; Leskinen MV, Yip KT, Valkonen A, Pihko PM. *J. Am. Chem. Soc.* 2012; 134:5750. [PubMed: 22424139]
9. For important examples, see: Andappan MMS, Nilsson P, von Schenck H, Larhed M. *J. Org. Chem.* 2004; 69:5212. [PubMed: 15287763] Andappan MMS, Nilsson P, Larhed M. *Chem. Commun.* 2004:218. Yoon CH, Yoo KS, Yi SW, Mishra RK, Jung KW. *Org. Lett.* 2004; 6:4037. [PubMed: 15496093] Enquist PA, Lindh J, Nilsson P, Larhed M. *Green Chem.* 2006; 8:338. Yoo KS, Yoon CH, Jung KW. *J. Am. Chem. Soc.* 2006; 128:16384. [PubMed: 17165795] Lindh J, Enquist P, Pilotti Å, Nilsson P, Larhed M. *J. Org. Chem.* 2007; 72:7957. [PubMed: 17887706] Yoo KS, Park CP, Yoon CH, Sakaguchi S, O'Neill J, Jung KW. *Org. Lett.* 2007; 9:3933. [PubMed: 17760452] Ruan J, Li X, Saidi O, Xiao J. *J. Am. Chem. Soc.* 2008; 130:2424. [PubMed: 18232688] Delcamp JH, Brucks AP, White MC. *J. Am. Chem. Soc.* 2008; 130:11270. [PubMed: 18671350] Werner EW, Sigman MS. *J. Am. Chem. Soc.* 2010; 132:13981. [PubMed: 20858011] Zheng CW, Wang D, Stahl SS. *J. Am. Chem. Soc.* 2012; 134:16496. [PubMed: 22998540]
10. For a review, see: Karimi B, Behzadnia H, Elhamifar D, Akhavan PF, Esfahani FK, Zamani A. *Synthesis.* 2010:1399.
11. For examples of oxidative Heck reactions with cyclohexenone, see ref. 9e and the following: Krishna TR, Jayaraman N. *Tetrahedron.* 2004; 60:10325. Fall Y, Doucet H, Santelli M. *Tetrahedron.* 2009; 65:489. Gottumukkala AL, Teichert JF, Heijnen D, Eisink N, Van Dijk S, Ferrer C, Minnaard AJ, Van Den Hoogenband A. *J. Org. Chem.* 2011; 76:3498. [PubMed: 21428446]
12. Other methods to synthesize 3-aryl cyclohexenones usually need pre-functionalized 3-substituted cyclohexenones for coupling partner. Please see selected examples: Zimmerman HE, Nesterov EE. *J. Am. Chem. Soc.* 2003; 125:5422. [PubMed: 12720456] Matsuo J, Aizawa Y. *Chem. Commun.* 2005:2399. Quasdorf KW, Riener M, Petrova KV, Garg NK. *J. Am. Chem. Soc.* 2009; 131:17748. [PubMed: 19928764]
13. Beletskaya IP, Cheprakov AV. *Chem. Rev.* 2000; 100:3009. [PubMed: 11749313]
14. For example, preliminary experiments revealed substantial biaryl product formation, arising from homocoupling of the boronic acid, or non-oxidative conjugate addition to the enone.
15. a) Campbell AN, White PB, Guzei IA, Stahl SS. *J. Am. Chem. Soc.* 2010; 132:15116. [PubMed: 20929224] b) Campbell AN, Meyer EB, Stahl SS. *Chem. Commun.* 2011:10257.
16. For previous use of AMS as a cocatalyst in a Pd-catalyzed oxidation reaction, see: Sheldon RA, Sobczak JM. *J. Mol. Catal.* 1991; 68:1.


17. Kathuria S, Gaetani S, Fegley D, Valiño F, Duranti A, Tontini A, Mor M, Tarzia G, La Rana G, Calignano A, Giustino A, Tattoli M, Palmery M, Cuomo V, Piomelli D. *Nat. Med.* 2002; 9:76. [PubMed: 12461523] b) Fegley D, Gaetani S, Duranti A, Tontini A, Mor M, Tarzia G, Piomelli D. *J. Pharmacol. Exp. Ther.* 2005; 313:352. [PubMed: 15579492] c) Piomelli D, Tarzia G, Duranti A, Tontini A, Mor M, Compton TR, Dasse O, Monaghan EP, Parrott JA, Putman D. *CNS Drug Reviews.* 2006; 12:21. [PubMed: 16834756]
18. For previous syntheses, see: Mor M, Rivara S, Lodola A, Plazzi PV, Tarzia G, Duranti A, Tontini A, Piersanti G, Kathuria S, Piomelli D. *J. Med. Chem.* 2004; 47:4998. [PubMed: 15456244]



**Scheme 1.**  
Strategy for the synthesis of *meta*-substituted phenols.




**Scheme 2.**  
Mechanistic steps highlighting the requirement for isomerization of the  $Pd^{II}$ -enolate intermediate in Heck reactions of cyclohexenone.



**Scheme 3.**  
Previously reported aerobic dehydrogenation conditions for the synthesis of phenols.



**Scheme 4.**

Application of one-pot oxidative Heck/dehydrogenation reactions in the synthesis of URB597.

Table 1



Dehydrogenation of 3-Cyclohexenones: Screening Results.<sup>[a]</sup><sup>[a]</sup>Reactions were performed on 1.0 mmol scale in DMSO (0.2 mL), 1 atm O<sub>2</sub>.<sup>[b]</sup>GC yield.<sup>[c]</sup>H<sub>2</sub>O (0.05 mL) and AMS (0.09 mmol) were added. Reaction time: 6.5 h. See Supporting Information for details.

Table 2

Dehydrogenation of Substituted Cyclohexenones,<sup>[a]</sup>

Entry	Cyclohexenone	Phenol	Time [h]	Yield [%] <sup>[b]</sup>
1			10	90 <sup>[c]</sup>
2			10	96 <sup>[c]</sup>
3			36	46
4			6.5	63
5			10	87
6			24	77
7			24	73
8			12	68
9			10	89
10			10	52 <sup>[c]</sup>
11			10	83

<sup>[a]</sup>Reactions were performed on 1.0 mmol scale in DMSO/H<sub>2</sub>O (v/v 0.2/0.05 mL).

<sup>[b]</sup>Isolated yield.

<sup>[c]</sup>Pd<sup>II</sup>/L5/AMS = 5%/5%/15%.

Table 3

Optimization of Conditions for the Oxidative Heck and one-pot Oxidative Heck/Dehydrogenation Reactions.<sup>[a]</sup>

Entry	Oxidative Heck			Oxidative Heck/ Dehydrogenation			
	Solvent	Conc. [M] <sup>[b]</sup>	t <sub>1</sub> [h]	Yield [%] <sup>[c]</sup>	Solvent	t <sub>2</sub> [h]	Yield [%] <sup>[d]</sup>
1	DMSO	1.3	8	65	-	24	64
2	DMF	1.3	3	95	-	24	25 <sup>[e]</sup>
3	NMP	1.3	3	90	-	-	-
4	Dioxane	1.3	3	93	-	-	-
5	CH <sub>3</sub> CN	1.3	3	73	-	-	-
6	DMF	2.5	3	96	-	-	-
7	DMF	5.0	3	80	-	-	-
8	DMF	2.5	3	-	DMSO	24	54
9	NMP	2.5	3	-	DMSO	24	72
10	NMP	3.1	3	-	DMSO	24	82
11 <sup>[f]</sup>	NMP	3.1	3	-	DMSO	32	82
12 <sup>[f,g]</sup>	NMP	3.1	4	95	DMSO	32	84

<sup>[a]</sup> Reaction conditions: Heck reactions [cyclohexenone (0.75 mmol), boronic acid (0.25 mmol), Pd(II) = Pd(CH<sub>3</sub>CN)<sub>4</sub>(BF<sub>4</sub>)<sub>2</sub>, 50 °C, 3–8 h]; Dehydrogenation: temp. increased to 80 °C, 24–32 h (for entries 8–15, 0.2 mL DMSO added).

<sup>[b]</sup> Conc. of boronic acid.

<sup>[c]</sup> GC yields.

<sup>[d]</sup> <sup>1</sup>H NMR yields.

<sup>[e]</sup> 65% of Heck product recovered.


[1] 10% AMS used.

[2] 0.5 mmol cyclohexenone.

Table 4

Oxidative Heck and One-Pot Oxidative Heck/Dehydrogenation Reactions to Prepare Substituted Cyclohexenones and Phenols.<sup>[47]</sup>

Entry	R	Oxidative Heck		Oxidative Heck/ Dehydrogenation	
		t <sub>1</sub> [h]	Yield <sup>[a]</sup> [%]	t <sub>2</sub> [h]	Yield <sup>[c]</sup> [%]
1	Me	4	97	32	91
2	<i>t</i> -Bu	4	98	36	85
3	MeO	4	94	32	80
4	Cl	4	85	32	68
5	Br	4	65	32	41
6	F	4	86	32	87
7	OH	4	85	32	64
8	Ph	13	62	36	63
9	COOMe	4	86	32	74
10	CF <sub>3</sub>	4	56	32	40
11	CN	13	41	36	35
12	H	4	76	32	75
13	MeO	13	83	36	47
14	Me	13	89	36	76
15	NO <sub>2</sub>	13	48	36	42
16	CF <sub>3</sub>	4	68	32	60



Entry	R	Oxidative Heck		Oxidative Heck/ Dehydrogenation	
		t <sub>1</sub> [h]	Yield <sup>[b]</sup> [%]	t <sub>2</sub> [h]	Yield <sup>[c]</sup> [%]
17	Me	13	73	36	58
18	F	13	75	36	66

<sup>[a]</sup> Reactions conditions: cyclohexenone (0.5 mmol), arylboronic acid (0.25 mmol) in NMP (0.2 mL for Heck and 0.08 mL for one-pot) at 50 °C for Heck reactions. For the one-pot reactions, DMSO was then added and the reactions were continued at 80 °C for 32 or 36 hours. See Supporting Information for details.

<sup>[b]</sup> Isolated yields of 3-substituted cyclohexenones.

<sup>[c]</sup> Isolated yields of phenols.