

# Residual beta cell function at diagnosis of type 1 diabetes in children and adolescents varies with gender and season

U. Samuelsson<sup>1\*</sup>

B. Lindblad<sup>2</sup>

A. Carlsson<sup>4</sup>

G. Forsander<sup>2</sup>

S. Ivarsson<sup>3</sup>

I. Kockum<sup>5</sup>

Å. Lernmark<sup>6</sup>

C. Marcus<sup>7</sup>

J. Ludvigsson<sup>1</sup>

Better Diabetes Diagnosis  
(BDD) study group

<sup>1</sup>Department of Clinical and  
Experimental Medicine,  
Division of Pediatrics and Diabetes,  
Research Center,  
Linköping University Hospital,  
Linköping, Sweden

<sup>2</sup>Department of Pediatrics,  
Queen Silvia Children's Hospital,  
Göteborg, Sweden

<sup>3</sup>Department of Pediatrics,  
University Hospital MAS,  
Malmö, Sweden

<sup>4</sup>Department of Pediatrics,  
Lund University Hospital,  
Lund, Sweden

<sup>5</sup>Department of Molecular Medicine,  
Karolinska Institute, Stockholm, Sweden

<sup>6</sup>Department of Clinical Sciences,  
Lund University, Malmö, Sweden

<sup>7</sup>Department of Pediatrics,  
Karolinska University Hospital,  
Huddinge, Sweden

\*Correspondence to: Ulf Samuelsson,  
Department of Clinical and  
Experimental Medicine, Division of  
Pediatrics and Diabetes Research  
Center, Linköping University Hospital,  
S-581 85 Linköping, Sweden.  
E-mail: ulf.samuelsson@lio.se

Received: 13 March 2012

Revised: 7 September 2012

Accepted: 11 October 2012

## Abstract

**Background** There are seasonal variations and gender differences in incidence of type 1 diabetes (T1D), metabolic control and responses to immune interventions at onset of the disease.

We hypothesized that there are seasonal and gender differences in residual insulin secretion already at diagnosis of T1D.

**Methods** In 2005, a national study, the Better Diabetes Diagnosis, was started to classify all newly diagnosed children and adolescents with diabetes. About 95% (3824/4017) of the patients were classified as T1D, and our analyses are based on the patients with T1D.

**Results** C-peptide was lower in younger children, 0–10 years of age ( $0.23 \pm 0.20$  nmol/L) than in older children, 11–18 years of age ( $0.34 \pm 0.28$  nmol/L) ( $p < 0.000$ ). There was a seasonal variation in non-fasting serum C-peptide, significantly correlated to the seasonal variation of diagnosis ( $p < 0.01$ ). Most children were diagnosed in January, February and March as well as in October when C-peptide was highest, whereas fewer patients were diagnosed in April and May when serum C-peptide was significantly lower ( $p < 0.01$ ). The seasonal variation of C-peptide was more pronounced in boys than in girls ( $p < 0.000$  and  $p < 0.01$ , respectively). Girls had higher C-peptide than boys ( $p < 0.05$ ), especially in early puberty.

**Conclusions** Both seasonal and gender differences in residual beta cell function exist already at diagnosis of T1D. These observations have consequences for treatment and for randomizing patients in immune intervention clinical trials. Copyright © 2012 John Wiley & Sons, Ltd.

**Keywords** C-peptide; children; type 1 diabetes; seasonal variation; gender; immune intervention

## Introduction

Immune intervention with glutamic acid decarboxylase (GAD) treatment of children and adolescents with recent onset of type 1 diabetes (T1D) seemed to delay the loss of endogenous C-peptide, at least in a phase II trial, and in some prespecified subgroups of a phase III trial [1,2]. In both phases II and III studies, there was a significant efficacy in patients treated during early spring [2], which raises the question whether there is a seasonal variation disease process and/or beta cell function. Seasonal variation in the incidence of T1D is well known [3], although there is no clear explanation of this

phenomenon. It has been speculated that the seasonal variation is related to virus infections [4]. Other possible explanations include hormonal variation, variation in physical activity with more physical activity and better insulin sensitivity during summer and therefore less insulin requirement, variation in sun exposure and vitamin D or a simple explanation could just be less active observation of symptoms such as thirst and polyuria during hot summer months.

In the European Phase III GAD trial, the efficacy was also different in men and women with a significant efficacy in men but not in women [2].

Regarding gender differences, we and others have earlier noticed that there were significant differences in the degree of metabolic control between girls and boys with T1D [5]. Thus, it is common that girls have higher HbA<sub>1c</sub> in the ages of 10–18 years, which often has been taken as a sign of less good care of teenage girls. There are differences in age of puberty between girls and boys, differences in physical activity with more physically active boys [6], girls have more subcutaneous fat and have also an increased tendency of overweight during adolescence [7] parallel to an increased risk of both bulimia and anorexia [8]. All factors mentioned previously may contribute to the less good metabolic control of girls with T1D, but are there differences also in disease process? Thus, we know that T1D is twice as common in boys as in girls after the age of 15 years [9], in contrast to other autoimmune diseases that tend to be more common in women than in men.

As both seasonal variation and gender seem to be involved in the course of T1D, we hypothesized that the disease process may be influenced leading to difference in residual insulin secretion already at diagnosis of T1D. Therefore, we decided to study these questions using data from almost 4000 newly diagnosed patients with T1D in a nationwide study in Sweden.

## Materials and methods

In 2005, a prospective national study, the Better Diabetes Diagnosis, was started in Sweden to classify all newly diagnosed children and adolescents with diabetes. Children below the age of 18 years with new onset diabetes are referred to a paediatric clinic. This cross-sectional prospective study is based on patients from all 43 Swedish paediatric clinics.

Diagnosis of diabetes is based on the American Diabetes Association criteria for diagnosis and classification of diabetes (i.e. casual plasma glucose > 11.1 mmol/L or a fasting plasma glucose > 7.0 mmol/L and symptoms of polyuria, polydipsia and weight loss) [10]. In total, 4017 patients were included in this study. Questions on family history regarding diabetes and autoimmune disorders among first-degree relatives, symptoms and signs as well as height and weight were registered in Swedish National Paediatric Diabetes Registry, a national incidence and quality control registry [11]. The diagnosis and classification of diabetes were initially based on clinical symptoms

and signs, later on strengthened by information on diabetes-related autoantibodies, human leukocyte antigen-types, C-peptide, and in some cases maturity onset diabetes of the young genetics [12]. About 95% (3824/4017) of the patients were classified as T1D (Table 1), and our analysis of difference in C-peptide between women and men is based on the 3824 patients with T1D.

The Karolinska Institute Research Ethics Board approved the study, and informed consent from the parents was obtained.

## Determination of C-peptide

Serum C-peptide from the random non-fasting blood sample, taken at diagnosis before the first insulin injection, was measured at Linköping University, Sweden, with a time-resolved fluoroimmunoassay (AutoDELFIA™ C-peptide kit, Wallac, Turku, Finland), with a detection level of 0.03 nmol/L. The sample was taken before the first insulin injection, mostly day 1. Each assay was validated by inclusion of a C-peptide control module containing a high, a medium and a low-level control (Immulate, DPC, UK). A 1224 MultiCalc® programme (Wallac) was used to calculate the levels of C-peptide.

## Statistical analysis

SPSS 18® (SPSS Inc., Chicago, IL, USA) was used for the analyses. Unpaired two-tailed Student's *t*-test and one-way analysis of variance was used. When there were indications of skewed distribution, Mann–Whitney *U*-test or Kruskal–Wallis test was used. Comparisons of groups were performed by crosstabs and chi-square ( $\chi^2$ ), or Fisher's exact test was used for proportions. To study whether there was a seasonal variation of diagnosis and/or a seasonal variation of C-peptide at diagnosis, an ordinary chi-square test [(observed (O) – expected (E))<sup>2</sup>/E] was used with 11 degrees of freedom. The mean C-peptide value each month was the observed value, and the mean value for all months was the expected value. Regarding the variation of diagnosis, the actual number of children was the observed value and the total number of children/12 was the expected value.  $p < 0.05$  was regarded as statistically significant. The results are expressed as mean  $\pm$  standard deviation.

## Results

C-peptide was lower in younger children, 0–10 years of age ( $0.23 \pm 0.20$  nmol/L) than in older children, 11–18 years of age ( $0.34 \pm 0.28$  nmol/L) ( $p < 0.000$ ) (Figure 1). There was a clear seasonal variation over the year in non-fasting serum C-peptide at diagnosis ( $\chi^2 = 25.4$ ,  $p < 0.01$ ) (Figure 2). This seasonal variation of C-peptide was significantly correlated to the seasonal variation of diagnosis (Figure 3) ( $p < 0.05$ , Spearman's two-tailed). Most children were diagnosed in January, February and March as well as in

October when C-peptide was highest, whereas few patients were diagnosed in April and May when serum C-peptide also was significantly lower ( $p < 0.01$ ). The seasonal




Figure 1. C-peptide in relation to age diagnosis of type 1 diabetes in girls and boys




Figure 2. Seasonal variation of non-fasting serum C-peptide in children and adolescents with type 1 diabetes




Figure 3. Seasonal variation of diagnosis of type 1 diabetes in boys and girls

variation was somewhat more pronounced for boys than for girls ( $X^2 = 49.9$ ,  $p < 0.000$  and  $X^2 = 27.3$ ,  $p < 0.01$ , respectively, in total  $X^2 = 71.2$ ,  $p < 0.000$ ) (Figure 3).

Girls had slightly higher C-peptide than boys ( $p < 0.05$ ) (Table 2). The lower C-peptide in boys was associated with differences between the two sexes in symptoms and signs at diagnosis (Table 2). These gender differences were similar in all age groups (Table 3), but the difference in C-peptide was especially pronounced from 9 years of age (Figure 1). The gender differences did not explain the seasonal variation in C-peptide as both genders had about the same seasonal variation of C-peptide at diagnosis (Figure 2).

C-peptide at diagnosis was related to symptoms and signs at diagnosis (Table 4). Thus, children without polyuria and polydipsia at diagnosis and children without weight loss at diagnosis had higher C-peptide values than children with such symptoms and signs.

Children with T1D and who had T1D in the family, in grandparents or in first-degree relatives in general had a higher mean C-peptide at onset than T1D children without T1D in the family or among relatives (Table 4). T2D in the family, in grandparents or in first-degree relatives had no such relation to C-peptide at diagnosis in the T1D patients. Furthermore, children with low pH ( $< 7.3$ ) and low body mass index standard deviation score at diagnosis had low mean C-peptide value. Co-morbidity with another autoimmune disease did not influence level of C-peptide (Table 4).

## Discussion

As shown already long time ago [13], younger patients had significantly lower C-peptide than patients diagnosed as teenagers. It is also reasonable that patients with T1D in the family are diagnosed a bit earlier with higher C-peptide. Type 2 diabetes in the family was not associated with higher C-peptide at diagnosis of T1D, nor an autoimmune disease beside T1D. As expected, there are also correlations between C-peptide, symptoms and signs at diagnosis.

The focus of this study was to elucidate if seasonal and gender differences of beta cell function might be part of the explanation why immune intervention studies have shown different results related to these parameters. Our results from this nationwide large unselected patient's population support the hypothesis that there, at least in

Table 1. Type of diabetes in a Swedish nationwide population of children and adolescents

Type of diabetes	Number	Percent
Type 1	3824	95
Type 2	86	2
Maturity onset diabetes of the young	40	1
Secondary diabetes	30	1
Unknown type	13	0.4
Antibody negative	17	0.5
Another type	7	0.1
	4017	100

Table 2. C-peptide and some clinical differences between boys and girls at diagnosis

	Boys			Girls		
	<i>n</i>	Mean	SD	<i>n</i>	Mean	SD
HbA <sub>1c</sub> (mmol/mol)	1913	92.2*	24.4	1528	96.2*	26.7
C-peptide (nmol/L, onset)	1838	0.28**	0.25	1426	0.30**	0.25
BMI-SDS	1838	-0.39	1.56	1448	-0.48	1.39
Age (year)	2020	10.2*	4.5	1588	9.4*	4.0
pH	1923	7.34	0.09	1511	7.34	0.10
BE (mmol/L)	1883	-4.1*	6.8	1492	-5.1*	7.5
<i>p</i> -glucose (mmol/L)	1950	27.3	9.1	1547	26.3	8.7
	<i>n</i>	Yes (%)			Yes (%)	
Weight loss	2131	1453 (68.2)		1693	1189 (70.2)	
Polydipsia	2131	1870 (87.8)		1693	1482 (87.5)	
Polyuria	2131	1897 (89)		1693	1482 (87.5)	
Other autoimmune disease	2131	80 (3.8)*		1693	110 (6.5)*	

BMI-SDS, body mass index standard deviation score; BE, base excess.

\*\**p* < 0.05.

\**p* < 0.01.

Table 3. C-peptide and some clinical differences between boys and girls at diagnosis in different age groups

	Sex	0–5 years		6–10 years		11–15 years		16–18 years	
		<i>n</i>	Mean ± SD	<i>n</i>	Mean ± SD	<i>n</i>	Mean ± SD	<i>n</i>	Mean ± SD
HbA <sub>1c</sub> (mmol/mol)	Boy	385	77.9 ± 18.5	551	87.9* ± 21.2	758	99.7* ± 24.8	221	101.8 ± 25.4
	Girl	318	80.1 ± 19.1	554	95.4* ± 24.4	523	104.8* ± 27.3	135	103.6 ± 30.6
C-peptide (nmol/L, onset)	Boy	364	0.23 ± 0.24	538	0.23* ± 0.16	721	0.31** ± 0.26	216	0.36 ± 0.32
	Girl	292	0.21 ± 0.15	513	0.26* ± 0.23	495	0.35** ± 0.29	128	0.42 ± 0.29
BMI-SDS	Boy	359	-0.66 ± 1.37	528	-0.20 ± 1.6	728	-0.32** ± 1.5	224	-0.72 ± 1.7
	Girl	295	-0.56 ± 1.24	517	-0.36 ± 1.4	510	-0.52** ± 1.4	129	-0.54 ± 1.5
pH	Boy	383	7.36 ± 0.09	559	7.36 ± 0.08	746	7.33 ± 0.09	228	7.33 ± 0.09
	Girl	318	7.35 ± 0.09	538	7.35 ± 0.09	522	7.32 ± 0.11	130	7.34 ± 0.10
BE	Boy	377	-4.3** ± 6.6	548	-3.3* ± 6.0	739	-4.5* ± 7.4	221	-4.1 ± 7.3
	Girl	317	-5.3** ± 6.8	534	-4.5* ± 7.2	515	-5.8* ± 8.2	129	-4.1 ± 7.8
<i>p</i> -glucose (mmol/L)	Boy	389	26.6 ± 7.8	560	26.9 ± 8.4	763	27.7 ± 9.6	231	28.5* ± 10.5
	Girl	324	27.0 ± 8.3	555	26.1 ± 8.0	529	26.6 ± 9.6	134	24.9* ± 8.9
		<i>n</i>	Yes (%)	<i>n</i>	Yes (%)	<i>n</i>	Yes (%)	<i>n</i>	Yes (%)
Weight loss	Boy	411	52.3**	580	69.1**	790	80)	239	83.7
	Girl	338	61.2**	569	75.4**	541	81.5	140	77.9
Polydipsia	Boy	411	90.3	580	92.4	790	93.2	239	92.5
	Girl	338	92.9	569	93.3	541	93.3	140	92.1
Polyuria	Boy	411	92.2	580	94.7	790	93.7	239	93.3
	Girl	338	93.2	569	93.3	541	93.2	140	92.1
Other autoimmune disease	Boy	411	3.6	580	5.6**	790	5.6 *	239	2.5**
	Girl	338	5.9	569	9.1**	541	9.1 *	140	7.9**

BMI-SDS, body mass index standard deviation score; BE, base excess.

\*\**p* < 0.05.

\**p* < 0.01.

Sweden, are seasonal variations in the disease process with differences in residual beta cell function already at diagnosis. Thus, we found not only a seasonal variation of diagnosis, which is well known [14], but also a clear seasonal variation of C-peptide at diagnosis with lower C-peptide in patients diagnosed in those months (April and May) when the incidence was lowest. We measured non-fasting C-peptide influenced by actual food intake, but there is no evidence that seasonal variation of food intake could explain our findings. Our results support that season may play a role for the precipitation of manifest T1D and/or course of T1D and may therefore also play a role for the effect of immune interventions. Why children diagnosed during April and May are fewer, but have significantly lower C-peptide, is unclear, but could, for instance, be related to infections [15,16]. There is also a

known seasonal variation in the immune system among children [17], which might to some extent be related to seasonal variation of vitamin D, associated with exposure to sun and light. The seasonal variation both in immune system and in residual beta cell function may be part of the explanation why the effect of GAD treatment in newly diagnosed T1D has shown best effect in patients diagnosed during early spring [1,2].

In light of the known gender differences in both incidence of T1D and metabolic control, the fact that immune intervention has shown different effects in women and men (2) is perhaps reasonable. It has earlier been shown that GAD antibodies response and its related C-peptide decline is more pronounced in women than in men [18]. Our results show that there is a difference between girls and boys already at diagnosis of diabetes. Difference in

**Table 4. Some symptoms and signs at diagnosis, as well as family history of diabetes in relation to C-peptide at diagnosis**

Symptoms/history	Number	C-peptide, diagnosis			
		Mean	SD	p-value	
Polyuria	Yes	3067	0.27	0.21	<0.000
	No	139	0.71	0.51	
Polydipsia	Yes	3043	0.26	0.21	<0.000
	No	154	0.70	0.51	
Weight loss	Yes	2397	0.25	0.19	<0.000
	No	701	0.42	0.37	
pH <7.3	Yes	513	0.18	0.15	<0.000
	No	2606	0.30	0.24	
BMI-SDS ≤0	Yes	1872	0.24	0.19	<0.000
	No	1125	0.37	0.32	
Other auto immune disease	Yes	173	0.32	0.31	n.s
	No	3268	0.29	0.26	
T1D in family	Yes	448	0.37	0.28	<0.000
	No	2993	0.27	0.26	
T1D in grandparents	Yes	220	0.34	0.25	0.01
	No	3221	0.29	0.27	
T2D in relatives	Yes	1121	0.31	0.27	0.01
	No	2320	0.28	0.27	

BMI-SDS, body mass index standard deviation score; T1D, type 1 diabetes; T2D, type 2 diabetes.

immune function between girls and boys [19–21] may partly explain why girls have significantly higher C-peptide at diagnosis. The difference exists in all ages but becomes especially pronounced in the ages 9–11 years when girls

go into puberty, somewhat earlier than boys, and therefore with increasing insulin resistance may get their manifest diabetes at a higher C-peptide production.

In conclusion, seasonal variation and gender differences in both immune function and T1D are well known, and evidently differences in beta cell function exist already at diagnosis of T1D. Even though it may be too early to tailor-cut individual treatments for patients with T1D, we have to be aware of whom we treat and in what situation when trying to improve the effect of immune interventions in T1D.

## Acknowledgements

We are grateful to all children, parents and the staff at the 43 paediatric clinics in Sweden and want to thank Lena Berglert, Division of Paediatrics, Linköping, who determined C-peptide. The study was supported by the Swedish Child Diabetes Foundation (Barndiabetesfonden).

## Conflict of interest

No authors have anything to disclose related to the content of this article.

## References

- Ludvigsson J, Faresjö M, Hjorth M, *et al.* GAD treatment and insulin secretion in recent-onset type 1 diabetes. *N Engl J Med* 2008; **359**: 1909–1920.
- Ludvigsson J, Krisky D, Casas R, *et al.* GAD65 antigen therapy in recently diagnosed type 1 diabetes mellitus. *N Engl J Med* 2012; **366**(5): 433–442.
- Moltchanova EV, Schreier N, Lammi N, Karvonen M. Seasonal variation of diagnosis of type 1 diabetes mellitus in children worldwide. *Diabet Med* 2009; **26**(7): 673–8.4.
- Gamble DR. The epidemiology of insulin dependent diabetes with particular reference to the relationship of virus infection to its etiology. *Epidemiol Rev* 1980; **2**: 49–70.
- Hanberger L, Samuelsson U, Lindblad B, Ludvigsson J. Swedish Childhood Diabetes Registry SWEDIABKIDS A1C in children and adolescents with diabetes in relation to certain clinical parameters: the Swedish Childhood Diabetes Registry SWEDIABKIDS. *Diabetes Care* 2008; **31**(5): 927–929.
- Bradley CB, McMurray RG, Harrell JS, Deng S. Changes in common activities of 3rd through 10th graders: the CHIC study. *Med Sci Sports Exerc* 2000; **32** (12): 2071–2078.
- Roemmich JN, Clark PA, Walter K, Patrie J, Weltman A, Rogol AD. Pubertal alterations in growth and body composition. V. Energy expenditure, adiposity, and fat distribution. *Am J Physiol Endocrinol Metab* 2000; **279**(6): E1426–E1436.
- Quiles-Marcos Y, Balaguer-Solá I, Pamies-Aubalat L, Quiles-Sebastián MJ, Marzo-Campos JC, Rodríguez-Marín J. Eating habits, physical activity, consumption of substances and eating disorders in adolescents. *Span J Psychol* 2011; **14**(2): 712–723.
- Pundziute-Lycká A, Dahlquist G, Nyström L, *et al.* The incidence of type I diabetes has not increased but shifted to a younger age at diagnosis in the 0–34 years group in Sweden 1983–1998. *Diabetologia* 2002; **45**: 783–791.
- The expert Committee on the diagnosis and classification of diabetes mellitus. Report of the expert committee on the diagnosis and classification of diabetes mellitus. *Diabetes Care* 2003; **26**(suppl 1): S5–S20.
- The Swedish pediatric diabetes quality register, (SWEDIABKIDS) (2010). <https://www.ndr.nu/ndr2/>
- Ludvigsson J, Carlsson A, Forsander G, *et al.* C-peptide in the classification of diabetes in children and adolescents. *Pediatr Diabetes* 2011 Sep 13. DOI: 10.1111/j.1399-5448.2011.00807.
- Ludvigsson J, Heding LG. C-peptide in children with juvenile diabetes. A preliminary report. *Diabetologia* 1976; **12**: 627–630.
- Christau B, Åkerblom H, Joner G, Dahlquist G, Ludvigsson J, Nerup J. Incidence of childhood diabetes mellitus in Denmark, Finland, Norway and Sweden. *Acta Endocrinol Suppl* 1981; **245**(Suppl): 68–77.
- Ludvigsson J, Afoke A. Seasonality of type I (insulin-dependent) diabetes mellitus: values of C-peptide, insulin antibodies and haemoglobin A1c show evidence of a more rapid loss of insulin secretion in epidemic patients. *Diabetologia* 1989; **32**: 8491.
- Afoke A, Ludvigsson J, Hed J, Lindblom B. Raised IgG and IgM in ‘epidemic’ IDDM suggest that infections are responsible for the seasonality of type I diabetes. *Diab Res* 1991; **16**: 11–17.
- Afoke A, Eeg-Olofsson O, Hed J, Kjellman N-IM, Lindblom B, Ludvigsson J. Seasonal variations and sex differences of circulating macrophages, immunoglobulins and lymphocytes in healthy school children. *Scand J Immunol* 1993; **37**: 209–215.
- Lindholm E, Hallengren B, Agardh CD. Gender differences in GAD antibody-positive diabetes mellitus in relation to age at onset, C-peptide and other endocrine autoimmune diseases. *Diabetes Metab Res Rev* 2004; **20**: 158–164.
- Okuyama K, Hamanaka Y, Kawano T, *et al.* T cell subsets related with a sex difference in IL-5 production. *Int Arch Allergy Immunol* 2011; **155**(Suppl 1): 21–26
- Dinesh RK, Hahn BH, Singh RP. PD-1, gender, and autoimmunity. *Autoimmun Rev* 2010; **9**(8): 583–587. Epub 2010 Apr 28
- Cohen-Solal JF, Jeganathan V, Hill L, *et al.* Hormonal regulation of B-cell function and systemic lupus erythematosus. *Lupus* 2008; **17**(6): 528–532.