

Published in final edited form as:

*Cell Mol Neurobiol.* 2013 May ; 33(4): 483–488. doi:10.1007/s10571-013-9929-4.

## Expression of NgR1-antagonizing proteins decreases with aging and cognitive decline in rat hippocampus

Heather D. VanGuilder Starkey<sup>a</sup>, Georgina V. Bixler<sup>a</sup>, William E. Sonntag<sup>b</sup>, and Willard M. Freeman<sup>a,\*</sup>

<sup>a</sup>Department of Pharmacology, Penn State University College of Medicine, 500 University Drive, R130, Hershey, PA, 17033, USA

<sup>b</sup>Donald W. Reynolds Department of Geriatric Medicine, University of Oklahoma Health Sciences Center, 975 NE 10<sup>th</sup> Street, BRC-1303, Oklahoma City, OK, 73103, USA

### Abstract

The myelin-associated inhibitor/Nogo-66 receptor 1 (NgR1) pathway directly functions in negative modulation of structural and electrophysiological synaptic plasticity. Previous work has established an important role of NgR1 pathway signaling in cognitive function, and we have demonstrated that multiple components of this pathway, including ligands, NgR1 co-receptors, and RhoA, are upregulated at the protein level specifically in cognitively impaired, but not age-matched cognitively intact aged rats. Recent studies have identified two novel endogenous NgR1 antagonists, LOTUS and LGI1, and an alternative co-receptor, ADAM22, which act to suppress NgR1 pathway signaling. To determine whether these endogenous NgR1-inhibiting proteins may play a compensatory role in age-related cognitive impairment by counteracting overexpression of NgR1 agonists and co-receptors, we quantified the expression of LOTUS, LGI1 and ADAM22 in hippocampal CA1, CA3 and DG subregions dissected from mature adult and aged rats cognitively phenotyped for spatial learning and memory by Morris water maze testing. We have found that endogenous inhibitors of NgR1 pathway action decrease significantly with aging and cognitive decline, and that lower expression levels correlate with declining cognitive ability, particularly in CA1 and CA3. These data suggest that decreased expression of NgR1-antagonizing proteins may exert a combinatorial effect with increased NgR1 signaling pathway components to result in abnormally strong suppression of synaptic plasticity in age-related cognitive impairment.

### Keywords

age-related cognitive decline; RhoA; ADAM22; LGI1; LOTUS/CRTAC1; plasticity; Nogo-66 receptor 1

### Introduction

NgR1 pathway signaling through RhoA, initiated by binding of myelin-associated inhibitors (MAIs) to NgR1 and mediated by multiple NgR1 co-receptors, suppresses neurite outgrowth and axon regeneration during development and following CNS damage. MAI/NgR1 pathway action also modulates synaptic plasticity in the mature, undamaged CNS by promoting structural rigidity and suppressing functional strengthening of synapses. Anatomical, biochemical and electrophysiological assessments demonstrate an inverse

\*Corresponding Author: Willard M. Freeman, wfreeman@hmc.psu.edu, Phone: 717-531-4037, Fax: 717-531-5013.

**Conflict of Interest:** The authors have no conflicts of interest to declare.

relationship between MAI/NgR1 pathway expression and hippocampal spine density, efficacy of activity-dependent synaptic plasticity, and spatial learning and memory (Zagrebelsky et al., 2005, Lee et al., 2008, Karlen et al., 2009, Raiker et al., 2010, Delekate et al., 2011). We have previously demonstrated the significant hippocampal upregulation of the MAI ligands Nogo-A, MAG, and OMgp, the NgR1 receptor and its signal-transducing co-receptors in a naturally occurring rat model of human age-related cognitive decline (VanGuilder et al., 2011b, 2012; VanGuilder et al., 2013) (Supplemental Table 1). Interestingly, induction of MAI/NgR1 pathway components occurs specifically in cognitively impaired, but not cognitively intact, aged rats phenotyped for hippocampal cognitive function, and is highly conserved within individual subjects, suggesting an important role of MAI/NgR1-mediated suppression of synaptic plasticity in impaired spatial learning and memory.

The MAI/NgR1 pathway has been well-characterized, but recently, endogenous NgR1 antagonists that compete with MAIs for NgR1 binding have been discovered, suggesting an additional level of complexity to NgR1 pathway regulation. LOTUS (lateral olfactory tract usher substance) is a transmembrane domain-containing secreted protein that antagonizes NgR1 to prevent Nogo-66-mediated growth cone collapse (Sato et al., 2011, Kurihara et al., 2012). LGI1 (leucine rich glioma inactivated 1) is a leucine rich repeat domain-containing secreted protein that competes with Nogo-66 for NgR1 binding, which effectively antagonizes the plasticity-suppressing action of the MAI/NgR1 pathway. Through interaction with ADAM22, a disintegrin and matrix metalloprotease and putative NgR1 co-receptor, LGI1 functions to enhance neuronal outgrowth. The known roles of LOTUS and LGI1 as endogenous NgR1-antagonizing ligands, and ADAM22 as an NgR1-interacting surface receptor, suggest a potential mechanism that may compensate for abnormal induction of MAI/NgR1 signaling in age related cognitive decline (Figure 1). The goal of the present study was to determine whether hippocampal expression of LOTUS, LGI1 and ADAM22 is regulated with cognitive impairment and to determine their potential relationship to spatial learning and memory ability.

## Materials and methods

### Animals: behavior and sample preparation

Behavioral stratification of subjects and dissection of CA1, CA3 and DG subregions has been described in detail elsewhere (VanGuilder et al., 2011a, 2012; VanGuilder Starkey et al., 2012, 2013.) All animal experiments were performed in accordance with IACUC and AALAC approved procedures. Briefly, mature adult (12 months) and aged (26 months) male Fischer 344 × Brown Norway (F1) hybrid rats were purchased from the National Institute on Aging rodent colony maintained by Harlan Industries (Indianapolis, IN) and housed singly in the OUHSC Reynolds Oklahoma Center on Aging barrier facility, with water and food (Purina Mills, Richmond, IN) freely available. Rats were tested for hippocampal spatial learning and memory ability by Morris water maze and aged rats were classified as cognitively intact or impaired relative to the mature adult group based on mean proximity to the escape platform location during probe trials (VanGuilder et al., 2011a, 2012; VanGuilder Starkey et al., 2012, 2013).

Rats were sacrificed by decapitation without anesthesia one week after conclusion of behavioral testing. CA1, CA3 and DG subregions were isolated individually from left and right hippocampi as previously described (Newton et al., 2005, VanGuilder et al., 2011, VanGuilder Starkey et al., 2013). Accuracy of our dissections has been verified by quantifying expression of subregion-specific genes (described in VanGuilder Starkey et al., 2013).

## Immunoblotting

As previously described (VanGuilder et al., 2012, VanGuilder Starkey et al., 2012), soluble protein from hippocampal CA1, CA3 and DG was extracted by homogenizing samples in 1% Tween20, 100mM NaCl, 20mM HEPES, 1mM EDTA, 1mM dithiothreitol, 1mM Na<sub>3</sub>VO<sub>4</sub>, and cOmplete Mini EDTA-free protease inhibitor tablets (Roche, Indianapolis, IN). Soluble protein concentrations were determined by BCA assay (Pierce, Rockford, IL), and samples were diluted in lysis buffer and LDS sample buffer (Life Technologies, Carlsbad, CA) and denatured/reduced with 50mM dithiothreitol at 95°C. Sample sizes were as follows: CA1: adult n=7, aged intact n=7, aged impaired n=10; CA3: adult n=7, aged intact n=7, aged impaired n=10; DG: adult: n=5, aged intact: n=5, aged impaired: n=6. Prepared samples were separated using 26-well Criterion TGX 4-20% acrylamide gradient gels (BioRad Hercules, CA) transferred to PVDF (GE Healthcare, Piscataway, NJ) and blocked with 3% bovine serum albumin in PBS/1% Tween20. Membranes were probed with goat polyclonal primary antibodies (anti-LOTUS: sc160254, anti-LGI1: sc9581, anti-ADAM22: sc25999; Santa Cruz Biotechnology, Santa Cruz, CA), washed in PBS/1% Tween20, incubated with HRP-conjugated mouse anti-goat secondary antibody (sc2768, Santa Cruz Biotechnology), developed with enhanced chemiluminescent substrate, visualized on film, and digitized at 800d.p.i. for quantitation by semi-automated digital densitometry (ImageQuant TL, GE Healthcare). Immunoreactive bands were eliminated by incubation of the primary antibody with their respective blocking peptide (5:1 w/w) (LOTUS: sc160254 P, LGI1: sc9581 P, ADAM22: sc25999 P) for two hours prior to addition to the blot (data not shown). Immunoblot data for each sample were normalized to corresponding wholelane densitometry of a total protein-stained gel (VanGuilder et al., 2011a, 2011b, 2012). No differences in protein banding, as determined by protein staining gels, or Actin immunoreactivity were observed, as described previously (VanGuilder et al., 2012).

## qPCR

Quantitation of gene expression was conducted as previously described (Bixler et al., 2011, VanGuilder Starkey et al., 2012) using RNA isolated and purified using standard phase separation/isopropanol precipitation and the RNeasy Mini kit (Qiagen, Valencia, CA). cDNA was synthesized using the High-capacity cDNA Reverse Transcription kit (Life Technologies), and qPCR was performed using TaqMan gene-specific primer/probe assays (LOTUS:Rn00592920\_m1, LGI1:Rn01454071\_m1, ADAM22: Rn01526955\_m1; Life Technologies) with a 7900HT Sequence Detection System (Life Technologies), with  $\beta$ -actin (Rn00667869\_m1) as the endogenous control. Sample sizes were as follows: CA1: adult n=7, aged intact n=7, aged impaired n=10; CA3: adult n=7, aged intact n=7, aged impaired n=10; DG: adult: n=7, aged intact: n=7, aged impaired: n=10. Data were analyzed using the  $2^{-\Delta\Delta C_t}$  method with automatic thresholding (SDS 2.2.2 software, Life Technologies).

## Statistical analysis

Morris water maze behavioral testing data, including statistical analysis of group acquisition and probe trial data, have been extensively described elsewhere (VanGuilder et al., 2011, 2012; VanGuilder Starkey et al., 2012, 2013) and are included to provide context for interpretation of protein data. Protein and mRNA data were evaluated by one-way ANOVA with Student Newman Keuls post hoc testing (ANOVA/SNK). Relationships between protein expression and water maze performance (mean proximity to platform location) were performed by Pearson product moment correlation analysis.

## Results

Decreased expression of LOTUS, LGI1 and ADAM22 with advanced aging was observed throughout the hippocampus, and was frequently exacerbated in aged rats with cognitive

deficits (Figure 2). Significant reductions in LGI1 were evident in CA1, CA3, and DG of aged impaired rats compared to mature adults, and ranged from 30% in CA1 ( $p < 0.01$ ) to greater than 50% in CA3 ( $p < 0.001$ ) and DG ( $p < 0.05$ ). LGI1 expression in aged impaired rats was also significantly reduced compared to aged intact rats in CA1 ( $p < 0.05$ ) and CA3 ( $p < 0.001$ ), but not DG. LOTUS followed a similar trend albeit to a lesser magnitude, and reached statistical significance in CA3 with significantly less LOTUS expression in aged impaired rats compared to aged intact and adult rats ( $p < 0.05$  and  $p < 0.01$ , respectively). Significantly decreased ADAM22 expression was evident in CA1 of both aged intact and aged impaired groups compared to adults ( $\sim 20\%$ ,  $p < 0.05$ ). In CA3, ADAM22 was reduced by 35% in aged intact rats ( $p < 0.01$ ) and by nearly 60% in aged impaired rats ( $p < 0.001$ ) compared to adults. In DG, a 30% reduction in ADAM22 was observed in aged impaired rats ( $p < 0.05$ ). Cognitive status-dependent decreases (i.e., aged impaired versus aged intact) in NgR1-antagonizing protein expression was observed in CA1 (LGI1), CA3 (LGI1, LOTUS), but not DG, while a combinatorial effect of aging and cognitive impairment (aged impaired versus adult) was observed more frequently [CA1 (LGI1, ADAM22), CA3 (LGI1, LOTUS, ADAM22) and DG (LGI1, ADAM22)]. These decreases were evident in the same samples for which cognitive status-dependent increases in MAI ligands (Nogo-A, MAG, and OMgp), NgR1 receptor and co-receptors (TROY, LINGO-1, p75) was evident (Supplemental Table 1) (VanGuilder et al., 2011b, 2012; VanGuilder et al., 2013).

We also observed an inverse relationship between protein expression and Morris water maze performance (Figure 3). Decreasing spatial cognitive ability (i.e., higher mean proximity values) correlated significantly with LGI1 expression in CA1 ( $r = -0.51$ ,  $p < 0.05$ ) and CA3 ( $r = -0.69$ ,  $p < 0.001$ ), and with LOTUS expression in CA3 ( $r = -0.051$ ,  $p < 0.05$ ). ADAM22 expression was consistently negatively correlated with spatial cognition in all three subregions assessed (CA1:  $r = -0.44$ ,  $p < 0.05$ ; CA3:  $r = -0.60$ ,  $p < 0.01$ ; DG:  $r = -0.61$ ,  $p < 0.05$ ).

Interestingly, protein expression changes were recapitulated at the level of LOTUS, LGI1 and ADAM22 mRNA only in CA1 and DG, but not CA3 (Figure S1).

## Discussion

We previously identified the plasticity-suppressing MAI/NgR1 pathway as upregulated in the hippocampus specifically in aged rats with deficits of spatial learning and memory (VanGuilder et al., 2011b, 2012; VanGuilder et al., 2013). LOTUS and LGI1 are recently-identified endogenous NgR1 antagonists that compete with MAIs for NgR1 binding, while ADAM22 interacts with NgR1 to facilitate LGI1/NgR1 binding (Thomas et al., 2010). The findings of our current work indicate that hippocampal expression of LOTUS, LGI1 and ADAM22 is significantly depleted from CA1, CA3 and DG subregions in aged, cognitively impaired rats, and identified a strong correlation of decreasing protein expression with decreasing cognitive ability. Interestingly, while decreased expression of these proteins was frequently observed in aged cognitively intact rats, this occurred to a lesser magnitude in all cases. Together, these data suggest that an endogenous mechanism of NgR1 pathway inactivation fails to compensate for NgR1 pathway induction in the hippocampus with age-related cognitive decline.

The proteins investigated here also function to modulate neuronal function in a variety of ways independent of NgR1 interaction. In addition to its known roles in epilepsy (Fukata et al., 2010), hippocampal LGI1 mediates postnatal pruning and functional maturation of glutamatergic synapses and facilitates AMPA receptor-mediated synaptic transmission (Zhou et al., 2009, Fukata et al., 2010). Interestingly LGI1, which is highly enriched in CA3 pyramidal and DG granule layers (Herranz-Perez et al., 2010) and transcribed almost exclusively by neurons (Senechal et al., 2005), forms a tripartite complex including

ADAM22 and PSD95 (Ogawa et al., 2010) in which ADAM22 is localized to synapses by LGI1 and anchored by PSD95, where it may modulate AMPA receptor aggregation through interactions with stargazin (Fukata et al., 2006). ADAM22 itself has profound effects dendritic maturation and clustering of potassium channels at axonal juxtaparanodes and initial segments (Zhou et al., 2009, Ogawa et al., 2010). LOTUS, on the other hand, impacts gross axonal morphology and fasciculation (Sato et al., 2011), and influences activity-dependent plasticity predominantly through its antagonism of NgR1. It is likely that concomitant upregulation of MAI/NgR1 signaling components and downregulation of NgR1 antagonists exerts a combinatorial effect on hippocampal synapses, leading to aberrant neurotransmission and abnormal inhibition of synaptic plasticity that is reflected in impaired spatial learning and memory (Figure 1).

## Supplementary Material

Refer to Web version on PubMed Central for supplementary material.

## Acknowledgments


This work was supported by National Institute on Aging grants R01AG026607 and P01AG11370 to WES, and the Donald W. Reynolds Foundation. The authors thank Yelena Rossini and Robert Brucklacher for technical assistance.

## References

- Bixler GV, VanGuilder HD, Brucklacher RM, Kimball SR, Bronson SK, Freeman WM. Chronic insulin treatment of diabetes does not fully normalize alterations in the retinal transcriptome. *BMC Med Genomics*. 2011; 4:40. [PubMed: 21575160]
- Delekate A, Zagrebelsky M, Kramer S, Schwab ME, Korte M. NogoA restricts synaptic plasticity in the adult hippocampus on a fast time scale. *Proc Natl Acad Sci*. 2011; 108:2569–2574. [PubMed: 21262805]
- Fukata Y, Adesnik H, Iwanaga T, Brecht DS, Nicoll RA, Fukata M. Epilepsy-related ligand/receptor complex LGI1 and ADAM22 regulate synaptic transmission. *Science*. 2006; 313:1792–1795. [PubMed: 16990550]
- Fukata Y, Lovero KL, Iwanaga T, Watanabe A, Yokoi N, Tabuchi K, Shigemoto R, Nicoll RA, Fukata M. Disruption of LGI1-linked synaptic complex causes abnormal synaptic transmission and epilepsy. *Proc Natl Acad Sci*. 2010; 107:3799–3804. [PubMed: 20133599]
- Herranz-Perez V, Olucha-Bordonau FE, Morante-Redolat JM, Perez-Tur J. Regional distribution of the leucine-rich glioma inactivated (LGI) gene family transcripts in the adult mouse brain. *Brain Res*. 2010; 1307:177–194. [PubMed: 19833108]
- Karlen A, Karlsson TE, Mattsson A, Lundstromer K, Codeluppi S, Pham TM, Backman CM, Ogren SO, Aberg E, Hoffman AF, Sherling MA, Lupica CR, Hoffer BJ, Spenger C, Josephson A, Brene S, Olson L. Nogo receptor 1 regulates formation of lasting memories. *Proc Natl Acad Sci*. 2009; 106:20476–20481. [PubMed: 19915139]
- Kurihara Y, Arie Y, Iketani M, Ito H, Nishiyama K, Sato Y, Nakamura F, Mizuki N, Goshima Y, Takei K. The carboxyl-terminal region of Crtac1B/LOTUS acts as a functional domain in endogenous antagonism to Nogo receptor-1. *Biochem Biophys Res Commun*. 2012; 418:390–395. [PubMed: 22281491]
- Lee H, Raiker SJ, Venkatesh K, Geary R, Robak LA, Zhang Y, Yeh HH, Shrager P, Giger RJ. Synaptic function for the Nogo-66 receptor NgR1: regulation of dendritic spine morphology and activity-dependent synaptic strength. *J Neurosci*. 2008; 28:2753–2765. [PubMed: 18337405]
- Newton IG, Forbes ME, Legault C, Johnson JE, Brunso-Bechtold JK, Riddle DR. Caloric restriction does not reverse aging-related changes in hippocampal BDNF. *Neurobiol Aging*. 2005; 26:683–688. [PubMed: 15708443]
- Ogawa Y, Oses-Prieto J, Kim MY, Horresh I, Peles E, Burlingame AL, Trimmer JS, Meijer D, Rasband MN. ADAM22, a Kv1 channel-interacting protein, recruits membrane-associated




- guanylate kinases to juxtaparanodes of myelinated axons. *J Neurosci.* 2010; 30:1038–1048. [PubMed: 20089912]
- Raiker SJ, Lee H, Baldwin KT, Duan Y, Shrager P, Giger RJ. Oligodendrocyte-myelin glycoprotein and Nogo negatively regulate activity-dependent synaptic plasticity. *J Neurosci.* 2010; 30:12432–12445.
- Sato Y, Iketani M, Kurihara Y, Yamaguchi M, Yamashita N, Nakamura F, Arie Y, Kawasaki T, Hirata T, Abe T, Kiyonari H, Strittmatter SM, Goshima Y, Takei K. Cartilage acidic protein-1B (LOTUS), an endogenous Nogo receptor antagonist for axon tract formation. *Science.* 2011; 333:769–773. [PubMed: 21817055]
- Senechal KR, Thaller C, Noebels JL. ADPEAF mutations reduce levels of secreted LGI1, a putative tumor suppressor protein linked to epilepsy. *Hum Mol Genet.* 2005; 14:1613–1620. [PubMed: 15857855]
- Thomas R, Favell K, Morante-Redolat J, Pool M, Kent C, Wright M, Daignault K, Ferraro GB, Montcalm S, Durocher Y, Fournier A, Perez-Tur J, Barker PA. LGI1 is a Nogo receptor 1 ligand that antagonizes myelin-based growth inhibition. *J Neurosci.* 2010; 30:6607–6612. [PubMed: 20463223]
- VanGuilder HD, Bixler GV, Brucklacher RM, Farley JA, Yan H, Warrington JP, Sonntag WE, Freeman WM. Concurrent hippocampal induction of MHC II pathway components and glial activation with advanced aging is not correlated with cognitive impairment. *J Neuroinflamm.* 2011a; 8:138.
- VanGuilder HD, Bixler GV, Sonntag WE, Freeman WM. Hippocampal expression of myelin-associated inhibitors is induced with age-related cognitive decline and correlates with deficits of spatial learning and memory. *J Neurochem.* 2012; 121:77–98. [PubMed: 22269040]
- VanGuilder HD, Farley JA, Yan H, Van Kirk CA, Mitschelen M, Sonntag WE, Freeman WM. Hippocampal dysregulation of synaptic plasticity-associated proteins with age-related cognitive decline. *Neurobiol Dis.* 2011b; 43:201–212. [PubMed: 21440628]
- VanGuilder Starkey, HD.; Sonntag, WE.; Freeman, WM. *Eur J Neurosci.* 2013. Increased hippocampal NgR1 signaling machinery in aged rats with deficits of spatial cognition. Feb 26th epub ahead of print
- VanGuilder Starkey HD, Van Kirk CA, Bixler GV, Imperio CG, Kale VP, Serfass JM, Farley JA, Yan H, Warrington JP, Han S, Mitschelen M, Sonntag WE, Freeman WM. Neuroglial expression of the MHCII pathway and PirB receptor is upregulated in the hippocampus with advanced aging. *J Mol Neurosci.* 2012; 48:111–126. [PubMed: 22562814]
- Zagrebelsky M, Holz A, Dechant G, Barde YA, Bonhoeffer T, Korte M. The p75 neurotrophin receptor negatively modulates dendrite complexity and spine density in hippocampal neurons. *J Neurosci.* 2005; 25:9989–9999. [PubMed: 16251447]
- Zhou YD, Lee S, Jin Z, Wright M, Smith SE, Anderson MP. Arrested maturation of excitatory synapses in autosomal dominant lateral temporal lobe epilepsy. *Nature Med.* 2009; 15:1208–1214. [PubMed: 19701204]




**Fig. 1. LOTUS, LGI1 and ADAM22 antagonize MAI/NgR1-mediated inhibition of plasticity**

The plasticity-suppressing ligands Nogo-A, MAG and OMgp bind a common receptor, NgR1. Two co-receptor complexes (NgR1/LINGO-1/TROY and NgR1/LINGO-1/p75) transduce MAI/NgR1 signals through intermediaries to activate the GTPase RhoA, which activates a cascade of plasticity-suppressing effectors and resulting in decreased structural remodeling and functional strengthening of synapses. The newly-discovered NgR1 antagonists LOTUS and LGI1 compete with MAIs for NgR1 binding sites and inhibit MAI/NgR1 pathway-mediated suppression of plasticity. ADAM22 interacts with NgR1 to create an LGI1-binding moiety that facilitates LGI1 antagonism of NgR1 to promote plasticity.



**Fig. 2. Expression of NgR1 antagonists decreases with aging and cognitive decline**  
 Protein expression of the antagonist ligands LOTUS and LGI1 and the NgR1 co-receptor ADAM22 decreases significantly in CA1, CA3 and DG subregions with advanced aging. Additional significant decreases were evident specifically between aged intact and aged impaired rats in CA1. Data are presented as % adult mean  $\pm$  standard error; representative blot images are included as insets. \* $p$ <0.05, \*\* $p$ <0.01, \*\*\* $p$ <0.005; ANOVA/SNK.





**Fig. 3. Expression of NgR1 antagonists correlates with spatial learning and memory ability**  
 There is a significant relationship between protein expression of LOTUS, LGI1 and ADAM22 and mean proximity to the escape platform location during Morris water maze probe trials of spatial learning and memory. Lower protein expression was associated with greater proximity values (i.e., indicating inferior targeting of the escape area), particularly in CA3. \* $p < 0.05$ , \*\* $p < 0.01$ , \*\*\* $p < 0.001$ ; Pearson correlation.