

# Quantitative estimation of diacerein in bulk and in capsule formulation using hydrotropic solubilizing agents by UV-spectrophotometry and the first order derivative using the area under curve method

## Abstract

**Ramchandra Pandey,  
Pravin O. Patil,  
Manohar U. Patil,  
Prashant K. Deshmukh,  
Sanjay B. Bari**

H.R. Patel Institute of  
Pharmaceutical Education and  
Research, Shirpur, Dist: Dhule,  
Maharashtra, India

### Address for correspondence:

Pravin O. Patil,  
H. R. Patel Institute of  
Pharmaceutical Education and  
Research, Shirpur Dist:  
Dhule 425 405,  
Maharashtra, India.  
E-mail: rpatilpravin@yahoo.co.in

### Access this article online

Website: [www.phmethods.org](http://www.phmethods.org)

DOI: 10.4103/2229-4708.97703

Quick response code



**Purpose:** This study was designed to develop and validate two simple, rapid, and economical UV-spectrophotometric and the first-order derivative methods using the area under curve method for estimation of diacerein in bulk and in capsule formulation. **Materials and Methods:** In this study, hydrotropic solution of 8 M urea and 0.5 M potassium citrate were employed as the solubilizing agent to solubilize a poorly water-soluble drug, diacerein. In the UV-spectrophotometry method, two wavelengths 252.0 nm and 266.2 nm and in the first-order derivative spectrophotometric methods two wavelengths 259.4 nm and 274.2 nm in 8 M urea and two wavelengths 247.8 nm and 267.4 nm in the UV-spectrophotometry method and in the first-order derivative spectrophotometric methods two wavelengths 259.2 nm and 274.2 nm in 0.5 M potassium citrate were selected for determination of areas. **Results:** Hydrotropic agents used did not interfere in spectrophotometric analysis of diacerein. Diacerein followed linearity in the concentration range of 2–12 µg/mL with a coefficient correlation of 0.999 for both methods. **Conclusion:** The amount of drugs estimated by both proposed methods are in good accord with label claim. The % RSD value in recovery, precision, and ruggedness studies are found to be less than 2 indicate that the method is accurate, precise, and rugged.

**Key words:** Diacerein, derivative spectroscopy, hydrotropic solubilization, potassium citrate, urea

## INTRODUCTION

Diacerein chemically is 4,5-bis(acetyloxy)-9,10-dioxo-2-anthracene carboxylic acid.<sup>[1-2]</sup> It is a selective inhibitor of interleukin-1 having a protective effect on granuloma-induced cartilage breakdown by a reduction in the concentrations of proinflammatory cytokines. It is used in the symptomatic treatment of osteoarthritis.<sup>[3]</sup> The literature survey revealed many analytical methods includes UV-spectroscopic methods,<sup>[4-7]</sup> RP-HPLC,<sup>[8-12]</sup> and HPTLC methods<sup>[13]</sup> for the estimation of diacerein in bulk, pharmaceutical formulation, and in biological samples. The spectrophotometric methods available for diacerein in literature reveal the use of dimethyl acetamide (DMA)<sup>[14]</sup> and dimethyl sulfoxide (DMSO)<sup>[15]</sup> to solubilize diacerein. Drawbacks of organic solvents include their higher cost, toxicity, and pollution. In addition, diacerein is poorly soluble in water. Therefore, hydrotropic solution may be a suitable alternative to exclude the use of organic solvents. Special systems are required to solubilize poorly water-soluble drugs. Hydrotrophy is one of such techniques. Hydrotrophy may be defined as a phenomenon in which the water solubility of poorly water-soluble compounds may be increased several folds by the use of diverse groups of hydrophilic solutes called hydrotropes.<sup>[16]</sup> Hydrotropes commonly used includes sodium benzoate,

sodium acetate, sodium salicylate, nicotinamide, urea, trisodium citrate, sodium ascorbate, piperazine, caffeine, potassium citrate, etc.<sup>[17]</sup> Hydrotropic agents have been observed to enhance the solubility of various substances in water.<sup>[18]</sup> The mixed hydrotropic solubilization process is used to increase the solubility of poorly water-soluble drugs by using blends of hydrotropic agents. There was a miraculous synergistic effect on enhancement in solubility of a poorly water-soluble drug by mixing two hydrotropic agents.<sup>[19]</sup> In the literature, several methods were reported for estimation of poorly water-soluble drugs using hydrotropic solubilizing agents.<sup>[20-22]</sup> Therefore, it was thought useful to utilize this hydrotropic solution to extract out the drug from fine powder of tablets to carry out spectrophotometric estimation.

The objective of the present investigation is to develop simple, precise, and accurate UV-spectrophotometric and first-order derivative methods for determination of diacerein in bulk and in the capsule dosage form using hydrotropic solubilizing agents. The developed methods were validated as per the ICH guidelines.<sup>[23-25]</sup>

## MATERIALS AND METHODS

### Chemicals

Diacerein supplied as a gift sample by Macleods Pharmaceuticals Ltd., Mumbai. Capsule formulation (Orcerein) was purchased from an Indian market, containing Diacerein 50 mg.

### Instrumentation

A UV-visible spectrophotometer (1700 Shimadzu, software UV Probe 2.21) with a spectral bandwidth of 1 nm was employed for all spectroscopic measurements, using a pair of 10 mm matched quartz cells.

### Selection of solvent

Urea and potassium citrate solution were selected as a solvent for developing spectral characteristics of a drug. The selection was made after assessing the solubility in different hydrotropic solvents.

### Preparation of stock standard solutions

Stock standard solutions of diacerein were prepared by dissolving 10 mg in a 100 mL volumetric flask containing 25 mL (8 M urea) solution and the volume was made up to the mark with water and again 10 mg diacerein was dissolved in a 100 mL volumetric flask containing 25 mL 0.5 M potassium citrate solution

and the volume was made up to the mark with water to obtain concentrations 100 µg/mL each. After appropriate dilutions, 10 µg/mL diacerein was scanned in the UV-region, i.e. 400–200 nm. Diacerein showed  $\lambda_{\max}$  258.2 nm in urea solution and 257.4 nm in potassium citrate solution.

### Area under curve method

The area under curve method is used when a broad spectrum of the drug is obtained. From the spectrum of diacerein, the area under the curve in the range of 252.0-266.2 nm for the zero-order spectra [Figure 1] and 259.4-274.2 nm for first-order derivative spectra [Figure 2] were selected for determination of areas using urea solution. In the case of potassium citrate solution, two wavelengths 247.80 and 267.40 nm in the UV spectrophotometric method [Figure 3] and in the first-order derivative spectrophotometric method two wavelengths 259.2 nm and 274.2 nm [Figure 4] were selected for determination of areas.




Figure 1: Zero-order spectra of diacerein showing  $\lambda_{\max}$  258.2 nm in 8 M urea






Figure 2: First derivative spectra of diacerein showing amplitude at 263.2 nm in 8 M urea



**Figure 3:** Zero-order spectra of diacerein showing  $\lambda_{\text{max}}$  257.4 nm in 0.5 M potassium citrate



**Figure 4:** First derivative spectra of diacerein showing amplitude at 264.2 nm in 0.5 M potassium citrate

### Study of linearities curve

An appropriate volume of diacerein in the range of 0.2-1.2 mL were transferred into six separate 10 mL volumetric flasks, from standard stock solutions of urea and potassium citrate and the volume was made up to mark with water to obtain concentration of 2-12  $\mu\text{g}/\text{mL}$  each. The area of diacerein was measured at selected wavelengths, and calibration curves were plotted as concentration *versus* area.

### Analysis of marketed formulation

An accurately weighed capsule content equivalent to 10 mg of diacerein was transfer into a 100 mL volumetric flask containing 25 mL (8 M urea) solution and the volume was made up to the mark with water, similarly 10 mg equivalent capsule content was transferred in a 100 mL volumetric flask containing 25 mL (0.5M potassium citrate) solution and again the volume was made up to the mark with water. This stock solution was filtered through a 0.45  $\mu\text{m}$  Whatmann filter paper. An appropriate volume of solution was further diluted with water to obtain the concentration 6  $\mu\text{g}/\text{mL}$  of diacerein. The area of this solution was measured at selected wavelengths and the concentration of the drug was determined using a linear regression equation. The analysis procedure was repeated for six times with capsule formulation. The results of analysis of the capsule formulation are reported in Table 1 and 2.

### Recovery study

The accuracy of the method was studied at three different levels, i.e. 80%, 100% and 120% levels. To the pre-analyzed sample solution (4  $\mu\text{g}/\text{mL}$  of diacerein), a known amount of drug standards of diacerein was

**Table 1: Results of the assay in 8 M urea**

Drug	UV-spectrophotometry		First-order derivative	
	Amount found (%) [n = 6]	% RSD	Amount found (%) [n = 6]	% RSD
Diacerein 50 mg capsule formulation	99.77	0.81	99.04	1.97

**Table 2: Results of the assay in 0.5 M potassium citrate**

Drug	UV-spectrophotometry		First-order derivative	
	Amount found (%) [n = 6]	% RSD	Amount found (%) [n = 6]	% RSD
Diacerein 50 mg capsule formulation	99.70	0.19	99.35	0.80

added. The solutions were reanalyzed by the proposed method.

## RESULTS AND DISCUSSION

Diacerein showed absorbance maximum at 258.2 nm in the UV-spectrophotometric and 263.2 nm amplitude in first-order derivative spectra in urea solution and 257.4 nm in UV-spectrophotometric and 264.20 nm amplitude in first-order derivative spectra in potassium citrate solution. Diacerein followed linearity in both methods in the concentration range of 2-12  $\mu\text{g}/\text{mL}$  at their respective wavelengths in zero-order and first-order spectra with the linear regression equations  $Y = 0.1228X + 0.2289$  in the UV-spectroscopic method ( $r^2 = 0.9993$ ) [Figure 5] and  $Y = 0.0196X + 0.0483$  in the first-order derivative method ( $r^2 = 0.999$ ) [Figure 6] in urea solution, and  $Y = 0.2428X + 0.4553$  in the UV-spectroscopic method ( $r^2 = 0.9996$ ) [Figure 7] and  $Y = 0.0216X + 0.0189$

in the first-order derivative method ( $r^2 = 0.9992$ ) [Figure 8] in potassium citrate solution. The amount of the drug estimated by using these two methods was found to be in good agreement with the label claim, which indicates that there was no interference from the excipients commonly present in the capsule formulation. These methods were validated for accuracy, precision, and ruggedness. These methods were found to be accurate indicated by low values ( $<2$ ) of % RSD. The precision of the methods were studied as repeatability, intra-day and inter-day precision. The % RSD values for precision studies of diacerein were found to be  $< 2$ , which indicates that the methods to be precise. Both these methods are simple, economical, and rapid and can suitably be used for determination of diacerein in bulk and in capsule formulation. The results are shown in Tables 3 and 4.

## CONCLUSION


The developed UV-spectrophotometric and first-order derivative methods for determination of diacerein were found to be simple, accurate, precise, and

**Table 3: Summary of validation parameters in 8 M urea**


Parameter	UV-spectrophotometry	First-order derivative
% Recovery ( $n = 9$ )	100.21	99.24
% RSD	0.35	0.28
Precision (% RSD)		
Intra-day ( $n = 3$ )	0.37-0.49	0.63-1.68
Inter-day ( $n = 3$ )	0.25-0.50	0.62-1.26
Repeatability [% RSD] ( $n = 6$ )	0.18	0.96
Ruggedness [% RSD]		
Analyst-I ( $n = 6$ )	0.021	0.036
Analyst-II ( $n = 6$ )	0.035	0.031

**Table 4: Summary of validation parameters in 0.5 M potassium citrate**


Parameter	UV-spectrophotometry	First-order derivative
% Recovery ( $n = 9$ )	99.93	99.71
% RSD	0.22	0.37
Precision [% RSD]		
Intra-day ( $n = 3$ )	0.04-0.30	0.33-0.67
Inter-day ( $n = 3$ )	0.05-0.25	0.30-0.45
Repeatability [% RSD] ( $n=6$ )	0.13	0.63
Ruggedness [% RSD]		
Analyst-I ( $n = 6$ )	0.41	0.86
Analyst-II ( $n = 6$ )	0.57	0.72




**Figure 5:** Linear regression equations  $Y = 0.1228X + 0.2289$  in the UV- spectroscopic method ( $r^2 = 0.9993$ ) in 8M urea



**Figure 6:** Linear regression equation  $Y=0.0196X + 0.0483$  in the first-order derivative method ( $r^2=0.999$ ) in 8M urea



**Figure 7:** Quantitative estimation of diacerein in bulk and in capsule formulation: Linear regression equations  $Y = 0.2428X + 0.4553$  in the UV-spectroscopic method ( $r^2 = 0.9996$ ) in 0.5 M potassium citrate



**Figure 8:** Linear regression equation  $Y = 0.0216X + 0.0189$  in the first-order derivative method ( $r^2 = 0.9992$ ) in 0.5 M potassium citrate



economical. These methods were proved to be rugged and can be used for routine analysis of diacerein in bulk and in capsule formulation.

## ACKNOWLEDGMENTS

The authors are thankful to H.R. Patel Institute of Pharmaceutical Education and Research, Shirpur (M.S.), India, for providing the required facilities to carry out this research work.

## REFERENCES

1. The Merck Index – An Encyclopedia of Chemicals, Drugs and Biologicals. 14<sup>th</sup> ed. Whitehouse Station, New Jersey, USA: Merck Research Laboratories; 2006. p. 2980.
2. Indian Pharmacopoeia, Govt. of India, Ministry of Health and Family Welfare. Vol. 2. Ghaziabad: Indian Pharmacopoeial Convention; 2010. p. 1191-3.
3. Tamura T, Shirai T, Kosaka N, Ohmori K, Takafumi K. Pharmacological studies of diacerein in animal models of inflammation, arthritis and bone resorption. *Eur J Pharmacol* 2002;448:81-7.
4. Deshpande MM, Gondane SJ, Mahajan MP, Deshpande AS, Sawant SD. Estimation of Diacerein by UV Spectrophotometry in Bulk and formulation. *Int J Chem Anal Sci* 2011;2:1-2.
5. Narade S, Patil S, Surve S, Shete D, Pore Y. Simultaneous UV spectrophotometric method for the determination of diacerein and aceclofenac in tablets. *J Pharm Sci Res* 2010;2:137-42.
6. Nyola NK, Kalra N. Spectrophotometric determination of diacerein in bulk and pharmaceutical formulation. *Int J Pharm Bio Sci* 2010;1:202-7.
7. Sreejith KR, Premalatha K. Novel spectrophotometric methods for estimation of diacerein from formulations. *Int J Res Pharm Biomed Sci* 2011;2:992-9.
8. Lalithaa KG, Venkatachalam T, Srinivasanb R, Kalaiselvi P, Kannappan N. A simple HPLC method for quantitation of diacerein in tablet dosage Form. *Euras J Anal Chem* 2010;5:81-8.
9. Gandhi SP, Dewani MG, Borole TC, Damle MC. Development and validation of stability indicating HPLC method for determination of diacerein and aceclofenac as bulk drug and in tablet dosage form. *Int J Res Pharm Chem* 2011;1:799-806.
10. Jagadeeswaran M, Gopal N, Sivakumar T. Development and validation of a RP-HPLC method for simultaneous determination of diacerein and aceclofenac in tablet dosage form. *Res Pharm Bio Chem Sci* 2010;1:418-24.
11. Kannappan N, Madhukar A, Srinivasan R, Srinivas RL, Naveen Kumar CH, Mannavalan R. Analytical method development and validation of diacerein tablets by RP-HPLC. *Int J Chem Tech Res* 2010;2:143-8.
12. Sekar V, Jayaseelan S, Udhaya Kumar E, Subash N, Prakash M, Perumal P. Development and validation of RP-HPLC method for the simultaneous estimation of diacerein and aceclofenac in dosage form. *Int J Chem Tech Res* 2010;2:168-71.
13. Chitlange SS, Pawbake GR, Mulla AI, Wankhede SB. Stability-indicating HPTLC method for estimation of diacerein in pharmaceutical dosage form. *Res Pharm Bio Chem Sci* 2010;1:226-34.
14. Gupta KR, Samrit VE, Thakur V, Hemke AT. UV-spectrophotometric estimation of diacerein in pharmaceutical formulation. *J Chem Pharm Res* 2010;2:467-72.
15. Sivakumar R, Nallasivan PK, Saranya KC, Sam Solomon WD, Akelesh T, Venkatnarayanan R. Visible spectrophotometric estimation of diacerein in bulk and pharmaceutical dosage forms. *Pharm Anal* 2010;2:414-6.
16. Neuberger C. Hydrotropy. *Biochemische Zeitschrift* 1916;76:107-76.
17. Coffman RE, Kildsig D. Hydrotropic solubilization- mechanistic studies. *Pharm Res* 1996;13:1460-3.
18. Maheshwari RK. Mixed hydrotropy in spectrophotometric analysis of aceclofenac. *Indian Pharm* 2007;64:67-9.
19. Jain NK, Agrawal RK, Singhai AK. Formulation of aqueous injection of carbamazepine. *Pharmazie* 1990;45:221.
20. Pareek V, Tambe SR, Bhalerao SB. Role of different hydrotropic agents in spectrophotometric and chromatographic estimation of Cefixime. *Int J Pharm Bio Sci* 2010;1:1-10.
21. Chandra M, Sharma S. Determination and validation of UV Spectrophotometric method for estimation of Paracetamol and Diclofenac Sodium in tablet dosage forms using hydrotropic solubilizing agents. *Int J Pharm Tech Res* 2011;3:244-7.
22. Jain N, Jain R, Jain A, Pandey SP, Jain DK. Spectrophotometric method development and validation for quantitative estimation of amlodipine besylate in bulk drug and their dosage forms by using hydrotropic agent. *Euras J Anal Chem* 2010;5:212-7.
23. ICH, Q2A: Text on Validation of Analytical Procedures, International Conference on Harmonization. October 1994.
24. ICH, Q3B: Validation of Analytical Procedures: Methodology, International Conference on Harmonization. November 1996.
25. ICH Harmonized Tripartite Guideline, Q2 (R1): Validation of Analytical Procedures: Text and Methodology. November 2005.

**How to cite this article:** Pandey R, Patil PO, Patil MU, Deshmukh PK, Bari SB. Quantitative estimation of diacerein in bulk and in capsule formulation using hydrotropic solubilizing agents by UV-spectrophotometry and the first order derivative using the area under curve method. *Pharm Methods* 2012;3:4-8.

**Source of Support:** Nil, **Conflict of Interest:** None declared.