



Published in final edited form as:

Hypertension. 2012 February ; 59(2): e8–e9. doi:10.1161/HYPERTENSIONAHA.111.187781.

Blood Pressure Threshold for Abnormal Ocular Fundus Findings is Lower than Expected

Beau B. Bruce, MS, MD,

Department of Ophthalmology, Neurology Emory University, Atlanta, GA

Cédric Lamirel, MD,

Department of Ophthalmology, Emory University, Atlanta, GA

David W. Wright, MD,

Department of Emergency Medicine, Emory University, Atlanta, GA

Valérie Biousse, MD, and

Department of Ophthalmology, Neurology Emory University, Atlanta, GA

Nancy J. Newman, MD

Department of Ophthalmology, Atlanta, Neurology, Neurological Surgery Emory, University, Atlanta, GA

Beau B. Bruce: bbbruce@emory.edu; Cédric Lamirel: cedric.lamirel@gmail.com; David W. Wright: dwwright@emory.edu; Valérie Biousse: vbiousse@emory.edu; Nancy J. Newman: ophntjn@emory.edu

To the Editor

Ocular fundus examination is a critical part of the physical examination in patients with severely elevated blood pressure (BP), which is defined by the Seventh Report of the Joint National Committee on Prevention, Detection, Evaluation, and Treatment of High Blood Pressure (JNC7) as a BP exceeding 180/120 mmHg.¹ Indeed, the presence or absence of severe, grade III/IV, hypertensive retinopathy helps differentiate hypertensive emergencies requiring intensive care, from less severe hypertensive urgencies.¹ As a secondary analysis in the Fundus photography vs. Ophthalmoscopy Trial Outcomes in the Emergency Department (FOTO-ED) study,^{2,3} we sought to explore potential risk factors, in particular BP, for the presence of ocular fundus abnormalities relevant to the care of emergency department (ED) patients. We found evidence of acute end organ ocular damage at lower blood pressures than the JNC7 criteria.

350 adult patients presenting to the Emory University ED with a chief complaint of headache, acute focal neurologic deficit, visual changes, or a diastolic BP (DBP) \geq 120 were prospectively enrolled during the FOTO-ED study (ClinicalTrials.gov NCT00873613) from March 2009 and January 2010. Photographs of the ocular fundus (optic disc and macula) were obtained from both eyes using a commercially available nonmydriatic ocular fundus camera (Kowa α -D, Torrance, CA) by nurse practitioners on eligible patients between 7 AM and 10 PM, 7 days a week. Photographs were reviewed by neuro-ophthalmologists for ocular fundus abnormalities within 24 hours. Presenting complaints, triage systolic (SBP)

Address correspondence and reprints to Dr. Beau B. Bruce, Neuro-Ophthalmology Unit, Emory Eye Center, 1365-B Clifton Rd. NE, Atlanta, GA 30322. Phone: (404)778-5360. Fax: (404)778-4849. bbbruce@emory.edu.

Publisher's Disclaimer: This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final citable form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

and diastolic BP (i.e., on arrival to the ED), heart rate, body mass index (BMI), age, race, and gender were recorded and evaluated as risk factors for ocular fundus abnormalities. Univariate and logistic regression analyses were performed.

Forty-four patients (13.6%) in the FOTO-ED study had ocular fundus abnormalities relevant to ED care: 13 optic disc edema, 4 optic disc pallor, 4 retinal vascular occlusion, 13 isolated intraocular hemorrhages, and 10 grade III/IV hypertensive retinopathy.^{2,3} Exploratory univariate analyses using quintiles of SBP and DBP suggested that the risk of ocular fundus abnormalities increased markedly within the highest quintile of SBP (169 mmHg, 23% abnormal vs. 10%, $p=0.004$) and within the highest two quintiles of DBP (86 mmHg, 19% vs. 9%, $p=0.005$). To avoid collinearity, these criteria were combined and only patients with both SBP 170 mmHg and DBP 90 mmHg were considered to have elevated BP. Obesity (BMI 30) increased the risk of abnormalities (18% vs. 10%, $p=0.049$), as did a chief complaint of visual changes (23 vs. 9%, $p=0.0005$). Heart rate, age, race, and gender were not associated with ocular fundus abnormalities.

Controlling for obesity and visual complaints, the odds ratio of a ocular fundus abnormality among patients with both SBP 170 mmHg and DBP 90 mmHg was 4.2 (95%CI:1.87-9.43) compared to those with a lower BP. Excluding the 21 of 350 (6%) patients who were enrolled on the basis of a DBP 120, this odds ratio was still significant: 2.8 (95%CI: 1.03-7.5, Table).

Seventeen of the 68 patients with elevated BP (25%) had fundus abnormalities. Seven of the seventeen (41%) with abnormalities were among patients enrolled for DBP 120 (all of whom met the JNC7 SBP criteria for hypertensive crisis/emergency). However, the remaining 10 patients (59%) with abnormalities had an elevated BP by the present study's definition (SBP 170 and DBP 90), but did not meet the JNC7 criteria. The ocular fundus findings among these 10 patients were 5 isolated intraocular hemorrhages, 2 grade III/IV hypertensive retinopathies (with BP 193/98 and 178/105), 1 anterior ischemic optic neuropathy, 1 branch retinal artery occlusion, and 1 optic neuritis.

We found that elevated BP (SBP 170 mmHg and DBP 90 mmHg) and obesity are risk factors for abnormal ocular fundus findings among patients presenting to the ED with headache or focal neurologic complaints. Notably, our BP threshold is substantially lower than that at which JNC7 proposes to examine patients for evidence of end organ damage suggestive of hypertensive emergency (i.e., $>180/120$ mmHg). Among our patients meeting this less stringent BP threshold, 90% of the ocular findings (all except optic neuritis) can be clinically associated with elevated BP. Disturbingly, there were even two cases of malignant hypertensive retinopathy among them. This data suggests that the BP threshold proposed by JNC7 for severely elevated blood pressure is likely too high. Furthermore, examining the ocular fundus of patients with both SBP 170 mmHg and DBP 90 mmHg who also have headache or focal neurologic complaints is particularly appropriate.

Future study is necessary to externally validate these proposed lower BP thresholds, determine the applicability of these lower BP thresholds to patients who otherwise have no classic indication for ocular fundoscopic examination (e.g., no headache or focal neurologic complaint), and determine the relevance of the ocular fundus abnormalities seen in patients with elevated BP to long-term outcomes.

Acknowledgments

Sources of Funding: This study was supported in part by an unrestricted departmental grant (Department of Ophthalmology) from Research to Prevent Blindness, Inc., New York, and by NIH/NEI core grant P30-EY06360 (Department of Ophthalmology). Dr. Bruce received research support from the NIH/PHS (KL2-RR025009, UL1-

RR025008), NIH/NEI (K23-EY019341), and the Knights Templar Eye Foundation; and received the American Academy of Neurology Practice Research Fellowship. Dr. Lamirel received research support from Institut Servier (Paris, France), Fondation Planiol (Varenes, France), and the Philippe Foundation, Inc. (New York, NY). Dr. Wright received research support from NIH/PHS (KL2-RR025009). Dr. Biousse received research support from NIH/PHS (UL1-RR025008). Dr. Newman is a recipient of the Research to Prevent Blindness Lew R. Wasserman Merit Award.¹

References

1. National High Blood Pressure Education Program. The Seventh Report of the Joint National Committee on Prevention, Detection, Evaluation, and Treatment of High Blood Pressure: Complete Report. Washington, DC: U.S. Department of Health and Human Services; 2004.
2. Bruce BB, Lamirel C, Wright DW, Ward A, Heilpern KL, Biousse V, Newman NJ. Non-mydratric ocular fundus photography reveals unrecognized relevant findings in the ED: the FOTO-ED study. *N Engl J Med*. 2011; 364:387–389. [PubMed: 21268749]
3. Bruce BB, Lamirel C, Biousse V, Ward A, Heilpern KL, Newman NJ, Wright DW. Feasibility of Nonmydratric Ocular Fundus Photography in the Emergency Department: Phase I of the FOTO-ED Study. *Acad Emerg Med*. 2011; 18:928–933. [PubMed: 21906202]

¹**Conflict of Interest/Disclosure Statement:** No funder had any role in the design or conduct of the study; in the collection, management, analysis, or interpretation of the data; or in the preparation, review, or approval of the manuscript. No conflicting relationship exists for any author.

Table

Multivariate logistic model for abnormal ocular fundus findings in 329 patients presenting to the emergency department with chief complaint of headache, acute focal neurologic deficit, or acute visual changes.

Variable	Odds ratio	95%CI	p-value
SBP 170 mmHg and DBP 90 mmHg	2.8	(1.03-7.5)	0.04
Obesity (BMI 30)	2.3	(1.1-4.8)	0.03
Acute visual chief complaint	5.1	(2.4-10.9)	<0.001

Intercept: -3.2, Hosmer-Lemeshow test, p=0.88, SBP=systolic blood pressure, DBP=diastolic blood pressure, BMI=body mass index