

MOLLUSCICIDAL ACTIVITY OF VARIOUS SOLVENT EXTRACTS FROM *SOLANUM NIGRUM* VAR. *VILLOSUM* L. AERIAL PARTS AGAINST *GALBA TRUNCATULA*

HAMMAMI H.*^a, MEZGHANI-JARRAYA R.**^{a1}, DAMAK M.** & AYADI A.*

Summary:

Molluscicidal activity of *Solanum nigrum* var. *villosum* (morelle velue) extracts and their fractions were tested against the mollusca gastropoda *Galba truncatula* intermediate host of *Fasciola hepatica*. The results indicated that the hydro-methanol (MeOH-H₂O) immature fruit extract possess the highest molluscicidal activity (LC₅₀ = 3.96 mg/L) against *Galba truncatula* compared with other tested compounds. After acido-basic treatment, the methanolic extract fraction isolated from the immature fruits and the richest in alkaloids was the most toxic (LC₅₀ = 1.65 mg/L). The fractions richest in saponosides obtained from the hydromethanolic and methanolic extracts of immature fruits showed interesting molluscicidal activities (LC₅₀ = 6.15 mg/L and LC₅₀ = 7.91 mg/L, respectively). The observed molluscicidal activity could be attributed to the presence of alkaloids or saponosides. So, the immature fruits of *Solanum nigrum* var. *villosum* could be substrates of choice for molluscicide activity. In addition, total alkaloids and saponosides present in this plant deserve further investigations in order to identify the active principles and demonstrate their activities on mollusks in their natural habitat. According to the World Health Organization's guidelines on screening for plant molluscicides, use of these fractions may add to the arsenal of methods to control snail transmitting fasciolosis in tropical and Third World countries where fasciolosis is a common disease.

KEY WORDS: molluscicidal activity, *Galba truncatula*, *Fasciola hepatica*, *Solanum nigrum* var. *villosum* L., alkaloids, saponosides.

Résumé : ACTIVITÉ MOLLUSCICIDE DES DIFFÉRENTS EXTRAITS DES PARTIES AÉRIENNES DE *SOLANUM NIGRUM* VAR. *VILLOSUM* L. CONTRE LE MOLLUSQUE *GALBA TRUNCATULA*

Les activités molluscicides des extraits au dichlorométhane, au méthanol et au méthanol-eau des différents organes (feuilles, fruits immatures et fruits mûres) de *Solanum nigrum* var. *villosum* (morelle velue) ont été testées contre le mollusque gastéropode *Galba truncatula* hôte intermédiaire de *Fasciola hepatica*. Les résultats ont montré que l'extrait hydrométhanolique (MeOH-H₂O) des fruits immatures était le plus actif contre le mollusque *Galba truncatula* (CL₅₀ = 3,96 mg/L) comparativement aux autres composés testés. Après traitement acido-basique, la fraction de l'extrait méthanolique, isolée à partir des fruits immatures et la plus riche en alcaloïdes, était la plus toxique (CL₅₀ = 1,65 mg/L). De plus, les fractions les plus riches en saponosides des extraits hydrométhanoliques et méthanoliques des fruits immatures ont montré des activités molluscicides intéressantes (CL₅₀ = 6,15 mg/L et CL₅₀ = 7,91 mg/L, respectivement). L'activité molluscicide observée pourrait donc être attribuée à la présence d'alcaloïdes ou de saponosides. Ainsi, les fruits immatures de *Solanum nigrum* var. *villosum* pourraient être des substrats de choix pour l'activité molluscicide. De plus, les alcaloïdes totaux et les saponosides présents méritent des investigations supplémentaires afin d'identifier les principes actifs et prouver leurs activités sur les mollusques dans leur habitat naturel. Selon l'OMS, l'utilisation de ces fractions pourrait s'ajouter à l'arsenal des méthodes de lutte contre les mollusques transmettant la fasciolose dans les pays tropicaux et le Tiers-Monde où la fasciolose est endémique.

MOTS-CLÉS : molluscicide, *Galba truncatula*, *Fasciola hepatica*, *Solanum nigrum* var. *villosum* L., alcaloïdes, saponosides.

Fasciolosis caused by *Fasciola hepatica* (*F. hepatica*), a parasite trematode is of considerable medical and veterinarian importance. This old disease has a great potential of expansion in view of the large colonization capacities of its fasciolid causal agents as well as the freshwater lymnaeid snail vector species (Mas-Coma *et al.*, 2001). The control of Fasciolosis depends on the understanding of the epidemiology of the disease, especially the biology, ecology

and distribution of the snails that act as intermediate hosts. It is generally considered that snail control by mollusciciding is of considerable medical and veterinarian importance and can be a rapid and effective means of reducing Fasciolosis transmission (Madsen, 1985). The search for local plants with molluscicidal activity was considered more sustainable than the use of synthetic molluscicides. Plants from Solanaceae family with several species of *Solanum* have been found to have a significant and potentially useful molluscicidal activity (Medina & Ritchi, 1980; Kloos *et al.*, 1987; Beccouche *et al.*, 2000; Wei *et al.*, 2002; Silva *et al.*, 2006). *Galba truncatula* (*G. truncatula*) has been identified as the main intermediate host of fasciolosis in southern Tunisia (Ayadi *et al.*, 1997, Hammami & Ayadi, 1999; Hammami *et al.*, 2007). Previous studies in our laboratories have shown the high molluscicidal

* Fungal and Parasitic Molecular Biology Laboratory, Faculty of Medicine, University of Sfax, 3029, Sfax, Tunisia.

** Laboratory of Chemistry of Natural Products, Faculty of Science, University of Sfax, Tunisia.

^a Co-authors.

Correspondence: Ali Ayadi.

Tel./Fax: 216 74 24 71 30.

E-mail: ali.ayadi@rns.tn

activity of alkaloids especially the N-methylisosalsoleine isolated from *Hammada scoparia*, Chenopodiaceae (Jarraya *et al.*, 2009). It was also reported in earlier studies that crude water extracts of *Solanum nigrum* var. *villosum* has molluscicidal and antiparasitic activity against the mollusca gastropoda *G. truncatula* (Hammami & Ayadi, 2008).

In our on going researches for the isolation and the purification of more plant molluscicidal active products, we aimed our study to explore molluscicidal activities of various solvent extracts and fractions from *Solanum nigrum* var. *villosum* aerial parts against *G. truncatula*.

MATERIAL AND METHODS

PLANT MATERIAL

Solanum nigrum var. *villosum* L. 1753, a common annual herbaceous weed (Solanaceae family) locally known as “Enab-El-Dheeb”, was described in many fields of the south- western of Tunisia by Pottier Alapetite (1981). Samples of this plant were collected in June 2006 from Tozeur’s traditional oases, Tunisia and botanically identified by Prof. Ridha Krichen, Botanist at “École d’Horticulture de Chott Mariem, Sousse, Tunisie”. Voucher specimens (LFPMB10) have been deposited at the “Fungal and Parasitic Molecular Biology Laboratory, Faculty of Medicine, Sfax, Tunisia”. Leaves, ripe red and unripe green fruits collected in Tozeur, Tunisia, were carefully detached from the fresh plant and air-dried.

EXTRACTIONS

Air-dried leaves, ripe red and unripe fruits of *Solanum nigrum* var. *villosum* were extracted (during 24 hours each time) successively in a Soxhlet apparatus by solvents of increasing polarity: dichloromethane, methanol and then lixiviated with methanol-water (8:2, v-v) at room temperature.

All extracts were filtered through paper Whatman No.1 and then concentrated under reduced pressure to afford the dichloromethane extract (DE) and the methanolic extract (ME). For the methanol-water (8:2, v-v) extract, methanol was removed under reduced pressure and the remaining aqueous phase was lyophilized to afford MeOH-H₂O extract. Each extract was weighed (Table I) and kept in dark at 4 °C until use.

PHYTOCHEMICAL SCREENING

The preliminary phytochemical screening was performed according to the Harborne (1964, 1973) methods. Plant extracts (dichloromethane, methanol and methanol-

water) were subjected to chemical tests for the presence of sterols, triterpenoids, carotenoids, tropolone, quinones, alkaloids, flavonoids and saponosides.

A sample of each extract or fraction obtained is dissolved in a minimum amount of solvent adequate for a solution E. This will be added to various reagents according to the protocols experimental T₁, T₂, T₃, T₄, T₅ and T₆.

T₁: reaction of Liebermann: 1 ml E + 0.2 ml of acetic anhydride + 4 drops of H₂SO₄ (cc). Violet changing to blue-green indicates the presence of sterols and (or) triterpenoids.

T₂: reaction of Carr and Price: 1 ml E + 4 drops of SbCl₃ 20 % in CHCl₃. Coloring fleeting blue to blue violet could be a sign of the presence of carotenoids and (or) triterpenoids.

T₃: reaction of Wiustater: 1 ml of E + 0.2 ml of MeOH + 1 drop of FeCl₃ (0.005 M) + 0.6 ml water + 0.4 ml of CHCl₃. Red coloration of the chloroform layer indicates the presence of a tropolone nucleus.

T₄: reaction of Borntraeger: 2 ml E + 2 ml of NaOH (0.1 M). If the aqueous phase is colored of red to violet, we can deduce the presence of free quinons.

T₅: test of flavonoids: 1 ml of E + 2 ml EtOH-H₂O (1-1) + some drops of HCl + Mg chips. Orange to red-violet coloration and blue indicates the presence of flavonoids.

T₆: Mayer reaction: 1 ml of E + 0.5 ml HCl (0.1 N) + 5 drops of Mayer reagent. Formation of a white precipitate indicates the presence of alkaloids.

The result with a single reagent being not absolute because there are false positive, we used a second reagent for alkaloids: the Dragendorff reagent revealing red-orange spots of alkaloids on thin layer chromatography plates.

. Mayer reagent: (solution A: 13.5 g HgCl₂ + 20 ml water and solution B: 49.8 g KI + 20 ml water). The two solutions are mixed and diluted with water to one liter.

. Dragendorff reagent: formula Munier and Macheboeuf for alkaloids: (solution I: dissolve 0.85 g of under bismuth nitrate in a mixture of 10 ml glacial acetic acid + 40 ml water; solution II: 8 g KI + 20 ml water). Immediately before use, mix 15 ml of solution I and 15 ml of solution II, and add 20 ml of glacial acetic acid and make up to 100 ml with water. On TLC, the total steroidal saponins appear red-purple after spraying a solution of chloroform saturated antimony trichloride, and few minutes after heating the plate to 100-110 °C. All steroidal nuclei are located without distinguishing saponosides and glycoalkaloids (GA). The glycoalkaloids (GA) were revealed as orange spots on TLC by Dragendorff reagent.

The concentration of the chemical groups investigated was scored: - (no reaction), + (weakly positive reaction), ++ (important positive reaction), +++ (very important positive reaction).

FRACTIONATION OF ACTIVE EXTRACTS

An acido-basic treatment of each active extract (except hydromethanolic extract of leaves) was performed as follows: m(g) of each extract were solubilized in hydrochloric solution (0.05 M) then extracted thrice by ethyl acetate (EtOAc). Following evaporation of the ethyl acetate, we obtained the extract A (m_A). The aqueous layer was alkalinized by addition of ammoniac then extracted by ethyl acetate until negative Mayer test. After evaporation of the ethyl acetate, we obtained the extract B (m_B) containing total alkaloids revealed as orange spots on TLC by Dragendorff reagent. Then, the remaining aqueous layer was extracted thrice by butanol. After evaporation of the butanol, we obtained the extract C (m_C) containing polar products such as saponins, which were revealed as red-purple spots on TLC by $SbCl_3/CHCl_3$ (Aubert *et al.*, 1989). Finally, the remaining aqueous layer was lyophilized to produce the extract D (m_D).

SNAILS

Uninfected adult *G. truncatula* (3-5 mm in length) were collected locally from a deep draining canal isolated from definitive host, located in the Tozeur's traditional oases in February 2007 and transferred in laboratory aquaria and acclimatized for a minimum period of four days, in holding tanks containing aerated, dechlorinated tap water and washed sand, before being exposed to the aqueous extracts. The toxicity of these preparations was also tested on *G. truncatula* control organisms.

MOLLUSCICIDAL TESTS

Evaluation of the molluscicidal activity of *Solanum nigrum* var. *villosum* extracts and $CuCl_2$ (used as positive control) against snails was investigated as recommended by the World Health Organization (1965). The $CuCl_2$ solution was prepared by adding dechlorinated water to 1 g of $CuCl_2$ up to 1 liter. The

extract or fraction of each plant part and of $CuCl_2$ was used to produce a series of concentrations so that the molluscicidal potency of each sample could be evaluated. The snails were exposed, in groups of ten (five replicates) for 24 to 48 hours, to 500 ml of each concentration of one of the tested materials: leaf, ripe red and unripe green fruit extracts, or $CuCl_2$ (positive control product) shown in Table III.

Galba truncatula negative control organisms were similarly immersed in dechlorinated water. After exposure, snails were rinsed thoroughly in dechlorinated water and left for 48 hours in dechlorinated water before mortality was evaluated.

Mortality was recorded after 24 to 48 hours and dead animals were removed immediately to avoid contamination of animals. Snail mortality was established by the contraction of the body within the shell; no response to a needle probe was taken as evidence of death.

STATISTICS

The concentrations that would kill 50 % (LC_{50}), 90 % (LC_{90}) of the exposed snails and the confidence interval (95 % CI), were determined by the R language analysis which is an integrated suite of software facilities for data manipulation, calculation and graphical display (Ihaka & Gentleman, 1996).

RESULTS

The mass and yields of different plant part extracts of *Solanum nigrum* var. *villosum* after 24 hours extraction with increasing polarity solvents were given in Table I.

The phytochemical tests performed for alkaloids and saponosides were positive for the methanolic extracts and important for the MeOH- H_2O (8:2, v-v) extracts. We detected sterols and terpenoids in dichloromethane extracts. The flavonoids were present in the hydromethanolic extracts (Table II).

The molluscicidal activities of *Solanum nigrum* var. *villosum* extracts and the cupric chloride against the mollusca *G. truncatula* were given in Table III. Toxic

Initial mass of plant parts (g)	Leaves (114 g)		Mature fruits (400 g)		Immature fruits (200 g)	
	Mass (g)	Yield (%)	Mass (g)	Yield (%)	Mass (g)	Yield (%)
Dichloromethane extract (CH_2Cl_2)	4.21	3.69	0.40	0.1	0.18	0.09
Methanol extract (MeOH)	18.06	15.84	21.72	5.43	8.31	4.15
MeOH- H_2O extract (8:2, v-v)	0.05	0.04	2.45	0.61	0.24	0.12

Table I. – Mass and yields of different plant part extracts of *Solanum nigrum* var. *villosum* after 24 hours extraction with increasing polarity solvents.

activity of this plant was determined for dichloromethane, methanolic and hydro-methanolic extracts.

The hydro-methanolic extract of the immature fruits, after 48 hours, showed the highest activity with values for $LC_{50} = 3.96$ mg/L and $LC_{90} = 7.49$ mg/L. By comparison, we obtained values of $LC_{50} = 15.19$ mg/L and $LC_{90} = 18.85$ mg/L for the methanolic extract and $LC_{50} = 34.41$ mg/L and $LC_{90} = 50.47$ mg/L for the dichloromethane extract.

For leaves, the hydro-methanolic extract ($LC_{50} = 15.92$ mg/L) and the methanolic extract ($LC_{50} = 17.51$ mg/L) showed interesting toxicity compared with the dichloromethane extract ($LC_{50} = 31.37$ mg/L).

Mature fruit extracts show the lowest toxicities with $LC_{50} = 50.84$ mg/L for the dichloromethane extract and $LC_{50} = 84.83$ mg/L for the hydro-methanolic extract. The methanolic extracts of mature fruits were inactive.

Cupric chloride, used as a reference substance, killed 90 % of mollusks at 11.34 mg/L after six days of

treatment. Control assays with dechlorinated water showed no effect on the snails (Table III). Haemolysis and hyper secretion of mucus were the common toxic responses of snails to the active tested materials.

The molluscicidal activity of methanolic and hydro-methanolic extracts of *Solanum nigrum* after acidobasic treatment were reported in Table IV.

The highest toxicity, after 48 hours, was shown with immature fruit fractions: the fractions B ($LC_{50} = 1.65$ mg/L) and C ($LC_{50} = 7.91$ mg/L) of the methanolic extract and the fraction C ($LC_{50} = 6.15$ mg/L) of the hydromethanolic extract.

Weaker molluscicidal activities were obtained for the fraction C of leaf methanol extract ($LC_{50} = 13.6$ mg/L) and mature fruit hydromethanolic extract ($LC_{50} = 14.29$ mg/L).

The molluscicidal activity disappeared for the fraction A of each fractionated extract.

Phytochemical test		T ₁	T ₂	T ₃	T ₄	T ₅	T ₆	
		(sterols; triterpenoids)	(carotenoids; triterpenoids)	(tropolone)	(quinone)	(flavonoids)	(alkaloids)	(saponosides)
Dichloromethane	Leaves	+	+	-	-	-	-	-
	Mature fruits	+	+	-	-	-	-	-
	Immature fruits	+	+	-	-	-	-	-
Methanol	Leaves	-	-	-	-	-	+	+
	Mature fruits	-	-	-	-	+	-	-
	Immature fruits	-	-	-	-	-	+	+
Methanol-H ₂ O (8:2, v-v)	Leaves	-	-	-	-	+	+	+
	Mature fruits	-	-	-	-	+	+	++
	Immature fruits	-	-	-	-	+	++	++

- : no reaction; + : weakly positive reaction; ++ : important positive reaction.

Table II. – Chemical groups present in the extracts from aerial parts of *Solanum nigrum* var. *villosum*.

Extract	Leaves		Mature fruits		Immature fruits	
	LC ₅₀ (mg/L) (95 % CI)	LC ₉₀ (mg/L) (95 % CI)	LC ₅₀ (mg/L) (95 % CI)	LC ₉₀ (mg/L) (95 % CI)	LC ₅₀ (mg/L) (95 % CI)	LC ₉₀ (mg/L) (95 % CI)
Dichloromethane (CH ₂ Cl ₂)	31.37 (27.50; 35.24)	46.53 (39.12; 53.94)	50.84 (45.53; 58.15)	64.8 (48.04; 81.21)	34.41 (30.64; 38.18)	50.47 (43.97; 56.97)
Methanol (MeOH)	17.51 (14.86; 20.16)	26.05 (21.97; 30.13)	0	0	15.19 (14.38; 96.19)	18.85 (17.44; 20.26)
MeOH-H ₂ O extract (8:2, v-v)	15.92 (13.43; 18.41)	23.45 (19.63; 27.27)	84.83 (43.28; 26.38)	152.36 (50.32; 254.39)	3.96 (3.11; 4.81)	7.49 (6.18; 8.8)
Dechlorinated water	0	0	0	0	0	0

LC₅₀: 50 % lethal concentration; LC₉₀: 90 % lethal concentration; CI: confidence interval.

CuCl₂ showed molluscicidal activity against the mollusca gastropoda *Galba truncatula* after 48 hours with $LC_{50} = 26.12$ (19.35; 31.69) mg/L and $LC_{90} = 62.71$ (49.83; 96.50) mg/L; and after 144 hours with $LC_{50} = 6.20$ (5.42; 6.94) mg/L and $LC_{90} = 11.34$ mg/L.

Table III. – Molluscicidal activities of *Solanum nigrum* var. *villosum* extracts, after 48 hours exposure time, against *Galba truncatula*.

Initial mass (mi) of extract (g)	Solvent / pH	Mass (mg)	Yields (%)	Activity after 24 H exposure time		Activity after 48 H exposure time		Phytochemical tests	
				LC ₅₀ (mg/L) (95 % CI)	LC ₉₀ (mg/L) (95 % CI)	LC ₅₀ (mg/L) (95 % CI)	LC ₉₀ (mg/L) (95 % CI)	Dragendorff	SbCl ₃
MeOH-H ₂ O / Mature fruits mi = 2.02	EtOAc / acid medium	m _A = 96	4.7	inactive	inactive	inactive	inactive	-	-
	EtOAc / alkali medium	m _B = 20	1	inactive	inactive	inactive	inactive	+	-
	Butanol / alkali medium	m _C = 288.2	14.25	15.54 (6.39 ; 37.75)	28.30 (20.42 ; 39.23)	14.29 (5.98 ; 34.10)	27.37 (14.79 ; 50.66)	-	++
MeOH / Leaves mi = 0.69	Aqueous lyophilized layer	m _D = 1,440	71.28	inactive	inactive	inactive	inactive	-	-
	EtOAc / acid medium	m _A = 182.1	26.38	inactive	inactive	inactive	inactive	-	-
	EtOAc / alkali medium	m _B = 3.6	0.52	inactive	inactive	inactive	inactive	+	-
MeOH-H ₂ O / Immatures fruits mi = 0.15	Butanol / alkali medium	m _C = 73	10.57	12.59 (2.61; 60.59)	22.76 (11.88; 43.59)	13.64 (2.28; 81.47)	24.36 (17.46; 34.00)	-	++
	Aqueous lyophilized layer	m _D = 340	49.27	inactive	inactive	inactive	inactive	-	-
	EtOAc / acid medium	m _A = 26.5	0.18	inactive	inactive	inactive	inactive	-	-
MeOH-H ₂ O / Immatures fruits mi = 0.15	EtOAc / alkali medium	m _B = 6.7	4.47	inactive	inactive	inactive	inactive	+	-
	Butanol / alkali medium	m _C = 23.6	15.73	10.96 (1.31; 91.61)	36.38 (19.32; 68.51)	6.15 (0.10; 347.89)	18.16 (2.83; 116.43)	-	+++
	Aqueous lyophilized layer	m _D = 90	60	inactive	inactive	inactive	inactive	-	-
MeOH / Immatures fruits mi = 0.65	EtOAc / acid medium	m _A = 81.8	12.58	inactive	inactive	inactive	inactive	-	-
	EtOAc / alkali medium	m _B = 11.1	1.7	2.56 (0.78; 8.46)	7.56 (3.44; 16.61)	1.65 (0.61; 4.45)	4.77 (0.83; 27.45)	+++	-
	Butanol / alkali medium	m _C = 181.6	27.94	14.69 (4.09; 52.71)	48.07 (21.10; 109.51)	7.91 (0.27; 234.79)	26.20 (7.99; 85.86)	-	+++
Dechlorinated water	Aqueous lyophilized layer	m _D = 340.2	52.34	24.59 (1.13; 536.74)	89.40 (5.44; 146.93)	20.46 (3.11; 134.43)	55.26 (17.89; 170.61)	-	+
				0	0	0	0		

LC₅₀: 50 % lethal concentration; LC₉₀: 90 % lethal concentration; CI: confidence interval. CuCl₂ showed molluscicidal activity against the mollusca gastropoda *Galba truncatula* after 48 hours with LC₅₀ = 26.12 (19.35; 31.69) mg/L and LC₉₀ = 62.71 (49.83; 96.50) mg/L; and after 144 hours with LC₅₀ = 6.20 (5.42; 6.94) mg/L and LC₉₀ = 11.34 mg/L.

Table IV. – Yields, phytochemical tests and molluscicidal activities of *Solanum nigrum* var. *villosum* fractions, after 24 and 48 hours exposure time, against *Galba truncatula*.

DISCUSSION

One of the major preventive steps against fasciolosis is the control of the vector snail populations. In recent years, several plant extracts have been studied against snail transmitted parasitic diseases and several species of *Solanum* have been found to have significant and potentially useful molluscicidal activity. Such activity varies greatly from species to species and even between different parts of the same plant (Silva *et al.*, 2005).

During our study, only immature fruit and leaf extracts showed significant molluscicidal activities with LC₅₀ values that fell well below the upper threshold of 40 mg/L set for a potential molluscicide by the World Health Organization (WHO, 1993). So that, the immature fruit extracts have the most toxicity against the snail *G. truncatula* correlating with previous studies (Hammami & Ayadi, 2008). By comparison with immature fruit crude water extract of the same plant (*Solanum nigrum* var. *villosum* collected from Tozeur's oases) obtained values (LC₅₀ = 41.2 mg/L and LC₉₀ = 66.6 mg/L) in our previous studies, the immature fruit hydro-methanolic extract was more efficacious and demonstrated the importance of the plant components separation.

This result was even supported by Ndamukong *et al.* (2006) when studying molluscicidal activity of *Solanum scabrum* against the snails *Bulinus camerunensis* and *B. truncatus* showing that methanolic extracts were more toxic than water extracts.

Methanolic extract was found even possessing a strong molluscicidal activity and exhibited potent effect on mollusks (Lahlou *et al.*, 2002).

In the present study, methanolic and hydro-methanolic extracts of the aerial green parts (leaves and immature fruits) were the most toxic from the whole extracts tested. These findings are in agreement with other data in the literature. For instance, aerial parts methanolic extracts of *Solanum jabrense* were described to have molluscicidal activity with LC₅₀ = 56.0 mg/L and LC₉₀ = 80.3 mg/L (Silva *et al.*, 2006).

It was even reported that the methanolic extract of the berries of *Solanum aculeastrum* also showed significant activity against host snails of schistosomiasis, the LC₁₀₀ = 50mg/L (Wanyonyi *et al.*, 2003).

According to other studies, the cercaricidal activity was related to the flavonoids, terpenoids, and saponosides (Chifundera *et al.*, 1993; Perret & Whitefield, 1995; Lahlou *et al.*, 2002). It was also reported that the toxicity in *Solanum* genera was accorded to the presence of flavonoids (Silva *et al.*, 2002) and steroidal alkaloids in the fruits (Bhattacharyya, 1984; Valverde *et al.*, 1993). Alkaloids of stems of *Solanum stipula-*

ceum were toxic against *Biomphalaria glabrata* LC₅₀ = 45.2 mg/L and LC₉₀ = 56.0 mg/L (Silva *et al.*, 2006).

In our study, the fractionation has concerned only active extracts showing significant LC₅₀ below the upper threshold of 40 mg/L (Table III) fixed by WHO (1993) and that were positive for alkaloid and saponosides phytochemical tests (Table II). Leaf hydromethanolic extract having a fewer yield (0.04 %) was not fractionated (Table I).

During our study, after acido-basic treatment, we noticed that fractions showing molluscicidal activity contained generally alkaloid and/or saponosides (Table IV).

Among the eight different total alkaloid and saponosides fractions tested, that from immature fruit methanolic extract was the most potent in *G. truncatula* (LC₅₀ = 1.65 mg/L) but it was presented in a weaker yield of 1.7 %. Even in other studies, it was reported that *S. nigrum* immature fruits are not a promising source of the alkaloids because of their small size and declining alkaloid content (Eltayeb *et al.*, 1997). Other workers have reported steroidal alkaloids in *Solanum incanum* fruits at levels ranging from traces (Lin *et al.*, 1990) to 0.031 % (Fukuhara & Kubo, 1991) on a fresh weight basis in mature fruits.

The fractions B less rich in alkaloids from mature fruit hydromethanolic (1 %) and leaf methanolic (0.52 %) extracts were not active against *G. truncatula* and that may be attributed to the fact that total alkaloids were not enough to produce molluscicidal activity.

The fraction C rich in saponosides was active in all our extracts tested and it presented important yields (ranging from 10.57 % to 52.34 %) and interesting molluscicidal activity. The fraction C from hydromethanolic immature fruit extract was found to be the next toxic against *G. truncatula* (LC₅₀ = 6.15 mg/L) and after that was classified the fraction C from immature fruit methanolic extract (LC₅₀ = 7.91 mg/L). We deduce that molluscicidal activity was related to polar fractions rich in alkaloids and /or saponosides and that the fraction richest in alkaloids from the immature fruits was more toxic than those rich in saponosides.

Saponosides fractions from leaves methanolic extract and mature fruits hydromethanolic extracts have shown weaker molluscicidal activity (LC₅₀ = 13.64 mg/L and LC₅₀ = 14.29 mg/L, respectively).

So, we concluded that immature fruits were best candidates for molluscicidal activity. This is in agreement with earlier studies carried out in our laboratories confirming the highest toxicity of crude water extracts of immature fruits of the same plant *Solanum nigrum* var. *villosum* (Hammami & Ayadi, 2008).

Immature fruits more rich in alkaloids were more toxic than mature fruits, this may be due to the high concentration of steroidal alkaloids. So, it was demonstrated

that small unripe fruits of *Solanum nigrum* had a high concentration of solasodine that decreased with fruit maturation (Eltayeb *et al.*, 1997).

Phytochemical tests indicated the steroidal nature of the total alkaloids of *Solanum nigrum*. Several steroidal alkaloids including solasodine, solasonine and solamargine were isolated and identified in this plant (Schreiber, 1968; Ripperger & Schreiber, 1981; Drewes & Staden, 1995; Eltayeb *et al.*, 1997). Steroidal saponins were previously isolated and identified as nigrumin I and II (Tsuyoshi *et al.*, 2000). Six new steroidal saponins were also isolated by Zhou *et al.* (2006).

Hence, the observed molluscicidal activity may be due to the presence of steroidal alkaloids. It was reported that steroidal alkaloids from *Solanum aculeastrum* possess strong molluscicidal properties (Wanyonyi *et al.*, 2002). In other studies, saponins and steroidal alkaloids were isolated from *Solanum melongena* L. fruits (Aubert *et al.*, 1989).

CONCLUSION

In the present study, molluscicidal activity of *Solanum nigrum* var. *villosum* varies between different parts of this plant and may be attributed to the nature of components. The analyses by thin layer chromatography achieved on the different extracts of this plant showed that the methanolic and hydro-methanolic ones were the richest in alkaloids and/or saponosides and the most active against *G. truncatula*. So, molluscicidal activity may be correlated with the content of total alkaloid and/or saponosides found in the methanolic and the hydro-methanolic immature fruit extracts. This finding is in agreement with other data in the literature confirming that alkaloids from methanolic extracts of *Solanum aculeastrum* showed significant activity against host snails of schistosomiasis (LC₅₀ = 50 ppm) (Wanyonyi *et al.*, 2003). However, molluscicidal activity of *Solanum nigrum* var. *villosum* collected in Tunisia was more active and especially the hydro-methanolic extract (LC₅₀ = 3.96 mg/L) and the methanolic extract fraction richest in alkaloids (LC₅₀ = 1.65 mg/L) isolated from the immature fruits.

Even the fractions richest in saponosides from the immature fruit hydromethanolic and methanolic extracts have shown interesting molluscicidal activity (LC₅₀ = 6.15 mg/L and LC₅₀ = 7.91 mg/L, respectively). In our knowledge, this is the first study proving that immature fruit methanolic or hydromethanolic fractions rich in alkaloids and/or saponosides of *Solanum nigrum* var. *villosum* have important molluscicidal activity against the snail gastropoda *G. truncatula*.

Green berries of *Solanum nigrum* var. *villosum* offers promised as a potential molluscicidal control agent

and the total alkaloids and saponosides from these plant merit further investigations to identify the active principal responsible of molluscicidal properties and to further assess their activity in *G. truncatula* and its associated fauna in their natural habitats.

According to the World Health Organization's guidelines on screening for plant molluscicides (WHO, 1983), use of these fractions may add to the arsenal of methods to control snail transmitting fasciolosis in tropical and third world countries where fasciolosis is a common disease.

ACKNOWLEDGEMENTS

This study was funded by the Ministry of Higher Education, Scientific Researches, Tunisia.

REFERENCES

- AUBERT S., DAUNAY M. C. & POCHARD E. Steroidic saponins of Egg-plant (*Solanum melongena* L.) I: Food interest, analytical methodology, localization in fruit. *Agronomie*, 1989, 9, 641-651.
- AYADI A., MAKNI F. & BEN SAÏD M. État actuel de la fasciolose en Tunisie. *Bulletin de la Société Française de Parasitologie*, 1997, 15, 27-32.
- BEKKOUCHE K., MARKOUK M., LARHSINI M., JANA M. & LAZREK H.B. Molluscicidal properties of glycoalkaloids extracts from Moroccan *Solanum* species. *Phytotherapy Research*, 2000, 14, 366-367.
- BHATTACHARYYA J. Isolation of solasodine from the fruits of *Solanum asperum* and *Solanum paludosum*. *Journal of Natural Products*, 1984, 47, 1059-1060.
- CHIFUNDERA K., BALUKA B. & MASHIMANGO B. Phytochemical screening and molluscicidal potency of some Zairian medicinal plants. *Pharmacology Research*, 1993, 28, 333-340.
- DREWES F.E. & STADEN J.V. Aspects of extraction and purification of solasodine from *Solanum aculeastrum* tissues. *Phytochemical analysis*, 1995, 6, 203-206.
- ELTAYEB E., AL-ANSARI A.S. & RODDICK J.G. Changes in the steroidal alkaloid solasodine during development of *Solanum nigrum* and *Solanum incanum*. *Phytochemistry*, 1997, 46, 489-494.
- FUKUHARA K. & KUBO I. Isolation of steroidal glycoalkaloids from *Solanum incanum* by two countercurrent chromatographic methods. *Phytochemistry*, 1991, 30, 685-688.
- HAMMAMI H. & AYADI A. Écologie de *Lymnaea truncatula* Muller, hôte intermédiaire de *Fasciola hepatica* Linné dans le microclimat de Tozeur (sud-ouest de la Tunisie). *Bulletin de la Société de Pathologie Exotique*, 1999, 92, 302-304.
- HAMMAMI H., HAMED N. & AYADI A. Epidemiological studies on *Fasciola hepatica* in Gafsa oases (South west of Tunisia). *Parasite*, 2007, 14, 261-264.

- HAMMAMI H. & AYADI A. Molluscicidal and antiparasitic activity of *Solanum nigrum villosum* against *Galba truncatula* infected or uninfected with *Fasciola hepatica*. *Journal of Helminthology*, 2008, 82, 235-239.
- HARBORNE J.B. Biochemistry of phenolic compounds. Academic press, London, 1964, 93-111.
- HARBORNE J.B. Phytochemical methods. Chapman & Hall, London, 1973.
- IHAKA R. & GENTLEMAN R.R. A language for data analysis and graphics. *Journal of Computational and Graphical Statistics*, 1996, 3, 299-314.
- JARRAYA R.M., HAMMAMI H., AYADI A. & DAMAK M. Molluscicidal activity of *Hammada scoparia* (Pomel) Iljin leaf extracts and the principal alkaloids isolated from them against *Galba truncatula*. *Memórias do Instituto Oswaldo Cruz*, 2009, 104, 1035-1038.
- KLOOS H., THIONGO F. W., OUMA J.H. & BUTTERWORTH A.E. Preliminary evaluation of some wild and cultivated plants for snail control in Machakos district, Kenya. *The Journal of Tropical Medicine and Hygiene*, 1987, 90, 197-204.
- LAHLOU M., EL MAHI M. & HAMAMOUCHE J. Evaluation of anti-fungal and molluscicidal activity of Moroccan *Zizyphus lotus* (L.) Desf. *Annales Pharmaceutiques Françaises*, 2002, 60, 410-414.
- LIN C.N., LU C.M., CHENG M.K. & GEN K.H. The cytotoxic principles of *Solanum incanum*. *Journal of Natural Products*, 1990, 53, 513-516.
- MADSEN H. Ecology and control of African freshwater pulmonate snails. Part I: Life cycle and methodology. Danish Bilharziasis Laboratory, Charlottenlund, Denmark, 1985.
- MAS-COMA S., FUNATSU I.R. & BARGUES M.D. *Fasciola hepatica* and lymnaeid snails occurring at very high altitude in South America. *Parasitology*, 2001, 123, 115-127.
- MEDINA F.R. & RITCHIE L.S. Molluscicidal activity of the Puerto Rican weed *Solanum nodiflorum* against snail hosts of *Fasciola hepatica*. *Economic Botany*, 1980, 34, 368-375.
- NDAMUKONG K.J., NTONIFOR N.N., MBUH J., ATEMNKENG A.F. & AKAM M.T. Molluscicidal activity of some Cameroonian plants on *Bulinus* species. *East African Medical Journal*, 2006, 83, 102-109.
- PERRET S. & WHITFIELD P.J. Currently available Molluscicides. *Parasitology Today*, 1995, 12, 156-159.
- POTTIER-ALAPETITE G. Flore de la Tunisie Angiospermes-Dicotylédones Gamopétales. Imprimerie Officielle de la République Tunisienne, Tunis, 1981, 821.
- RIPPERGER H. & SCHREIBER K. *Solanum* steroid alkaloids, in: The alkaloids. Chemistry and Physiology. Manske R.H.F. & Rodrigo R.G.A. (Eds), New York, Academic Press, 1981, Vol. XIX, 81-192.
- SCHREIBER K. Steroid alkaloids: the *Solanum* group, in: The alkaloids. Chemistry and Physiology. Manske R.H.F. (Ed.). New York, Academic Press, 1968, Vol. X, 1-192.
- SILVA T.M., BRAZ-FILHO R., CARVALHO M.G. & AGRA M.F. Flavonoids and an alkamide from *Solanum paludosum* Moric. *Biochemical Systematics and Ecology*, 2002, 30, 479-481.
- SILVA T.M., BATISTA M., CAMARA C.A. & AGRA M.F. Molluscicidal activity of some Brazilian *Solanum* spp. (Solanaceae) against *Biomphalaria glabrata*. *Annals of Tropical Medicine and Parasitology*, 2005, 99, 419-425.
- SILVA T.M., CAMARA C.A., AGRA M. DE F., DE CARVALHO M.G., FRANA M.T., BRANDOLINE S.V., DA SILVA PASCHOAL L. & BRAZ-FILHO R. Molluscicidal activity of *Solanum* species of the Northeast of Brazil on *Biomphalaria glabrata*. *Fitoterapia*, 2006, 77, 449-452.
- TSUYOSHI I., HIDETSUGU T. & TOSHIHIRO N. Steroidal oligoglycosides from *Solanum nigrum*. *Chemical & Pharmaceutical Bulletin*, 2000, 48, 1062-1064.
- VALVERDE M., LAVAUD C., BOUSTIE J., EL BADAOUI H., MUGUET B. & HENRY M. Solamargine: the main glycoalkaloid from the fruits of *Solanum paludosum*. *Planta Medica*, 1993, 59, 483-484.
- WANYONYI A.W., CHHABRA S.C., MKOJI G., EILERT U. & NJUE W.M. Bioactive steroidal alkaloid glycosides from *Solanum aculeastrum*. *Phytochemistry*, 2002, 59, 79-84.
- WANYONYI A.W., CHHABRA S.C., MKOJI G., NJUE W. & TARUS P.K. Molluscicidal and antimicrobial activity of *Solanum aculeastrum*. *Fitoterapia*, 2003, 74, 298-301.
- WEI F.H., XU X.J., LIU J.B., DAI Y.H., DUSSART G. & TRIGWELL J. Toxicology of a potential molluscicide derived from the plant *Solanum xanthocarpum*: a preliminary study. *Annals of Tropical Medicine and Parasitology*, 2002, 96, 325-331.
- WORLD HEALTH ORGANIZATION. Molluscicide screening and evaluation. *Bulletin of the World Health Organization*, 1965, 33, 567-581.
- WORLD HEALTH ORGANIZATION. Report of the Scientific Working Group on Plant Molluscicides. Document WHO/TDR/SCH-SWE 4, 83.3, WHO, Geneva, 1983, 1-11.
- WORLD HEALTH ORGANIZATION. Tropical Disease Research. *Bulletin of the World Health Organization*, 1993, 830, 1-86.
- ZHOU X., HE X., WANG G., GAO H., ZHOU G., YE W. & YAO X. Steroidal saponins from *Solanum nigrum*. *Journal of Natural Products*, 2006, 69, 1158-1163.

Received on November 4th, 2009Accepted on June 9th, 2010