

Unwinding the Wnt action of casein kinase 1

Cell Research (2013) 23:737-738. doi:10.1038/cr.2013.51; published online 9 April 2013

The casein kinase 1 (CK1) family, a major intracellular serine/threonine kinase, is implicated in multiple pathways; however, understanding its regulation has proven challenging. A recent study published in *Science* identifying allosteric activation of CK1 by the DEAD-box RNA helicase DDX3 expands our understanding of the control of this abundant kinase family.

The human CK1 protein kinase family is encoded by six genes (α , $\gamma 1$, $\gamma 2$, $\gamma 3$, δ , and ϵ) and regulates diverse biochemical processes including hedgehog signaling, circadian rhythms and the p53 tumor suppression [reviewed in 1]. In the Wnt/ β -catenin pathway, all CK1 family members are involved, each with a distinct role. To carry out their functions, CK1 family members achieve specificity by several mechanisms, but how their kinase activity is regulated has been less clear. Here, we discuss the findings of the Niehrs lab [2] in the context of what is known about CK1 control in the Wnt pathway.

CK1 γ proteins are membrane bound due to C-terminal S-palmitoylation and phosphorylate the Wnt co-receptor LRP5/6 in the presence of Wnts and Disheveled to activate the pathway [3, 4]. One mechanism of activation may be via 'priming' by upstream phosphorylation of LRP5/6, a common characteristic of CK1 substrate recognition [5]. CK1 δ and CK1 ϵ bind to and phosphorylate Disheveled, an activity regulated by Wnt signaling and protein phosphatases [6, 7]. CK1 α interacts with and phosphorylates APC, Axin and Ser45 of β -catenin in an apparently unregulated reaction. The CK1 α -catalyzed phosphorylation

primes β -catenin for further phosphorylation by GSK3 and subsequent degradation. How does CK1 accomplish so many different jobs in the Wnt pathway and how is it controlled?

A key mechanism for regulation is CK1s' differential interaction with scaffolds and membranes. CK1 δ and CK1 ϵ bind to substrates including Disheveled, Period and NFAT1; CK1 α interacts with Axin, and CK1 γ localizes to membranes where it phosphorylates LRP6. These interactions take place at protein motifs distinct from the phosphorylation sites. However, binding and co-localization alone are probably not sufficient for precise biological control.

Each CK1 isoform is likely to be regulated differently. CK1 α is the smallest member of the family (~38 kDa), and has been thought to be constitutively active. CK1 δ and CK1 ϵ have closely-related C-terminal domains (148-184 aa) that are actively autophosphorylated, resulting in a kinase-phosphotail interaction that restricts access of protein substrates to the active site of the kinase. CK1 δ and CK1 ϵ can be relieved of this auto-inhibition by the action of protein phosphatases that in turn can be stimulated by extracellular signals such as glutaminergic and Wnt signaling [1, 6]. The regulation of CK1 γ is not well understood.

Although the kinase domains between CK1s are highly conserved, subtle differences govern their binding to scaffolds. For example, two key residues determine the differential binding of CK1 α and CK1 ϵ to Disheveled and Period [8]. Motifs on the scaffolds also facilitate binding to CK1. CK1 ϵ binds to an F-X-X-X-F motif on PER2 and

NFAT1 that is quite distal from the phosphorylation sites [9]. The F-X-X-X-F motif is also present on additional CK1 partners including DDX3, although its importance has not yet been tested. The presence of kinase-binding motifs can greatly enhance the phosphorylation of the substrate. Thus, regulating the affinity of CK1 for scaffold-binding sites can have profound effects on rates of phosphorylation.

Protein kinase activity can be controlled by diverse mechanisms, the most commonly studied being phosphorylation, addition or removal of regulatory subunits, and targeting to scaffolds (Figure 1). An additional, under-explored mechanism is allosteric regulation. While allosteric regulation has a proud history in enzymology, there are only a few examples (e.g., AMP-kinase, phospholase kinase) of small-molecule allosteric regulation of protein kinases [reviewed in 10]. Notably, a recent screen for inhibitors of the Wnt/ β -catenin pathway identified the drug pyrvinium pamoate as an allosteric activator of CK1 α [11]. As a clue to mechanism, pyrvinium bound to but did not activate other CK1 isoforms. However, it could activate CK1 δ lacking its C-terminal regulatory domain. This suggests that there is a conserved site in the CK1 family to which pyrvinium binds that allosterically activates the kinases. Additional inhibitory mechanisms, such as the C-terminal phosphodomains of CK1 δ and CK1 ϵ , may be able to override the small-molecule activation. The finding of allosteric activation by pyrvinium suggests that endogenous allosteric regulators of the CK1 family may also exist.

Cruciat *et al.* [2] now provide evidence for protein allosteric activators of CK1. In a cell-based RNAi screen for new regulators of the Wnt/ β -catenin signaling, they identified the DEAD-box RNA helicase DDX3. DDX3 is required both for Wnt/ β -catenin signaling in human cells as well as Wnt-dependent apical-posterior neural patterning of the central nervous system in *Xenopus* and neuroblast migration in *C. elegans*. Epistatic and biochemical analysis place DDX3 at the level of LRP6 and Disheveled phosphorylation. DDX3 cooperates with CK1 ϵ in phosphorylating Disheveled, and physically interacts with CK1 ϵ after Wnt stimulation. Kinetic analysis revealed that DDX3 is an allosteric activator of all CK1 family members tested.

The DDX genes encode a family of DEAD-box RNA helicases, so named for the “DEAD” (Asp-Glu-Ala-Asp) amino acid sequence in their motif II. The 37 members of the DDX family of proteins have the defining function of unwinding double-stranded RNA in an ATP-dependent manner. DDX proteins are multifunctional, playing roles in translation initiation, mRNA transport and spliceosome assembly [12]. Stimulating CK1 activity is a ‘moonlighting’ function of DDX3, as key helicase domains and the helicase activity of DDX3 are not required to stimulate Wnt signaling and activate CK1 ϵ .

DDX3 and pyrvinium are similar in that both are allosteric activators of CK1. Whether the underlying mechanisms of activation are similar remains to be determined. The C-terminal regulatory domain of CK1 δ blocks pyrvinium action [11], whereas DDX3 is able to activate full-length CK1 δ . Pyrvinium and DDX3 might interact with the same region on CK1, but the protein activators may function more efficiently than small molecules. Future structural studies may provide us with more definitive answers.

While DDX3 was identified as a Wnt/ β -catenin activator via CK1 ϵ , it also activates CK1 α and CK1 γ , raising




Figure 1 Regulation of the CK1 family. As described in the text, diverse mechanisms exist to regulate the activity of CK1.

the question of why one activity predominates over the other in the cell- and animal-based assays used by Cruciat *et al.* Furthermore, the authors found that a handful of other DDX proteins activate CK1. While this suggests redundancy, they found strong phenotypes with loss-of-function DDX3 alone. This may be due to tissue-specific expression. It is also possible that other binding partners may modulate the interaction of DDX proteins with CK1.

As always, more questions remain. Does DDX3 regulation of CK1 extend beyond Disheveled signaling? For example, could DDX proteins activate CK1 δ/ϵ to phosphorylate Period proteins during circadian rhythms? How does Wnt signaling stimulate the interaction between CK1 ϵ and DDX3? Is DDX3 also a substrate for Wnt-activated CK1 ϵ , and what effect does that have on DDX3’s other functions? How do all these regulatory mechanisms on the various CK1s work together temporally and spatially? Further work will illuminate what happens when the DEAD-box family brings CK1 to life.

Daniel GR Yim^{1,2},
David M Virshup¹

¹Program in Cancer and Stem Cell Biology, Duke-NUS Graduate Medical School Singapore, Singapore; ²Graduate School for Integrative

Sciences and Engineering, National University of Singapore, Singapore
Correspondence: David M Virshup
E-mail: david.virshup@duke-nus.edu.sg

References

- Cheong JK, Virshup DM. *Int J Biochem Cell Biol* 2011; **43**:465-469.
- Cruciat CM, Dolde C, de Groot RE, *et al. Science* 2013; **339**:1436-1441.
- Davidson G, Wu W, Shen J, *et al. Nature* 2005; **438**:867-872.
- Zeng X, Tamai K, Doble B, *et al. Nature* 2005; **438**:873-877.
- Flotow H, Graves PR, Wang AQ, *et al. J Biol Chem* 1990; **265**:14264-14269.
- Swiatek W, Tsai IC, Klimowski L, *et al. J Biol Chem* 2004; **279**:13011-13017.
- Gao ZH, Seeling JM, Hill V, *et al. Proc Natl Acad Sci USA* 2002; **99**:1182-1187.
- Dahlberg CL, Nguyen EZ, Goodlett D, *et al. PLoS One* 2009; **4**:e4766.
- Okamura H, Garcia-Rodriguez C, Martinson H, *et al. Mol Cell Biol* 2004; **24**:4184-4195.
- Lindsley JE, Rutter J. *Proc Natl Acad Sci USA* 2006; **103**:10533-10535.
- Thorne CA, Hanson AJ, Schneider J, *et al. Nat Chem Biol* 2010; **6**:829-836.
- Schröder M. *Biochem Pharmacol* 2010; **79**:297-306.