

Novel Tumor Suppressor Function of Glucocorticoid-Induced TNF Receptor GITR in Multiple Myeloma

Yang Liu¹, Phong Quang¹, Esteban Braggio², Hai Ngo¹, Gayane Badalian-Very¹, Ludmila Flores¹, Yong Zhang¹, Antonio Sacco¹, Patricia Maiso¹, Abdel Kareem Azab¹, Feda Azab¹, Ruben Carrasco¹, Barrett J. Rollins¹, Aldo M. Roccaro¹, Irene M. Ghobrial^{1*}

1 Department of Medical Oncology, Dana-Farber Cancer Institute, Harvard Medical School, Boston, Massachusetts, United States of America, **2** Department of Medicine, Division of Haematology, Mayo Clinic College of Medicine, Scottsdale, Arizona, United States of America

Abstract

Glucocorticoid-induced TNF receptor (GITR) plays a crucial role in modulating immune response and inflammation, however the role of GITR in human cancers is poorly understood. In this study, we demonstrated that GITR is inactivated during tumor progression in Multiple Myeloma (MM) through promoter CpG island methylation, mediating gene silencing in primary MM plasma cells and MM cell lines. Restoration of GITR expression in GITR deficient MM cells led to inhibition of MM proliferation *in vitro* and *in vivo* and induction of apoptosis. These findings were supported by the presence of induction of p21 and PUMA, two direct downstream targets of p53, together with modulation of NF- κ B in GITR-overexpressing MM cells. Moreover, the unbalanced expression of GITR in clonal plasma cells correlated with MM disease progression, poor prognosis and survival. These findings provide novel insights into the pivotal role of GITR in MM pathogenesis and disease progression.

Citation: Liu Y, Quang P, Braggio E, Ngo H, Badalian-Very G, et al. (2013) Novel Tumor Suppressor Function of Glucocorticoid-Induced TNF Receptor GITR in Multiple Myeloma. PLoS ONE 8(6): e66982. doi:10.1371/journal.pone.0066982

Editor: Zhengqi Wang, Emory University, United States of America

Received: March 13, 2013; **Accepted:** May 13, 2013; **Published:** June 13, 2013

Copyright: © 2013 Liu et al. This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

Funding: This work was supported by NIH R01CA125690 and R01CA133799

Competing interests: There is no competing interest related to this article.

* E-mail: irene_ghobrial@dfci.harvard.edu

Introduction

Tumor Necrosis Factor receptor superfamily members (TNFRSFs) play an important role in immune responses and inflammatory reactions [1–5]. It has been recently shown that TNFRSFs are also associated with the pathogenesis of Multiple Myeloma (MM) [6,7]. For example, CD40 (*TNFSF5*) mediates MM cell survival and proliferation, as well as migration through the NF- κ B pathway [8]. Other members of the family such as APRIL (*TNFRSF13B*) and BAFF (*TNFRSF17*) were shown to be involved in the protection of MM cells from apoptosis via NF- κ B activation [9], while loss-of-function mutations of Fas antigen (*TNFRSF6*) could inhibit Fas ligand induced apoptosis in MM cells [10]. These studies suggest that TNFRSFs could play multiple roles in the pathogenesis of MM. Glucocorticoid-induced TNFR-related gene (GITR) is a member of the TNFR super family (*TNFRSF18*) that is considered a key regulator in a multitude of immune functions. GITR is

expressed and further upregulated on most immune cell types like regulatory T cells (T-regs), naïve T cells and natural killer cells (NKs) [11]. GITR plays a pivotal role in inflammation processes and autoimmune diseases. It is triggered by its ligand (GITRL), mainly expressed in antigen-presenting cells and endothelial cells [12]. Conversely, GITR engagement in NK cells induces an inhibitory effect, even though a separate study provides opposite results, demonstrating that co-stimulation by GITR/GITRL interaction is found either to activate or to inhibit NK cells [13,14]. GITR expression in tumor infiltrating lymphocytes has been found to be associated with cancer progression in patients suffering from esophageal adenocarcinomas. However, the role of GITR as a direct regulator of tumor progression in MM has not been previously described.

Multistep tumor progression occurs as a succession of clonal expansions, which is triggered by acquisition of an enabling mutant genotype. Clonal expansion may occur due to acquired mutations or epigenetic

alterations such as DNA methylation and histone modifications affecting the regulation of gene expression. In general, cancers are characterized by a global DNA hypomethylation and locus-specific hypermethylation of tumor suppressor genes.

In this study, we demonstrated that GITR is inactivated during tumor progression in MM through promoter CpG island methylation, mediating gene silencing in primary MM plasma cells and MM cell lines. Restoration of GITR expression in GITR deficient MM cells led to inhibition of MM proliferation *in vitro* and *in vivo* and induction of apoptosis. Notably, GITR/GITRL interaction increased the level of p53-regulated genes, such as *CDKN1A* (p21) and *BBC3* (PUMA) in a ligand dependent manner. Mechanistically, we demonstrated that GITR negatively regulates the NF- κ B signaling pathway in MM cells leading to apoptosis in response to TNF- α . These findings imply that GITR acts as a potential tumor suppressor gene in MM, and its epigenetic silencing facilitates NF- κ B activation and tumor proliferation in MM.

Materials and Methods

Ethics statement

Bone marrow samples from patients with MM were obtained under Dana-Farber Cancer Institute IRB approval with written informed consent and according to the declaration of Helsinki. In *in vivo* studies, mice were treated, monitored, and sacrificed in accordance with approved protocol of the Dana-Farber Cancer Institute Animal Care and Use Committee.

Cultured cell lines and primary tumor samples

Five human myeloma cell lines were used: MM1.S, U266, RPM1 (ATCC, Manassas, VA); OPM1 and INA6 (kind gift of Dr. K. Anderson, Dana-Farber Cancer Institute, Boston, MA) (15). Cells were cultured in RPMI1640 medium with 10% FBS. The umbilical vein endothelial HUVEC cell line (Cambrex, Walkersville, MD) was cultured in EGM-2 MV media (Cambrex) reconstituted according to the manufacturer. Plasma cells from patients with multiple myeloma were obtained using anti-human CD138 microbead selection (Miltenyi Biotec, Auburn, CA). GFP+/GITR+ MM.1S cells were generated using lentivirus based transfection approach.

Methylated DNA immunoprecipitation (MeDIP) assay

25 mg of gDNA sample were diluted in TE buffer (10 mM Tris-HCl, pH 7.5, 1 mM EDTA) and sheared to between 300 to 800 bp. 4 μ g DNA of each sample was saved as input and the rest heated to 95°C for 10 min and immediately placed on ice. Immunoprecipitation was performed using 10 μ g anti-5MeCyt monoclonal antibody for (Eurogentec, Bi-MECY-0100) sheared gDNA

in IP buffer (20 mM Na-Phosphate, pH 7.0, 1 M NaCl, 2% Triton-X100). MeDIP was purified and precipitated using phenol and chloroform: isoamyl alcohol. The extent of methylated DNA enrichment in our MeDIP samples was verified by qPCR on an Applied Biosystems 7500 Real-Time PCR System using the following primers:

TNFRSF18F: 5'-TTCAAGAGCCCCACAGCCAGTTG-3'
 TNFRSF18R: 5'-ATTCTCAGGTCATGAACGGTCCC-3'.
 TNFRSF11AF: 5'-GAGTTTTGGGGAGCGGAGGC-3'
 TNFRSF11AR: 5'-GTTATCTGCCGCAAGTGAGG-3'
 TNFRSF9F: 5'-CATTGAGCAGCTGGAGAACCCTC-3'
 TNFRSF9R: 5'-CGTCCGCATCTGTCCGCATC-3'
 TNFRSF19F: 5'-CACTCATCCGCTTCAGTAGCC-3'
 TNFRSF19R: 5'-GCACACCGTGCACCTCCAG-3'
 TNFRSF21F: 5'-GAGAGGTCCCCATGGCTGAAC-3'
 TNFRSF21R: 5'-GTGCTGAGCGCCCTAGAGC-3'
 TNFRSF8F: 5'-GCTGGACTGATTCATTCATTC-3'
 TNFRSF8R: 5'-CGGACGACCAACCTCCCATG-3'
 TNFRSF11BF: 5'-CTTACCACGAGCGCGCAGC-3'
 TNFRSF11BR: 5'-CTGATCAAAGGCAGGCGATAC-3'

The relative changes in the extent of promoter methylation were determined by measuring the amount of promoter in immunoprecipitated DNA after normalization to the input DNA $\%(\text{MeDNA-IP}/\text{Input}) = 2^{-(\text{Ct}(\text{input})-\text{Ct}(\text{MeDNA-IP})) \times 100}$.

Bisulfite sequencing and methylation specific PCR. Genomic DNA was prepared and bisulfite converted by EpiTect Plus DNA Bisulfite Kit (Qiagen, Valencia, CA). PCR products were purified and cloned into PGEM-T-easy vector (Promega, San Luis Obispo, CA), then sent for DNA automatic sequencing. Primer sequences for GITR are Forward 5'-GGGTTATTGGTTGGAGTTGGTAT-3', reverse 5'-CCTATCCCCATTCTATAAAATTCC-3'. Primer sequences for MSP are

MF: 5'-GCGGCAGCAGCGCGTCGTGTGAACCCG-3', MR: 5'-CGCGTTTTCGAACCCTATACGA-3' UF: 5'-GTGGTAGTAGTGTGTGTGTGAACCTG-3' UR: 5'-CACATTTTCAGGCCCTGTGCAG-3'

Bone marrow Tissue Immunohistochemistry. Bone marrow tissues were obtained from both healthy individuals and MM patients. Samples were embedded in paraffin-decalcified in Rapid-Cal-Immuno (BBC, Stanwood, WA) for 1 hour, washed, and then paraffin embedded. For GITR staining, 3- to 4- μ m tissue sections were mounted on plus slides, dried for 2 hours in a 60°C oven, and then stained using GITR polyclonal antibody (R&D system, Minneapolis, MN, cat# AF689; 1:200 dilution) according to established protocols in a BenchMark automated immunostainer (Ventana Medical Systems, Tucson, AZ). Heat-induced epitope retrieval was performed with the EDTA-buffered cell conditioning (CC1) retrieval solution. Immunostaining was completed with the iView diaminobenzidine tetrahydrochloride (DAB) detection kit (Ventana). According to the description of GITR antibody,

immersion fixed paraffin-embedded sections of human tonsil was subjected to G1TR antibody and considered as positive control.

Lentiviral infection. PCDH-ef1-CMV-EGFP lentiviral vector was purchased from SBI Company. cDNA clone of G1TR was purchased from Origene company. cDNA of G1TR was amplified and subcloned into PCDH-ef1-CMV-EGFP using BamHI and XbaI restriction enzyme sites. The PCDH-empty control and PCDH-G1TR plasmid were co-transfected with pPACK™ packaging plasmid mix (SBI, LV050A-1) into 293T cells, which is available in the frozen stock of the lab. Viral supernatant was obtained after 72 hrs post-transfection. MM1.S and OPM1 cell lines were infected by PCDH-control and PCDH-G1TR recombinant lentivirus for 72 hours. Both GFP positive and GFP negative MM1.S and OPM1 were sorted by FACS.

Competition assay and cell proliferation assay. The GFP competition assay was used to assess the ability of proliferation of two days after infection. MM cells were split into replicate wells of 500,000 cells in 12-well plates. After 24-h treatments with a range of drug doses, the GFP-positive percentage was quantified in the surviving cell population by using a BD Biosciences LSRII flow cytometry. GFP+ and GFP-cells were sorted and combined as 1:1 ratio. The percentage of GFP+ cells was examined everyday by flow cytometry. Proliferation rate and cytotoxicity were measured by DNA synthesis using BrdU (R&D) assay and by 3-(4,5-dimethylthiazol-2-yl)-2,5-diphenyltetrazolium bromide (MTT; Chemicon International, Temecula, CA) dye absorbance, respectively.

Cell cycle analysis. For cell cycle analysis by flow cytometry, cells were seeded in a 10 cm dish at 60% confluency. The cells were concomitantly treated with RNase A (Sigma Aldrich, San Luis, MO) and stained with 10ng/ml propidium iodide. Cell cycle status was determined using a FACS caliber flow cytometer (Becton Dickinson, Oxford, UK) and analyzed using FlowJo7.6.5 software.

Immunoblotting. Whole-cell lysates were subjected to sodium dodecyl sulfate-polyacrylamide gel electrophoresis (SDS-PAGE) and transferred to polyvinylidene fluoride (PVDF) membrane (Bio-Rad Laboratories, Hercules, CA). The antibodies used for immunoblotting included anti-caspase3, p-Ikk- β , Ikk- β , IkkB, P65, P50, (Cell Signaling Technology, Danvers, MA), -p53 and actin (Santa Cruz Biotechnology, Santa Cruz, CA). Nuclear extracts of the cells were prepared using the Nuclear Extraction Kit (Panomics, Redwood City, CA) and subjected to immunoblotting with anti-p65, -p50/p105, (Cell Signaling Technology, CA), and anti-nucleolin (Santa Cruz Biotechnology, CA) antibodies. In addition, the G1TR antibody (R&D Cell Biology, CA) was used for immunoblotting and immunohistochemistry on tissue microarray sections.

Active Motif for NF- κ B activity. NF- κ B activity was investigated using the Active Motif TransAM NF- κ B Family Kit, a DNA-binding enzyme-linked immunosorbent assay (ELISA)-based assay (Active Motif, North America, Carlsbad, CA). Briefly, MM1.S cells (empty control MM1.S and G1TR-transfected MM1.S) were cultured in presence or absence of TNF- α (10ng/ml) for 20 minutes. NF κ B-p65, -p50 transcription factors binding to the related consensus sequence on the plate-bound oligonucleotide were studied from nuclear extracts, following the manufacturer's procedure.

In vivo tumor progression of MM cells. Seven SCID mice for each group were injected with 5×10^6 MM1.S cells. After 4 weeks of tail vein injection, the bone marrow from the femur and skull, as well as peripheral blood were isolated from the two groups of mice. Mice bone marrow tissues were crushed and filtered into single cell suspension. Anti-human CD138 monoclonal antibody (BD bioscience, San Jose, CA) as used to examine MM cells growth by both flow cytometry and immunofluorescence.

Immunofluorescence. Immunocytochemical analysis was performed using an epifluorescence microscope (Nikon Eclipse E800; Nikon, Avon, MA) and a Photometrics, Cool snap CF color camera (Nikon, Lewisville, TX). Briefly cells were fixed with 4% paraformaldehyde in phosphate-buffered saline and permeabilized with 0.2% Triton X-100. Cells were blocked in 1X PBS / 5% normal goat serum PBS and incubated with primary antibody at 4°C overnight. Cells were incubated with Alexa Fluor 488 goat anti-rabbit secondary antibody for 2hrs.

cDNA expression profiling. Total RNA of control_MM1.S and G1TR_MM1.S was isolated using an RNeasy kit (QIAGEN, Valencia, CA). Quality control of RNA was done using RNA6000 Nano assay on the Agilent 2100 Bioanalyzer (Santa Clara, CA). Hybridization to the Human U133A 2.0 microarray was performed in Dana-Farber Cancer Institute core facility. In the experiment comparing the overexpressing G1TR versus empty control, the list of filtered probe-set was analyzed looking for differentially expressed genes between experiments (>2-fold change).

Analysis of expression profiling of available datasets. G1TR expression was analyzed in publicly available datasets comprising bone marrow plasma cells from healthy donors (NPC) and different stages of the MM disease. NPC, MG, US, SMM data was obtained from GEO dataset GSE5900 and MM data from GSE2658 and the MMRC portal (data available at www.broadinstitute.org/mmgp/home). All the intensity files were MAS5 transformed and the data was normalized to the median. Comparisons between two groups were performed using t-test (2 tailed) and for more than 2 groups using ANOVA. The comparison of G1TR expression level between molecular MM subgroups was performed using the two most relevant

molecular classifications in MM: the TC and the UAMS classifications (16,17). Results were considered significant when $p < 0.05$.

Statistical Analysis. Statistical significance of differences in the different groups was determined using Student's *t*-test. The minimal level of significance was $p < 0.05$. Experiments were repeated in triplicates. Error bars reported in the figures represent standard deviations.

Results

GITR is aberrantly methylated in both primary MM cells and MM cell lines

We hypothesized that deregulation of TNFR super family members may play a pivotal role in modulating MM pathogenesis. We therefore evaluated the expression of TNFRSFs in primary MM cells by analyzing GEO dataset GSE2658 and found that 7 members of the TNFRSF family including *TNFRSF18* (GITR), *TNFRSF11A*, *TNFRSF9*, *TNFRSF19*, *TNFRSF8*, *TNFRSF21* and *TNFRSF11B* exhibited lower expression levels in MM cells compared to their normal cellular counterpart (Figure S1a). We next investigated the mechanisms responsible for down-regulation of TNFRSFs members; and profiled DNA methylation status of the promoter CpG islands (CGI) of the related genes using MeDIP assay in 5 MM cell lines. Primers were designed within 500bp region around the CpG islands at the promoter region (Figure 1a). MM cells presented with significant methylation in the promoter CGI of GITR (*TNFRSF18*), with specific higher methylation in OPM1, MM.1S and U266 cells, compared to RPMI.8226 and INA6, where lower methylation levels were documented (Figure 1b). We further confirmed that hypermethylation of GITR promoter occurred in MM cells, by performing MeDIP assay using MM1.S and OPM1 cell lines treated with the demethylating agent, 5' azacytidine. Untreated cells were used as control, (Figure S2a).

To further evaluate the methylation status of GITR in MM cells, we performed methylation-specific PCR (MSP) for GITR in MM cell line and primary MM bone marrow CD138+ plasma cells (Figure S2b). The MSP result was further validated by bisulfate conversion. All the identified methylated CpG sites were represented as dark circles, whereas unmethylated sites in that same region were represented as blank circles across the entire CpG island (Figure 1.c). Further confirmation was performed by sanger sequencing (Figure S2.c). Based on the results of MSP and bisulfate sequencing, we found that primary clonal plasma cells presented with promoter GITR methylation in 3 out of the 5 MM patients studied, compared to the related normal cellular counterpart, where an unmethylated status was observed. Notably, primary MM cells and MM cell lines presented with a similar methylation pattern: specifically, the hypermethylation was mainly

demonstrated within the first 130 base pairs of CpG island located at the promoter region. In addition, we found that expression of GITR negatively correlated with the promoter CGI methylation status, indicating that loss of GITR expression correlates with aberrant DNA methylation in MM cells (Figure 1e). These observations were further confirmed by exposing MM cells to 5' azacytidine treatment that led to increase of GITR levels, both in MM.1S and OPM1 cells (Figure 1d), while no relatively significant increase of GITR expression was obtained by performing qPCR in INA6, U266 and RPMI8226 cell lines, which were characterized by lower methylation pattern of GITR promoter (Figure S1d). Furthermore, these data were also confirmed at protein level; GITR expression significantly increased upon 5' azacytidine treatment in a dose dependent manner, as detected by flow cytometry (Figure S3b). These findings suggest that deregulation of GITR may result from an aberrant promoter CGI methylation in MM cells.

The expression of GITR was next evaluated using immunohistochemistry in bone marrow specimens of both MM patients and healthy individuals. 82% of the normal bone marrows evaluated showed GITR positivity, compared to MM specimens that were GITR positive only in 33% of the cases (Figure 1e).

Deregulation of GITR correlates with MM progression, prognosis and survival

We next evaluated the GITR mRNA expression levels in MM patients-derived CDC138 cells, using publically available gene expression data sets (GSE5900; GSE2658); and found a significant reduction of GITR expression in MM samples, compared to healthy individuals, and interestingly, we found a sustained reduced expression of GITR with disease progression, as shown by comparing MG, US to smoldering MM to active symptomatic MM patients (Figure 2a; $P < 0.001$). Further expression analysis was performed in the subclasses of MM patients, showing that the lowest expression of GITR was observed in PR group, which has an advanced and more proliferative form of myeloma (Fig. 2bP < 0.01). We next evaluated the expression of GITR according to prognosis and survival; and found that lower expression of GITR correlated with poor prognosis and survival in MM patients, further suggesting the functional role of GITR in modulating MM pathogenesis and disease progression (Figure 2c, $P < 0.001$; Figure 2d, $P 0.03$).

Functional role of GITR in MM cells

We next dissected the functional relevance of GITR expression in modulating MM cell behavior, both *in vitro* and *in vivo*. We performed GFP competition assay by introducing lentivirus-based vector with GFP labeling into MM1.S and OPM1 cell lines. Both lenti-CMV-GFP and lenti-GITR-GFP were selected by flow cytometry and combined with GFP negative MM cells in 1:1 ratio.




Figure 1. Promoter CpG island Hypermethylation leads to repression of GTR in MM cells. A) Primer location for Medip, MSP and bisulfite sequencing assay. Bar arrow represents 1 kb region on the genome. TSS represents transcription starting site. B) Analysis of DNA methylation status of 7 TNFRSF members promoter CpG islands in 5 MM cell lines. Methylated DNA immunoprecipitation assay was performed to profile promoter methylation status. Promoter of housekeeping gene GAPDH was used as unmethylated control. Fold enrichment was calculated as described in the method and normalized to GAPDH. C) Bisulfite sequencing at the CpG island of GTR promoter, with circles denoting the methylation status of each selected clone. Black and white circles are methylated CG dinucleotides, and non-methylated CpG dinucleotides and TG dinucleotides, respectively. D) Re-expression of GTR mRNA level was examined by real-time PCR in the presence of 5' azacytidine (5 μM) for 4 days. Total RNA was extracted and reverse transcribed with oligo_dT. The mRNA level of GTR was normalized to 18s. Mean±SD. E) Expression of GTR was determined in 5 MM cell lines by real-time PCR. The mRNA level was normalized to 18s. Mean±SD. DNA methylation status of GTR in 5 MM cell lines was calculated by counting the methylated CG sites in the promoter of GTR and divided by total CG number. F) Expression of GTR examined in 9 primary MM and 11 normal bone marrow by tissue array using anti-human GTR monoclonal antibody. 4 slides of each group were shown. GTR expression was represented as brown color. Staining of tonsil for GTR staining was considered as positive control.

doi: 10.1371/journal.pone.0066982.g001




Figure 2. Expression of GTR correlates with MM clinical pathogenesis. A) Expression of GTR was evaluated by Gene expression profiling analysis (GEO 2658 datasets). The average of GTR expression in MG, US and smoldering MM was labeled as red bar. B) Expression of GTR was evaluated by Gene expression profiling analysis (GEO 2658 datasets). The average of GTR expression in every MM subtype was labeled as red bar. >C) Patients belonging to Total Therapy 2 (TT2) and Total Therapy 3 (TT3) trials with available gene expression data were divided in poor and good prognosis groups based on the 70-gene index generated by the University of Arkansas for Medical Sciences (UAMS). The cutoffs for the poor prognosis groups were 0.66 and 0.79 for TT2 and TT3, respectively. Median normalized TNFRSF18 expression level was compared between good and poor prognosis groups. D) Kaplan-Meier curve showing survival of 414 MM patients in GTR high and GTR low groups. The two groups were divided by the median of GTR expression in GEP datasets (GSE2658 and GSE5900) and analyzed by graphpad software (P=0.03).

doi: 10.1371/journal.pone.0066982.g002

Expression of GFP was measured by flow cytometry (Figure S4a). We found that the GTR+/GFP+ cells exhibited a competitive disadvantage compared to

GTR-/GFP+ cells in MM1.S, OPM1 cell lines, indicating that expression of GTR could lead to increased cytotoxicity of MM cells (Figure 3a). Cell proliferation

was next evaluated in GITR expressing MM1.S cells (GITR+), compared to control cells (GITR-); we showed that inhibition of proliferation was observed in GITR-overexpressing cells (Figure 3b). We next evaluated how GITR loss of function would affect proliferation of MM cells that present with higher GITR levels, such as RPMI8226 cells; and demonstrated that reduced expression of GITR in RPMI8226 cells led to increased cell proliferation (Figure 3c). GITR knockdown efficiency in RPMI8226 cells has been determined by real-time PCR (Figure S4b). Due to the importance of neo-angiogenesis to support MM disease progression and due to the presence of GITR specific ligand (GITRL) [15] on the surface of endothelial cells (HUVECs), we next investigated the role of GITR in MM cells in the presence of endothelial cells. *In vitro* co-culture of GITR-overexpressing MM1.S with HUVECs led to significant inhibition of cell growth, thus providing a further evidence of the tumor suppressor role of GITR in MM (Figure S4c).

We next validated our *in vitro* findings by using an *in vivo* xenograft MM mouse model. GITR expression was enhanced in MM1.S cells by using lentiviral vector. 5×10^6 GITR+ MM1.S cells and GITR-empty vector infected control MM1.S cell were injected into SCID-bg mice via tail vein respectively. After 4 weeks, bone marrow tissues were extracted from both control and experimental mice groups. We observed significant reduction of growth of MM cells in mice that received GITR-overexpressing cells, compared to control groups, suggesting that GITR is a critical regulator of tumor growth in MM cells both *in vitro* and *in vivo* (Figure 3d-3e). Taken together, these studies suggest that GITR acts as a potential tumor suppressor in MM.

GITR induces expression of p21 and PUMA in a ligand dependent manner

To understand the molecular basis of the role of GITR as a potential tumor suppressor in MM, we analyzed the specific gene expression signature in GITR overexpressing cells compared to cells transfected with empty vector. Gene annotation analysis showed that the p53 signaling pathway represents the most significantly enriched gene set, ($P < 0.001$), together with up-regulation of MDM2, CDKN1a (p21), E2F1, Fas, IGF-BP3 and CD82 genes (Table. S1). Meanwhile, apoptosis response genes including Tumor Necrosis Factor Receptor Type 1-Associated Death Domain protein (TRADD) and Death Associated Protein (DAP) were also upregulated in GITR-overexpressing MM cells. Expression of these genes was validated in both GITR+ MM1.S and GITR knockdown RPMI8226 cells by real time PCR; showing overexpression of GITR could increase the level of p21, Fas, TRADD and DAP, in contrast knockdown of GITR could lead to decreased mRNA level of these genes (Figure S5a). Based on these results, we examined the expression of p53 in MM cells in response to stimulation with GITRL using

qRT-PCR and western blot. We found that expression of GITR could lead to both increased mRNA and protein level of p53 in 12hrs after stimulation with 10ng/ml GITRL (Figure 4a). The elevated p53 protein could subsequently lead to the induction of p21, which is another downstream target of p53 and a regulator of mitochondrial-mediated apoptosis (Figure 4b, and Figure S5b). Next, we evaluated whether elevated levels of p21 and puma proteins could affect MM cell cycle progression or apoptosis. We demonstrated that GITR expressing cells presented with increased numbers of cells in subG1/G1 phase, together with S and G2 phase inhibition (Figure 4c). Induction of apoptosis by GITR expression was also confirmed using flow cytometry with PI/Annexin V dual staining combined with immunoblotting for caspase cleavage in the presence of GITRL (Figure 4d-e). Our results indicate that GITR could modulate p53-mediated p21 and puma expression, thus mediating cell cycle progression and apoptosis in MM cells.

GITR negatively regulates NF- κ B pathway in MM cells

NF- κ B transcription factors play a key role in the survival and proliferation of several hematologic malignancies, including MM [16]. It has been shown that GITR may negatively regulate NK cell activation through suppression of the NF- κ B pathway [13]. We therefore dissected the role of GITR in modulating the canonical NF- κ B pathway in both GITR- and GITR+ cell lines. Cells were exposed to TNF- α (10ng/mL) and nuclear DNA binding activity and p50/p65 units was studied by using DNA binding ELISA assay and immunoblot at 0, 15, 30 and 60 minutes. We showed that TNF- α was able to activate NF- κ B pathway in both control and GITR expressing groups at 15 min (Figure 5a). However, NF- κ B activation was abolished at early 30 to 60 min in GITR+ cells after the stimulation, whereas the NF- κ B signal remained elevated up to 60 min in the GITR-cells, suggesting a role of GITR in terminating TNF- α -induced NF- κ B activation (Figure 5a). In addition, we also demonstrated inhibition of TNF- α -dependent p50 nuclear translocation in GITR+ cells, as shown by immunofluorescence (Figure 5b). Previous studies showed that IKK- β represents a critical upstream regulator of P65/p50 activation [17]; we therefore examined the phosphorylation of IKK- β stimulated by TNF- α by immunoblotting. Consistently, the phosphorylation of IKK- β was markedly diminished by the increased expression of GITR at early time points. IKK- α -degradation was also measured in TNF- α stimulated control and GITR overexpressing cells. We found that IKK- α protein decreased back to the basal level at 30 to 60 min time points in GITR expressing MM1.S cells, suggesting that TNF- α stimulated NF- κ B activation is IKK- α -dependent (Figure 5c).

To further confirm that GITR induces inhibition of TNF- α -induced NF- κ B signaling, we evaluated the




Figure 3. Effect of G1TR on MM tumor proliferation in vitro and in vivo. A) MM cells (MM.1S and OPM1) were transfected with PCDH empty and PCDH-G1TR with GFP labeled lentiviral vector (Cat# CD511B-1, SBI Inc.) respectively. Cell proliferation has been evaluated by using GFP competition assay. Expression of GFP and G1TR has been examined by flow cytometry by using anti-human G1TR PE labeled primary antibody. GFP+ and GFP-MM cell were sorted by BD laser II flow machine. Ratio of GFP+/GFP- was recorded everyday after combination. B) Effect of G1TR-overexpression on MM1.S cell line proliferation. 96 wells plate coated with MM1.S cells were read after 48 and 72 hours respectively. Anti-BrdU-pod was used to detect the absorbance. Mean±SD. *, P<0.05. C) Effect G1TR-knockdown on RPMI.8226 cell line proliferation (RPMI.8226). 96 wells plate coated with MM1.S cells were read after 48 and 72 hours respectively. Anti-BrdU-pod was used to detect the absorbance. Mean±SD. *, P<0.05. D) SCID mice were injected i.v. with 5 million MM1.S cells, transfected with either empty vector (control C1, C2) or G1TR (G1TR+, G1, G2). In vivo tumor progression has been evaluated by using immunofluorescence staining with anti-human CD138 monoclonal antibody after 4 weeks injection, on bone marrow femurs. E) Detection of MM cells from tissues of mice injected with either empty vector (control) or G1TR (G1TR+). MM cells have been detected by using flow cytometry analysis for CD138. Mean±SD. *, P<0.05; **, P<0.01.

doi: 10.1371/journal.pone.0066982.g003




Figure 4. GITR induces up-regulation of p21 and PUMA in MM cells. A) Total protein and RNA was extracted in 0, 3, 6, 12, 24 and 48 hours after treatment with GITRL (10ng/mL) from GITR- and GITR+ MM1.S cells. Both Protein and mRNA level of p53 have been evaluated by western blot and qRT-PCR. Actin was considered as the protein loading control. CT value of qRT-PCR was normalization to 18s. Mean \pm SD. B) GITR- and GITR+ cells were exposed to GITR-L (0-5-10ng/mL) for 24 hrs. whole cellular protein was extracted and subjected to Western blot using anti-p21, -PUMA and -actin antibodies. C) GITR- and GITR+ MM1.S cells were exposed to GITR-L (5-10 ng/mL) for 24 hours respectively. Cell cycle has been evaluated by using PI staining and flow cytometry analysis. D) GITR- and GITR+ MM1.S cells were exposed to GITR-L (5-10ng/mL) for 24 hours. Whole protein lysates have been subjected to Western blot using anti-caspase-3, 9 and -tubulin antibodies. E) GITR- and GITR+ MM1.S cells were exposed to GITR-L (5-10ng/mL) for 24 hours. Cells were stained by PI/Annexin V to evaluate apoptosis effect using flow cytometry. Mean \pm SD.

doi: 10.1371/journal.pone.0066982.g004




Figure 5. Effect of G1TR on canonical NF- κ B pathway. A) Effect of G1TR on TNF- α induced NF- κ B activation. NF- κ B activity has been evaluated by using DNA binding ELISA assay in G1TR- and G1TR+ MM1.S cells. Cells were exposed to TNF- α (10ng/mL) for 15-30-60 minutes. NF- κ B p65 and p50 transcription factor binding to its consensus sequence on the plate-bound oligo nucleotide was examined from nuclear extracts. Data represent mean plus or minus SD of triplicate experiments. Nuclear protein lysates were also subjected to Western blot using anti-p50, -p65 and -nucleolin antibodies. B) G1TR- and G1TR+ were harvested at 24 hours after treatment with and without TNF- α (10 ng/mL) for 60 minutes; Immunocytochemical analysis was assessed using anti-phospho-NF- κ B-p50 antibody, with DAPI used to stain nuclei. C) G1TR- and G1TR+ cells were exposed to TNF- α (10ng/mL) for 15-30-60 minutes. Whole cellular protein lysates have been subjected to Western blot using anti-p65, -phospho(p)-I κ B beta, -I κ B beta, -I κ B alpha, and -actin antibodies. D) G1TR- and G1TR+ cells were exposed to TNF- α (2.5-5-10ng/mL) for 16 hours. Whole cellular protein lysates have been subjected to Western blot using anti-BCL-2, -survivin and -actin antibodies. E) G1TR- and G1TR+ cells were exposed to TNF- α (2.5-10ng/mL) for 12 hours. Apoptosis was evaluated by Annexin/PI staining and flow cytometry analysis. Mean \pm SD.

doi: 10.1371/journal.pone.0066982.g005

expression p65/p50-downstream targets, such as BCL-2 and survivin. As demonstrated at protein level, BCL2 and survivin expression was upregulated in GITR-MM cells, upon stimulation with TNF- α . In contrast, no significant changes of BCL-2 and survivin were observed in GITR overexpressing cells in response to TNF- α (Figure 5d).

Previous studies showed that TNF- α alone may be responsible for NF- κ B nuclear translocation, cIAP-1 and cIAP-2 up-regulation, thus leading to increase in MM cell proliferation [18]. To explore the effect of GITR on apoptosis induced by TNF- α in MM cells, we performed PI/Annexin dual staining assay. We found that GITR was able to counteract NF- κ B mediated anti-apoptosis signals and facilitate the apoptosis induced by TNF- α in GITR expressing MM cells (Figure 5e). Taken together, these findings further support our hypothesis that GITR negatively regulates canonical NF- κ B pathway activation by inhibiting phosphorylation of IKK- β in MM cells at early time points.

Discussion

MM is a plasma cell neoplasia characterized by activation of the NF- κ B pathway through cell autonomous mutations or external signaling from the tumor microenvironment [19,20]. In this study, we demonstrated that epigenetic silencing of *GITR* is involved in MM cell proliferation, is associated with an anti-apoptosis effect and is contributing to NF- κ B-mediated MM pathogenesis. Notably, using bisulfite sequencing, we observed that all of the methylated cytosines are distributed at the first 130 base pair of promoter CGI region in both MM cell lines and primary bone marrow CD138+ MM cells, suggesting that the aberrant methylation of *GITR* promoter in MM cells is not a random epigenetic event. To link the function of *GITR* to MM cell behavior, we sought to investigate if the expression of *GITR* correlates with MM cells proliferation. Indeed, we demonstrated that expression of *GITR* could inhibit MM cells proliferation *in vitro* and *in vivo*. These results indicate that *GITR* acts as a potential tumor suppressor gene on MM cells and we confirmed that its re-expression induces apoptosis in MM cells. These results are consistent with previous reports indicating that deletion of *GITR* could lead to increased proliferation in T lymphocytes [21].

GITR is located on chromosome 1p36. Deletion of 1p36 is associated with increased risk for neoplasia, including neuroblastoma, prostate cancer, lung cancer, melanoma and non-Hodgkin lymphomas [22–27]. It has been previously shown that heterozygous deletions at 1p36 were also found in 8% of MM patients, indicating that this region may contain putative tumor suppressor genes [28]. We analyzed published aCGH data from 238 MM patients and 60 human myeloma cell lines and found that about 8% MM patients have monoallelic deletion of 1p, which is consistent with the previous

reports. The DNA methylation-mediated loss of *GITR* in MM may also provide evidence of alternative contribution of 1p36 abnormality to MM tumorigenesis.

Recent studies in Chronic Lymphocytic Leukemia (CLL) have shown pronounced expression of *GITRL* and the *GITR* receptor was expressed at significantly higher levels on NK cells of CLL patients compared with healthy controls. In addition, *GITRL* contributed to resistance of CLL cells to rituximab therapy, indicating that *GITR/GITRL* contributes to disease progression and resistance to Rituximab-induced NK reactivity in CLL [29]. In another study, constitutive expression of *GITRL* by tumor cells diminished NK cell antitumor immunity [30]. Moreover, in *GITR*^{-/-} mouse models, the lack of expression of *GITR* was shown to modestly contribute to mature B cell homeostasis [31]. Our study demonstrated that the tumor suppressor properties of *GITR* are characterized by inhibition of proliferation and induction of apoptosis as well as induction of downstream targets of p53, such as p21, MDM2, Fas, IGF-BP3 and CD82. This data support the presence of a putative crosstalk between *GITR* and the p53 pathway. Therefore we attempted to elucidate the molecular basis that link p53 signal to the *GITR* pathway. We observed significantly increased p53 level in *GITR*⁺, leading to induction of p21 and puma in MM cells in response to *GITRL*. Further studies are needed to better define the interaction between *GITR* and the p53-related signaling cascade.

NF- κ B transcription factors play a key role in the survival and proliferation of plasma cell malignancies, including multiple myeloma [32–34]. It was shown that mutations involved in the NF- κ B pathway are present in at least 17% of MM tumors and 40% of MM cell lines [35–37]. These mutations can lead to activation of the canonical NF- κ B pathway via modulating the level of NIK, TRAFs and CD40. Therefore, targeting the NF- κ B pathway is an attractive therapy for MM [20,38]. Our data indicates that *GITR* negatively regulates the canonical NF- κ B pathway by terminating the canonical NF- κ B activation as early as 30 minutes post-stimulation with TNF- α . The inhibitory role of *GITR* in the NF- κ B pathway was further supported by inhibition of phosphorylation of IKK- β and protection of degradation of I κ B. We speculate that this effect could possibly result from either binding of TNF- α to *GITR* or blocking the intracellular cell signaling transduction of TNF- α by *GITR* (Figure 6). Further elucidation of *GITR* mediated disruption of the NF- κ B pathway will be critical for understanding the role of *GITR* in the pathogenesis of MM and the development of future therapeutic agents.

Overall, our results demonstrate that *GITR* acts as a potential tumor suppressor in MM. Furthermore, our data links *GITR* to p53-associated modulation of p21 and puma in myeloma cells. Most importantly, our findings reveal an inhibitory role of *GITR* in regulating NF- κ B pathway, providing a novel insight into the role




Figure 6. GITR inhibits NF-κB pathway. A proposed model to summarize the ability of GITR to modulate TNF-α induced canonical NF-κB pathway. The presence of GITR in MM cells could disrupt TNF-α/TNFR induced NF-κB activation by blocking phosphorylation of IKK-beta, subsequently leading to early termination of nuclear translocation of NF-κB p65/p50 and impaired induction of BCL-2 and Survivin.

doi: 10.1371/journal.pone.0066982.g006

of TNFRSFs family in MM pathogenesis and disease progression, thus supporting the therapeutic use of NF-κB inhibitors in MM patients who present with increased GITR methylation.

Supporting Information

Figure S1. Deregulation of TNFRSFs in multiple myeloma. A) Expression of TNFRSFs member was assessed by analyzing GEO dataset GSE2658. Expression value was used to calculate the p-value by T-test. The heat map was generated for all of the TNFRSFs members. Significance of differential expression of TNFRSFs members, which were selected according to the p-value(<0.05), is shown by the intensity of red (up-regulation) versus blue (down-regulation).

Figure S2. Hypermethylation of GITR in MM cells. A) Loss of methylation enrichment of GITR promoter in the presence of demethylation agent 5' azacytidine in MM cell line by meDIP assay. Fold enrichment was normalized to GAPDH. B) Methylation specific PCR was performed to evaluate the methylation status of primers MM cells from 5 patients. According to the BSP results, MM1.S were considered as positive control and ddH₂O as negative control. C) DNA sequencing was done in primary MM DNA samples after bisulfate conversion followed by subcloning into T-easy vector. C > T represents unmethylated C, whereas C>C represents methylated C. D) expression of GITR mRNA level was examined by real-time PCR in the presence of 5' azacytidine (5 μM) for 4 days. Total RNA was extracted and reverse transcribed with oligo_dT from INA6, U266 and RPMI8226 cell lines. The mRNA level of GITR was normalized to 18s. Mean ±SD.

Figure S3. Promoter DNA methylation leads to GITR silencing in MM cells. Re-expression of GITR protein level was determined by flow cytometry exposed to 5' azacytidine in a dose dependent manner. Cells were harvested after 96 hrs of incubation with 5' azacytidine and stained with anti-GITR-PE labeled antibody.

Figure S4. Expression of GITR affects MM cell proliferation. A) MM.1S cells have been transfected with either empty vector (contrl) or GITR (GITR+). Expression of GITR has been evaluated by flow cytometry. Anti-GITR-PE conjugated antibody has been used. B) Knockdown of GITR in RPMI8226 cell lines by siRNA was measured by real-time PCR. mRNA level was normalized to 18s. C) MM cells have been transfected with either empty vector (contrl) or GITR (GITR+) and cultured in presence or absence of HUVECs for 48 hours. Cell proliferation has been evaluated by BrdU assay. Mean \pm SD.

Figure S5. Overexpression of GITR induced the expression of p21 and puma in a ligand dependent way. F) MM1.S cells were transfected with both GITR overexpressing vector (GITR+) and

empty control vector (GITR-) respectively. RPMI8226 cells were transfected with si-scramble and pooled si-GITR oligo. Total RNA was extracted in 24 hours after the transfection. Expression of p21, TRADD, Fas and DAP has been evaluated by qRT-PCR, with normalization to 18s. Mean \pm SD. G) MM cells were transfected with either empty vector (contrl) or GITR (GITR+). Total RNA was extracted in 12 hours after treatment with GITRL (5-10ng/mL). Expression of p21 and puma was evaluated by qRT-PCR, with normalization to 18s. Mean \pm SD.

Table S1. Overexpression of GITR is associated with p53 pathway. Genes involved in p53 pathway are shown (cut off = 2 fold, $p < 0.05$).

Author Contributions

Conceived and designed the experiments: YL, PQ, EB, HN, RC, BR, AR, IG. Performed the experiments: YL, PQ, EB, HN, GB, LF, YZ, AS, PM, AA, FA, RC. Analyzed the data: YL, PQ, EB, AR, IG. Contributed reagents/materials/analysis tools: EB, GB, LF, RC, BR. Wrote the manuscript: YL, EB, AR, IG.

References

1. Hehlgans T, Pfeffer K (2005) The intriguing biology of the tumor necrosis factor/tumour necrosis factor receptor superfamily: players, rules and the games. *Immunology* 115: 1-20. doi:10.1111/j.1365-2567.2005.02143.x. PubMed: 15819693.
2. Cheng X, Kinoshita M, Murali R, Greene MI (2003) The TNF receptor superfamily: role in immune inflammation and bone formation. *Immunol Res* 27: 287-294. doi:10.1385/IR:27:2-3:287. PubMed: 12857975.
3. Kwon B, Kim BS, Cho HR, Park JE, Kwon BS (2003) Involvement of tumor necrosis factor receptor superfamily(TNFRSF) members in the pathogenesis of inflammatory diseases. *Exp Mol Med* 35: 8-16. doi:10.1038/emmm.2003.2. PubMed: 12642898.
4. Zhou T, Mountz JD, Kimberly RP (2002) Immunobiology of tumor necrosis factor receptor superfamily. *Immunol Res* 26: 323-336. doi:10.1385/IR:26:1-3:323. PubMed: 12403370.
5. Locksley RM, Killeen N, Lenardo MJ (2001) The TNF and TNF receptor superfamilies: integrating mammalian biology. *Cell* 104: 487-501. doi:10.1016/S0092-8674(01)00237-9. PubMed: 11239407.
6. Silke J, Brink R (2010) Regulation of TNFRSF and innate immune signalling complexes by TRAFs and cIAPs. *Cell Death Differ* 17: 35-45. doi:10.1038/cdd.2009.114. PubMed: 19680262.
7. Tamada K, Chen L (2006) Renewed interest in cancer immunotherapy with the tumor necrosis factor superfamily molecules. *Cancer Immunol Immunother* 55: 355-362. doi: 10.1007/s00262-005-0081-y. PubMed: 16187084.
8. Tai YT, Podar K, Mitsiades N, Lin B, Mitsiades C et al. (2003) CD40 induces human multiple myeloma cell migration via phosphatidylinositol 3-kinase/AKT/NF-kappa B signaling. *Blood* 101: 2762-2769. doi:10.1182/blood-2002-09-2813. PubMed: 12433678.
9. Shen X, Zhu W, Zhang X, Xu G, Ju S (2011) A role of both NF- κ B pathways in expression and transcription regulation of BAFF-R gene in multiple myeloma cells. *Mol Cell Biochem* 357: 21-30. doi:10.1007/s11010-011-0871-9. PubMed: 21607696.
10. Dimberg LY, Dimberg AI, Ivarsson K, Strömberg T, Osterborg A et al. (2005) Ectopic and IFN-induced expression of Fas overcomes resistance to Fas-mediated apoptosis in multiple myeloma cells. *Blood* 106: 1346-1354. doi:10.1182/blood-2004-04-1322. PubMed: 15860671.
11. Shimizu J, Yamazaki S, Takahashi T, Ishida Y, Sakaguchi S (2002) Stimulation of CD25(+)CD4(+) regulatory T cells through GITR breaks immunological self-tolerance. *Nat Immunol* 3: 135-142. doi:10.1038/nri759. PubMed: 11812990.
12. Vecchiarelli A, Pericolini E, Gabrielli E, Agostini M, Bistoni F et al. (2009) The GITRL-GITR system alters TLR-4 expression on DC during fungal infection. *Cell Immunol* 257: 13-22. doi: 10.1016/j.cellimm.2009.02.001. PubMed: 19272591.
13. Liu B, Li Z, Mahesh SP, Pantanelli S, Hwang FS et al. (2008) Glucocorticoid-induced tumor necrosis factor receptor negatively regulates activation of human primary natural killer (NK) cells by blocking proliferative signals and increasing NK cell apoptosis. *J Biol Chem* 283: 8202-8210. doi:10.1074/jbc.M708944200. PubMed: 18230609.
14. Hanabuchi S, Watanabe N, Wang YH, Ito T, Shaw J et al. (2006) Human plasmacytoid dendritic cells activate NK cells through glucocorticoid-induced tumor necrosis factor receptor-ligand (GITRL). *Blood* 107: 3617-3623. doi:10.1182/blood-2005-08-3419. PubMed: 16397134.
15. Tuytaerts S, Van Meirvenne S, Bonehill A, Heirman C, Corthals J et al. (2007) Expression of human GITRL on myeloid dendritic cells enhances their immunostimulatory function but does not abrogate the suppressive effect of CD4+CD25+ regulatory T cells. *J Leukoc Biol* 82: 93-105. doi:10.1189/jlb.0906568. PubMed: 17449724.
16. Demchenko YN, Kuehl WM (2010) A critical role for the NF κ B pathway in multiple myeloma. *Oncotarget* 1: 59-68. PubMed: 20890394.
17. Yang F, Tang E, Guan K, Wang CY (2003) IKK beta plays an essential role in the phosphorylation of RelA/p65 on serine 536 induced by lipopolysaccharide. *J Immunol* 170: 5630-5635. PubMed: 12759443.
18. Mitsiades N, Mitsiades CS, Poulaki V, Chauhan D, Richardson PG et al. (2002) Biologic sequelae of nuclear factor-kappaB blockade in multiple myeloma: therapeutic applications. *Blood* 99: 4079-4086. doi:10.1182/blood.V99.11.4079. PubMed: 12010810.

19. Tai YT, Li XF, Breitzkreutz I, Song W, Neri P et al. (2006) Role of B-cell-activating factor in adhesion and growth of human multiple myeloma cells in the bone marrow microenvironment. *Cancer Res* 66: 6675-6682. doi: 10.1158/0008-5472.CAN-06-0190. PubMed: 16818641.
20. Chapman MA, Lawrence MS, Keats JJ, Cibulskis K, Sougnez C et al. (2011) Initial genome sequencing and analysis of multiple myeloma. *Nature* 471: 467-472. doi:10.1038/nature09837. PubMed: 21430775.
21. Ronchetti S, Nocentini G, Riccardi C, Pandolfi PP (2002) Role of GITR in activation response of T lymphocytes. *Blood* 100: 350-352. doi:10.1182/blood-2001-12-0276. PubMed: 12070049.
22. White PS, Maris JM, Beltinger C, Sulman E, Marshall HN et al. (1995) A region of consistent deletion in neuroblastoma maps within human chromosome 1p36.2-36.3. *Proc Natl Acad Sci U S A* 92: 5520-5524. doi:10.1073/pnas.92.12.5520. PubMed: 7777541.
23. Biegel JA, White PS, Marshall HN, Fujimori M, Zackai EH et al. (1993) Constitutional 1p36 deletion in a child with neuroblastoma. *Am J Hum Genet* 52: 176-182. PubMed: 8434586.
24. Best CJ, Gillespie JW, Yi Y, Chandramouli GV, Perlmutter MA et al. (2005) Molecular alterations in primary prostate cancer after androgen ablation therapy. *Clin Cancer Res* 11: 6823-6834. doi:10.1158/1078-0432.CCR-05-0585. PubMed: 16203770.
25. Nomoto S, Haruki N, Tatematsu Y, Konishi H, Mitsudomi T et al. (2000) Frequent allelic imbalance suggests involvement of a tumor suppressor gene at 1p36 in the pathogenesis of human lung cancers. *Genes Chromosomes Cancer* 28: 342-346. doi:10.1002/1098-2264(200007)28:3. PubMed: 10862041.
26. Poetsch M, Dittberner T, Woenckhaus C (2002) Frameshift mutations of RIZ, but no point mutations in RIZ1 exons in malignant melanomas with deletions in 1p36. *Oncogene* 21: 3038-3042. doi:10.1038/sj.onc.1205457. PubMed: 12082534.
27. Katzenberger T, Kalla J, Leich E, Stöcklein H, Hartmann E et al. (2009) A distinctive subtype of t(14;18)-negative nodal follicular non-Hodgkin lymphoma characterized by a predominantly diffuse growth pattern and deletions in the chromosomal region 1p36. *Blood* 113: 1053-1061. PubMed: 18978208.
28. Schultheis B, Krämer A, Willer A, Hegenbart U, Goldschmidt H et al. (1999) Analysis of p73 and p53 gene deletions in multiple myeloma. *Leukemia* 13: 2099-2103. doi:10.1038/sj.leu.2401609. PubMed: 10602435.
29. Buechele C, Baessler T, Wirths S, Schmolh JU, Schmiedel BJ et al. (2012) Glucocorticoid-induced TNFR-related protein (GITR) ligand modulates cytokine release and NK cell reactivity in chronic lymphocytic leukemia (CLL). *Leukemia* 26: 991-1000. doi:10.1038/leu.2011.313. PubMed: 22064350.
30. Baltz KM, Krusch M, Bringmann A, Brossart P, Mayer F et al. (2007) Cancer immunoediting by GITR (glucocorticoid-induced TNF-related protein) ligand in humans: NK cell/tumor cell interactions. *FASEB J* 21: 2442-2454. doi:10.1096/fj.06-7724com. PubMed: 17360848.
31. Teodorovic LS, Riccardi C, Torres RM, Pelanda R (2012) Murine B cell development and antibody responses to model antigens are not impaired in the absence of the TNF receptor GITR. *PLOS ONE* 7: e31632. doi:10.1371/journal.pone.0031632. PubMed: 22328941.
32. Rauert H, Stühmer T, Bargou R, Wajant H, Siegmund D (2011) TNFR1 and TNFR2 regulate the extrinsic apoptotic pathway in myeloma cells by multiple mechanisms. *Cell Death Dis* 2: e194. doi:10.1038/cddis.2011.78. PubMed: 21850048.
33. Chauhan D, Tian Z, Zhou B, Kuhn D, Orłowski R et al. (2011) In vitro and in vivo selective antitumor activity of a novel orally bioavailable proteasome inhibitor MLN9708 against multiple myeloma cells. *Clin Cancer Res* 17: 5311-5321. doi: 10.1158/1078-0432.CCR-11-0476. PubMed: 21724551.
34. Berenson JR, Ma HM, Vescio R (2001) The role of nuclear factor-kappaB in the biology and treatment of multiple myeloma. *Semin Oncol* 28: 626-633. doi:10.1053/sonc.2001.29542. PubMed: 11740821.
35. Demchenko YN, Glebov OK, Zingone A, Keats JJ, Bergsagel PL et al. (2010) Classical and/or alternative NF-kappaB pathway activation in multiple myeloma. *Blood* 115: 3541-3552. doi: 10.1182/blood-2009-09-243535. PubMed: 20053756.
36. Annunziata CM, Davis RE, Demchenko Y, Bellamy W, Gabrea A et al. (2007) Frequent engagement of the classical and alternative NF-kappaB pathways by diverse genetic abnormalities in multiple myeloma. *Cancer Cell* 12: 115-130. doi:10.1016/j.ccr.2007.07.004. PubMed: 17692804.
37. Keats JJ, Fonseca R, Chesi M, Schop R, Baker A et al. (2007) Promiscuous mutations activate the noncanonical NF-kappaB pathway in multiple myeloma. *Cancer Cell* 12: 131-144. doi: 10.1016/j.ccr.2007.07.003. PubMed: 17692805.
38. Gilmore TD (2007) Multiple myeloma: lusting for NF-kappaB. *Cancer Cell* 12: 95-97. doi:10.1016/j.ccr.2007.07.010. PubMed: 17692798.