

## INSECTS, INSECTICIDES AND HORMESIS: EVIDENCE AND CONSIDERATIONS FOR STUDY

### G. Christopher Cutler □ Nova Scotia Agricultural College

□ Insects are ubiquitous, crucial components of almost all terrestrial and fresh water ecosystems. In agricultural settings they are subjected to, intentionally or unintentionally, an array of synthetic pesticides and other chemical stressors. These ecological underpinnings, the amenability of insects to laboratory and field experiments, and our strong knowledgebase in insecticide toxicology, make the insect-insecticide model an excellent one to study many questions surrounding hormesis. Moreover, there is practical importance for agriculture with evidence of pest population growth being accelerated by insecticide hormesis. Nevertheless, insects have been underutilized in studies of hormesis. Where hormesis hypotheses have been tested, results clearly demonstrate stimulatory effects on multiple taxa as measured through several biological endpoints, both at individual and population levels. However, many basic questions are outstanding given the myriad of chemicals, responses, and ecological interactions that are likely to occur.

*Keywords: insects; hormesis; hormoligosis; sublethal insecticide exposure; pest resurgence*

### INSECT EXPOSURE TO INSECTICIDES

Management of insect pests the past 70 years has been achieved mainly through application of synthetic pesticides. Since the discovery of the insecticidal properties of DDT by Paul Müller in 1939, hundreds of insecticidal compounds have been developed, accompanied by a relatively steady increase in insecticide use. Insecticide use has waned in some crops with the recent advent of transgenic varieties that express insecticidal toxins (Benbrook 2004; Cattaneo *et al.* 2006), but synthetic insecticide use remains high in most commodities. Approximately 560 million kg of insecticide were used globally in 2001, over three-quarters of which was for agricultural purposes (Kiely *et al.* 2004). The Environmental Protection Agency has approved the use of about 225 insecticidal active ingredients, and there are typically multiple formulations of each used in a variety of applications (Yu 2008).

Insect populations in agriculture and forestry are thus potentially exposed to great amounts of pesticide. Exposures may occur through direct contact (i.e. topical application of the spray), ingestion or residual contact. Unborn progeny or gametes may be affected through exposed adults. Although many individuals will be killed by these compounds, oth-

Address correspondence to G. Christopher Cutler, Department of Environmental Sciences, Nova Scotia Agricultural College, Truro, NS, Canada B2N 5E3; Tel.: 902-896-2471; Fax: 902-893-1404; Email: ccutler@nsac.ca

ers will be subject to a number of sublethal effects (Croft and Brown 1975; Haynes 1988; Stark and Banks 2003). Effects depend on a number of factors but dose is a key determinant of elicited response. In a field situation, the pesticide dose to which the insect is exposed will differ greatly over space and time. Growers attempt to apply sprays evenly to their crops, but even a small breeze can cause drift, resulting in deposition of variable amounts of solution to plants throughout a field. Volatilization of pesticides, which is particularly prominent during applications on dry, hot days, can significantly reduce the amount of product that remains at the target. Even within a plant penetration of the spray through the canopy can vary significantly, whether comparing the top vs. bottom of the plant, or the upper vs. lower surface of a leaf. The addition of time will further alter the exposure. Microbial and chemical degradation in or on soil and foliage are important processes that change the toxicity of an applied solution, and these vary with temperature, moisture, pH and adsorption. Similarly, the rate of insecticide photodegradation will vary with light intensity. While these processes usually render the insecticide less effective, in some cases metabolites of the parent compound may be more toxic to the target insect (e.g. Nauen *et al.* 1998). Systemic insecticides that are applied to soil or seeds are expected to reach concentrations in the leaves that are lethal to insects, but sublethal concentrations are present in the plant during accumulation and degradation of the toxicant. Further, concentrations of systemic insecticide can vary through a plant, as well as in old and new foliage over time (Olson *et al.* 2004). Thus, although growers try to apply pesticides evenly at concentrations intended to kill target pests, many biotic and abiotic processes will spatially and temporally change the dose of pesticide to which an insect is actually exposed in the field. Very often these will be a range of sublethal concentrations.

## **HORMESIS AND INSECT PEST MANAGEMENT**

Although the study of dose-response relationships has traditionally been guided by the threshold and/or linear non-threshold models, the hormetic dose-response model – a biphasic model characterized by low-dose stimulation and high-dose inhibition – is now widely recognized as a general, real and reproducible biological phenomenon (Calabrese 2005a; 2005b; 2010). Hormesis has been observed in a myriad of single-cell and multicellular organisms, and for many biological measures including growth, longevity, numerous metabolic and molecular processes, cognitive function and immune response (Calabrese and Baldwin 2003a; Calabrese and Baldwin 2003b; Calabrese and Blain 2005). Hormetic effects are not limited to chemical stressors, such as pesticides and heavy metals, and may manifest following mild temperature stress (Luckey 1968; Stolzing *et al.* 2006; Galbadage and Hartman 2008; Gomez

*et al.* 2009), induced radiation (Luckey 1991; Azzam *et al.* 1996; Feinendegen 2005), reduced caloric intake or exercise (Le Bourg and Rattan 2008; Mattson and Calabrese 2010). Further, the hormetic dose-response appears to be the most fundamental dose-response, outperforming the threshold and linear non-threshold models in head-to-head comparisons using numerous criteria (Calabrese and Baldwin 2003a; Calabrese and Blain 2005). Hormesis has attracted attention from a variety of disciplines, particularly in risk assessment for chemical and radiation protection (Luckey 1991; Calabrese and Cook 2005; Sanders 2010) and in the biomedical sciences (Le Bourg and Rattan 2008; Mattson and Calabrese 2010). As one might predict, evolutionary biologists have also become interested in the phenomenon, questioning whether hormesis of a trait is adaptive, resulting in increased fitness for genotypes exhibiting the response, maladaptive (i.e. associated with decreased fitness), or selectively neutral (Forbes 2000; Parsons 2001; Mattson 2009).

Insect toxicology and the study of insecticide dose-response have a rich scientific history. In the very first issue of the *Annual Review of Entomology* in 1956, no less than six papers examined topics in insecticide chemistry (Martin 1956), insecticide mode of action (Kearns 1956), insecticide resistance (Hoskins and Gordon 1956), plant-insecticide residues (Gunther and Blinn 1956), application techniques (Brann 1956) and impacts of pesticides on arthropod populations (Ripper 1956). This trend continues today with hundreds of papers published every year on these very same subjects and more in emerging related fields. However, studies in insect toxicology with pest and beneficial insects have traditionally focused mostly on high doses and lethal effects, i.e. LD<sub>50</sub>/LC<sub>50</sub> data, as have toxicology studies in other disciplines (Calabrese 2005a). Further, although the importance of low dose insecticide exposures and sublethal effects have long been realized (Ripper 1956; Croft and Brown 1975; Haynes 1988; Stark and Banks 2003), these are generally presented within the context of deleterious impacts of low doses on insect fecundity, longevity, behavior and similar endpoints. The alternative response – insecticide-induced stimulation of biological processes via hormetic mechanisms – has received far less attention.

Why is it important to study hormesis in insects? I would suggest that there are at least two broad reasons: (1) the pursuit of basic knowledge on hormesis; and (2) practical importance in pest management. In the first case, it is only over the past decade or so that hormesis has been widely accepted in scientific circles and a large number of fundamental questions concerning the phenomenon remain to be answered. Insects are, of course, tremendously useful scientific models. They are complex multicellular organisms, and their widespread diversity and abundance, crucial ecological status, amenability to cheap and rapid mass production in controlled environments, and the lack regulatory/ethical hurdles concern-

ing experimentation make them attractive experimental subjects in toxicology and in an array of other biological disciplines. Great effort has been dedicated to the study of insect toxicology, biochemistry, physiology, molecular biology, genetics, reproduction and behavior, and this work serves as an excellent foundation for basic dose-response investigations. The genomes of many insect species have been mapped completely or in part (NCBI 2012) and the functions of a large number of insect genes are known. Thus, opportunities abound to study the basic workings of hormesis using insects as model organisms.

In the second case, the study of hormesis in insects is important because of its potential implications in agriculture and pest management (Morse 1998; Cohen 2006; Cutler *et al.* 2009; Guedes *et al.* 2009). It has long been noted that following an insecticide application, there is sometimes a surge in insect or mite population growth at a rate greater than what would have been observed without the application. This may be observed in the primary pest targeted with the insecticide (known as ‘pest resurgence’), or in a secondary pest species that is initially of lesser economic importance (known as a ‘secondary pest outbreak’) (Ripper 1956; Hardin *et al.* 1995). These surges are usually attributed to natural enemy disturbance. In such a scenario the application of a non-selective insecticide reduces both the pest and natural enemy populations, but the natural enemies are slower to recover. The reasons for this are mainly twofold: (1) natural enemy population growth lags as their food supply (the pest) recovers; and (2) natural enemies, particularly parasitoids, are often more susceptible to insecticides than their host and thus are slower to recover from toxicological effects (Croft and Brown 1975). Free of its natural enemies the pest population is able to quickly exceed its previous level of infestation.

Natural enemy decline is without question an important cause of pest resurgence, but it is usually the default or assumed cause. In reality, empirical data to support such claims is usually lacking (Hardin *et al.* 1995) and there are other possible mechanisms for an insect outbreak following an insecticide application. These include reduced competition with other herbivores, changes in pest behavior, altered host-plant nutrition, increased attractiveness of the plant host, or direct stimulation of the pesticide on the insect, factors that may operate singly or in tandem to give additive or synergistic effects (Hardin *et al.* 1995; Cohen 2006). Experiments do indeed demonstrate that hormesis could be an alternate or additional mechanism contributing to the pest resurgence phenomenon. Such resurgences could not only result in increased crop/commodity damage, but could lead to additional pesticide treatments, potentially exacerbating non-target impacts, insecticide resistance development and environmental contamination. The problem might be especially relevant in insecticide-resistant pest populations where a typical field rate might

expose individuals to the ‘hormetic-zone’ of the dose-response curve, boosting reproduction of resistant populations and increasing the frequency of the resistance alleles (Guedes *et al.* 2010).

### EVIDENCE OF HORMESIS IN INSECTS

Cohen (2006) provided discussion and a list of papers from the literature demonstrating pesticide-induced hormesis-like effects in arthropods. Table 1 and the discussion below include many of the works cited by Cohen (2006) plus additional citations, but focuses on studies reporting stimulatory effects of insecticides on insects; reports of stimulatory effects of pesticides on mites (Acari) are cited by Cohen (2006) and are generally not considered here. Also not discussed here are a number of recent biomedical articles reporting hormesis in *Drosophila* spp. when exposed to various temperature, oxidative, chemical and radiation stressors (e.g. Le Bourg 2010).

Pest resurgences following pesticide applications were reported from the beginning of the insecticide era and even then it was suspected that a favorable influence of the pesticide on arthropods may in part explain these resurgences (Ripper 1956; Peterson 1963). However, an awareness of the hormetic dose-response phenomenon seems to have been lacking among entomologists and no toxicological explanation for such occurrences was provided. In one of the earliest papers, Sun (1945) observed that while high concentrations of rotenone dust were toxic to adult female bean aphids, *Aphis rumicis*, females treated with low concentrations of the poison produced more young than control aphids. Subsequently, sublethal doses of dieldrin were found to increase longevity of *Drosophila melanogaster* (Knutson 1955) and increase weight and fecundity of house flies, *Musca domestica* (Afifi and Knutson 1956). Other early studies with *M. domestica* found stimulated reproduction when this insect was exposed to sublethal doses of various insecticides (Kilpatrick and Schoof 1956; Ouye and Knutson 1957; Hunter *et al.* 1958). Kuenen (1958) found that when the weevil *Sitophilus granarius* was fed wheat spiked with sublethal concentrations of DDT, about 20% more offspring were produced compared to unexposed weevils. Kuenen pondered whether stimulation in insects exposed to low-doses of insecticide could be a universal phenomenon. He stated that given the widely different physiological mechanisms of poisons, one “... cannot possibly expect a general rule to exist for all toxicants”, but then admits that “... a parallel with many other toxicants and DDT forces itself upon us when we see that low doses have a stimulating effect (on reproduction), while high doses kill the organism” (Kuenen 1958). Others reported that DDT stimulated oviposition in beneficial insect predators (Fleschner and Scriven 1957; Atallah and Newson 1966) and a parasitic wasp (Grosch and Valcovic 1967). In 1968, Luckey published

TABLE 1. Evidence of insecticide induced hormesis in insects.

Target insect	Chemical stressor <sup>1</sup> (sublethal doses)	Reported effects	Reference
<i>Acheta domesticus</i> (Orthoptera; house cricket)	12 different insecticides	Growth stimulation (weight)	Luckey (1968)
<i>Aphis citricola</i> (Homoptera; spirea aphid)	aldicarb	Reduced developmental time; increased longevity, fecundity, and weight	Neubauer <i>et al.</i> (1983)
<i>Aphis rumicis</i> (Homoptera; bean aphid)	rotenone	Enhanced reproduction	Sun (1945)
<i>Bemisia tabaci</i> (Homoptera; white-fly)	fenvalerate, acephate	Stimulated oviposition	Abdullah and Joginder (2004)
<i>Blaberus craniifer</i> (Blattaria; death's head cockroach)	charybdotoxin	Increased vitellogenesis	Coudey-Perribre <i>et al.</i> (1997)
<i>Bracon hebetor</i> (Hymenoptera; braconid wasp)	DDT	Stimulated oviposition	Grosch and Valcovic (1967)
<i>Ceratitis capitata</i> (Diptera; Mediterranean fruit fly)	<i>Metarhizium anisopliae</i> crude extract	Decreased pupal mortality	Ortiz-Urquiza <i>et al.</i> (2010)
<i>Choristoneura fumiferana</i> (Lepidoptera; spruce budworm)	fenitrothion, phosphamidon	Increased weight, Ca <sup>++</sup> , and total protein production	Smirnov (1983)
<i>Choristoneura rosaceana</i> (Lepidoptera; obliquebanded leafroller)	pyriproxyfen	Increased weight of pupae and adults	Sial and Brummer (2010)
<i>Chrysopa californica</i> (Neuroptera; lacewing)	DDT	Stimulated oviposition	Fleschner and Scriven (1957)
<i>Coleomegilla maculata</i> (Coleoptera; lady bird beetle)	DDT	Stimulated oviposition	Atallah and Newson (1966)
<i>Cydia pomonella</i> (Lepidoptera; codling moth)	azinphos-methyl	Stimulated copulation and oviposition	Abivardi <i>et al.</i> (1998)
<i>Diabrotica virgifera</i> (Coleoptera; western corn rootworm)	carbofuran, carbaryl	Stimulated oviposition, increased longevity	Ball and Su (1979)
<i>Drosophila melanogaster</i> <sup>2</sup> (Diptera; fruit fly)	dieldrin	Enhanced longevity	Knutson (1955)
<i>Dysdercus koenigi</i> (Heroptera; red cotton bug)	eucalyptus oil volatiles	Reduced duration of post-embryonic development	Srivastava <i>et al.</i> (1995)

continued...

TABLE 1. Continued

Target insect	Chemical stressor <sup>1</sup> (sublethal doses)	Reported effects	Reference
<i>Folsomia candida</i> (Collembola; springtail)	nonylphenol	Stimulated body length and fecundity	Widarto <i>et al.</i> (2007)
<i>Leptinotarsa decemlineata</i> (Coleoptera; Colorado potato beetle)	novaluron	Increased larval weight	Cutler <i>et al.</i> (2005)
<i>Leptopilina heterotoma</i> (Hymenoptera; <i>Drosophila</i> parasitoid)	chlorpyrifos	Stimulated host searching	Rafalimanana <i>et al.</i> (2002); Delpuech <i>et al.</i> (2005)
<i>Musca domestica</i> (Diptera; house fly)	dieldrin, DDT, BHC, chlordane, aldrin, diazinon, malathion	Increased fecundity, fertility and/ or weight	Affi and Knutson (1956); Ouye and Knutson (1957); Hunter <i>et al.</i> (1958)
<i>Myzus persicae</i> (Homoptera; green peach aphid)	azinphosmethyl, fenvalerate, imidacloprid	Stimulated reproduction	Gordon and McEwen (1984); Jackson and Wilkins (1985); Lowery and Sears (1986a; 1986b); Cutler <i>et al.</i> (2009); Yu <i>et al.</i> (2010)
<i>Nilaparvata lugens</i> (Homoptera; brown planthopper)	decamethrin, methyl parathion	Enhanced fecundity	Chilliah <i>et al.</i> (1980); Chilliah and Heimrichs (1980)
<i>Phormia regina</i> (Diptera; queen blow fly)	cadmium	Enhanced pupation	Nascarella <i>et al.</i> (2003)
<i>Phutella xylostella</i> (Lepidoptera; diamondback moth)	fenvalerate, methomyl	Increased fecundity	Sota <i>et al.</i> (1998); Fujiwara <i>et al.</i> (2002)
<i>Podisus distinctus</i> (Heteroptera; predatory stinkbug)	permethrin	Increased egg viability, increased net reproduction	Magalhães <i>et al.</i> (2002); Guedes <i>et al.</i> (2009); Zanuncio <i>et al.</i> (2011)
<i>Podisus nigripinus</i> (Heteroptera; predatory stinkbug)	gammacyhalothrin	Increased survival and longevity	Pereira <i>et al.</i> (2009)
<i>Poncilio scaber</i> (Isopoda; woodlouse)	dimethoate	Increased hyperactivity (locomotion)	Bayley and Baatrup (1996)
<i>Pyrrhocoris apterus</i> (Heteroptera; fire bug)	thiometon	Stimulated oviposition	Honěk and Novák (1977)

continued...

TABLE 1. Continued

Target insect	Chemical stressor <sup>1</sup> (sublethal doses)	Reported effects	Reference
<i>Rhyacionis kumarii</i> (Heteroptera; reduviid predator)	endosulfan	Increased fecundity and longevity	George and Ambrose (1999)
<i>Scirtothrips citri</i> (Thysanoptera; citrus thrips)	dicofof, fluralinate, formetanate, malathion	Stimulated reproduction	Morse and Zareh (1991)
<i>Sitophilus granarius</i> (Coleoptera; stored product weevil)	DDT	Stimulated reproduction	Keunen (1958)
<i>Sitophilus zeamais</i> (Curculionidae; maize weevil)	deltamethrin	Population growth	Guedes <i>et al.</i> (2010)
<i>Spodoptera littoralis</i> (Lepidoptera; Egyptian cotton leafworm)	carbaryl, methyl parathion, deltamethrin	Stimulated oviposition	Esaac <i>et al.</i> (1972)
<i>Suppoptius cincticeps</i> (Heteroptera; predatory stinkbug)	permethrin	Enhanced number of ovarioles, oviposition and life span of adult females	Kanaoka <i>et al.</i> (1996)
<i>Tribolium castaneum</i> (Coleoptera; red flour beetle)	azadirachtin	Reduced pre-oviposition period, increased survival and weight	Zanuncio <i>et al.</i> (2003, 2005)
<i>Zabrotes subfasciatus</i> (Coleoptera; Mexican bean weevil)	azadirachtin <i>Tetradenia riparia</i> extract	Enhanced survival, adult emergence, larval weight and fecundity Stimulated esterase production Increased oviposition and adult emergence	Ramachandran <i>et al.</i> (1988) Mukherjee <i>et al.</i> (1993) Weaver <i>et al.</i> (1992)

<sup>1</sup> Entomopathogens and insecticidal botanical extracts are included.

<sup>2</sup> *Drosophila melanogaster* has been used extensively as a model to study hormesis in biomedical research. Only one citation is mentioned here as the main focus is agriculturally based pesticide-induced hormesis.



his well-known article on insect hormoligosis, where he showed that weight gain of house crickets, *Acheta domesticus*, was stimulated when exposed to sublethal concentrations of 14 different insecticides (Luckey 1968). This was one of the first papers to propose a mechanism for the low-dose stimulation. Luckey (1968) suggested that insecticides in minute quantities might act as noncompetitive inhibitors of polymeric allosteric enzymes, shifting the equilibria toward the more active of two species of the polymeric molecule and resulting in the formation of other more active enzymes in the system.

Despite these intriguing papers, study of insect hormesis/hormoligosis was slow to catch on, or was dismissed. For example, Moriarty (1969) suggested that the results presented by Luckey (1968) were of “doubtful validity”. There were some reports in the 1970s describing spikes in arthropod populations following applications of pesticides in the field (e.g. Dittrich *et al.* 1974), and a few others that more closely explored low-dose stimulation by insecticides in the laboratory (Esaac *et al.* 1972; Honěk and Novák 1977), but the decade was relatively uneventful in terms of hormesis investigations. However, in the 1980s interest in the stimulatory effects of insecticides on insects picked up. Chelliah *et al.* (1980) showed that topical applications insecticides increased reproductive outputs and increased longevity of brown planthopper, *Nilaparvata lugens*, but that response varied depending on the dose and the active ingredient. Insecticide-induced stimulation of growth and/or reproduction was later observed in the flour beetle, *Tribolium castaneum* (Ramachandran *et al.* 1988), and in several aphid species (Neubauer *et al.* 1983; Gordon and McEwen 1984; Jackson and Wilkins 1985; Lowery and Sears 1986a). Smirnoff (1983) observed spruce budworm, *Choristoneura fumiferana*, larvae that survived organophosphorus and carbamates insecticide treatments developed into heavier pupae and contained more calcium and total proteins as compared to untreated larvae. In the early 1990s and years following, study of hormesis/hormoligosis accelerated with biological stimulation in insects due to low doses of insecticides being reported in many species across several orders, including thrips, wasps, woodlice, cockroaches, collembolans, and multiple species of bugs, beetles, flies, and moths (Table 1).

Several things are apparent about agricultural insect hormesis research to date.

- (i) Study of hormesis/hormoligosis per se in insects has increased, particularly in the past decade as the terms and phenomenon have become more familiar to scientists (Table 2). This pattern parallels the general ballooning interest in hormesis and the exponential increase in the number of peer-review studies citing the term *hormesis*

**TABLE 2.** Citations of hormoligosis and hormesis with insect- or pesticide-related terms.<sup>1</sup>

Search Terms	Citations per decade					Totals
	1960-69	1970-79	1980-89	1990-99	2000-12	
Hormoligosis <sup>2</sup>	2	0	4	4	16	26
Hormesis AND Coleoptera	0	0	0	0	3	3
Hormesis AND Diptera	0	0	0	0	3	3
Hormesis AND Hemiptera	0	0	0	0	1	1
Hormesis AND Heteroptera	0	0	0	0	5	5
Hormesis AND Lepidoptera	0	0	0	0	2	2
Hormesis AND Insect	0	0	1	2	18	21
Hormesis AND insecticide	0	0	1	7	21	29
Hormesis AND pesticide	0	0	3	2	35	40
Hormesis AND <i>Drosophila</i> <sup>3</sup>	0	0	2	1	63	66

<sup>1</sup> From the SciVerse Scopus database.

<sup>2</sup> Use of this term has generally, but not exclusively, been confined to insects and arthropods

<sup>3</sup> Predominantly biomedical research.

(E. Calabrese, pers. comm.).<sup>1</sup> However, in many instances biological stimulation due to low doses of insecticide is not reported as hormesis, with no mention of the terms “hormesis” or “hormoligosis” appearing in these articles (e.g. Ge *et al.* 2010; Hu *et al.* 2010; Martoub *et al.* 2011). This emphasizes that the hormesis concept is not comprehended by all insect toxicologists, even those who may be currently working on the problem.

- (ii) Documentation of hormesis in insects spans across many taxonomic orders, occurring in ametabolous, hemimetabolous and holometabolous groups, supporting arguments that hormesis is a general phenomenon (Calabrese 2010).
- (iii) Effects are seen when insects are exposed to stressors as eggs, larvae, pupae or adults, and usually stimulatory effects will carryover across life stage (but see Nascarella *et al.* 2003). This inter-life stage carry-over effect is intriguing considering the massive disassembly and restructuring of the insect DNA blueprint during metamorphosis in holometabolous taxa. Inter-generation effects over time have been less studied.
- (iv) Many different insecticidal active ingredients can elicit hormetic effects, again illustrating the generality of the phenomenon. However, most studies to date have focused on insecticidal neurotoxins. The occurrence and nature of hormesis due to, for example,

<sup>1</sup> *Hormesis Update 2010*. International Hormesis Society Conference Hormesis: Implications for Toxicology, Medicine, and Risk Assessment. University of Massachusetts, Amherst MA, 27 April 2010

insect growth regulators, pathogens or parasites that occur in agricultural systems has received little study.

- (v) Relative concentrations that induce stimulation in insects are quite variable and sometimes seem to deviate from the quantitative features typically observed in a hormetic dose response (Calabrese 2010). Mainly, in some cases we see reports of insecticide-induced stimulation with concentrations well above the No Observable Effects Concentration (NOEC).
- (vi) The large majority of studies report some sort of reproductive stimulation, although some authors have measured influences on other toxicological endpoints such as weight and some physiological or behavioral measures. Few have considered biochemical, hormonal or molecular changes in insects during hormesis, although this is likely to change in the coming years (e.g. Ge *et al.* 2010; Hu *et al.* 2010; Yu *et al.* 2010).
- (vii) With the exception of a few field and semi-field studies (Chelliah *et al.* 1980; Lowery and Sears 1986b; Morse and Zareh 1991), almost all work documenting hormesis in insects has been done in the lab, despite important implications for insecticide-induced hormesis in the field.

## INSECT HORMESIS SEMANTICS

As covered by Rozman and Doull (2003), the concept of hormesis has evolved substantially over the years and a number of terms have been used to describe the hormetic dose-response (Calabrese and Baldwin 2002). In literature related to insects, mainly three terms have been used: (1) hormesis; (2) hormoligosis; and (3) pesticide-mediated homeostatic modulation. The definition and principles of hormesis have been thoroughly covered by others (e.g. Calabrese and Baldwin 2002; e.g. Rozman and Doull 2003; Calabrese 2005a; Mattson and Calabrese 2010) and will not be repeated here. The terms ‘hormesis’ and ‘hormoligosis’ have been compared previously (Morse 1998; Cohen 2006; Guedes *et al.* 2009), and Cohen (2006) proposed the additional concept of ‘pesticide-mediated homeostatic modulation’. Here I provide a more in-depth examination of the terms ‘hormoligosis’ and ‘pesticide-mediated homeostatic modulation’, as they relate to hormesis.

### Hormoligosis

The terms ‘hormoligosis’ and ‘hormoligant’ were coined by T.D. Luckey in 1955 at the First International Conference on Antibiotics in Agriculture (Luckey 1956; 2008). The hormoligosis concept originally, and erroneously, encompassed all substances that were stimulatory at

small doses, i.e. not only stressors or toxic substances, with a hormoligant being the entity that excites or stimulates in small quantities. It was noted that the stimulation was more pronounced in individuals that were under stress and it seems that the definition was subsequently modified bearing this in mind. Luckey (1963) described hormoligosis as a situation where "... minute quantities of any stressing agent (chemical, physical, psychological or social) would be stimulatory to an organism under a wide variety of conditions, whereas larger quantities of stressing agent would be harmful to the same organism." Here the principle of low-dose stimulation and high-dose inhibition becomes apparent. In 1968, the paper *Insect Hormoligosis* was published (Luckey 1968) and in describing hormoligosis it is stated that "... subharmful quantities of many stress agents may be helpful when presented to organisms in suboptimal environments". This definition for hormoligosis seems to have been the most enduring, though ironically most contemporary authors using the term have tended to ignore the criterion of "suboptimal environment" (Cohen 2006; Guedes *et al.* 2009). It has been suggested that processes initiated by a stimulatory signal followed by homeostatic overcompensation in the other direction could give hormoligotic responses, being delineated from hormetic responses that are initiated by an inhibitory signal followed by homeostatic overcompensation (Rozman and Doull 2003), although broadly defined hormesis could apply to both scenarios (Calabrese 2010). Luckey's definition suggests that hormoligosis applies to situations where an organism is subjected to stress but no stimulation is observed until a small amount of a second stressor is presented to the system.

From an evolutionary perspective, hormoligosis can be seen as a small (but not too small) dose of stress optimizing the fitness of an individual to deal with subsequent stresses (Rozman and Doull 2003). It is not clear in this definition whether or not the effect of the first stress acts additively or synergistically with the second, or if stimulation can occur with either stressor alone if its dose was increased. Differentiating between these potential interactive effects is important since it has been shown that mixtures of stressors at low doses can produce hormetic responses and that the magnitude of stimulation can be roughly predicted in mixtures with knowledge of the concentration–response relationships of the individual stressors (Belz *et al.* 2008). That is, is hormoligosis simply a form of mixture hormesis, or is it something different? Since not all stressors may induce hormesis (Belz *et al.* 2008), the later may be true. This should be a key research question for insect toxicologists and toxicologists in general with an interest in hormesis. At present, except where conceptual and mechanistic differences between hormesis and hormoligosis are specified (e.g. Rozman and Doull 2003; Cohen 2006; Guedes *et al.* 2009), these terms have generally been treated synonymously in the insect toxicology literature.

### **Pesticide-mediated homeostatic modulation**

Cohen (2006) argued that hormesis does not apply in situations where stimulatory effects are observed in an arthropod pest which had neither been targeted nor controlled by a pesticide. He distinguished ‘acaricides’, stressors targeting mites, from ‘insecticides’, chemicals that are exclusively toxic to insects, and ‘pesticides’, chemicals that affect both types of arthropods. Examples are presented mainly as cases where stimulation of mite reproduction is seen following exposure to pesticides such as DDT, carbaryl, insecticidal pyrethroids or imidacloprid. In terms of their use patterns these compounds are not considered acaricides and are not intended to control mite pests. But mites and insects often cohabit the same environment and in some cases when the aforementioned compounds are used to manage insect pests, mite population surges have been observed. Cohen (2006) rightfully points out that to establish hormesis, dose-response endpoints such as LD or LC values should first be clearly established. He contends that stimulatory effects observed with pesticides that are non-toxic to arthropods at high doses (e.g. DDT stimulation of two-spotted spider mite, *Tetranychus urticae*, reproduction in the field) cannot be attributed to hormesis and proposed *pesticide-mediated homeostatic modulation* as a broader term to include both hormesis and stimulatory effects of pesticides on non-target pests.

However, pesticide-mediated homeostatic modulation is a dubious concept. Cohen’s (2006) arguments are not toxicologically driven, but rather are semantic in nature and rest mainly on differentiating stimulatory effects in ‘target’ vs. ‘non-target’ arthropods due to exposure to ‘insecticides’ vs. ‘acaricides’. Such anthropocentric distinctions are of no matter to the underlying mechanisms that cause observed stimulations. Rather, it is the toxicological nature of the dose-response that is fundamental in delineating a response as hormetic. The central tenant of toxicology is Paracelsus’ maxim that any chemical is potentially toxic (poisonous) and it is only the dose that renders it so. Many common substances are not considered insecticides per se, but they can kill insects depending on the dose.

Indeed, several of the compounds that Cohen (2006) suggests are non-acaricidal can in fact be lethal to mites. For example, DDT, methyl parathion and permethrin are not considered acaricides or practical pesticides for management of *T. urticae*, but they undoubtedly may kill *T. urticae* depending on the dose (e.g. Attiah and Boudreaux 1964; Ayyappatii *et al.* 1997). A dose-response of *T. urticae* to either of these chemicals may therefore be established, along with the possibility of high-dose inhibition and low-dose stimulation as hormesis. Hormesis may very well apply to situations where a pest insect is not controlled by a pesticide if lethal or “high” doses exceed field rates. For instance, several researchers have observed that *M. persicae* reproduction was greatly stim-

ulated by field rates of azinphosmethyl (Peterson 1963; Gordon and McEwen 1984; Lowery and Sears 1986a; 1986b), but this compound is also clearly lethal to this insect (Lowery 1985; Raman 1988). Further doubt of the validity of the pesticide-mediated homeostatic modulation concept is raised when it is suggested that it and hormesis are different but that the underlying physiological/biochemical mechanisms are presumably identical (Cohen 2006).

It seems that “hormesis” should suffice *in lieu* of the terms “hormoligosis” and “pesticide-mediated homeostatic modulation”, at least to explain stimulatory responses encountered thus far in insect toxicology. Pesticide-mediated homeostatic modulation does not propose a dose-response or mechanistic underpinning any different from hormesis. It is a concept based on semantics and is not biologically demarcated from hormesis. Hormoligosis is a term with strong historical significance in insect toxicology and one that continues to be used by some authors when describing insecticide-induced stimulation in insects. By definition it may be delineated from hormesis in that the insect (or any organism/cell) is to be experiencing sub-optimal conditions prior to any observed biological stimulation by a low dose of insecticide (or any stressor). This may in fact be a case of mixture hormesis, where two stressors are employed with or without a temporal separation in exposure.

## **CONSIDERATIONS FOR STUDY**

Hormesis is a burgeoning field of study. However, despite the potential implications of hormesis-rooted stimulations in pest management, and the opportunities insect-insecticide models can offer in basic investigations, the phenomenon has received relatively scant attention from insect toxicologists. Cohen (2006) offered several important suggestions for further research and below I briefly describe additional problems that researchers may wish to consider in future hormesis investigations. Some problems are specific to entomology, but others may be of interest in other disciplines and with alternative models.

### **General experimental design**

Most experiments exploring chemical-induced hormesis in insects have suffered from the same shortcoming as similar studies in other disciplines, mainly that too few doses are used, no or few sub-NOEC are used, too many high doses are used, there is inadequate replication, and there is no time component (Calabrese and Baldwin 1998; Calabrese 2005b). When examining stimulatory effects of insecticides on insects, insect toxicologists have usually chosen 3-4 test concentrations below the LC50, e.g. LC5, LC10 or LC25, but little or no rationale for their treatment choices is given. In some cases reports of hormesis are *a posteriori*

and the experimenter may have had no knowledge of the phenomenon. However, where there is *a priori* knowledge of hormesis, experiments that aim to study stimulatory or sublethal effects of insecticides on insects should incorporate many replicates and doses, including several in the theoretical hormetic zone below the NOEC (but see below), derived from previous experiments if necessary (Cutler *et al.* 2009).

### **Insecticide-plant-insect interactions**

Insecticides have been shown to alter plant photosynthesis, respiration, transpiration, and metabolism of nitrogen, carbohydrates and minerals (NAS 1968), which may make the plant more attractive and susceptible to attack by phytophagous insects. When conducting dose-response/hormesis experiments using insects and plants, it is therefore important to be mindful of the exposure route and consider whether the observed stimulatory effects are due to direct interaction of the insecticide with the insect, or some indirect influence of the compound on the plant tissue on which the insect is feeding (Cohen 2006). Neubauer *et al.* (1983) found that stimulated growth and reproduction in the aphid *A. citricola* when exposed to leaves systemically treated with aldicarb, but not when this compound was administered in an artificial diet, suggesting the hormetic effects observed may have been plant mediated. Similarly, Ferguson and Chapman (1993) found that carbaryl applications in potato caused *M. persicae* population surges and after excluding reductions in natural enemies as a reason for this, concluded that stimulation in aphid reproduction was due to the insecticide treatment. However, they also noted in laboratory experiments that carbaryl increased total nitrogen content in leaves, which was correlated with increased nymph production. It has been suggested that any agent that disturbs nitrogen metabolism in plant tissues will favor increased survival and abundance of herbivores feeding on those tissues (White 1984).

Insecticides may also change physical characteristics of a plant that may increase their attractiveness to insects (NAS 1968). Such phenomena should not be misinterpreted as hormesis. For example, Hari and Mahal (2008a) reported that application of sub-lethal concentrations of the insecticides acephate and cypermethrin reduced cotton height, plant spread and upper canopy leaf area, and this resulted in increased oviposition of *Helicoverpa armigera* (cotton bollworm).

### **Stimulatory concentrations**

Meta-analyses reveal that hormetic stimulation typically peaks at concentrations below the NOEC (Calabrese and Baldwin 1998; 2001; Calabrese 2005a; 2005b). However, in studies with insects, stimulation is sometimes reported at concentrations well above the NOEC. Stimulation

above control levels following exposure to concentrations in the LC25 range is not uncommon (e.g. Sota *et al.* 1998; Bao *et al.* 2009), and stimulation at the LC50 has been observed (Esaac *et al.* 1972; Chelliah *et al.* 1980; Sota *et al.* 1998). Examination of variances in response among treated (e.g. insects exposed to an insecticide LC25 concentration) and control groups in these studies suggests that individuals within and across treatment groups respond homogeneously, so results are not skewed by a few individuals that have abnormally high reproductive outputs. It is important to keep in mind that in a field situation, although reproduction in some individuals may be stimulated by exposure to a LC25 insecticide concentration, 25% of the population would be killed by this exposure, possibly negating stimulatory effects on the population as a whole. Nonetheless, apparent insecticide-induced stimulatory effects at concentrations well above NOEC levels is an important deviation from the hormetic dose-response defined by Calabrese (2005a; 2005b) and is worth investigating further.

### **The nature of the stressor**

The occurrence of hormesis may not be absolute among all chemicals that induce a toxicological response (e.g. Belz *et al.* 2008). This may be true in particular for agents that induce hormesis via direct stimulation of a receptor mediated pathway. Even where the molecular structure of chemicals is highly similar, the ability of these different chemicals to induce hormesis may differ (Calabrese 2010). It has sometimes been observed that not all insecticides in low doses (not necessarily doses around the NOEC) induce stimulation. For example, Chelliah *et al.* (1980) found that while reproductive stimulation occurred in *N. lugens* at LC25 and LC50 doses of methyl parathion (an organophosphorus insecticide) and decamethrin (a synthetic pyrethroid insecticide), respectively, no stimulation was found when the insect was exposed to perthane (a chlorinated hydrocarbon) at several similar concentrations below the LC50. Neubauer *et al.* (1983) observed statistically significant hormetic effects in spirea aphids, *Aphis citricola*, exposed to sublethal concentrations of aldicarb, but not with ethiofencarb or dimethoate. When exposed to 1/10 LC50 concentrations of several insecticides, endosulfan increased fecundity and longevity of *Rhynocoris kumari*, where the other compounds tested decreased fecundity and longevity (George and Ambrose 1999). Similarly, LC30 concentrations of endosulfan resulted in reduced developmental times for *Helicoverpa armigera* but the same concentrations of spinosad, chlorpyrifos, acephate and cypermethrin had deleterious impacts on the insect's development (Hari and Mahal 2008b). Experiments should easily be able to determine whether such results are simply contingent on the experimental design (i.e. picking the best suite of doses to detect potential hormesis), or whether hormetic



dose-response curves vary with chemical mode of action or structure. In contrast to neurotoxins, hormesis in insects exposed to low doses of insect growth regulators or entomopathogens has been less studied (Ortiz-Urquiza *et al.* 2010).

### **Peak-dip-peak phenomenon?**

Luckey (1963) studied the response of nematodes under heat stress to various concentrations of antibiotic. He noted that with several antibiotics there were two peaks of stimulation whereas normal response or inhibition of survival was found at concentrations between the peaks. Similar results were seen in some experiments with crickets exposed to insecticides, with two peaks of stimulated weight gain observed, along with intermediate doses resulting in reduced or no significant weight gain (Luckey 1968). In our own experiments we too have seen this peak-dip-peak phenomenon (Cutler *et al.* 2009; Mohan *et al.* unpublished). We have found its occurrence to be real and reproducible, and variable with exposure scenario, dose selection and life stage. Stimulation of different molecular pathways at different doses may be at play, but this needs to be explored further. Smooth J-shaped or inverted-U shaped hormetic curves may not always be observed in insects during stimulatory responses to low levels of stress.

### **Fitness tradeoffs**

It has been suggested that hormesis results from a temporary reallocation of resources, which will be followed by a measurable fitness tradeoff in some other biological process (Forbes 2000). There is evidence of such tradeoffs in the insect-hormesis literature. For example, Sial and Brunner (2010) found that although *Choristoneura rosaceana* (oblique-banded leafroller) pupae and adult weights significantly increased following larval exposure to sublethal concentrations of pyriproxyfen, subsequent fecundity and fertility of the insect was reduced at these doses. Enhanced pupation was observed in queen blowfly following exposure of maggots to cadmium but this was followed by reduced adult emergence from pupae, reflecting stage specific hormesis (Nascarella *et al.* 2003). Fujiwara *et al.* (2002) observed that although egg laying was stimulated in diamondback moths exposed to LC25 treatments, eggs laid were smaller and offspring survival was reduced, resulting in no change in reproductive effort compared to untreated insects. Thus, tradeoffs are apparent but seem to vary depending on the exposure regime. It would be worthwhile determining if there are consistent trends in how tradeoffs manifest under different exposure regimes and among different taxa, and how these exposures affect biological fitness occur across generations (Widarto *et al.* 2007). Stepping beyond traditional hormesis endpoint

assessments for insects (e.g. fecundity, fertility, longevity) is encouraged. Entomologists should look more closely at molecular, physiological, morphological, behavioral and population markers for hormesis or counter tradeoffs.

### **Hormesis in insecticide resistant populations**

Insecticide resistance continues to be a major issue for the management of insect pests in food production (Denholm *et al.* 1998), and the control of disease vectors (Hemingway and Ranson 2000; Labbé *et al.* 2007). Study of hormesis within the context of insecticide resistance is of interest. As Guedes *et al.* (2010) suggested, if field doses of an insecticide do not control an insecticide resistant population, such sublethal exposures might very well boost their population growth through stimulated reproduction and increase the frequency of resistance alleles within the population, exacerbating the resistance problem. Insecticide-induced hormesis therefore may be important for the evolution of pesticide resistance and design of resistance management programs (Guedes *et al.* 2010), but this has not been well studied.

### **Hormesis and beneficial insects**

Mass rearing and management of beneficial insects is a multi-billion dollar industry. Insects are reared for pollination, honey and honey bee hive products, biological control, sterile insect release programs, pet food, human food, biological and medical research, silk, and many other end uses. Potential beneficial implications for hormesis for human health have been well documented (Mattson and Calabrese 2010), and we might be able to apply hormetic principles during mass culture of insects to, for example, improve insect longevity, immunity or reproductive outputs. For example, Wojda *et al.* (2009) found that exposure of *Galleria mellonella* – a common experimental model and source of food for animals – to mild heat shock for 30 minutes decreased the insect's susceptibility to a common entomopathogenic pathogenic fungus, possibly due to increased expression of antimicrobial peptides and higher antifungal and lysozyme activities. Guedes *et al.* (2009) showed that reproductive outputs of the beneficial predatory bug *Podisus distinctus* increased and generation time was reduced with exposure to a single sublethal dose of permethrin. Similar results were seen with the insect predator *Supputius cincticeps* (Zanuncio *et al.* 2005). Long-term experiments should be pursued to determine if these types of hormesis-based observations can translate into economic benefits during mass culturing of beneficial insects.

## Field studies

Although hormesis has been implicated as a potentially important factor in insect pest resurgences and resistance development (Guedes *et al.* 2009, 2010), the prevalence of the phenomenon in agricultural settings is generally not known, and very few agricultural studies have tested insect hormesis hypotheses under field or greenhouse conditions. This is ironic since observed pesticide-induced resurgences in the field were often the principle driver for initial studies of insect hormoligosis and hormesis (Dittrich *et al.* 1974; Chelliah *et al.* 1980; Lowery and Sears 1986b; Morse and Zareh 1991). Field experiments with specific pests, crops and pesticide exposure regimes are needed to determine if and under what circumstances insect hormesis occurs. This will clarify whether or not pesticide-induced insect hormesis has economic repercussions for crop production and pest management.

## ACKNOWLEDGEMENTS

I thank the Natural Sciences and Engineering Research Council (NSERC) for funding my research on hormesis, and comments from an anonymous reviewer that helped improve the manuscript.

## REFERENCES

- Abdullah NMM and Joginder S. 2004. Effect of insecticides on longevity of the whitefly, *Bemisia tabaci* (Gennadius) on cotton. *Univ Aden J Nat Appl Sci* 8:261-268
- Abivardi C, Weber DC, and Dorn S. 1998. Effects of azinphos-methyl and pyrifenoxy on reproductive performance of *Cydia pomonella* L. (Lepidoptera: Tortricidae) at recommended rates and lower concentrations. *Annals Appl Biol* 132:19-33
- Affi SED and Knutson H. 1956. Reproductive potential, longevity, and weight of house flies which survived one insecticidal treatment. *J Econ Entomol* 49:310-313
- Atallah YH and Newson LL. 1966. Ecological and nutritional studies on *Coleomegilla maculata* De Geer (Coleoptera: Coccinellidae). III. The effects of DDT, toxaphene, and endrin on the reproductive and survival potentials. *J Econ Entomol* 59:1181-1187
- Attiah HH and Boudreaux HB. 1964. Population dynamics of spider mites influenced by DDT. *J Econ Entomol* 57:53-57
- Ayyappathi R, Witicowski JF, and Higley LG. 1997. Ovipositional responses of two species of spider mites (Acari: Tetranychidae) to sublethal concentrations of permethrin and methyl parathion on corn. *Environ Entomol* 26:489-496
- Azzam EI, de Toledo SM, Raaphorst GP, and Mitchel REJ. 1996. Low-dose ionizing radiation decreases the frequency of neoplastic transformation to a level below the spontaneous rate in C3H 10T1/2 cells. *Radiat Res* 146:369-373
- Ball H and Su PP. 1979. Effect of sublethal dosages of carbofuran and carbaryl on fecundity and longevity of the female western corn rootworm. *J Econ Entomol* 72:873-876
- Bao HB, Liu SH, Gu JH, Wang XZ, Liang XL, and Liu ZW. 2009. Sublethal effects of four insecticides on the reproduction and wing formation of brown planthopper, *Nilaparvata lugens*. *Pest Manag Sci* 65:170-174
- Bayley M and Baatrup E. 1996. Pesticide uptake and locomotor behaviour in the woodlouse: An experimental study employing video tracking and <sup>14</sup>C-labelling. *Ecotoxicol* 5:35-45
- Belz RG, Cedergreen N and Sorensen H. 2008. Hormesis in mixtures - Can it be predicted? *Sci Total Environ* 404:77-87

- Benbrook CM. 2004. Genetically engineered crops and pesticide use in the United States: the first nine years. BioTech InfoNet, Technical Paper No. 7 (available at [http://organic.insightd.net/reportfiles/Full\\_first\\_nine.pdf](http://organic.insightd.net/reportfiles/Full_first_nine.pdf))
- Brann JL. 1956. Apparatus for application of insecticides. *Annu Rev Entomol* 1:241-260
- Calabrese EJ. 2005a. Toxicological awakenings: the rebirth of hormesis as a central pillar of toxicology. *Toxicol Appl Pharmacol* 204:1-8
- Calabrese EJ. 2005b. Paradigm lost, paradigm found: the re-emergence of hormesis as a fundamental dose response model in the toxicological sciences. *Environ Poll* 138:378-411
- Calabrese EJ. 2010. Hormesis is central to toxicology, pharmacology and risk assessment. *Hum Exp Toxicol* 29:249-261
- Calabrese EJ and Baldwin LA. 1998. Hormesis as a biological hypothesis. *Environ Health Pers* 106:357-362
- Calabrese EJ and Baldwin LA. 2001. Hormesis: A generalizable and unifying hypothesis. *Crit Rev Toxicol* 31:353-424
- Calabrese EJ and Baldwin LA. 2002. Applications of hormesis in toxicology, risk assessment and chemotherapeutics. *Trends Pharmacol Sci* 23:331-337
- Calabrese EJ and Baldwin LA. 2003a. The hormetic dose-response model is more common than the threshold model in toxicology. *Toxicol Sci* 71:246-250
- Calabrese EJ and Baldwin LA. 2003b. Toxicology rethinks its central belief. *Nature* 421:691-692
- Calabrese EJ and Blain R. 2005. The occurrence of hormetic dose response in the toxicological literature, the hormesis database: an overview. *Toxicol Appl Pharmacol* 202:289-301
- Calabrese EJ and Cook RR. 2005. Hormesis: how it could affect the risk assessment process. *Human Exp Toxicol* 24:265-270
- Cattaneo MG, Yafuso C, Schmidt C, Huang CY, Rahman M, Olson C, Ellers-Kirk C, Orr BJ, Marsh SE, Antilla L, Dutilleu P, and Carriere Y. 2006. Farm-scale evaluation of the impacts of transgenic cotton on biodiversity, pesticide use, and yield. *Proc Nat Acad Sci* 103:7571-7576
- Chelliah S, Fabellar LT, and Heinrichs EA. 1980. Effect of sub-lethal doses of three insecticides on the reproductive rate of the brown planthopper, *Nilaparvata lugens*, on rice. *Environ Entomol* 9:778-780
- Chelliah S and Heinrichs EA. 1980. Factors affecting insecticide-induced resurgence of the brown planthopper, *Nilaparvata lugens*, on rice. *Environ Entomol* 9:773-777
- Cohen E. 2006. Pesticide-mediated homeostatic modulation in arthropods. *Pestic Biochem Physiol* 85:21-27
- Croft BA and Brown AWA. 1975. Response of arthropod natural enemies to insecticides. *Annu Rev Entomol* 20:285-335
- Cutler GC, Ramanaidu K, Astatkie T, and Isman MB. 2009. Green peach aphid, *Myzus persicae* (Hemiptera: Aphididae), reproduction during exposure to sublethal concentrations of imidacloprid and azadirachtin. *Pest Manag Sci* 65:205-209
- Cutler GC, Scott-Dupree CD, Tolman JH, and Harris CR. 2005. Acute and sublethal toxicity of novaluron, a novel chitin synthesis inhibitor, to *Leptinotarsa decemlineata* (Coleoptera : Chrysomelidae). *Pest Manag Sci* 61:1060-1068
- Delpuech JM, Bardon C, and Boulétreau M. 2005. Increase of the behavioral response to kairomones by the parasitoid wasp *Leptopilina heterotoma* surviving insecticides. *Arch Environ Contam Toxicol* 49:186-191
- Denholm I, Cahill M, Dennehy TJ, and Horowitz AR. 1998. Challenges with managing insecticide resistance in agricultural pests, exemplified by the whitefly *Bemisia tabaci*. *Phil Trans R Soc Lond B* 353:1757-1767
- Dittrich V, Streibert P, and Bathe PA. 1974. An old case reopened: mite stimulation by insecticide residues. *Environ Entomol* 3:534-540
- Esaac EG, El-Gogary S, Abdel-Fatah MS, and Ali AM. 1972. Effect of carbaryl, methyl parathion, and dieldrin on egg production and percent pupation of the Egyptian cotton leafworm, *Spodoptera littoralis* (Boisd.). *J Appl Entomol* 71:263-70
- Feinendegen LE. 2005. Evidence for beneficial low level radiation effects and radiation hormesis. *Brit J Radiol* 78:3-7
- Ferguson JS and Chapman RK. 1993. Factors involved in carbaryl-induced population buildups of *Myzus persicae* (Sulzer) (Homoptera, Aphididae) on potato. *J Agricult Entomol* 10:51-64
- Fleschner CA and Scriven GT. 1957. Effect of soil-type and DDT on ovipositional response of *Chrysopa californica* (Coq.). *J Econ Entomol* 50:221-222

- Forbes VE. 2000. Is hormesis an evolutionary expectation? *Funct Ecol* 14:12-24
- Fujiwara Y, Takahashi T, Yoshioka T, and Nakasuji F. 2002. Changes in egg size of the diamondback moth *Plutella xylostella* (Lepidoptera : Yponomeutidae) treated with fenvalerate at sublethal doses and viability of the eggs. *Appl Entomol and Zool* 37:103-109
- Galbadage T and Hartman PS. 2008. Repeated temperature fluctuation extends the life span of *Caenorhabditis elegans* in a daf-16-dependent fashion. *Mech Ageing Dev* 129:507-514
- Ge L-Q, Wu J-C, Zhao K-F, Chen Y, and Yang G-Q. 2010. Induction of *Nlvg* and suppression of *Nljhe* gene expression in *Nilaparvata lugens* (Stål) (Hemiptera: Delphacidae) adult females and males exposed to two insecticides. *Pestic Biochem Physiol* 98:279-288
- George PJE and Ambrose DP. 1999. Insecticidal impact on the post-embryonic development of *Rhynocoris kumarii* Ambrose and Livingstone (Het., Reduviidae). *J Appl Entomol* 123:498-401
- Gomez FH, Bertoli CI, Sambucetti P, Scannapieco AC, and Norry FM. 2009. Heat-induced hormesis in longevity as correlated response to thermal-stress selection in *Drosophila buzzatii*. *J Therm Biol* 34:17-22
- Gordon PL and McEwen FL. 1984. Insecticide-stimulated reproduction of *Myzus persicae*, the green peach aphid (Homoptera: Aphididae). *Can Entomol* 116:7783-784
- Goudey-Perrière F, Dahmani F, Perrière C, Brousse-Gamy P, and Menez A. 1997. Enhancement of oocyte growth in the cockroach *Blaberus cranzifer* by a scorpion toxin, charybdotoxin. *Life Sci* 60:199-205
- Grosch DS and Valcovic LR. 1967. Chlorinated hydrocarbon insecticides are not mutagenic in *Bracon hebetor* tests. *J Econ Entomol* 60:1177-1179
- Guedes RNC, Magalhaes LC, and Cosme LV. 2009. Stimulatory sublethal response of a generalist predator to permethrin: hormesis, hormoligosis, or homeostatic regulation? *J Econ Entomol* 102:170-176
- Guedes NMP, Tolledo J, Corrêa AS, and Guedes RNC. 2010. Insecticide-induced hormesis in an insecticide-resistant strain of the maize weevil, *Sitophilus zeamais*. *J Appl Entomol* 134:142-148
- Gunther FA and Blinn RC. 1956. Persisting insecticide residues in plant materials. *Annu Rev Entomol* 1:167-180
- Hardin MR, Benrey B, Coll M, Lamp WO, Roderick GK, and Barbosa P. 1995. Arthropod pest resurgence: an overview of potential mechanisms. *Crop Prot* 14:3-18
- Hari NS and Mahal MS. 2008a. Sub-lethal influences of different insecticides on oviposition preference of *Helicoverpa armigera* (Hubner) (Lepidoptera : Noctuidae) on cotton in relation to altered plant morphology. *Int J Pest Manag* 54:181-187
- Hari NS and Mahal MS. 2008b. Effect of repeated sub-lethal applications of different insecticides on egg laying preference by *Helicoverpa armigera* (Hubner) on cotton. *Pestic Res J* 20:95-98
- Haynes KF. 1988. Sublethal effects of neurotoxic insecticides on insect behavior. *Annu Rev Entomol* 33:149-168
- Hemingway J and Ranson H. 2000. Insecticide resistance in insect vectors of human disease. *Annu Rev Entomol* 45:371-39
- Honěk A and Novák I. 1977. Effect of sublethal doses of Intra 50 on reproduction in *Pyrrhocoris apterus* L. (Heteroptera, Pyrrhocoridae). *J Appl Entomol* 83:364-370
- Hoskins WM and Gordon HT. 1956. Arthropod resistance to chemicals. *Annu Rev Entomol* 1:89-122
- Hu J-H, Wu J-C, Yin J-L, and Gu H-N. 2010. Physiology of insecticide-induced stimulation of reproduction in the rice brown planthopper (*Nilaparvata lugens* (Stål)): Dynamics of protein in fat body and ovary. *Int J Pest Manag* 56:23-30
- Hunter PE, Cutkomp LK, and Kolkaila AM. 1958. Reproduction in DDT- and diazinon-treated house flies. *J Econ Entomol* 11:579-582
- Jackson AEM and Wilkins RM. 1985. The effect of sub-lethal dosages of the synthetic pyrethroid fenvalerate on the reproductive rate of the aphid *Myzus persicae*. *Pest Sci* 16:364-368
- Kanaoka A, Yamaguchi R, and Konno T. 1996. Effect of buprofezin on oviposition of brown planthopper, *Nilaparvata lugens*, at sub-lethal dose. *J Pestic Sci* 21:153-157
- Kearns CW. 1956. The mode of action of insecticides. *Annu Rev Entomol* 1:123-148
- Keunen D. 1958. Influence of sublethal doses of DDT upon the multiplication rate of *Sitophilus granarius* (Coleopt. Curculionidae). *Entomol Exp Appl* 1:147-152
- Kiely T, Donaldson D, and Grube A. 2004. Pesticides Industry Sales and Usage: 2000 and 2001 Market Estimates. EPA's Biological and Economic Analysis Division, Office of Pesticide Programs, and Office of Prevention, Pesticides, and Toxic Substances. [www.epa.gov/pesticides](http://www.epa.gov/pesticides).

- Kilpatrick JW and Schoof JH. 1956. Fly production in treated and untreated privies. *Public Health Rep* 71:787-796
- Knutson H. 1955. Modifications in fecundity and lifespan of *Drosophila melanogaster* Meigen following sublethal exposure to an insecticide. *Ann Entomol Soc Amer* 48:35-39
- Kuenen DJ. 1958. Influence of sublethal doses of DDT upon the multiplication rate of *Sitophilus granarius* (Coleopt.: Curculionidae). *Entomol Exp Appl* 1:147-152
- Labbé P, Berticat C, Berthomieu A, Unal S, Bernard C, Weill M, and Lenormand T. 2007. Forty years of erratic insecticide resistance evolution in the mosquito *Culex pipiens*. *PLoS Genet* 3: e205
- Le Bourg E. 2011. Using *Drosophila melanogaster* to study the positive effects of mild stress on aging. *Exp Gerontol* 46:345-348
- Le Bourg E and Rattan SIS. 2008. *Mild Stress and Healthy Aging: Applying Hormesis in Aging Research and Interventions*. Springer, New York
- Lowery DT. 1985. Effect of azinphosmethyl, an organophosphorus insecticide, on reproduction of the green peach aphid (Homoptera: Aphididae). M.Sc. Thesis, University of Guelph, Guelph, Ontario, Canada
- Lowery DT and Sears MK. 1986a. Effect of exposure to the insecticide azinphosmethyl on reproduction of green peach aphid (Homoptera: Aphididae). *J Econ Entomol* 79:1534-1538
- Lowery DT and Sears MK. 1986b. Stimulation of reproduction of the green peach aphid (Homoptera: Aphididae) by azinphosmethyl applied to potato. *J Econ Entomol* 79:1530-1533
- Luckey TD. 1956. Mode of action of antibiotics - evidence from germfree birds. *Use of Antibiotics in Agriculture*. National Academy of Sciences, Washington, D.C. p. 135-145.
- Luckey TD. 1963. Antibiotic action in adaptation. *Nature* 198:263-265
- Luckey TD. 1968. Insecticide hormoligosis. *J Econ Entomol* 61:7-12
- Luckey TD. 1991. *Radiation Hormesis*. CRC Press, Boca Raton
- Luckey TD. 2008. Sir Samuraj T. D. Luckey, PhD. *Dose-Response* 6:97-112
- Magalhães LC, Guedes RNC, Oliveira EE, and Tuelher ES. 2002. Development and reproduction of the predator *Podisus distinctus* (Stal) (Heteroptera: Pentatomidae) exposed to sublethal doses of permethrin. *Neotrop Entomol* 31:445-448
- Martin H. 1956. The chemistry of insecticides. *Annu Rev Entomol* 1:149-166
- Martoub M, Couty A, Giordanengo P, and Amelin A. 2011. Opposite effects of different mineral oil treatments on *Macrosiphum euphorbiae* survival and fecundity. *J Pest Sci* 84:229-233
- Mattson MP. 2009. The fundamental role of hormesis in evolution In: Mattson MP, Calabrese EJ (eds), *Hormesis: A Revolution in Biology, Toxicology and Medicine*, pp 53-68. Springer, New York
- Mattson MP and Calabrese EJ (eds). 2010. *Hormesis: A Revolution in Biology, Toxicology and Medicine*. Springer, New York
- Mohan M, Cutler GC, and Scott-Dupree CD. 2012. Inter-generational and life-stage differences in imidacloprid induced hormesis in green peach aphid. (unpublished data)
- Moriarty F. 1969. The sublethal effects of synthetic insecticides on insects. *Biol Rev* 44:321-357
- Morse JG. 1998. Agricultural implications of pesticide-induced hormesis of insects and mites. *Hum Exp Toxicol* 17:266-269
- Morse JG and Zareh N. 1991. Pesticide-induced hormoligosis of citrus thrips (Thysanoptera: Thripidae) fecundity. *J Econ Entomol* 84:1169-1174
- Mukherjee SN, Rawal SK, Ghumare SS, and Sharma RN. 1993. Hormetic concentrations of azadirachtin and isoesterase profiles in *Tribolium castaneum* (Herbst) (Coleoptera, Tenebrionidae). *Experientia* 49:557-560
- NAS. 1968. *Effects of Pesticides on Fruit and Vegetable Physiology*. Principles of Plant and Animal Pest Control, Vol. 6. National Academy of Sciences, Washington.
- Nascarella MA, Stoffolano JG, Stanek EJ, Kostecki PT, and Calabrese EJ. 2003. Hormesis and stage specific toxicity induced by cadmium in an insect model, the queen blowfly, *Phormia regina* Meig. *Environ Poll* 124:257-262
- Nauen R, Tietjen K, Wagner K, and Elbert A. 1998. Efficacy of plant metabolites of imidacloprid against *Myzus persicae* and *Aphis gossypii* (Homoptera: Aphididae). *Pestic. Sci.* 52:53-57
- NCBI. 2012. Basic Local Alignment Search Tool, National Center for Biotechnology Information. Available at <http://blast.ncbi.nlm.nih.gov/Blast.cgi>
- Neubauer I, Raccach B, Aharonson N, Swirski E, and Ishaaya I. 1983. Systemic effect of aldicarb, dimethoate and ethiofencarb on mortality and population dynamics of the spirea aphid, *Aphis citricola* Van der Goot. *Crop Prot* 2:211-218

- Olson ER, Dively GP, and Nelson JO. 2004. Bioassay determination of the distribution of imidacloprid in potato plants: Implications to resistance development. *J Econ Entomol* 97:614-620
- Ortiz-Urquiza A, Vergara-Ortiz A, Santiago-Alvarez C, and Quesada-Moraga E. 2010. Insecticidal and sublethal reproductive effects of *Metarhizium anisopliae* culture supernatant protein extract on the Mediterranean fruit fly. *J Appl Entomol* 134:581-591
- Ouye MT and Knutson H. 1957. Reproductive potential, longevity, and weight of house flies following treatments of larvae with malathion. *J Econ Entomol* 50:490-493
- Parsons PA. 2001. The hormetic zone: an ecological and evolutionary perspective based upon habitat characteristics and fitness selection. *Q Rev Biol* 76:459-67
- Pereira AIA, de Sousa Ramalho F, de Medeiros Bandeira C, Malaquias JB, and Zanuncio JC. 2009. Age-dependent fecundity of *Podisus nigrispinus* (Dallas) (Heteroptera: Pentatomidae) with sublethal doses of gamma-cyhalothrin. *Braz Arch Biol Tech* 52:1157-1166
- Peterson AG. 1963. Increases of the green peach aphid following the use of some insecticides on potatoes. *Am Potato J* 40:121-129
- Rafalimanana H, Kaiser L, and Delpuech J-M. 2002. Stimulating effects of the insecticide chlorpyrifos on host searching and infestation efficacy of a parasitoid wasp. *Pest Manag Sci* 58:321-328
- Ramachandran R, Mukherjee SN, and Sharma RN. 1988. Hormetic effect of azadirachtin on *Tribolium castaneum* Herbst (Coleoptera: Tenebrionidae). *Indian J Exp Biol* 26:913-914
- Raman KV. 1988. Insecticide toxicity to three strains of green peach aphid (*Myzus persicae* Sulzer) reared on resistant and susceptible potato cultivars. *Crop Prot* 7:62-65
- Ripper WE. 1956. Effects of pesticides on balance of arthropod populations. *Annu Rev Entomol* 1:403-438
- Rozman KK and Doull J. 2003. Scientific foundations of hormesis. Part 2. Maturation, strengths, limitations, and possible applications in toxicology, pharmacology, and epidemiology. *Crit Rev Toxicol* 33:451-462
- Sanders CL. 2010. Radiation Hormesis and the Linear-No-Threshold Assumption. Springer, New York
- Sial AA and Brunner JF. 2010. Lethal and sublethal effects of an insect growth regulator, pyriproxyfen, on obliquebanded leafroller (Lepidoptera: Tortricidae). *J Econ Entomol* 103:340-347
- Smirnov WA. 1983. Residual effects of *Bacillus thuringiensis* and chemical insecticide treatments on spruce budworm (*Choristoneura fumiferana* Clemens). *Crop Prot* 2:225-230
- Sota N, Motoyama N, Fujisaki K, and Nakasuji F. 1998. Possible amplification of insecticide hormoligosis from resistance in the diamondback moth, *Plutella xylostella* (Lepidoptera: Yponomeutidae). *Appl Entomol Zool* 33:435-440
- Srivastava RK, Gurusubramanian G, and Krishna SS. 1995. Postembryonic development and reproduction in *Dysdercus koenigii* (F) (Heteroptera, Pyrrhocoridae) on exposure to Eucalyptus oil volatiles. *Biol Agric Hort* 12:81-88
- Stark JD and Banks JE. 2003. Population-level effects of pesticides and other toxicants on arthropods. *Annu Rev Entomol* 48:505-519
- Stolzinger A, Sethe S, and Scutt AM. 2006. Stressed stem cells: Temperature response in aged mesenchymal stem cells. *Stem Cells Dev* 15:478-487
- Sun Y-P. 1945. Effect of rotenone and Velsicol (AR-60) dusts on the control and reproduction of bean aphids. *J Econ Entomol* 38:124-125
- Weaver DK, Dunkel FV, Cusker JL, and Vanpuylvelde L. 1992. Oviposition patterns in two species of bruchids (Coleoptera: Bruchidae) as influenced by the dried leaves of *Tetradenia riparia*, a perennial mint (Lamiales: Lamiaceae) that suppresses population size. *Environ Entomol* 21:1121-1129
- White TCR. 1984. The abundance of invertebrate herbivores in relation to the availability of nitrogen in stressed food plants. *Oecologia* 63:90-105
- Widarto TH, Krogh PH, and Forbes VE. 2007. Nonylphenol stimulates fecundity but not population growth rate ( $\lambda$ ) of *Folsomia candida*. *Ecotoxicol Environ Safe* 67:369-377
- Wojda I, Kowalski P, and Jakubowicz T. 2009. Humoral immune response of *Galleria mellonella* larvae after infection by *Beauveria bassiana* under optimal and heat-shock conditions. *J Insect Phys* 55:525-531
- Yu SJ. 2008. *The Toxicology and Biochemistry of Insecticides*. CRC Press, Boca Raton
- Yu Y, Shen G, Zhu H, and Lu Y. 2010. Imidacloprid-induced hormesis on the fecundity and juvenile hormone levels of the green peach aphid *Myzus persicae* (Sulzer). *Pestic Biochem Phys* 98:38-242

- Zanuncio TV, Serrao JE, Zanuncio JC, and Guedes RNC. 2003. Permethrin-induced hormesis on the predator *Supputius cincticeps* (Stal, 1860) (Heteroptera: Pentatomidae). *Crop Prot* 22:941-947
- Zanuncio TV, Zanuncio JC, Serrao JE, Medeiros RS, Pinon TBM, and Sedyama CA. 2005. Fertility and life expectancy of the predator *Supputius cincticeps* (Heteroptera: Pentatomidae) exposed to sublethal doses of permethrin. *Biol Res* 38:31-39
- Zanuncio JC, Jusselino-Filho P, Ribeiro RC, Zanuncio TV, de S. Ramalho F, and Serrão JE. 2011. Hormetic responses of a stinkbug predator to sublethal doses of pyrethroid. *Bull Environ Contam Toxicol* 8:608-614