

FORMIDABEL: The Belgian Ants Database

Dimitri Brosens¹, François Vankerkhoven², David Ignace³, Philippe Wegnez⁴,
Nicolas Noé⁵, André Heughebaert⁵, Jeannine Bortels⁶, Wouter Dekoninck⁷

1 Instituut voor Natuur en Bosonderzoek, Kliniekstraat 25, B-1070, Brussels, Belgium **2** Polyergus, Wolvenstraat 9, B-3290, Diest, Belgium **3** FourmisWalBru, Rue Winston Churchill 91, B-6180, Courcelles, Belgium **4** FourmisWalBru, Rue de la Grotte, 23, B-4651, Herve, Belgium **5** ULB, Université Libre de Bruxelles, Campus de la Plaine CP257, Brussels, Belgium **6** ULG, Université de Liège, Gembloux Agro-Bio Tech, Passage des Déportés, 2, B-5030 Gembloux - Belgium **7** Royal Belgian Institute of Natural Sciences (RBINS), Vautierstraat 29, B-1000, Brussels, Belgium

Corresponding author: Wouter Dekoninck (Wouter.Dekoninck@naturalsciences.be)

Academic editor: V. Chavan | Received 25 February 2013 | Accepted 22 May 2013 | Published 3 June 2013

Citation: Brosens D, Vankerkhoven F, Ignace D, Wegnez P, Noé N, Heughebaert A, Bortels J, Dekoninck W (2013) FORMIDABEL: The Belgian Ants Database. ZooKeys 306: 59–70. doi: 10.3897/zookeys.306.4898 Resource ID: GBIF key: <http://gbrds.gbif.org/browse/agent?uuid=b528799a-2d52-4023-aa02-9ce081e3ca5f>

Resource citation: Polyergus Working Group and FourmisWalBru Working Group (2013 -). FORMIDABEL: The Belgian Ants Database, 27264 records. Contributed by Brosens D, Vankerkhoven F, Ignace D, Wegnez P, Noé N, Heughebaert A, Bortels J and W Dekoninck. Online at <http://ipt.biodiversity.be/resource.do?r=formidabel> and, www.formicidae-atlas.be, Version 1.0 (last updated on 2013-04-19), GBIF key: <http://gbrds.gbif.org/browse/agent?uuid=b528799a-2d52-4023-aa02-9ce081e3ca5f>. Data Paper ID: doi: 10.3897/zookeys.306.4898

Abstract

FORMIDABEL is a database of Belgian Ants containing more than 27.000 occurrence records. These records originate from collections, field sampling and literature. The database gives information on 76 native and 9 introduced ant species found in Belgium. The collection records originated mainly from the ants collection in Royal Belgian Institute of Natural Sciences (RBINS), the ‘Gaspar’ Ants collection in Gembloux and the zoological collection of the University of Liège (ULG). The oldest occurrences date back from May 1866, the most recent refer to August 2012. FORMIDABEL is a work in progress and the database is updated twice a year.

The latest version of the dataset is publicly and freely accessible through this url: <http://ipt.biodiversity.be/resource.do?r=formidabel>. The dataset is also retrievable via the GBIF data portal through this link: <http://data.gbif.org/datasets/resource/14697>




Figure 1. www.formicidae-atlas.be

A dedicated geo-portal, developed by the Belgian Biodiversity Platform is accessible at: <http://www.formicidae-atlas.be>

Purpose: FORMIDABEL is a joint cooperation of the Flemish ants working group “Polyergus” (<http://formicidae.be>) and the Wallonian ants working group “FourmisWalBru” (<http://fourmiswalbru.be>). The original database was created in 2002 in the context of the preliminary red data book of Flemish Ants (Dekoninck et al. 2003). Later, in 2005, data from the Southern part of Belgium; Wallonia and Brussels were added. In 2012 this dataset was again updated for the creation of the first Belgian Ants Atlas (Figure 1) (Dekoninck et al. 2012). The main purpose of this atlas was to generate maps for all outdoor-living ant species in Belgium using an overlay of the standard Belgian ecoregions. By using this overlay for most species, we can discern a clear and often restricted distribution pattern in Belgium, mainly based on vegetation and soil types.

Keywords

Formicidae, Belgium, Flanders, Wallonia, Brussels Capital Region, ecological data, grid mapping, UTM, historical data, literature, collections, observations, trapping, ants

Taxonomic coverage

General taxonomic coverage description

The taxonomic coverage (Figure 2) of this database spans the full range of ants pertaining to Belgium (indigenous ant species and exotic introduced species). The determination level is species level and, if appropriate, hybrid level. For some species, information on micro- and macrogynes is available. Key milestones of FORMIDABEL from conception till date are described in the “Dataset” section of this manuscript.




Figure 2. Taxonomic range of the FORMIDABEL database subfamilies

The taxonomic authorities followed are: Radchenko and Elmes (2010) for the genus *Myrmica* and Seifert (2007) for the other genera. The dataset contains occurrences of 76 native and 9 introduced species.

As depicted in Figure 2, the most abundant subfamily in the database is the Formicinae (49,4%) followed by the Myrmicinae (48,7%), the Dolichoderinae (0,9%) and the Ponerinae (0,8%). The top five most recorded species are *Lasius niger* (2846 records), *Myrmica rubra* (2601 records), *Myrmica scabrinodis* (1626 records), *Formica fusca* (1467 records) and *Myrmica sabuleti* (1202 records).

Taxonomic ranks

Phylum: Arthropoda

Subphylum: Hexapoda

Class: Insecta

Order: Hymenoptera

Suborder: Apocrita

Superfamily: Vespoidea

Family: Formicidae

Subfamily: Dolichoderinae, Formicinae, Myrmicinae, Ponerinae

Genera: *Anergates*, *Aphaenogaster*, *Camponotus*, *Dolichoderus*, *Formica*, *Formicoxenus*, *Harpagoxenus*, *Hypoponera*, *Lasius*, *Leptothorax*, *Linepithema*, *Manica*, *Monomorium*, *Myrmecina*, *Myrmica*, *Plagiolepis*, *Polyergus*, *Ponera*, *Solenopsis*, *Stenamma*, *Strongylognathus*, *Tapinoma*, *Technomyrmex*, *Temnothorax* and *Tetramorium*.

Common names: Ants

Spatial coverage

General spatial coverage

Belgium is a small country in Western Europe. To the west, its 70 km coastline fronts the North Sea; to the north lies the Netherlands; to the east, Germany, and to the south, France and Luxembourg. Biogeographically, the fauna of eastern Belgium belongs to the Central European Province of the Eurasian (Palaeartic) region. By contrast, the rest of the country primarily consists of an Atlantic fauna plus a few Central European relict species.

Politically and geographically, the country is divided into three parts: Flanders, Wallonia and the Brussels Capital Region (Figure 3). In Flanders (13,522 km² and population about 6 million people), to the north, soils are mainly sandy to loamy. Here, the most important habitats for ants are heathlands and dry grasslands. The Brussels Capital Region is a small region (162 km²) entirely situated in the sandy loam area. In Wallonia (17,006 km² and about 3,5 million people), to the south, soils and habitats are more diverse, ranging from forests to rocky and calcareous grasslands on loam and chalky soils. Eastern Wallonia, near the German border, includes the Hautes Fagnes, a large area of bogs and peat with some typical ant species.

Geographical method

The Universal Transverse Mercator Projection (UTM), an adaptation of the standard Mercator projection, uses a two dimensional Cartesian co-ordinate system to identify locations on the surface of the Earth (Wikipedia).

The UTM 5 Km (Universal Transverse Mercator Projection) raster projection divides Belgium in approximately 1200 25 km² squares (Figure 3). A representative number of UTM squares has been sampled (1125 UTM 5×5 km squares of which 659 squares with more than 10 records: see Figures 3 and 4) to complete the dataset. All the records in FORMIDABEL are georeferenced through the centroid coordinates of the corresponding UTM 5 km square. Therefore, the uncertainty on these coordinates is 3.500 meters, the distance between the centre and the corner of the UTM square.

Ecocodes

More than half of the records are provided with a description of the microhabitat of the record locality. This allows us to give details on ecological preferences of all Belgian ant species. In FORMIDABEL we created a list of potential microhabitats for ants in Belgium. For each of these microhabitats we use a code called the “ecocode”. This code thus gives information on the habitat where the occurrence was made. Nine types of habitat and landscape were defined to collect accurate information on the habi-




Figure 3. The UTM 5 Km grid of Belgium




Figure 4. Projection of the number of records per UTM 5x5 km square (grey dots= 1-20 records, pink dots= 21-50 records, red dots= 51-150 records, dark red dots=151-300 records and black dots=301-644 records).

tat preference of all ant species (Dekoninck et al. 2005). These nine types are based on the EIS-code and the Flemish nature types (Vandenbussche, 2002; Zwaenepoel et al. 2002). When no habitat description was available for a record (for example, with some older records) the habitat was coded as 'Not known or not observed'. The nine

main habitats we defined were: i) anthropogenic habitats, ii) dry grasslands, iii) moist grasslands, iv) forests, v) chalk grasslands, stony slopes and other rocky xerothermic habitats, vi) shrubs, vii) heathlands, viii) fens and highland bogs and ix) coastal and inland dunes.

Coordinates

49°27'0"N and 51°32'24"N Latitude; 2°28'12"E and 6°27'36"E Longitude

Temporal coverage

The oldest record in the database goes back to May 5, 1866 and the most recent records are from August 2012. The largest part of the records were obtained after 1991 (Figure 5).

Methods

Method step description:

A large portion of the occurrence data have been collected by volunteers, other records originated from several projects and research programs. The data and specimens were sent to the Belgian ant curators, and after validation, the information was incorporated in the database. The collection records “dry specimen” originate from the Gembloux “Ant” collection and the Charles Gaspar collection, the collection of the “Cercle des entomologistes Liégeois”, the RBINS collection and the private collection “Roland Vannieuwenhuysse”. After revision and validation, this information was also included in the database. The literature-based records were retrieved from van Boven 1970; van




Figure 5. Temporal distribution of the records

Boven and Mabelis 1986; Dekoninck et al. 2006 and references therein. How the database evolved is described in the Database history section.

Sampling description: Most occurrence records originate from hand/nest sampling (42,3% of all records and mainly from Wallonia) and pitfall sampling (36,7% mainly from Flanders). The followed procedure differs from region to region. This is due to historical reasons. Some very interesting occurrence records were obtained by sifting, coloured water traps and Malaise traps (all less than 3 % of the total sampling). An extensive description of the sampling methods is provided by Schauff (2001).

Quality control description: All the records were validated by the dataset curators before being added to the FORMIDABEL database. The dataset curators also checked the determinations of the collection specimens. If needed, the determination was adapted and made consistent with modern taxonomy; Radchenko and Elmes (2010) for the genus *Myrmica* and Seifert (2007) for the other genera. Before the final publication of the database all the records were tested on geographical consistency by the Belgian Biodiversity Platform and corrected if necessary.

Dataset

Dataset history

At the beginning of 2001 all available records of ants in Flanders (northern part of Belgium) were brought together for the first time and several inventories were started. More than 20.000 records (for the most part gathered after 1990) were assembled in the FORMIDABEL (FORMIcidaeDATABELgium) database resulting in the ‘Verspreidingsatlas en voorlopige Rode lijst van de mieren van Vlaanderen; Dekoninck et al. 2003. [Distribution atlas and preliminary Red list of ant species in Flanders, Belgium]. This report contains all available distribution data for Flanders. In the southern part of Belgium (Wallonia) intensive inventorying started in 2005. Until then knowledge on the distribution of ants in Wallonia was limited. Only a few areas (Haute Fagnes, Famenne and the Viroin valley) had already been inventoried. Thanks to the good cooperation between the Polyergus and the FourmisWalBru working groups; many Belgian ant records were brought together in the FORMIDABEL database. Since then, FORMIDABEL is updated with data originating from FourmisWalBru twice a year. The FORMIDABEL dataset was then used for the creation of the “Belgian Ant Atlas”, (Dekoninck et al. 2012). In 2010 a cooperation agreement between the authors of the atlas and the Belgian Biodiversity Platform (www.biodiversity.be) was made. Together with the publication of the book, the data was published in Darwin Core Archives to GBIF ([url: http://data.gbif.org/datasets/resource/14697/](http://data.gbif.org/datasets/resource/14697/)) and a dedicated data portal was created: www.formicidae-atlas.be.

The original FORMIDABEL database was created in Microsoft Access. Later, this database was completely imported in a relational SQL database. During this process




Figure 6. Formidabel flowchart

additional data cleaning was performed; see section Quality control description. The dataflow is illustrated in Figure 6.

Dataset description

The FORMIDABEL Darwin Core Archive is a custom made SQL view on the original version of the FORMIDABEL access database which is in the custody of the Belgian Ant working groups Polyergus and fourmisWalBru. Mind that every record in FORMIDABEL represent at least one occurrence, but primarily contains information on the presence of a species. The view only shows data that are accepted for publication. Fields given are:

id, decimalLatitude, family, basisOfRecord, stateProvince, identifiedBy, eventDate, modified, country, individualCount, scientificName, kingdom, order, geodeticDatum, genus, collectionCode, decimalLongitude, samplingProtocol, catalogNumber, phylum, recordNumber, countryCode, coordinatePrecision, language, coordinateUncertaintyInMeters, locality, specificEpithet, recordedBy, institutionCode, nomenclaturalCode, class.

The dataset contains primary biodiversity data, mostly occurrence data (Figure 7). Some records hold an indirect link to collection specimens. This link is only available in the original database.

Dataset preview

Object name: Darwin Core Archive Formidabel; Belgian Ants Database

Character encoding: UTF-8 **Format name:** Darwin Core Archive format

Format version: 1.0

Distribution: <http://ipt.biodiversity.be/resource.do?r=formidabel>

Publication date of data: 2013-08-02

Language: English

Licenses of use: This work is licensed under a Creative Commons Attribution-Non-Commercial-ShareAlike 3.0 Unported License. <http://creativecommons.org/licenses/by-nc-sa/3.0/>

Metadata language: English

Date of metadata creation: 2013-02-18

Hierarchy level: Dataset

Norms for data use and publication:

Based on <http://www.canadensys.net/norms>

Give credit where credit is due

As is common practice in scientific research, cite the data you are using.

Be responsible

Use the data responsibly. The data are published to allow anyone to better study and understand the world around us, so please do not use the data in any way that is unlawful, harmful or misleading. Understand that the data are subject to change, errors and sampling bias. Protect the reputation of the data publisher and clearly indicate any changes you may have made to the data.

Share knowledge

Let us know if you have used the data. It helps our participants to showcase their efforts and it helps you reach a wider audience. Inform the data publisher(s) if you have comments about the data, notice errors, or want more information.

Respect the data license

Understand and respect the data license or waiver under which the data are published. It is indicated in the rights field of every record and in the dataset metadata.

Collection data: Ant Collection Gembloux ([urn:lsid:biocol.org:col:33368](http://rsb2012.org/urn:lsid:biocol.org:col:33368)), Collection Charles Gaspar, Collection “Cercle des entomologistes Liégeois”, RBINS Belgian Formicidae Collection ([urn:lsid:biocol.org:col:35271](http://rsb2012.org/urn:lsid:biocol.org:col:35271)), Private collection “Van Nieuwenhuysse”. All collections are dry prepared insect collections. The dataset does not contain unique identifiers for specimens. To track a collection specimen, the corresponding author should be contacted.

id	basisOfRecord	eventDate	scientificName	order	locality	class	decimalLatitude	decimalLongitude
KBVE-SRB PreservedSpecim		36353	<u>Anergates atratulus</u> (Scher Hymenopte Zonhoven)			Insecta	51	5.387
KBVE-SRB HumanObservatic		36776	<u>Anergates atratulus</u> (Scher Hymenopte Meeuwen-Gr.)			Insecta	51.085	5.606
KBVE-SRB HumanObservatic		36710	Anergates atratulus (Scher Hymenopte Neerpelt)			Insecta	51.269	5.401
KBVE-SRB HumanObservatic		36663	<u>Anergates atratulus</u> (Scher Hymenopte Zonhoven)			Insecta	50.998	5.458
KBVE-SRB HumanObservatic		36693	Anergates atratulus (Scher Hymenopte Zonhoven)			Insecta	50.998	5.458
KBVE-SRB HumanObservatic		36757	<u>Anergates atratulus</u> (Scher Hymenopte Neerpelt)			Insecta	51.269	5.401
KBVE-SRB HumanObservatic		36757	Anergates atratulus (Scher Hymenopte Neerpelt)			Insecta	51.269	5.401
KBVE-SRB HumanObservatic		21385	<u>Anergates atratulus</u> (Scher Hymenopte As)			Insecta	50.995	5.601
KBVE-SRB HumanObservatic		2721	Anergates atratulus (Scher Hymenopte Rotselaar)			Insecta	50.966	4.745
KBVE-SRB PreservedSpecim		36318	<u>Anergates atratulus</u> (Scher Hymenopte Genk)			Insecta	50.998	5.458
KBVE-SRB PreservedSpecim		38972	Anergates atratulus (Scher Hymenopte Adinkerke)			Insecta	51.068	2.536
KBVE-SRB HumanObservatic		40029	<u>Anergates atratulus</u> (Scher Hymenopte Diest)			Insecta	51.006	5.031
KBVE-SRB HumanObservatic		40029	Anergates atratulus (Scher Hymenopte Diest)			Insecta	51.006	5.031
KBVE-SRB PreservedSpecim		39201	<u>Anergates atratulus</u> (Scher Hymenopte Peer)			Insecta	51.088	5.463
KBVE-SRB PreservedSpecim		39215	Anergates atratulus (Scher Hymenopte Peer)			Insecta	51.043	5.461
KBVE-SRB HumanObservatic		38507	<u>Anergates atratulus</u> (Scher Hymenopte Kalmthout)			Insecta	51.42	4.402
KBVE-SRB HumanObservatic		38507	Anergates atratulus (Scher Hymenopte Kalmthout)			Insecta	51.375	4.401
KBVE-SRB HumanObservatic		38639	<u>Anergates atratulus</u> (Scher Hymenopte Kalmthout)			Insecta	51.42	4.402
KBVE-SRB PreservedSpecim		40427	Anergates atratulus (Scher Hymenopte Lombardsijde)			Insecta	51.159	2.75
KBVE-SRB PreservedSpecim		38589	<u>Anergates atratulus</u> (Scher Hymenopte Visç.)			Insecta	50.769	5.659
KBVE-SRB PreservedSpecim		39694	Anergates atratulus (Scher Hymenopte Oostmalle)			Insecta	51.281	4.756
KBVE-SRB PreservedSpecim		39694	<u>Anergates atratulus</u> (Scher Hymenopte Oostmalle)			Insecta	51.281	4.756
KBVE-SRB HumanObservatic		27302	Anergates atratulus (Scher Hymenopte Marchin)			Insecta	50.463	5.219
KBVE-SRB PreservedSpecim		40020	<u>Anergates atratulus</u> (Scher Hymenopte Ampsin)			Insecta	50.552	5.294
KBVE-SRB PreservedSpecim		30491	Aphaenogaster subterrane Hymenopte Marche-les-Deux)			Insecta	50.513	4.939
KBVE-SRB PreservedSpecim		23543	<u>Aphaenogaster subterrane</u> Hymenopte Neuville-sous-Origny)			Insecta	50.552	5.294
KBVE-SRB HumanObservatic		37771	Aphaenogaster subterrane Hymenopte Nismes)			Insecta	50.069	4.572
KBVE-SRB HumanObservatic		37771	<u>Aphaenogaster subterrane</u> Hymenopte Treignes)			Insecta	50.113	4.643
KBVE-SRB PreservedSpecim		7561	Aphaenogaster subterrane Hymenopte Trooz)			Insecta	50.589	5.649
KBVE-SRB HumanObservatic		39327	<u>Aphaenogaster subterrane</u> Hymenopte Treignes)			Insecta	50.113	4.643
KBVE-SRB PreservedSpecim		3873	Aphaenogaster subterrane Hymenopte Beez)			Insecta	50.468	4.938
KBVE-SRB PreservedSpecim		3867	<u>Aphaenogaster subterrane</u> Hymenopte Samson)			Insecta	50.467	5.008
KBVE-SRB PreservedSpecim		3886	Aphaenogaster subterrane Hymenopte Samson)			Insecta	50.467	5.008
KBVE-SRB PreservedSpecim		7561	<u>Aphaenogaster subterrane</u> Hymenopte Trooz)			Insecta	50.588	5.72

Figure 7. A small preview of the Darwin Core-Archive

Additional information

This dataset was originally created to develop the Belgian Ants Atlas. However, the dataset can be reused for a variety of purposes. Since the link between individual data records and underlined specimens (stored in multiple collections) is not recorded, we doubt if the dataset can be used for taxonomic or systematic studies. However, this being an occurrence dataset, it can be used for understanding species richness, distribution pattern and modeling studies such as ecological niche modeling. In order to enhance the confidence of use, we have documented the metadata as well as subjected the data records to a series of quality assessment and enhancement processes as described in the earlier section quality control description.

Acknowledgments

The authors would like to thank all the contributors to this data paper. Especially all the volunteers who spend hours in the field collecting ants. We would also like to thank the Entomology Department of RBINS (Dr. Patrick Grootaert), the Cercle des Entomologistes Liégeois (P. Lays) and Gembloux Agro-Bio Tech, Unité d'Entomologie

fonctionnelle et évolutive for giving us access to the collections and the Belgian Biodiversity Platform and GBIF to make this work possible. Last but not least we would like to thank Thomas Little for making sure that the English language was respected during the creation of this paper.

References

1) References cited within the metadata

- Schauff ME (2001) Collecting And Preserving Insects And Mites Techniques And Tools Update and modified WWW version of: Steyskal GC, Murphy WL, Hoover EM (Eds)(1986) Insects and mites: techniques for collection and preservation. Agricultural Research Service, USDA, Miscellaneous Publication, Vol. 1443
- Dekoninck W, Vankerkhoven F, Maelfait J-P (2003) Verspreidingsatlas en voorlopige Rode Lijst van de mieren van Vlaanderen. Rapport van het Instituut voor Natuurbehoud IN.R.2003.07. Brussel 191 pp.
- Dekoninck W, Maelfait J-P, Vankerkhoven F, Grootaert P (2005) Remarks on the distribution and use of a provisional red list of the ants of Flanders (Formicidae, Hymenoptera). In: Procter D, Harding PY (Eds) JNCC Report No. 367 Proceedings of *IN* Cardiff 2003, Red Lists for Invertebrates: their application at different spatial scales – practical issues, pragmatic approaches, 74–85.
- Dekoninck W, Maelfait J-P, Vankerkhoven F, Baugnée J-Y, Grootaert P (2006) – An update of the checklist of the Belgian ant fauna with comments on new species for the country (Hymenoptera, Formicidae). *Belgium Journal of Entomology* 8: 27–41.
- Dekoninck W, Igance D, Vankerkhoven F, Wegnez P (2012) Verspreidingsatlas van de mieren van België/Atlas des fourmis de Belgique. *Bulletin SRBE/KBVE*, 148 (III): 95–186pp.
- Seifert B (2007) Die Ameisen Mittel-und Nordeuropas - *Lutra Verlags- und Vertriebsgesellschaft*. Klitten, 368 pp.
- Radchenko AG, Elmes GW (2010) *Myrmica* ants (Hymenoptera: Formicidae) of the Old World. *Fauna Mundi*, Vol. 3, 789 pp.
- Vandenbussche V (2002) Systematiek van natuurtypes voor de biotopen heide, moeras, duin, slik en schor: deel 1: Inleiding. Brussels, Instituut voor Natuurbehoud.
- Zwaenepoel A, T’Jollyn F, Vandenbussche V, Hoffmann M (2002) Systematiek van Natuurtypes voor de biotoop grasland. Brugge & Brussels, WVI & Instituut voor Natuurbehoud.

2) References used to develop the dataset

- Bondroit J (1909) Les fourmis de Belgique. *Annales de la Société royale belge d’Entomologie* 1909, 53: 479–500.
- Bondroit J (1910) Compte rendu d’une excursion aux environs de Hockai. *Annales de la Société royale belge d’Entomologie*, 54: 231–232.
- Bondroit J (1911) Contribution à la fauna de Belgique, Notes diverses. *Annales de la Société royale belge d’Entomologie*, 55: 8–13.
- Bondroit J (1912) Fourmis des Hautes Fagnes. *Bulletin et Annales de la Société royale belge d’Entomologie*, 56: 351–352.
- Bondroit J (1918) Les Fourmis de France et de Belgique. *Annales de la Société d’Entomologie de France*, 87: 1–174.

- Cammaerts R, Cammaerts M-C (1988) Four ants (Hym.: Formicidae) new to the Belgian Fauna. *Entomologist's Record*, 100: 37–38.
- De Biseau J-C, Couvreur J-M (1994) Fourmis (Formicidae). *Faune de Belgique*. Institut royal des Sciences naturelles de Belgique, 56 pp.
- Dessart P, Cammaerts R (1995) Recaptures d'*Hypoponera punctatissima* en Belgique (Hymenoptera Formicidae Ponerinae). *Bulletin S.R.B.E./K.B.V.E.*, 131: 487–489.
- Gaspar C (1970) Hymenoptera Formicidae: cartes 143–164. In *Atlas Provisoire des Insectes de Belgique*, cartes 101–200. Edité par J. Leclercq, Faculté des Sciences Agronomiques de l'Etat, Zoologie générale et Faunistique, Gembloux.
- Gaspar C (1970) Hymenoptera Formicidae: cartes 15–30. In *Atlas Provisoire des Insectes de Belgique*, cartes 1–100. Edité par J. Leclercq, Faculté des Sciences Agronomiques de l'Etat, Zoologie générale et Faunistique, Gembloux.
- Gaspar C (1971) Hymenoptera Formicidae: cartes 203–216. In *Atlas Provisoire des Insectes de Belgique*, cartes 201–300. Edité par J. Leclercq, Faculté des Sciences Agronomiques de l'Etat, Zoologie générale et Faunistique, Gembloux.
- Schoeters E, Vankerhoven F (2001) *Onze mieren*. Educatie Limburgs Landschap, Heusden-Zolder, 175 pp.
- van Boven JKA (1947) Liste de détermination des principales espèces de Fourmis belges (Hymenoptera: Formicidae). *Bulletin S.R.B.E./K.B.V.E.*, 83: 163–190.
- van Boven JKA (1949) Notes sur la faune des Hautes Fagnes en Belgique. *Bulletin S.R.B.E./K.B.V.E.*, 85: 135–143.
- van Boven, JKA (1970) Vliesvleugelige insecten - Hymenoptera, Angeldragers - Aculeata, Mieren - Formicidae. *Wetenschappelijke Mededelingen van de Koninklijke Nederlandse Natuurhistorische Vereniging*, 30: 52 pp.
- van Boven JKA, Mabelis AA (1986) De mieren van de Benelux. *Wetenschappelijke Mededelingen van de KNNV* 173, 64 pp.

3) Publications based on this dataset

- Dekoninck W, Igance D, Vankerhoven F, Wegnez P (2012) *Verspreidingsatlas van de mieren van België/Atlas des fourmis de Belgique*. *Bulletin SRBE/KBVE*, 148 (II): 94 pp.
- Dekoninck W, Maelfait J-P, Vankerhoven F, Bagnée J-Y, Grootaert P (2006) An update of the checklist of the Belgian ant fauna with comments on new species for the country (Hymenoptera, Formicidae). *Belgium Journal of Entomology*, 8: 27–41.