

Published in final edited form as:

Br J Nutr. 2010 November ; 104(9): 1288–1296. doi:10.1017/S0007114510002217.

***In vivo* modulation of 4E binding protein 1 (4E-BP1) phosphorylation by watercress: a pilot study**

Sharifah S. Syed Alwi¹, Breeze E. Cavell¹, Urvi Telang², Marilyn E. Morris², Barbara M. Parry³, and Graham Packham^{1,*}

¹Cancer Sciences Division, School of Medicine, Cancer Research UK Centre, Southampton General Hospital, The Somers Cancer Research Building (MP824), University of Southampton, Southampton SO16 6YD, UK

²Department of Pharmaceutical Sciences, School of Pharmacy and Pharmaceutical Sciences, University at Buffalo, State University of New York, Amherst, NY 14260-1200, USA

³Winchester and Andover Breast Unit, Royal Hampshire County Hospital, Winchester SO22 5DG, UK

Abstract

Dietary intake of isothiocyanates (ITC) has been associated with reduced cancer risk. The dietary phenethyl ITC (PEITC) has previously been shown to decrease the phosphorylation of the translation regulator 4E binding protein 1 (4E-BP1). Decreased 4E-BP1 phosphorylation has been linked to the inhibition of cancer cell survival and decreased activity of the transcription factor hypoxia-inducible factor (HIF), a key positive regulator of angiogenesis, and may therefore contribute to potential anti-cancer effects of PEITC. In the present study, we have investigated the *in vitro* and *in vivo* effects of watercress, which is a rich source of PEITC. We first demonstrated that, similar to PEITC, crude watercress extracts inhibited cancer cell growth and HIF activity *in vitro*. To examine the effects of dietary intake of watercress, we obtained plasma and peripheral blood mononuclear cells following the ingestion of an 80 g portion of watercress from healthy participants who had previously been treated for breast cancer. Analysis of PEITC in plasma samples from nine participants demonstrated a mean maximum plasma concentration of 297 nM following the ingestion of watercress. Flow cytometric analysis of 4E-BP1 phosphorylation in peripheral blood cells from four participants demonstrated significantly reduced 4E-BP1 phosphorylation at 6 and 8 h following the ingestion of watercress. Although further investigations with larger numbers of participants are required to confirm these findings, this pilot study suggests that flow cytometry may be a suitable approach to measure changes in 4E-BP1 phosphorylation following the ingestion of watercress, and that dietary intake of watercress may be sufficient to modulate this potential anti-cancer pathway.

Keywords

Isothiocyanates; Cancer; 4E binding protein 1 (4E-BP1); Phosphorylation

Epidemiological studies have suggested that a diet rich in cruciferous vegetables such as broccoli, cabbage and watercress is associated with reduced risk of multiple cancer types⁽¹⁾. The potential anti-cancer effect of high cruciferous vegetable intake has been linked to the

© The Authors 2010

*Corresponding author: Professor G. Packham, fax +44 23 8079 5152, g.k.packham@soton.ac.uk.

The authors declare no conflicts of interest.

presence of glucosinolates within these foods. Following the release of the plant enzyme myrosinase by chewing or cutting, hydrolysis of glucosinolates gives rise to a number of products including isothiocyanates (ITC), indoles, thiocyanates and nitriles^(2,3). Over 100 glucosinolates which give rise to chemically distinct hydrolysis products have been identified. Following absorption, ITC are rapidly conjugated to glutathione (GSH) via the action of glutathione-*S*-transferase enzymes and metabolised predominantly via the mercapturic acid pathway^(2,3). Several studies have indicated that the potential cancer-protective effects of a high-cruciferous vegetable diet are modulated by sequence variations within glutathione-*S*-transferase enzymes, most notably GSTM1 and GSTT1^(4–7). These variants may be linked to the enhanced protective effects of a high-cruciferous vegetable diet via potential effects on ITC metabolism⁽⁸⁾. The mechanisms of action of the potential anti-cancer activity of ITC are complex, and at present, are incompletely understood. However, alterations of carcinogen metabolism via inhibition of phase I enzymes and induction of phase II enzymes, as well as direct modulation of pathways controlling key cancer hallmarks, such as proliferation, resistance to apoptosis and angiogenesis, are thought to be involved^(2,3,9,10).

Following uptake into cells, ITC conjugate rapidly with the free thiol of glutathione^(11,12). Cycles of ITC conjugate efflux, regeneration of ITC by extracellular hydrolysis and reuptake of ITC lead to a very marked accumulation of ITC within cells and concomitant depletion of intracellular GSH. Decreased intracellular GSH leads to increased levels of reactive oxygen species, which may play an important role in suppressing growth and survival of transformed cells⁽¹³⁾. In the absence of GSH-mediated defence, intracellular ITC are thought to conjugate with various cellular proteins, predominantly via reactive cysteine thiols^(2,14). One arm of this response, driven by covalent modification of Keap1, leads to the stabilisation of Nrf2, a master regulator of antioxidant gene expression, and induction of protective antioxidant proteins^(15–18). Similar events are thought to contribute to the direct anti-cancer effects of ITC on proliferation, survival and angiogenesis⁽²⁾. Conjugation to α - and β -tubulins may contribute to the mitotic arrest that is frequently observed in ITC-treated cells, although it is likely that ITC exert their biological effects via modulation of multiple downstream effectors^(14,19–22). Metabolites of ITC are also likely to be involved, potentially via effects on chromatin remodelling and gene expression⁽²³⁾.

We have previously investigated the effects of phenethyl ITC (PEITC) on the response of cancer cells to hypoxia⁽²⁴⁾. PEITC is derived from the glucosinolate gluconasturtiin, which is found at particularly high levels in watercress (*Nasturtium officinale* or *Rorippa nasturtium-aquaticum*). We demonstrated that PEITC interfered with the ability of hypoxia to activate hypoxia-inducible factor (HIF)⁽²⁴⁾, a key transcription factor that mediates cellular responses to low pO₂^(25,26). Activation of HIF leads to increased transcription of a wide range of genes involved in angiogenesis (e.g. vascular endothelial growth factor), apoptosis (e.g. Bcl-2/adenovirus E1B 19-kDa protein-interacting protein 3) and metabolism (e.g. GLUT1). Inhibition of HIF may be important for the anti-cancer effects of PEITC, since PEITC and other ITC have been shown to possess anti-angiogenic activity *in vitro* and *in vivo*^(27–29), and angiogenesis plays a role in tumourigenesis, enabling growth of nascent tumours beyond a small size limit dictated by the perfusion distance of O₂ away from blood vessels⁽³⁰⁾.

The mechanism by which PEITC inhibits HIF activity appears to involve inhibition of HIF1 α mRNA translation⁽²⁴⁾. We⁽²⁴⁾ and others⁽³¹⁾ have demonstrated that PEITC decreases the levels of phosphorylation of 4E binding protein 1 (4E-BP1), and this may play an important role in the modulation of HIF1 α mRNA translation. The translation of HIF1 α mRNA is highly dependent on the eIF4E translation factor, which is, in turn, regulated by 4E-BP1⁽³²⁾. Dephosphorylation of 4E-BP1 facilitates its interaction with 4E-BP1, leading to

decreased translation of RNA such as HIF1 α mRNA. Hu *et al.*⁽³¹⁾ have also demonstrated that PEITC-induced cell death is reversed by overexpression of eIF4E. Thus, 4E-BP1 may be a key target for PEITC-associated anti-cancer effects, leading to the loss of both growth- and angiogenesis-promoting pathways. PEITC decreases the phosphorylation of 4E-BP1 on multiple sites, including Thr⁷⁰, Ser⁶⁵ and Thr^{37/46}^(24,31).

A small number of studies have investigated the *in vivo* effects of watercress consumption, generally, on pathways of carcinogen metabolism and oxidative stress. Pharmacokinetic analysis has demonstrated rapid absorption of PEITC into the blood with a mean maximal plasma concentration (C_{max}) of 928 nM after the ingestion of 100 g watercress⁽³³⁾. Hecht *et al.*^(34,35) have demonstrated that dietary intake of watercress increased urinary metabolites of the tobacco-specific lung carcinogens 4-(methylnitrosamino)-1-(3-pyridyl)-1-butanone and cotinine in smokers. Gill *et al.*⁽³⁶⁾ have demonstrated reduced levels of basal and hydrogen peroxide-induced DNA damage in peripheral blood lymphocytes following daily intake of 85 g of watercress for 8 weeks, associated with modest increases in erythrocyte superoxide dismutase and glutathione peroxidase 1 activity in specific cohort individuals carrying the *GSTM1*1* allele⁽³⁷⁾. A very recent study⁽²¹⁾ demonstrated reduced immunoreactivity of the proinflammatory cytokine MIF in plasma following the ingestion of a single 50 g portion of watercress.

Since decreased 4E-BP1 modulation has been functionally linked to *in vitro* anti-cancer effects of PEITC^(24,31), we investigated the effects of watercress extract on cancer cell growth inhibition and HIF activity. We also performed a small pilot study to determine whether dietary intake of watercress was sufficient to modulate 4E-BP1 phosphorylation *in vivo*.

Experimental methods

Cell culture

Human MCF7 breast cancer cells were obtained from the American Type Culture Collection (Manassas, VA, USA), and were maintained in Dulbecco's Modified Eagle's medium (Lonza Group Limited, Basel, Switzerland) supplemented with 10 % (v/v) fetal calf serum (PAA Laboratories, Yeovil, UK), 1 mM L-glutamine and penicillin/streptomycin (Lonza group Limited). PEITC and desferrioxamine (DFO) were obtained from Sigma Chemicals (Poole, UK). HIF reporter assays were performed as described previously⁽²⁴⁾. Peripheral blood mononuclear cells (PBMC) were isolated using Lymphoprep (Axis-Shield Diagnostics, Dundee, UK) according to the manufacturer's instructions. Growth inhibition assays were performed in duplicate as described previously⁽²⁴⁾. LY294002 was obtained from Sigma Chemicals.

Watercress extracts

Watercress samples were snap frozen in liquid N₂ before being ground to a fine powder using a pestle and mortar. Ground watercress (1 g) was decanted into a QIAshredder homogeniser (Qiagen, Crawley, UK), and incubated at room temperature for 1 h. Samples were centrifuged at 16 000 g for 6 min to collect the crude watercress extract.

Analysis of 4E binding protein 1 phosphorylation

The analysis of 4E-BP1 phosphorylation was done by single cell flow cytometry. PBMC were washed in ice-cold Roswell Park Memorial Institute 1640 medium (Invitrogen Limited, Paisley, UK) and resuspended in 1 ml of Roswell Park Memorial Institute 1640 medium. Cytotfix buffer (BD Biosciences, Oxford, UK; 1 ml) was added, and the cells were incubated for 10 min at 37°C before storage at -80°C, as per the manufacturer's

instructions. On the day of the analysis, samples were thawed, and the cells were washed with flow cytometry buffer (BD Biosciences). Cells were resuspended in 1 ml of Phosflow Permeabilisation buffer III (BD Biosciences), and were incubated on ice for 30 min. Cells were then washed twice with Stain Buffer (BD Biosciences), collected by centrifugation and resuspended in 1 ml of Stain Buffer containing 100 µl of phycoerythrin-conjugated anti-4E-BP1 antibody (Thr^{37/46} phospho-specific) (BD Biosciences). Unstained cells were analysed as controls. Cells were incubated in the dark at room temperature for 30 min, washed with Stain Buffer and resuspended in 500 µl of the same buffer before flow cytometry. Flow cytometry was performed using the FL2 channel (585 nm) on a BD Biosciences FACSCalibur. An average of 785 000 total events was collected for each sample. For 4E-BP1 fluorescence, we measured the proportion of monocytes with fluorescence values greater than those of unstained controls.

The analysis of 4E-BP1 phosphorylation was performed on cells gated on the basis of their forward scatter (FSC)/side scatter (SSC) properties. We found considerable variation in the FSC/SSC profiles of isolated PBMC following fixation and immunostaining. The impact of fixatives used for intra-cellular staining with phospho-specific antibodies on scatter profiles has been described previously⁽³⁸⁾. We separately gated on lymphocytes (abundant population with low FSC and SSC; Fig. 2(a)) and a population with increased FSC/SSC (see oval gate in Figs. 2(a) and 4) that we tentatively described as 'monocytes' (see 'Discussion'). In some analyses, we detected a third population of cells (to the right of the lymphocytes) that we believe is due to variation in fixation. When present, these were excluded from the analysis. Some samples also contained relatively high amounts of dead cells with very low FSC (see Fig. 4). Again, these were excluded from the analysis.

In vivo study

The watercress feeding study was based on the previous work done by Ji *et al.*⁽³³⁾, who studied the plasma pharmacokinetics of PEITC following the ingestion of 100 g of watercress in four normal participants. The present study was conducted according to the guidelines laid down in the Declaration of Helsinki, and all procedures involving human participants were approved by the Southampton and South West Research Ethics Committee (Ref 08/H0504/86). Written informed consent was obtained from all the participants. Potential participants were identified using the Winchester and Andover Breast Unit database of patients, where clinical and diagnostic data are routinely compiled for audit purposes using criteria defined by the British Association for Surgical Oncology. Although the participants had previously been treated for breast cancer, they were considered free of clinically detectable disease for a minimum of 2 years and were otherwise healthy. None of the women was taking any pharmacological medications or herbal supplements.

Participants were requested to avoid the following foods for 3 d before the study day to exclude known sources of glucosinolates from the diet: cabbage, brussels sprouts, broccoli, calabrese, cauliflower, turnip, swede, rutabaga, kohlrabi, kale, chinese kale, sea kale, curly kale, collard, pak choi, radish, horseradish, mustard, mustard greens, mustard leaf, wasabi, salad rocket, cress, watercress, capers, papaya (pawpaw) and nasturtium (Indian cress). The work of McNaughton & Marks⁽³⁹⁾ informed the compilation of the list of food items to be excluded from the diet. Food and drink diaries were completed to confirm avoidance of glucosinolate-containing food items; dietary histories were also taken to describe participants' usual glucosinolate intakes, although these data are not presented here. The participants fasted from midnight on the day of the study. Approximately, 8 h later the volunteers were cannulated, and a baseline blood sample (5 ml) was obtained. The participants then ate 80 g of watercress, obtained from a local grower and packager (Vitacress Salads Limited, Andover, UK). Eighty grams is the weight of a portion of fruit and vegetables defined by the WHO, and is equivalent to one of the five portions of fruit and

vegetables that describes one of UK's nutrition public health messages^(40,41). Blood samples were then obtained at approximately 7.5 min, 15 min, 30 min, 45 min, 1 h, 1.5 h, 2 h, 3 h, 4 h, 6 h and 8 h following the ingestion of the watercress meal. A 24 h blood sample was obtained by venipuncture. Blood samples were collected in heparinised glass tubes, and stored on ice before processing. Participants were free to take water throughout the study, and were allowed to eat and drink freely from 3 h after the ingestion of watercress meal, although glucosinolate-containing foods were eliminated until the end of the study.

We enrolled twelve women into the study, and analysed plasma PEITC concentrations and/or 4E-BP1 phosphorylation in samples from nine of these women (mean age 58 years, median age 56 years and range 48–82). For eight participants, we prepared both PBMC and plasma; samples were overlaid onto Lymphoprep and centrifuged at 800 g for 20 min. The plasma was recovered and stored at –80°C. For one participant (participant 5 in Table 1), we prepared only plasma; samples were centrifuged at 486 g for 15 min, and the plasma was recovered and stored at –80°C. Plasma PEITC concentrations were determined as described previously⁽³³⁾, with the exception that analyses were performed using 0.5 ml of plasma. PBMC were recovered from the interface, washed twice in ice-cold Roswell Park Memorial Institute 1640 medium and fixed using Cytotfix (BD Biosciences) as described earlier. Cells were frozen before the analysis, and all the samples from an individual participant were analysed in parallel. The analysis of 4E-BP1 phosphorylation was done as described earlier. The analysis was performed on samples from five participants. However, two were excluded due to technical failures, and another two were excluded because of poor viability of cells.

Results

Effect of watercress extract on cancer cell growth and hypoxia-inducible factor activity

PEITC has been shown to decrease cancer cell growth and to inhibit HIF activity, actions which have been linked to decreased 4E-BP1 phosphorylation^(24,31). Watercress is a particularly rich source of the PEITC precursor glucosinolate, gluconasturtiin, and we therefore investigated whether watercress extract could exert similar effects. Extracts were prepared by grinding watercress leaf, which releases myrosinase, the enzyme which catalyses the conversion of gluconasturtiin to PEITC. Watercress extract inhibited the growth of MCF7 breast cancer cells with an half maximal inhibitory concentration (IC₅₀) of 24.6 (SD 12.3) µl/ml (Fig. 1(a)).

To analyse the transcriptional activity of HIF, MCF7 cells were transfected with a HIF reporter construct and treated with the hypoxia mimetic DFO (Fig. 1(b)). DFO displaces Fe from prolyl hydroxylases leading to the stabilisation of HIF α proteins. DFO caused a strong induction of the HIF reporter construct, and this was inhibited in a dose-dependent manner by watercress extract, although this was statistically significant only at the highest concentration. Watercress extract also decreased the basal activity of the HIF reporter in the absence of DFO. As a control, we also analysed the effects of DFO and watercress extract on the activity of the SV40-promoter-based reporter plasmid pGL3-promoter. The activity of the control promoter was reduced following treatment with DFO, and was not affected by watercress extract under any condition. Therefore, similar to PEITC⁽²⁴⁾, watercress extract inhibits HIF activity in MCF7 breast cancer cells.

Phenethyl isothiocyanates decreases 4E binding protein 1 phosphorylation in peripheral blood mononuclear cells

Since PEITC decreases 4E-BP1 phosphorylation and this has been linked to HIF inhibition⁽²⁴⁾ and growth inhibition⁽³¹⁾, we selected 4E-BP1 phosphorylation as a potential biomarker to monitor *in vivo* exposure to PEITC. To explore the potential utility of 4E-BP1

phosphorylation as a biomarker, we first examined the levels of 4E-BP1 phosphorylation in PBMC and its modulation by PEITC.

PBMC were obtained from healthy donors and analysed by single cell flow cytometry. For these studies, we used an antibody that selectively reacted with 4E-BP1 phosphorylated on Thr^{37/46}, since it gave stronger staining compared with a 4E-BP1 Thr⁷⁰ phospho-specific antibody (data not shown). 4E-BP1 phosphorylation varied among different cell populations in peripheral blood based on FSC/SCC properties (see 'Experimental methods' for gating strategy). We readily detected phosphorylated 4E-BP1 in a population of cells that we tentatively described as 'monocytes', whereas the levels of phosphorylation were much lower in lymphocytes (Fig. 2). To confirm that immunostaining was specific for phosphorylated 4E-BP1, cells were treated with LY294002. LY294002 decreases 4E-BP1 phosphorylation via inhibition of Phosphoinositide 3 kinase, a key upstream positive regulator of 4E-BP1 phosphorylation via the phosphoinositide 3 kinase-AKT-mammalian target of rapamycin pathway⁽⁴²⁾. Treatment of cells with LY294002 reduced 4E-BP1 phosphorylation by 63 ± 13 % (mean \pm range of two experiments; Fig. 2). Consistent with previous data^(24,31), there was also a significant reduction in 4E-BP1 phosphorylation following *in vitro* treatment with PEITC (Fig. 2). Thus, 4E-BP1 phosphorylation appears to be a suitable biomarker to monitor *in vivo* responses to PEITC in PBMC.

Watercress consumption down-regulates 4E binding protein 1 phosphorylation *in vivo*

We performed a small feeding study to determine whether the ingestion of watercress was sufficient to modulate 4E-BP1 phosphorylation *in vivo*. We first analysed plasma concentrations of PEITC (Table 1). Similar to a previous study⁽³³⁾, there was a rapid increase in plasma PEITC, which, on average, reached a maximal concentration at 3 h (Fig. 3). The mean C_{\max} was 297 nM, although there was a wide interindividual variation (range 61–656 nM). Background PEITC concentrations before the ingestion of watercress were very low (generally <1 nM).

The analysis of 4E-BP1 phosphorylation was done in four participants (Fig. 4). All the participants showed a marked reduction in 4E-BP1 phosphorylation at 6 and 8 h following the ingestion of watercress compared with pre-watercress meal values. Twenty-four hour data were available for three participants; in one participant, 4E-BP1 phosphorylation was maintained at a low level until this time point, whereas recovery of 4E-BP1 phosphorylation was observed in the other two participants. The decreases in 4E-BP1 phosphorylation at 6 and 8 h were highly statistically significant ($P=0.001$ and 0.002 , respectively; Student's *t* test compared to pre-watercress samples). There was a considerable variation in the levels of 4E-BP1 phosphorylation before 6 h within individual subjects. It is possible that this is due to technical variation in transport and sample processing, and the average level of 4E-BP1 phosphorylation was not significantly altered at these time points. Thus, the analysis of 4E-BP1 Thr^{37/46} phosphorylation suggests a significant decrease at 6 and 8 h after the ingestion of watercress.

Discussion

Numerous epidemiological studies have suggested that high dietary intake of cruciferous vegetables is associated with reduced cancer risk, and *in vitro* studies have indicated that reduced 4E-BP1 phosphorylation may be an important mechanism contributing to anti-cancer effects of PEITC^(24,31). In this work, we have performed a pilot study to determine whether the ingestion of watercress, a rich dietary source of PEITC, is sufficient to modulate 4E-BP1 phosphorylation levels *in vivo*. The mechanisms by which PEITC decreases 4E-BP1 phosphorylation are not known. Modulation of upstream regulators, including

mammalian target of rapamycin and phosphatase and tensin homologue (PTEN), both of which contain redox-sensitive cysteine residues, may play a role^(43,44).

We selected 4E-BP1 phosphorylation as a potential molecular biomarker for several reasons. First, modulation of 4E-BP1 phosphorylation by PEITC has been demonstrated in multiple cell types^(24,31). Secondly, modulation of 4E-BP1 phosphorylation has been mechanistically linked to both growth-inhibitory and anti-angiogenic effects of PEITC^(24,31). Thirdly, it was possible to measure 4E-BP1 phosphorylation on a single cell basis using a quantitative flow cytometry assay. Finally, we were able to measure 4E-BP1 phosphorylation in PBMC, a relatively accessible tissue source suitable for repeat sampling. However, it is possible that other watercress-derived ITC, PEITC metabolites or unrelated bioactives may also modulate 4E-BP1 phosphorylation. We therefore cannot exclude the possibility that other compounds are involved in the *in vitro* and *in vivo* effects of watercress. Although we consider it unlikely, all the participants ate the watercress at the same time of day, and we therefore also cannot exclude the possibility that decreased 4E-BP1 phosphorylation is unrelated to watercress consumption, and may reflect diurnal modulation of activity.

Although flow cytometric analysis of 4E-BP1 phosphorylation appeared to be a promising approach for the evaluation of *in vivo* responses to ITC, this pilot study identified a number of key technical challenges that should be addressed in future studies. First, intracellular antibody staining requires cell fixation, and this resulted in a large variation in the scatter properties of the cells. The difficulties caused by variation in scatter properties, specifically in the context of staining with phospho-specific antibodies, have been discussed previously⁽³⁸⁾. Our gating strategy excluded lymphocytes and encompassed a population that we tentatively identified as monocytes. However, in future studies, it would be important to combine intracellular staining with specific surface markers to unambiguously identify specific cell subpopulations, and to improve the flow cytometric analyses. Secondly, we observed significant levels of cell death in some samples. The clinical and laboratory sites were geographically separate, and transportation between these sites may have resulted in increased cell death. For technically demanding analyses, such as phospho-specific flow cytometry, we recommend that samples be processed with the minimum of delay.

Flow cytometric analysis demonstrated a statistically significant reduction in 4E-BP1 Thr^{37/46} phosphorylation at 6 and 8 h following the consumption of watercress in 4/4 participants studied. Caution is required in interpreting our data, since there was significant variability in the levels of 4E-BP1 phosphorylation within individual samples before this time point. However, these differences disappeared when results from the four individuals were combined, whereas the consistent down-regulation of 4E-BP1 phosphorylation at 6 and 8 h was highly statistically significant. This suggests that dietary intake of a single 80 g portion of watercress is sufficient to modulate this potential anti-cancer pathway. These results are consistent with those reported previously from studies investigating the effects of watercress consumption on carcinogen metabolism/oxidative stress, although it should be noted that these studies involved repeated ingestion of watercress over a period of 3 d–8 weeks^(34–37). Recently, Brown *et al.*^(20,21) have shown that the ingestion of a single 50 g portion of watercress is associated with reduced plasma immunoreactivity of the proinflammatory cytokine macrophage migration inhibitory factor, which is a direct target for covalent modification by PEITC.

As part of the present study, we also analysed the plasma concentrations of PEITC. These results are generally in line with those reported previously by Ji *et al.*⁽³³⁾, who analysed the plasma concentrations of PEITC in four participants following the consumption of 100 g of watercress. Both the studies showed a rapid increase in plasma PEITC concentrations,

peaking at 2–3 h, followed by a decline to near background levels at 24 h. However, the C_{\max} values in this study were generally lower (297 v. 929 nM on average), and showed much more interindividual variation in the present study. Multiple variables are likely to contribute to these differences.

The lower average C_{\max} at least partly reflects the smaller portion size selected for the present study (80 v. 100 g), and although the difference in intake is small, it is not clear whether PEITC accumulation in the plasma is proportionate to dose. It is also possible that differences in glucosinolate content of the crop will have contributed to the variation in mean C_{\max} concentrations between the studies, since it is known that differences in sunlight exposure, temperature and added fertilisers can all influence gluconasturtiin production in watercress^(45,46). The average age of the participants in the present study was relatively high, and age-related changes in absorption may have also impacted on the overall C_{\max} and contributed to interindividual variation. Finally, glutathione-*S*-transferase variants have previously been demonstrated to modulate ITC metabolism and potential chemopreventive effects, and it is also possible that genetic variation may have contributed to some of the differences. Such age-, genetic- and crop-related differences are all ‘real world’ variables that are likely to interact to determine exposure, and thus complicate the analysis of biological effects of plant-derived agents.

A key question is why dietary intake of watercress may be sufficient to modulate 4E-BP1 phosphorylation, although plasma concentrations may not achieve the concentrations that are required to effect this pathway *in vitro* (typically 1–5 μM ^(24,31)). One possible explanation lies in the interaction of PEITC with GSH. Efflux of ITC conjugates, extracellular hydrolysis and reuptake of ITC lead to marked accumulation of intracellular ITC^(11,12). Thus, the intracellular concentration of PEITC in monocytes may be much higher than what is predicted from the plasma concentrations. Similar interactions may account for the time lag between peak plasma concentration and inhibition of 4E-BP1 phosphorylation. It will be important to investigate further how differences in glutathione levels and metabolism alter cellular accumulation of ITC and their metabolites, since GSH levels are altered in many cancer cells⁽⁴⁷⁾. As discussed earlier, it is also important to bear in mind the alternate possibilities. For example, *in vivo* modulation of 4E-BP1 phosphorylation may be unrelated to or only partially dependent on PEITC, and may be due to other bioactives derived from watercress. For example, Brown *et al.*⁽²¹⁾ recently reported that the mean plasma concentration of total ITC and dithiocarbamate metabolites reached approximately 1.5 μM at 2 h following the ingestion of a 50 g portion of watercress. At present, it is not known whether dithiocarbamate metabolites may contribute to the modulation of 4E-BP1 phosphorylation, either directly or following conversion to ITC.

In summary, we have performed a small pilot study to investigate the feasibility of measuring 4E-BP1 phosphorylation as a biomarker to monitor *in vivo* effects of PEITC. We tentatively conclude that flow cytometry may be a suitable approach to measure changes in 4E-BP1 phosphorylation following the ingestion of watercress. However, further studies are required with larger sample sizes to test this more rigorously.

Acknowledgments

We thank the medical assistants and ward staff at the Royal Hampshire County Hospital for their assistance and expertise. We particularly thank the women who volunteered for the *in vivo* study. This work was supported by the Watercress Alliance and BBSRC. G. P. and B. M. P. were the principal investigators of the study, and were responsible for the study design and supervision. M. E. M. was responsible for the design and supervision of LC/MS/MS analysis. S. S. A. A. was responsible for the analysis of 4E-BP1 phosphorylation. B. E. C. was responsible for the analysis of HIF activity and cell growth inhibition assays. U. T. was responsible for LC/MS/MS analysis. G. P. wrote the manuscript, which was reviewed by all the authors.

Abbreviations

C_{\max}	maximal plasma concentration
DFO	desferrioxamine
4E-BP1	4E binding protein 1
FSC	forward scatter
HIF	hypoxia-inducible factor
ITC	isothiocyanates
PBMC	peripheral blood mononuclear cells
PEITC	phenethyl ITC
SSC	side scatter

References

- Higdon JV, Delage B, Williams DE, et al. Cruciferous vegetables and human cancer risk: epidemiologic evidence and mechanistic basis. *Pharmacol Res.* 2007; 55:224–236. [PubMed: 17317210]
- Zhang Y, Yao S, Li J. Vegetable-derived isothiocyanates: anti-proliferative activity and mechanism of action. *Proc Nutr Soc.* 2006; 65:68–75. [PubMed: 16441946]
- Zhang Y. Cancer-preventive isothiocyanates: measurement of human exposure and mechanism of action. *Mutat Res.* 2004; 555:173–190. [PubMed: 15476859]
- Brennan P, Hsu CC, Moulán N, et al. Effect of cruciferous vegetables on lung cancer in patients stratified by genetic status: a mendelian randomisation approach. *Lancet.* 2005; 366:1558–1560. [PubMed: 16257343]
- London SJ, Yuan JM, Chung FL, et al. Isothiocyanates, glutathione *S*-transferase M1 and T1 polymorphisms, and lung-cancer risk: a prospective study of men in Shanghai, China. *Lancet.* 2000; 356:724–729. [PubMed: 11085692]
- Zhao B, Seow A, Lee EJ, et al. Dietary isothiocyanates, glutathione *S*-transferase -M1, -T1 polymorphisms and lung cancer risk among Chinese women in Singapore. *Cancer Epidemiol Biomarkers Prev.* 2001; 10:1063–1067. [PubMed: 11588132]
- Spitz MR, Duphorne CM, Detry MA, et al. Dietary intake of isothiocyanates: evidence of a joint effect with glutathione *S*-transferase polymorphisms in lung cancer risk. *Cancer Epidemiol Biomarkers Prev.* 2000; 9:1017–1020. [PubMed: 11045782]
- Steck SE, Hebert JR. GST polymorphism and excretion of heterocyclic aromatic amine and isothiocyanate metabolites after Brassica consumption. *Environ Mol Mutagen.* 2009; 50:238–246. [PubMed: 19197987]
- Clarke JD, Dashwood RH, Ho E. Multi-targeted prevention of cancer by sulforaphane. *Cancer Lett.* 2008; 269:291–304. [PubMed: 18504070]
- Hecht SS. Inhibition of carcinogenesis by isothiocyanates. *Drug Metab Rev.* 2000; 32:395–411. [PubMed: 11139137]
- Zhang Y. Role of glutathione in the accumulation of anticarcinogenic isothiocyanates and their glutathione conjugates by murine hepatoma cells. *Carcinogenesis.* 2000; 21:1175–1182. [PubMed: 10837007]
- Xu K, Thornalley PJ. Involvement of glutathione metabolism in the cytotoxicity of the phenethyl isothiocyanate and its cysteine conjugate to human leukaemia cells *in vitro*. *Biochem Pharmacol.* 2001; 61:165–177. [PubMed: 11163331]
- Trachootham D, Zhou Y, Zhang H, et al. Selective killing of oncogenically transformed cells through a ROS-mediated mechanism by beta-phenylethyl isothiocyanate. *Cancer Cell.* 2006; 10:241–252. [PubMed: 16959615]

14. Mi L, Wang X, Govind S, et al. The role of protein binding in induction of apoptosis by phenethyl isothiocyanate and sulforaphane in human non-small lung cancer cells. *Cancer Res.* 2007; 67:6409–6416. [PubMed: 17616701]
15. Dinkova-Kostova AT, Holtzclaw WD, Cole RN, et al. Direct evidence that sulfhydryl groups of Keap1 are the sensors regulating induction of phase 2 enzymes that protect against carcinogens and oxidants. *Proc Natl Acad Sci U S A.* 2002; 99:11908–11913. [PubMed: 12193649]
16. Keum YS, Owuor ED, Kim BR, et al. Involvement of Nrf2 and JNK1 in the activation of antioxidant responsive element (ARE) by chemopreventive agent phenethyl isothiocyanate (PEITC). *Pharm Res.* 2003; 20:1351–1356. [PubMed: 14567627]
17. McMahon M, Itoh K, Yamamoto M, et al. Keap1-dependent proteasomal degradation of transcription factor Nrf2 contributes to the negative regulation of antioxidant response element-driven gene expression. *J Biol Chem.* 2003; 278:21592–21600. [PubMed: 12682069]
18. Xu C, Yuan X, Pan Z, et al. Mechanism of action of isothiocyanates: the induction of ARE-regulated genes is associated with activation of ERK and JNK and the phosphorylation and nuclear translocation of Nrf2. *Mol Cancer Ther.* 2006; 5:1918–1926. [PubMed: 16928811]
19. Mi L, Xiao Z, Hood BL, et al. Covalent binding to tubulin by isothiocyanates. A mechanism of cell growth arrest and apoptosis. *J Biol Chem.* 2008; 283:22136–22146. [PubMed: 18524779]
20. Cross JV, Rady JM, Foss FW, et al. Nutrient isothiocyanates covalently modify and inhibit the inflammatory cytokine macrophage migration inhibitory factor (MIF). *Biochem J.* 2009; 423:315–321. [PubMed: 19723024]
21. Brown KK, Blaikie FH, Smith RA, et al. Direct modification of the pro-inflammatory cytokine MIF by dietary isothiocyanates. *J Biol Chem.* 2009; 284:32425–32433. [PubMed: 19776019]
22. Cross JV, Foss FW, Rady JM, et al. The isothiocyanate class of bioactive nutrients covalently inhibit the MEKK1 protein kinase. *BMC Cancer.* 2007; 7:183. [PubMed: 17894894]
23. Myzak MC, Karplus PA, Chung FL, et al. A novel mechanism of chemoprotection by sulforaphane: inhibition of histone deacetylase. *Cancer Res.* 2004; 64:5767–5774. [PubMed: 15313918]
24. Wang XH, Cavell BE, Syed Alwi SS, et al. Inhibition of hypoxia inducible factor by phenethyl isothiocyanate. *Biochem Pharmacol.* 2009; 78:261–272. [PubMed: 19376091]
25. Rankin EB, Giaccia AJ. The role of hypoxia-inducible factors in tumorigenesis. *Cell Death Differ.* 2008; 15:678–685. [PubMed: 18259193]
26. Weidemann A, Johnson RS. Biology of HIF-1 α . *Cell Death Differ.* 2008; 15:621–627. [PubMed: 18259201]
27. Xiao D, Singh SV. Phenethyl isothiocyanate inhibits angiogenesis *in vitro* and *ex vivo*. *Cancer Res.* 2007; 67:2239–2246. [PubMed: 17332354]
28. Bertl E, Bartsch H, Gerhauser C. Inhibition of angiogenesis and endothelial cell functions are novel sulforaphane-mediated mechanisms in chemoprevention. *Mol Cancer Ther.* 2006; 5:575–585. [PubMed: 16546971]
29. Jackson SJ, Singletary KW, Venema RC. Sulforaphane suppresses angiogenesis and disrupts endothelial mitotic progression and microtubule polymerization. *Vascul Pharmacol.* 2007; 46:77–84. [PubMed: 16938492]
30. Carmeliet P, Jain RK. Angiogenesis in cancer and other diseases. *Nature.* 2000; 407:249–257. [PubMed: 11001068]
31. Hu J, Straub J, Xiao D, et al. Phenethyl isothiocyanate, a cancer chemopreventive constituent of cruciferous vegetables, inhibits cap-dependent translation by regulating the level and phosphorylation of 4E-BP1. *Cancer Res.* 2007; 67:3569–3573. [PubMed: 17440067]
32. Yee Koh M, Spivak-Kroizman TR, Powis G. HIF-1 regulation: not so easy come, easy go. *Trends Biochem Sci.* 2008; 33:526–534. [PubMed: 18809331]
33. Ji Y, Morris ME. Determination of phenethyl isothiocyanate in human plasma and urine by ammonia derivatization and liquid chromatography–tandem mass spectrometry. *Anal Biochem.* 2003; 323:39–47. [PubMed: 14622957]
34. Hecht SS, Carmella SG, Murphy SE. Effects of watercress consumption on urinary metabolites of nicotine in smokers. *Cancer Epidemiol Biomarkers Prev.* 1999; 8:907–913. [PubMed: 10548320]

35. Hecht SS, Chung FL, Richie JP, et al. Effects of watercress consumption on metabolism of a tobacco-specific lung carcinogen in smokers. *Cancer Epidemiol Biomarkers Prev.* 1995; 4:877–884. [PubMed: 8634661]
36. Gill CI, Haldar S, Boyd LA, et al. Watercress supplementation in diet reduces lymphocyte DNA damage and alters blood antioxidant status in healthy adults. *Am J Clin Nutr.* 2007; 85:504–510. [PubMed: 17284750]
37. Hofmann T, Kuhnert A, Schubert A, et al. Modulation of detoxification enzymes by watercress: *in vitro* and *in vivo* investigations in human peripheral blood cells. *Eur J Nutr.* 2009; 48:483–491. [PubMed: 19636603]
38. Krutzik PO, Irish JM, Nolan GP, et al. Analysis of protein phosphorylation and cellular signaling events by flow cytometry: techniques and clinical applications. *Clin Immunol.* 2004; 110:206–221. [PubMed: 15047199]
39. McNaughton SA, Marks GC. Development of a food composition database for the estimation of dietary intakes of glucosinolates, the biologically active constituents of cruciferous vegetables. *Br J Nutr.* 2003; 90:687–697. [PubMed: 13129476]
40. World Health Organization. Joint WHO/FAO Expert Consultation. Geneva: WHO; 2003. Diet, Nutrition and the Prevention of Chronic Diseases. WHO Technical Report Series no. 916
41. Department of Health. [accessed 19 October 2009] 5 A DAY general information. 2009. <http://www.dh.gov.uk/en/Publichealth/Healthimprovement/FiveADay/FiveADaygeneralinformation/index.htm>
42. Proud CG. Signalling to translation: how signal transduction pathways control the protein synthetic machinery. *Biochem J.* 2007; 403:217–234. [PubMed: 17376031]
43. Dames SA, Mulet JM, Rathgeb-Szabo K, et al. The solution structure of the FATC domain of the protein kinase target of rapamycin suggests a role for redox-dependent structural and cellular stability. *J Biol Chem.* 2005; 280:20558–20564. [PubMed: 15772072]
44. Salmeen A, Barford D. Functions and mechanisms of redox regulation of cysteine-based phosphatases. *Antioxid Redox Signal.* 2005; 7:560–577. [PubMed: 15890001]
45. Engelen-Eigles G, Holden G, Cohen JD, et al. The effect of temperature, photoperiod, and light quality on gluconasturtiin concentration in watercress (*Nasturtium officinale* R. Br.). *J Agric Food Chem.* 2006; 54:328–334. [PubMed: 16417287]
46. Kopsell DA, Barickman TC, Sams CE, et al. Influence of nitrogen and sulfur on biomass production and carotenoid and glucosinolate concentrations in watercress (*Nasturtium officinale* R. Br.). *J Agric Food Chem.* 2007; 55:10628–10634. [PubMed: 18052091]
47. Estrela JM, Ortega A, Obrador E. Glutathione in cancer biology and therapy. *Crit Rev Clin Lab Sci.* 2006; 43:143–181. [PubMed: 16517421]




Fig. 1. Inhibition of cancer cell growth and hypoxia-inducible factor (HIF)-dependent transcription by watercress extract. (a) Representative growth inhibition experiment. MCF7 cells were incubated with the indicated concentrations of watercress extract. After 6 d, relative cell numbers were determined using the CellTiter 96[®] AQueous One Solution reagent. Results are derived from means of duplicate wells. (b) MCF7 cells were transfected with (i) pGL2-9TK-HRE or (ii) control pGL3-promoter reporter constructs, and were treated with the indicated concentrations of watercress extract (■) or with equivalent amounts of water (□). HIF activity was induced by treating the cells with desferrioxamine (DFO) (100 μM), and luciferase activity was analysed after 24 h. Data indicate mean with their standard errors for luciferase activity relative to control cells (i.e. cell in the absence of DFO and with water equivalent to the lowest concentration of watercress extract tested was set to 1.0) derived from three independent experiments, each performed in triplicate. Statistically significant differences (Student's *t* test) between dimethyl sulphoxide- and watercress-treated cells are shown. All other differences were not statistically significant.




Fig. 2. Analysis of 4E binding protein 1 (4E-BP1) phosphorylation in peripheral blood mononuclear cells (PBMC). PBMC were isolated from healthy individuals and analysed by flow cytometry using a Thr^{37/46} 4E-BP1 phosphorylation-specific antibody. (a) Forward scatter (FSC)/side scatter (SSC) plot showing gating of lymphocytes (lower left) and a population of cells with increased FSC and SSC, which we tentatively classed as monocytes (oval gate). (b–e) Fluorescence intensity of ‘monocytes’ (b)–(d) and lymphocytes (e). In (b) and (e), black line indicates unstained control, and the grey line indicates antibody-stained cells. In (c), the grey line indicates antibody-stained cells, and the black line indicates cells treated with LY294002 (100 μ M) for 2 h before staining with the 4E-BP1 antibody. In (d), the grey line indicates antibody-stained cells, and the black line indicates antibody-stained cells treated with phenethyl isothiocyanates (20 μ M) for 2 h before staining with the 4E-BP1 antibody. Data are representative of at least two independent experiments. FL2-H, fluorescence pulse height.




Fig. 3. Analysis of plasma phenethyl isothiocyanates (PEITC) concentrations following the consumption of watercress. Plasma concentration of PEITC was determined at various time points following the consumption of 80 g watercress. Data obtained from four representative subjects are shown.




Fig. 4. Analysis of 4E binding protein 1 (4E-BP1) phosphorylation following the consumption of watercress. 4E-BP1 phosphorylation was analysed by flow cytometry in peripheral blood-derived monocytes at various time points following the consumption of 80 g watercress. (a) Representative data obtained from two participants showing forward scatter (FSC)/side scatter (SSC) plots with ‘monocyte’ gate (a) and fluorescence intensity (b) of unstained control cells and stained cells before (T_0) and 8 h (T_{480}) after the consumption of watercress. (b) Graphs showing (i and ii) the levels of 4E-BP1 phosphorylation in two representative participants (1 and 4) and (iii) the mean with their levels of 4E-BP1 phosphorylation in all four subjects following the consumption of watercress. In (iii), the level of 4E-BP1 phosphorylation at T_0 was set to 1.0 to allow comparison between individuals. FL2-H, fluorescence pulse height. Mean values were significantly different compared to T_0 are indicated (Student’s *t* test): * $P=0.001$, ** $P=0.002$.

Table 1

Analysis of plasma phenethyl isothiocyanates concentrations following the consumption of watercress

Participant	C_{\max} (nM)	t_{\max} (min)
1	104	120
2	61	125
3	117	180
4	164	480
5	375	185
6	384	90
7	209	120
8	656	140
9	604	180
Mean	297	180

C_{\max} , maximal plasma concentration; t_{\max} , time to reach C_{\max} .