

Draft Genome Sequence of the Ascomycete *Phaeoacremonium aleophilum* Strain UCR-PA7, a Causal Agent of the Esca Disease Complex in Grapevines

Barbara Blanco-Ulate,^{a,b} Philippe Rolshausen,^c Dario Cantu^a

Department of Viticulture and Enology, University of California, Davis, Davis, California, USA^a; Department of Plant Sciences, University of California, Davis, Davis, California, USA^b; Department of Botany & Plant Sciences, University of California, Riverside, Riverside, California, USA^c

Grapevine infections by *Phaeoacremonium aleophilum* in association with other pathogenic fungi cause complex and economically important vascular diseases. Here we present the first draft of the *P. aleophilum* genome sequence, which comprises 624 scaffolds with a total length of 47.5 Mb (L₅₀, 45; N₅₀, 336 kb) and 8,926 predicted protein-coding genes.

Received 10 May 2013 Accepted 14 May 2013 Published 27 June 2013

Citation Blanco-Ulate B, Rolshausen P, Cantu D. 2013. Draft genome sequence of the ascomycete *Phaeoacremonium aleophilum* strain UCR-PA7, a causal agent of the esca disease complex in grapevines. *Genome Announc.* 1(3):e00390-13. doi:10.1128/genomeA.00390-13.

Copyright © 2013 Blanco-Ulate et al. This is an open-access article distributed under the terms of the [Creative Commons Attribution 3.0 Unported license](https://creativecommons.org/licenses/by/3.0/).

Address correspondence to Dario Cantu, dacantu@ucdavis.edu.

The esca disease complex of grapevines refers to five syndromes: brown wood-streaking, petri disease, young esca, esca, and esca proper (1), which are caused by the ascomycetes *Phaeoacremonium aleophilum* W. Gams, P. W. Crous, M. J. Wingfield, and L. Mugnai (teleomorph, *Togninia minima*), *Phaeoacremonium aleophilum* (1–3). Disease symptoms include, internally, wood discoloration, streaking, and vascular necrosis (3) and, externally, leaf chlorosis and necrotic stripes, berry black spots, decline in vigor and yield, and in severe cases plant death (3, 4).

The effective colonization of the host tissues appears to depend on *P. aleophilum* competence to produce phytotoxic metabolites (5, 6), overcome host preformed and inducible barriers (7, 8), degrade the plant cell wall (CW) (8, 9), and cooperate with other pathogens during infection (1, 3).

P. aleophilum strain UCR-PA7 was collected from the margin of a grapevine (*Vitis vinifera* cv. “Thomson”) wood canker in a commercial vineyard (Fresno County, CA) in October 2011. UCR-PA7 was hyphal-tip purified and taxonomically characterized using DNA markers (10). DNA was extracted using a modified cetyltrimethylammonium bromide (CTAB) protocol (11) and sequenced to an estimated median coverage of 103× using the Illumina HiSeq2000 platform. Using CLC Genomics Workbench v6.0, we assembled 99% of the 4.7 × 10⁷ trimmed (Q ≥ 30) and contaminant-filtered reads into 624 scaffolds (N₅₀, 336 kb; L₅₀, 45; G + C content, 49.66%; gaps, 83 kb) with a total length of 47.5 Mb. Assembly parameters were optimized based on the degree of completeness of the gene space estimated with CEGMA (12). The UCR-PA7 genome was estimated to be >97% complete based on the mapping of 248 low-copy and conserved core eukaryotic genes (CEGs) (12).

The gene finder Augustus (13) was trained using the CEG structures determined by CEGMA and identified *ab initio* 8,926 complete protein-coding genes on repeat-masked scaffolds (RepeatMasker [14]). A total of 97% of these genes have homologs in other ascomycetes (BLASTp, E value ≤ 10⁻³). We identified 658

potentially secreted proteins (SignalP-v4.0 [15]), of which at least 23% consist of putative plant CW-degrading enzymes based on homology to proteins in the CAZy database (16). Among these, 17 cellulases (GH3s, GH5s, GH6s, GH7s, and GH45s), 12 hemicellulases (GH10s, GH11s, GH31s, GH29s, GH67s, and GH115s), 21 pectin-degrading enzymes (GH28s, GH78s, PL1s, PL3s, PL4s, PL9s, CE8s, and CE12s), 12 callose-degrading enzymes (GH55s), and 1 cutinase (CE5) might play important roles during tissue colonization and systemic infection.

We also detected 79 cytochrome P450 monooxygenases, 2 laccases, and 2 lignin peroxidases, supporting the ability of *P. aleophilum* to degrade lignocellulose (2, 3, 8). However, the number of putative lignin-degrading proteins in the *P. aleophilum* genome is smaller than that in other wood-decay fungi previously described (e.g., *Neofusicoccum parvum*, 212 P450s [17]; *Eutypa lata*, 205 P450s [18]; and *Phanerochaete carnosa*, 266 P450s [19]), which suggests that synergism with other vascular pathogens during plant infection may favor the effective breakdown of lignified tissues (4, 8).

Nucleotide sequence accession numbers. This Whole-Genome Shotgun project has been deposited at DDBJ/EMBL/GenBank under the accession no. [AORD000000000](https://www.ncbi.nlm.nih.gov/nuccore/AORD000000000). The version described in this paper is the first version, [AORD010000000](https://www.ncbi.nlm.nih.gov/nuccore/AORD010000000).

ACKNOWLEDGMENTS

This work was supported by funding to D.C. from the College of Agricultural and Environmental Sciences (UC, Davis) and to P.R. from the College of Natural and Agricultural Sciences (UC, Riverside). Support to B.B.-U. was provided by the Consejo Nacional de Ciencia y Tecnología (Ministerio de Ciencia y Tecnología, Costa Rica).

We thank Henriette O’Geen (UC Davis Genome Center) and Abraham Morales for technical assistance.

REFERENCES

1. Surico G, Mugnai L, Marchi G. 2008. The esca disease complex, p 119–136. In Ciancio A, Mukerji KG (ed), *Integrated management of dis-*

- eases caused by fungi, phytoplasma and bacteria. Springer Verlag, Dordrecht, The Netherlands.
2. Mostert L, Groenewald JZ, Summerbell RC, Gams W, Crous PW, Samson RA. 2006. Taxonomy and pathology of *Togninia* (Diaporthales) and its phaeoacremonium anamorphs, vol. 54. Centraalbureau voor Schimmelcultures, Utrecht, The Netherlands.
 3. Bertsch C, Ramírez-Suero M, Magnin-Robert M, Larignon P, Chong J, Abou-Mansour E, Spagnolo A, Clément C, Fontaine F. 2013. Grapevine trunk diseases: complex and still poorly understood. *Plant Pathol.* **62**: 243–265.
 4. Mugnai L, Graniti A, Surico G. 1999. Esca (black measles) and brown wood-streaking: two old and elusive diseases of grapevines. *Plant Dis.* **83**:404–418.
 5. Luini E, Fleurat-Lessard P, Rousseau L, Roblin G, Berjeaud J-M. 2010. Inhibitory effects of polypeptides secreted by the grapevine pathogens *Phaeoconiella chlamydospora* and *Phaeoacremonium aleophilum* on plant cell activities. *Physiol. Mol. Plant Pathol.* **74**:403–411.
 6. Bruno G, Sparapano L, Graniti A. 2007. Effects of three esca-associated fungi on *Vitis vinifera*: IV. Diffusion through the xylem of metabolites produced by two tracheiphilous fungi in the woody tissue of grapevine leads to esca-like symptoms on leaves and berries. *Physiol. Mol. Plant Pathol.* **71**:106–124.
 7. Bruno G, Sparapano L. 2006. Effects of three esca-associated fungi on *Vitis vinifera*: II. Characterization of biomolecules in xylem sap and leaves of healthy and diseased vines. *Physiol. Mol. Plant Pathol.* **69**:195–208.
 8. Bruno G, Sparapano L. 2006. Effects of three esca-associated fungi on *Vitis vinifera*: III. Enzymes produced by the pathogens and their role in fungus-to-plant or in fungus-to-fungus interactions. *Physiol. Mol. Plant Pathol.* **69**:182–194.
 9. Valtaud C, Larignon P, Roblin G, Fleurat-Lessard P. 2009. Developmental and ultrastructural features of *Phaeoconiella chlamydospora* and *phaeoacremonium aleophilum* in relation to xylem degradation in esca disease of the grapevine. *J. Plant Pathol.* **91**:37–51.
 10. Groenewald M, Kang JC, Crous PW, Gams W. 2001. ITS and beta-tubulin phylogeny of *Phaeoacremonium* and *Phaeoconiella* species. *Mycol. Res.* **105**:651–657.
 11. Möller EM, Bahnweg G, Sandermann H, Geiger HH. 1992. A simple and efficient protocol for isolation of high molecular weight DNA from filamentous fungi, fruit bodies, and infected plant tissues. *Nucleic Acids Res.* **20**:6115–6116.
 12. Parra G, Bradnam K, Ning Z, Keane T, Korf I. 2009. Assessing the gene space in draft genomes. *Nucleic Acids Res.* **37**:289–297.
 13. Stanke M, Diekhans M, Baertsch R, Haussler D. 2008. Using native and syntetically mapped cDNA alignments to improve de novo gene finding. *Bioinformatics* **24**:637–644.
 14. Smit AFA, Hubley R, Green P. 2003, posting date. RepeatMasker. Institute for Systems Biology, Seattle, WA. <http://repeatmasker.org>.
 15. Petersen TN, Brunak S, von Heijne G, Nielsen H. 2011. SignalP 4.0: discriminating signal peptides from transmembrane regions. *Nat. Methods* **8**:785–786.
 16. Cantarel BL, Coutinho PM, Rancurel C, Bernard T, Lombard V, Henrissat B. 2009. The carbohydrate-active enzymes database (CAZy): an expert resource for glycogenomics. *Nucleic Acids Res.* **37**:D233–D238.
 17. Blanco-Ulate B, Rolshausen P, Cantu D. 2013. Draft genome sequence of *Neofusicoccum parvum* isolate UCR-NP2, a fungal vascular pathogen associated with grapevine cankers. *Genome Announc.* **1**(3):e00339-13. doi: [10.1128/genomeA.00339-13](https://doi.org/10.1128/genomeA.00339-13).
 18. Blanco-Ulate B, Rolshausen PE, Cantu D. 2013. Draft genome sequence of the grapevine dieback fungus *Eutypa lata* UCR-EL1. *Genome Announc.* **1**(3):e00228-13. doi: [10.1128/genomeA.00228-13](https://doi.org/10.1128/genomeA.00228-13).
 19. Suzuki H, MacDonald J, Syed K, Salamov A, Hori C, Aerts A, Henrissat B, Wiebenga A, vanKuyk P, Barry K, Lindquist E, LaButti K, Lapidus A, Lucas S, Coutinho P, Gong Y, Samejima M, Mahadevan R, Abou-Zaid M, de Vries R, Igarashi K, Yadav J, Grigoriev I, Master E. 2012. Comparative genomics of the white-rot fungi, *Phanerochaete carnosae* and *P. chrysosporium*, to elucidate the genetic basis of the distinct wood types they colonize. *BMC Genomics* **13**:444.