



Published in final edited form as:

Nat Genet. 2013 June ; 45(6): 621–631. doi:10.1038/ng.2610.

Identification of heart rate–associated loci and their effects on cardiac conduction and rhythm disorders

A full list of authors and affiliations appears at the end of the article.

Abstract

Elevated resting heart rate is associated with greater risk of cardiovascular disease and mortality. In a 2-stage meta-analysis of genome-wide association studies in up to 181,171 individuals, we identified 14 new loci associated with heart rate and confirmed associations with all 7 previously established loci. Experimental downregulation of gene expression in *Drosophila melanogaster* and *Danio rerio* identified 20 genes at 11 loci that are relevant for heart rate regulation and highlight a role for genes involved in signal transmission, embryonic cardiac development and the pathophysiology of dilated cardiomyopathy, congenital heart failure and/or sudden cardiac death. In addition, genetic susceptibility to increased heart rate is associated with altered cardiac conduction and reduced risk of sick sinus syndrome, and both heart rate–increasing and heart rate–decreasing variants associate with risk of atrial fibrillation. Our findings provide fresh insights into the mechanisms regulating heart rate and identify new therapeutic targets.

A high resting heart rate has been associated with increased incidence of cardiovascular disease, as well as with cardiovascular and all-cause mortality, independent of traditional risk factors^{1–3}. There are several potential mechanisms by which higher heart rate may contribute to greater cardiovascular risk. For example, higher heart rate entails elevated myocardial oxygen requirement and a shift in cardiac control from parasympathetic to sympathetic dominance, which may increase the likelihood of myocardial ischemia and electrical instability⁴. In addition, experimental alteration of heart rate by sinoatrial node ablation has been shown to influence the progression of atherosclerosis induced by an atherogenic high-cholesterol diet in cynomolgus monkeys^{5,6}. In humans, selective reduction of heart rate using ivabradine was shown to reduce clinical events in individuals with heart failure, suggesting that elevated heart rate is a clinically relevant and modifiable risk factor⁷. However, whether the association of higher heart rate with cardiovascular risk is causal remains to be clarified.

© 2013 Nature America, Inc. All rights reserved.

Correspondence should be addressed to R.J.F.L. (ruth.loos@mssm.edu).

COMPETING FINANCIAL INTERESTS

The authors declare competing financial interests: details are available in the online version of the paper.

Reprints and permissions information is available online at <http://www.nature.com/reprints/index.html>.

URLs. PANTHER, <http://www.pantherdb.org/>; Gene Ontology (GO), <http://www.geneontology.org/>; Molecular Signatures Database (MsigDB), <http://www.broad.mit.edu/gsea/msigdb/collections.jsp>; Mascot Search algorithm, <http://www.matrixscience.com>; Ensembl, <http://www.ensembl.org/index.html>; SNIPPER, <http://csg.sph.umich.edu/boehnke/snipper/>.

Note: Supplementary information is available in the online version of the paper.

Large twin studies with electrocardiogram (ECG) data have shown that genetic factors contribute to interindividual variation in heart rate, with heritability estimates ranging from 55 to 77% (refs. 8–10). So far, 3 genome-wide association studies (GWAS)^{11–13}, incorporating data from up to 38,991 individuals each, have identified variants in 7 loci that show evidence of association with heart rate. These variants are common in the general population (minor allele frequency (MAF) ~10%) and together explain ~0.7% of the variance in heart rate¹². To gain more comprehensive insight into the genetic regulation of heart rate, we performed a 2-stage meta-analysis of GWAS in data from up to 181,171 individuals. Loci convincingly associated with heart rate were subsequently tested for association with cardiac conduction, rhythm disorders and cardiovascular disease to elucidate potential mechanisms underlying the association between heart rate and cardiovascular disease and mortality. Furthermore, we undertook experimental studies in *D. melanogaster* and *D. rerio* models as a first step toward identifying the causal genes within the associated loci.

RESULTS

Stage 1 GWAS identifies five new loci associated with heart rate

We performed a meta-analysis of the associations between 2,516,789 SNPs and heart rate in data from up to 85,787 individuals of European ancestry from 36 GWAS, including data from up to 11,207 individuals described previously¹³ and 6,568 individuals of Indian Asian ancestry (Online Methods, Supplementary Figs. 1–3 and Supplementary Tables 1–4). All studies included have been approved by local ethics committees, and all participants have provided their consent in writing. Our stage 1 meta-analysis showed associations with heart rate at genome-wide significance ($P < 5 \times 10^{-8}$) for variants in 12 loci (Table 1). These 12 loci included all 7 previously identified loci (in *MYH6*, *CD46* and *FADS1* and near *GJAI*, *ACHE*, *SLC35F1* and *LINC00477* (also known as *C12orf67*))^{11–13} and 5 additional loci (in *KIAA1755*, *CCDC141*, *SYT10* and *FLRT2* and near *HCN4*).

To validate associations of the loci that were significantly associated with heart rate in stage 1 and to identify additional loci (Supplementary Fig. 1), lead SNPs at 42 loci (associated at $P < 3 \times 10^{-5}$) were selected for follow-up (Online Methods and Supplementary Table 5). Conditional analyses based on summary statistics of stage 1 meta-analysis results¹⁴ identified two loci with secondary associations that remained significant ($P < 5 \times 10^{-8}$) after adjusting for the association of the lead SNP. These secondary associations were also selected for follow-up (Online Methods and Supplementary Table 6).

Stage 2 follow-up identifies nine additional new loci

In stage 2, we examined associations between the 42 loci identified in stage 1 and heart rate in data from up to 88,823 additional individuals of European descent from 27 GWAS, including data from up to 38,991 individuals from 15 GWAS described previously¹², as well as 11 studies with Metabochip and 1 study with Cardiochip data (Online Methods, Supplementary Fig. 1 and Supplementary Tables 7–10).

In a joint analysis of results from stage 1 and stage 2, variants in 21 loci had associations that reached $P < 5 \times 10^{-8}$ in data from up to 181,171 individuals (Fig. 1, Table 1 and Supplementary Table 5). Among the 21 loci were all 12 loci with association $P < 5 \times 10^{-8}$ after stage 1, as well as 9 additional loci (in *CHRM2*, *RFX4*, *CPNE8* and *TFPI* and near *GNB4*, *NKX2-5*, *GNG11*, *B3GNT7* and *FNDC3B*). Hence, our study confirms the 7 previously identified loci^{11–13} and identifies 14 new loci robustly associated with heart rate.

Impact of the 21 confirmed loci on heart rate

The frequency of the heart rate-increasing alleles ranged from 10 to 85% for the 21 confirmed associations. Effect sizes of associations ranged from 0.21 to 0.74 beats per minute (bpm) per effect allele (mean \pm s.d., 0.41 ± 0.15 bpm per effect allele) (Fig. 2 and Table 1).

To estimate the combined effect of the 21 loci on heart rate, we constructed a genetic predisposition score (GPS) by summing the number of heart rate-increasing alleles of the 21 associations. We examined associations between the GPS and heart rate in data from 5,053 adults from LifeLines2 (data for 19 loci available) and 4,000 12-year-old children from ALSPAC (data for 21 loci available) (Online Methods). The difference in average heart rate between individuals in the lowest and highest 5% of the GPS distribution was 4.1 bpm in adults (66.1 versus 70.2 bpm) and 4.9 bpm in children (73.7 versus 78.6 bpm) (Fig. 3a,b), differences that were previously shown to be clinically relevant¹⁵. The GPS explained 0.9% of the variance in heart rate in adults from LifeLines2 and 0.8% of the variance in children from ALSPAC.

Conduction, rhythm disorders and cardiovascular disease

An altered heart rate reflects sinoatrial function and may reflect disturbed electrophysiological properties that are also present in other compartments of the heart. Such properties include atrial and atrioventricular nodal conduction (PR duration), ventricular depolarization (QRS duration) and myocardial repolarization (QT duration), which can be quantified on a 12-lead ECG. We examined whether the heart rate-associated loci showed evidence of association with cardiac conduction in data from previously reported GWAS for PR¹⁶, QRS¹⁷ and QT duration (QT-IGC Consortium (C.N.-C.), personal communication). Furthermore, we examined the association of the 21 loci with the risk of several conduction-related disorders, including atrial fibrillation, advanced (second- and third-degree) atrioventricular block and sick sinus syndrome (SSS, also known as sinus node dysfunction), as well as pacemaker implantation and sudden cardiac death^{13,18}. Finally, elevated resting heart rate is a well-recognized precursor of increased blood pressure and hypertension, independent of initial blood pressure levels^{19,20}, and predicts the incidence of coronary heart disease during up to 10 years of follow-up, independent of other major risk factors^{1,2}. We therefore also examined associations of the heart rate loci with systolic blood pressure, diastolic blood pressure and the prevalence of hypertension, coronary artery disease (CAD) and myocardial infarction in data from the Global BPgen²¹ and CARDIoGRAM consortia²² (Online Methods).

For each of the ECG traits, we found a significant association with individual heart rate loci ($P < 0.002$). Heart rate-increasing alleles of these loci were associated with prolonged PR duration (near *LINC00477* and *NKX2-5*) and reduced QT duration (near *GJA1*, *FADS1*, *SLC35F1* and *NKX2-5*), independent of heart rate, as well as with both reduced (near *GJA1*, *FADS1*, *SLC35F1* and *NKX2-5*) and prolonged (in *CCDC141*) QRS duration (Table 2). Common variants of the loci in or near *GJA1*, *FADS1*, *CCDC141* and *NKX2-5* were not previously identified as being associated with these cardiac conduction traits (Supplementary Table 11). In addition, stronger genetic susceptibility for increased heart rate as conferred by the GPS of 21 loci was associated with prolonged PR duration ($P = 1.3 \times 10^{-4}$) and reduced QT duration ($P = 1.1 \times 10^{-17}$), independent of heart rate, as well as with reduced QRS duration ($P = 1.8 \times 10^{-5}$) (Table 2 and Supplementary Fig. 4). These results suggest that, to some extent, similar cellular processes control heart rate and cardiac conduction through the atria and ventricles.

Five of the 21 heart rate loci are associated with atrial fibrillation ($P < 0.002$). Heart rate-increasing alleles of these loci were associated with both increased (near *SLC35F1*, *LINC00477* and *NKX2-5*; odds ratio (OR) = 1.06–1.13) and decreased (near *GJA1* and *HCN4*; OR = 0.86–0.90) risk of atrial fibrillation (Table 2). Common variants of the loci in or near *GJA1*, *SLC35F1* and *NKX2-5* were not previously identified as being associated with atrial fibrillation (Supplementary Table 11). Stronger genetic susceptibility for increased heart rate in the 21 loci combined was not associated with atrial fibrillation, which reflects the bidirectionality of the associations in the individual loci (Table 2).

None of the heart rate loci showed evidence of association with the risk of atrioventricular block, SSS, pacemaker implantation or sudden cardiac death individually (Supplementary Table 12). However, a higher GPS was associated with reduced risk of SSS ($P = 2.3 \times 10^{-4}$) and pacemaker implantation ($P = 3.6 \times 10^{-4}$) (Table 2 and Supplementary Fig. 4). SSS encompasses a group of sinus rhythm disorders, including pathological sinus bradycardia (slow heart rate), sinus arrest, sinoatrial block and paroxysmal tachycardias (bradycardia-tachycardia syndrome). SSS is the most common indicator for permanent pacemaker implantation²³, and ~80% of individuals with SSS in our data set had undergone pacemaker implantation¹⁸. Hence, the association between the heart rate loci and pacemaker implantation is likely secondary to the association with SSS in this study population.

None of the heart rate loci showed evidence of association with blood pressure or prevalent hypertension, CAD or myocardial infarction, either individually or when combined in the GPS of 21 loci (Supplementary Fig. 4 and Supplementary Tables 13 and 14). In addition, we showed, at most, limited evidence of association with heart rate for loci previously identified as being associated with blood pressure or prevalent hypertension, CAD or myocardial infarction (Supplementary Tables 15 and 16).

Pathway analyses

The 21 confirmed loci contain 234 genes that are located within 500 kb of the associations with heart rate (Supplementary Fig. 5 and Supplementary Table 17). To systematically identify biological connections between these genes and to identify new pathways associated with heart rate, we tested whether biological processes or molecular functions that were

predefined in five databases were enriched for multiple modest heart rate associations using MAGENTA²⁴ (Online Methods). We found evidence of enrichment of associations in pathways involved in dilated, hypertrophic and arrhythmogenic right ventricular cardiomyopathy, (cardiac) muscle contraction, regulation of heart contraction, integration of energy metabolism, positive regulation of cell adhesion and Alzheimer's disease ($P < 2 \times 10^{-3}$, false discovery rate (FDR) < 0.1) (Supplementary Tables 18 and 19).

Follow-up in *D. melanogaster* and *D. rerio*

Variants identified by GWAS typically implicate genomic regions rather than individual genes. We used a range of approaches to identify promising candidate genes for heart rate regulation within the 21 loci, including proteomics experiments aimed at identifying genes expressed at the protein level in mouse heart that are phosphorylated upon stimulation of the $\beta 1$ adrenergic receptor ($\beta 1$ AR), gene expression quantitative trait locus (eQTL) analysis in blood, *in silico* search for potentially functional variants in high linkage disequilibrium (LD) with lead variants ($r^2 > 0.8$), an automated literature search using the program SNIPPER and biological candidacy (Online Methods and Supplementary Tables 20–23). These approaches labeled 49 of the 234 genes located within the 21 loci as candidate genes for heart rate regulation (Supplementary Table 24).

To examine whether some of the 49 candidate genes are likely to underlie the associations identified by GWAS, we performed 2 series of experiments using animal models. First, we compared heart rate and risk of arrhythmia in *D. melanogaster* control pupae and pupae in which orthologs of the candidate genes were downregulated using RNA-mediated gene interference (RNAi), both at rest and after 20 min of tachypacing (Online Methods and Supplementary Tables 25 and 26). Second, we compared heart rate and fractional shortening of the ventricular chamber in control embryos of the zebrafish *D. rerio* and embryos in which orthologs of the candidate genes were downregulated using morpholino oligonucleotides (Online Methods and Supplementary Table 27).

Results were available for the orthologs of 25 candidate genes from 13 loci in *D. melanogaster* pupae and for orthologs of 12 genes from 7 loci in *D. rerio* embryos; results from orthologs of 6 genes were available in both species (Supplementary Fig. 6 and Supplementary Table 24). Results from these experiments support a role in heart rate regulation for 20 of the 31 candidate genes tested across the 2 models: *ACHE*, *UFSP1*, *TRIP6*, *EPHB4* and *PCOLCE* (locus 3), *PLXNA2* (locus 4), *FADS1*, *FADS2*, *FADS3* and *BEST1* (locus 5), *TTN* (locus 9), *MFN1* (locus 12), *CHRM2* (locus 14), *GNG11* (locus 15), *NCL* and *HTR2B* (locus 17), *PLD1* and *GHSR* (locus 18), *CRY1* (locus 19) and *CALCRL* (locus 21) ($P < 2 \times 10^{-3}$ in *D. melanogaster* pupae and $P < 4 \times 10^{-3}$ in *D. rerio* embryos) (Fig. 4 and Supplementary Tables 28 and 29).

The most convincing results were observed for orthologs of *MFN1* and *PLD1* (Supplementary Note), for which downregulated gene expression reduced resting heart rate in both models ($P < 1 \times 10^{-5}$). Moreover, *D. melanogaster* pupae with downregulated expression of the *MFN1* ortholog were characterized by reduced heart rate after 20 min of tachypacing ($P = 9.5 \times 10^{-5}$) and by increased risk of arrhythmia, both at rest ($P = 4.3 \times 10^{-4}$) and after tachypacing ($P = 2.0 \times 10^{-6}$). In *D. rerio* embryos, in addition to reduced

heart rate, downregulation of *mfn1* (the ortholog of *MFN1*) was accompanied by edema in 73% of embryos, whereas 51% of embryos with reduced expression of *pld1a* (the ortholog of *PLD1*) had an unlooped heart (Supplementary Table 29).

In addition to reduced heart rate, downregulated gene expression of orthologs of *ACHE*, *PCOLCE* and *FADS3* was associated with increased risk of arrhythmia after 20 min of tachypacing in *D. melanogaster* pupae. Furthermore, *D. rerio* embryos with reduced gene expression of *D. rerio* orthologs that was accompanied by reduced heart rate were also characterized by edema (*BEST1* and *MFN1*), blood pooling (*FADS2*), an unlooped heart (*HTR2B*, *NCL*, *PLD1* and *CALCRL*) and atrioventricular canal malformation (*CALCRL*). Downregulated expression of the *CRY1* ortholog was accompanied by a highly penetrant developmental malformation that likely mediates the heart rate effect (Supplementary Fig. 7 and Supplementary Table 29). Reduced expression of the *COLIA2* ortholog did not affect heart rate in *D. rerio* embryos ($P = 0.07$) but resulted in reduced fractional shortening of the ventricular chamber ($P = 1.8 \times 10^{-3}$).

In summary, results from experiments in *D. melanogaster* and *D. rerio* models support a role in heart rate regulation for 20 genes found within 11 loci associated with heart rate (Supplementary Table 30). Notably, results from animal models confirmed the eQTL associations observed in humans for all available orthologs, with a consistent direction of effect across species for all genes except *TRIP6* (Supplementary Tables 21 and 30).

DISCUSSION

Using a 2-stage meta-analysis of GWAS in up to 181,171 individuals, we identified 14 loci previously unknown to be robustly associated with heart rate and confirmed the 7 previously established loci, increasing the total number of heart rate loci to 21. Results from experiments in *D. melanogaster* and *D. rerio* models support a role in heart rate regulation for 20 candidate genes from 11 loci. These experiments highlight a role for genes that are essential for embryonic cardiovascular development and signal transmission, as well as for genes with a role in the pathophysiology of dilated cardiomyopathy, congestive heart failure and/or sudden cardiac death (Supplementary Note). In addition, stronger genetic susceptibility to higher heart rate is associated with prolonged PR duration and reduced QT duration, both independent of heart rate, as well as with reduced QRS duration and lower risk of SSS, a group of sinus rhythm disorders that result from sinus node dysfunction and are characterized by bradycardia.

Prevalent SSS is unlikely to explain the association between common variants and heart rate shown by GWAS, as the associations were essentially unchanged with a priori exclusion of individuals with prevalent cardiovascular disease, heart rate outside the range of 50–100 bpm and/or using heart rate-altering medication. This suggests that the confirmed loci have subtle effects on sinus node function in the general population, which manifest themselves in higher heart rate and reduced risk of SSS, showing the clinical relevance of our findings. Future studies should address whether such effects on sinus node function also affect the risk of mortality. The associations with higher heart rate do not translate into significantly higher risk of CAD or myocardial infarction, either individually or in combination, which

may reflect low statistical power given the effect sizes for heart rate and the number of available CAD and myocardial infarction cases.

Heart rate-increasing alleles of the loci near *GJAI*, *SLC35F1* (*PLN*) and *NKX2-5* show unidirectional associations with reduced QRS and QT duration but bidirectional associations with atrial fibrillation. These findings suggest that both tails of the ventricular depolarization and myocardial repolarization distributions are associated with increased risk of atrial fibrillation and that altered heart rate associated with genetic predisposition in these loci may reflect adaptations to disturbed electrophysiological properties (compartments) of the heart. Results from experiments in a *D. melanogaster* model highlight genes in additional loci that show suggestive evidence of a role in both heart rate regulation and arrhythmia susceptibility (*ACHE*, *PCOLCE*, *FADS3* and *MFNI*). Taken together, these results may enable the discovery of new druggable targets for the prevention and treatment of cardiovascular endpoints by selective reduction of heart rate and arrhythmia susceptibility, similar to the way ivabradine likely exerts its effects through targeting of *HCN4* (refs. 7,25).

Results from experiments in *D. melanogaster* and *D. rerio* models support a role in heart rate regulation for genes that are essential for embryonic cardiovascular development (*EPHB4*, *PLXNA2*, *PLD1* and *CALCRL*), as well as for genes with a role in the pathophysiology of dilated cardiomyopathy, congestive heart failure and/or sudden cardiac death (*TTN*, *MFNI*, *CHRM2* and *PLD1*). In congruence, we show that zebrafish embryos with downregulated expression of orthologs of these genes have edema (*MFNI*), an unlooped heart (*PLD1* and *CALCRL*) and atrioventricular canal malformation (*CALCRL*). Such defects in cardiovascular development can be hypothesized to mediate the reduced heart rate that we observe in these embryos. Future studies are required to determine whether individuals with genetic susceptibility for reduced heart rate in these loci are enriched for mild forms of such cardiovascular phenotypes.

In conclusion, our results provide new insights into the mechanisms that regulate or modulate heart rate in health and disease and provide a new perspective on the well-recognized association of heart rate with cardiovascular disease and mortality.

ONLINE METHODS

Stage 1 genome-wide association meta-analysis

The discovery sample encompassed 36 studies with data on heart rate in up to 85,787 individuals of European ancestry. Heart rate was derived from ECG in 12 studies (32% of the total sample) and peripheral pulse rate in 22 studies (49%) and was self-reported in 2 studies with data from health professionals (19%) (Supplementary Table 1). All studies included have been approved by local ethics committees, and all participants have provided their consent in writing.

Samples were genotyped using Affymetrix and Illumina genome-wide genotyping arrays (Supplementary Tables 2 and 3) and were imputed for polymorphic HapMap Phase 2 European CEU SNPs using MACH²⁶, IMPUTE²⁷, BIMBAM²⁸ or Beagle²⁹ (Supplementary Table 2).

Each study performed SNP association analyses with heart rate using an additive genetic model implemented in MACH2QTL³⁰, Merlin³¹, SNPTEST²⁷, ProbABEL³², GenABEL³³, LME in R, MMAP, Matlab or PLINK³⁴ (Supplementary Table 2). Associations were adjusted for age, age², body mass index (BMI) and study-specific covariates when appropriate (for example, principal components). Analyses were stratified by sex and case status for samples ascertained for diseases or conditions. To allow for relatedness in the deCODE, HAPI Heart, Heritage, Korcula, NBS, NSPHS and SPLIT studies, regression coefficients were estimated in the context of a variance component model that took into account relatedness in men and women, with sex as an additional covariate. Before meta-analysis of the results from the 36 GWAS, we excluded SNPs with poor imputation quality score (r^2 hat < 0.3 in MACH, proper_info < 0.4 in IMPUTE and BIMBAM, info < 0.8 in Beagle), low minor allele count ($n \times \text{MAF} < 3$) and/or extreme effect size ($\beta > \pm 50$ bpm per effect allele, that is, ~ 5 times the standard deviation of heart rate as typically observed in the contributing studies) in each sex- and case-specific stratum. Individual GWAS were corrected by genomic control before meta-analysis when appropriate ($\lambda_{\text{GC}} > 1.000$) to adjust for population stratification.

We performed the stage 1 fixed-effects meta-analysis using the inverse variance method in METAL³⁵. Before SNPs were selected for follow-up, a final genomic control correction of the meta-analysis results was performed ($\lambda_{\text{GC}} = 1.106$), giving conservative association estimates.

Lead SNPs at 42 independent loci were selected for follow-up in stage 2 ($P < 3 \times 10^{-5}$) (Supplementary Table 5). Loci were considered to be independent if pairwise r^2 for LD was less than 0.2 and if they were separated by at least 1 Mb. We subsequently performed conditional analyses using summary statistics of stage 1 results¹⁴ to examine whether any of the 42 loci contained secondary associations with heart rate that remained significant after adjusting for the association of the lead SNP ($P < 5 \times 10^{-8}$). Before embarking on the follow-up analysis using all available data, we made sure that the inclusion of subgroups of the population did not affect the results (Supplementary Table 4).

Stage 2 follow-up

We tested for association of the 42 lead SNPs and 3 secondary associations in data from up to 88,823 individuals of European descent from 37 *in silico* replication studies with heart rate in stage 2. Heart rate was derived from ECG in 22 studies (57% of the total sample) and from peripheral pulse rate in 15 studies (43%) (Supplementary Table 7). GWAS data were available for up to 60,396 individuals of European descent from 2 sources: a previously reported meta-analysis of 15 GWAS for RR interval (RRgen Consortium)¹² and 12 GWAS that have not been described previously in this context (Supplementary Tables 7–9). Additional data were available for 24,334 individuals of European ancestry from 11 studies who were genotyped using the Metabochip³⁶ and from 5,171 individuals of European descent from 1 study who were genotyped using the Cardiochip (Supplementary Tables 8 and 10).

Samples and SNPs that did not meet the quality control criteria described by each individual study and for stage 1 were excluded. Minimum genotyping quality control criteria were

defined as Hardy-Weinberg equilibrium $P > 1 \times 10^{-6}$ and call rate $> 95\%$ in each of the follow-up studies.

We tested the association of the 42 lead SNPs and 3 secondary associations with heart rate in each stage 2 study separately as described for stage 1 studies. Missing SNPs in GWAS of stage 2 were replaced by one of up to three proxies selected a priori ($r^2 > 0.8$) (if available). Prioritizing of proxies was based on (in order of importance) (i) availability on the Metabochip and/or Cardiochip with $r^2 > 0.8$; (ii) r^2 for LD; and (iii) proximity to the lead SNP. This resulted in the inclusion of three proxies in LifeLines2 and one proxy in ACTS, all of which had $r^2 > 0.9$ with the lead SNP at the locus (Supplementary Table 10). None of the loci for which these proxies were included has association reaching $P < 5 \times 10^{-8}$ after meta-analysis of stages 1 and 2 together.

We performed meta-analysis on summary statistics from the stage 1 meta-analysis and stage 2 studies using the weighted z -score method in data from up to 181,171 individuals (Supplementary Table 5). Genomic control-adjusted P values were used throughout stages 1 and 2 for GWAS. For studies with data from Metabochip and Cardiochip, little evidence for population stratification was previously observed for associations with other cardiovascular and metabolic traits, and, hence, no correction of P values was applied.

For loci with secondary associations, the SNP with the lowest P value for association with heart rate after combined meta-analysis of stage 1 and 2 results was considered the most representative for the locus. For loci with confirmed associations after meta-analysis of stage 1 and 2 results, an estimate of the effect size was obtained by fixed-effects meta-analysis of summary statistics from the stage 1 meta-analysis and stage 2 studies with heart rate, using the inverse variance method.

Additional analyses and functional follow-up experiments

Cumulative effects of confirmed loci and interindividual variation in heart rate

—To estimate the cumulative effect of the 21 heart rate-associated loci, we calculated the GPS in 5,053 adults from the LifeLines2 study and 4,000 12-year-old children from the ALSPAC study by summing the number of heart rate-increasing alleles carried by an individual at the lead SNP of each heart rate locus. The number of heart rate-increasing alleles ranged from 9 to 26 for the 19 available loci in LifeLines2 (data from rs6127471 and rs2340782 were not available) and from 10 to 29 for the 21 loci in ALSPAC.

We compared the explained variance (r^2) between covariate-adjusted models with and without the GPS to assess the variance in heart rate that can be explained by these loci in the LifeLines2 and ALSPAC studies.

Association analyses with related traits—Associations between the 21 heart rate loci and related cardiovascular intermediates and endpoints were extracted from GWAS data of the CHARGE Consortium (PR duration)¹⁶, the QRS GWAS Consortium (QRS duration)¹⁷ and the QT-IGC Consortium (QT duration; QT-IGC Consortium (C.N.-C.), personal communication) (Table 2), as well as from deCODE Genetics (prevalent advanced (second- or third-degree) atrioventricular block, SSS, pacemaker implantation and sudden cardiac

death)^{13,18} (Table 2 and Supplementary Table 12), deCODE Genetics¹³ and the CHARGE-AF Consortium²⁵ (atrial fibrillation) (Table 2), the Global BPgen Consortium (systolic and diastolic blood pressure, as well as prevalent hypertension)²¹ (Supplementary Table 13) and the CARDIoGRAM Consortium (prevalent CAD and myocardial infarction)²² (Supplementary Table 14). All associations were adjusted for covariates as described previously^{13,16–18,21,22,25,37}.

We calculated multi-SNP predisposition scores for each trait to examine the association of the 21 heart rate loci combined with each of the before-mentioned traits, on the basis of single-SNP summary statistics and weighting by effect sizes for association with heart rate after meta-analysis of stages 1 and 2 together.

Associations of the 21 heart rate loci, both individually and in combination in a multi-SNP predisposition score, with related traits were considered statistically significant at $P < 0.002$, that is, $\alpha = 0.05$ with Bonferroni correction for 21 independent tests.

Enrichment analysis of heart rate associations in biological pathways—We used MAGENTA²⁴ to test whether predefined biological processes or molecular functions were enriched for multiple modest heart rate associations, aiming to discover new pathways associated with heart rate and to test whether the 21 heart rate loci cluster near genes that constitute specific biological connections (Supplementary Tables 18 and 19). First, we calculated a corrected gene association P value for each gene in the genome and grouped genes into pathways using annotations from the Kyoto Encyclopedia of Genes and Genomes (KEGG), BIOCARTA, Protein Analysis THrough Evolutionary Relationships (PANTHER)³⁸, Biological Processes (PANTHER, BP) and Molecular Functions (PANTHER, MF), REACTOME, Gene Ontology (GO) and Ingenuity databases. Finally, for each pathway, we evaluated potential enrichment of highly ranked gene scores by comparing the fraction of genes within each gene set whose corrected P value was more significant than the 95th percentile of all gene P values to that of 10,000 randomly sampled gene sets of identical size from the genome²⁴. In significantly enriched gene sets, in addition to genes in validated association regions, the top ranked genes above the enrichment cutoff may suggest new modest associations for follow-up (Supplementary Table 19).

Proteomics experiments in mouse heart and genetic enrichment analysis—We used results from proteomics experiments to identify genes located within the heart rate loci that are expressed at the protein level in mouse heart and that are phosphorylated upon stimulation of β 1AR (A.L., M.N. Andersen, A.B. Steffensen, H. Horn, C.D. Kelstrup *et al.*, unpublished data). Briefly, male C57BL/6 mice were either treated with β 1AR- and β 2AR-specific antagonists (control group, $n = 3$) or with a β 2AR-specific antagonist followed by β 1AR-specific agonist (test group, $n = 3$). Cardiac proteins were extracted and digested, enriched for phosphopeptides by TiO₂ chromatography and analyzed by nanoflow liquid chromatography tandem mass spectrometry, as described previously³⁹.

A total of 8,518 phosphorylation sites that could be mapped to a specific residue were identified. Mice in the test and control groups were compared using a two-sided t test with permutation-based FDR < 0.01 . The number of genes encoding proteins identified in the

experiments was 4,096. Forty-one of these were located within 500 kb of confirmed heart rate associations, four of which were regulated by β 1AR stimulation (*MYH6*, *PLN*, *TTN* and *NCL*).

eQTL analyses—We examined associations between each of the heart rate loci and expression of genes in *cis* in 1,469 whole-blood samples (PAXgene), reflecting primary leukocyte gene expression⁴⁰ (Supplementary Table 21). Transcriptional components were applied to reduce a substantial proportion of interindividual non-genetic expression variation. An eQTL meta-analysis was subsequently performed on the residual expression variation. We used FDR < 0.05 to correct for multiple testing. We removed 50 principal components by linear regression to remove non-genetic variation in gene expression. In addition, we performed conditional analyses to examine to what extent each heart rate-associated SNP explains the association between the gene transcript and the SNP most significantly associated with the gene transcript.

Significant *cis* associations were observed between five heart rate-associated SNPs and the levels of nine nearby transcripts in blood (Table 1 and Supplementary Table 21). The heart rate-associated SNPs explained a substantial proportion of the association with the most significant SNP for the gene transcript in conditional analyses (adjusted $P > 0.05$) for *TRIP6*, *TMEM258(C11orf10)*, *FADS1*, *BEST1-FTH1*, *FADS2*, *CEP85L(C6orf204)* and *NCL-SNORD20*.

Potentially functional variants within the 21 loci—To identify SNPs in the confirmed loci that may be causal for the association with heart rate, we explored whether the heart rate-associated SNPs were in strong LD ($r^2 > 0.8$) with variants in transcription factor binding sites, nonsynonymous SNPs or copy number variants (deletion variants and mobile element insertion polymorphisms) identified in the 1000 Genomes Project CEU Pilot 1 or HapMap CEU reference panels⁴¹. For nonsynonymous SNPs, PANTHER³⁸ was used to assess whether the variant was likely to have a detrimental effect on protein function, based on alignment of evolutionarily related proteins (Supplementary Table 20).

One association tagged a variant in a transcription factor binding site near *CEP85L(C6orf204)*; near *SLC35F1*). Nonsynonymous variants in strong LD with heart rate-associated SNPs were present in six genes (Supplementary Table 22), with the p.Arg1045Trp alteration encoded in *KIAA1755* likely having a deleterious effect on protein function³⁸. Of interest, the rs180242 allele that was associated with lower heart rate tagged a common 723-bp deletion variant located 8 kb upstream of *GNG11* (ref. 42) (1000 Genomes Pilot ID P2_M_061510_7_474; $r^2 = 0.96$) and was additionally associated with lower expression of *GNG11* in blood (Supplementary Table 21).

Candidate genes based on the literature—To identify additional candidate genes in the heart rate loci, we identified all genes within 500 kb of the 21 heart rate-associated SNPs and performed an automated literature search using the search term ‘heart’ in the program SNIPPER (Supplementary Table 23). We identified many genes with established connections to embryonic cardiac development, cardiac conduction, cardiac contractile proteins, calcium regulation, angiogenesis and endothelial function (Supplementary Note).

Many of the loci harbored genes in which mutations lead to dilated and hypertrophic cardiomyopathy (in *MYH6* and *MYH7*, *PLN* (near *SLC35F1*), *TTN* (near *CCDC141*), *MFN1* (near *GNB4*) and *CHRM2*).

Experimental follow-up of positional candidate genes in *D. melanogaster* and *D. rerio*

We used *D. melanogaster* and the zebrafish *D. rerio* as models to examine whether candidate genes within the heart rate loci were likely to underlie the associations identified by GWAS. Forty-nine positional candidate genes were identified on the basis of results from proteomics experiments and genetic enrichment analysis in mouse heart and eQTL analyses, as well as by the presence of functional variants and results from the automated literature search. In addition, we searched the genes located within 500 kb of associations for biological candidates (Supplementary Table 24).

Experiments in *D. melanogaster*—BLAST searches were performed to identify obvious *D. melanogaster* orthologs of positional candidate genes (Supplementary Table 25). We subsequently used RNAi^{43,44} to downregulate orthologs of candidate genes (Vienna *Drosophila* RNAi Center). Expression of RNAi was induced by crossing with a *D. melanogaster* line expressing GAL4 driven by an actin promoter (stock 4414, Bloomington *Drosophila* Stock Center) (Supplementary Table 25).

D. melanogaster stocks were kept at 25°C on standard medium. Pre-pupae were selected for tachypacing, an established *D. melanogaster* model for atrial fibrillation, as previously described⁴⁵. We recorded videos through a microscope at 10× magnification before and after tachypacing to visualize heart contractions in triplicate periods of 10 s. Heart rate was subsequently quantified using ImageJ software. An arrhythmia index was calculated as the ratio of arrhythmic periods and total measurement duration using the same software.

The number of positional candidate genes for which we analyzed results was reduced from 49 to 25 owing to the absence of orthologs ($n = 13$ genes), lack of RNAi lines ($n = 3$), non-viability of offspring ($n = 1$) and reduced viability, defined as the generation of fewer than 5 live offspring ($n = 7$) (Supplementary Fig. 6 and Supplementary Table 25). For the 25 remaining genes, represented by 23 orthologs, we compared heart rate and risk of arrhythmia in 11 ± 5 (mean \pm s.d.) pupae with downregulated gene expression and 30 6000V controls. Ten additional pupae were available with downregulation of *stwl* (stonewall), a gene that is not anticipated to have a role in heart rate regulation and which can thus be interpreted as an extra control group. We targeted 13 orthologs using multiple independent RNAi lines.

We compared differences in heart rate between the offspring of RNAi-treated *D. melanogaster* and 6000V controls using a multilevel approach, adjusting for the dependence of repeated measures within pupae as a random effect. We compared RNAi lines with 6000V controls at baseline and after tachypacing. Differences in heart rate after tachypacing were examined with and without adjusting for average heart rate at baseline as a fixed effect. Results of analyses from multiple independent RNAi lines targeting the same ortholog were combined using the fixed-effects meta-analysis with inverse variance method (Supplementary Table 26).

We examined the risk of arrhythmia by comparing the number of arrhythmic cases and controls between RNAi-targeted orthologs and 6000V controls, both before and after tachypacing, using Fisher's exact test. Differences were considered statistically significant at $P < 0.002$, that is, $\alpha = 0.05$ with Bonferroni correction for 23 independent tests.

Experiments in *D. rerio*—In the zebrafish *D. rerio*, we excluded genes in eight loci that mapped in or near positional candidate genes with extensive a priori evidence of a role in cardiovascular processes (Supplementary Table 27). A maximum of two positional candidate genes per locus were selected in the remaining loci, which together with a lack of zebrafish orthologs in 12 genes resulted in the selection of 12 positional candidate genes for follow-up in zebrafish experiments (Supplementary Fig. 6 and Supplementary Tables 25 and 27).

Wild-type *D. rerio* stocks from Ekwill Fish Farm were maintained using standard procedures. Morpholino oligonucleotides (GeneTools) were designed against orthologs of the 12 positional candidate gene primary transcripts targeting the first exon-intron boundaries, except for *tfpia* (the ortholog of *TFPI*), which was designed to target the intron 1–exon 2 boundary (Supplementary Table 27). Embryos were injected at the single-cell stage and were scored and analyzed 48 h later. Downregulation of candidate genes was confirmed using quantitative PCR (Supplementary Table 29).

Heart rate analysis was performed as previously described⁴⁶. For each ortholog, we performed the procedure twice on different days, comparing heart rate in embryos from the same embryonic aliquot that were injected with either morpholino oligonucleotides or PBS. Heart rate was measured in 26 ± 4 embryos injected with morpholino oligonucleotides and in 27 ± 5 embryos injected with PBS.

Measures of heart rate alone do not provide information on cardiac contractility. We therefore additionally measured ventricular fractional shortening in a subsample of embryos (6 ± 1 embryos injected with morpholino oligonucleotides and 6 ± 2 embryos injected with PBS) as previously described⁴⁷.

Differences in heart rate and fractional shortening of the ventricular chamber were examined using linear regression and were adjusted for variation in the timing of the heart rate measurement.

For each group of embryos injected with morpholino oligonucleotides (42 ± 18 embryos), we assessed whether downregulation of positional candidate gene expression resulted in visible phenotypes that distinguished treated embryos from those injected with PBS.

Supplementary Material

Refer to Web version on PubMed Central for supplementary material.

Authors

Marcel den Hoed^{1,2}, Mark Eijgelsheim³, Tõnu Esko^{4,5,6}, Bianca J J M Brundel⁷, David S Peal⁸, David M Evans⁹, Ilja M Nolte¹⁰, Ayellet V Segrè^{11,12}, Hilma Holm¹³, Robert E Handsaker^{12,14}, Harm-Jan Westra¹⁵, Toby Johnson^{16,17}, Aaron Isaacs^{3,18}, Jian Yang¹⁹, Alicia Lundby^{20,21,22}, Jing Hua Zhao¹, Young Jin Kim²³, Min Jin Go²³, Peter Almgren²⁴, Murielle Bochud²⁵, Gabrielle Boucher^{26,27}, Marilyn C Cornelis²⁸, Daniel Gudbjartsson¹³, David Hadley^{29,30}, Pim Van Der Harst³¹, Caroline Hayward³², Martin Den Heijer³³, Wilmar Igl³⁴, Anne U Jackson^{35,36}, Zoltán Kutalik^{37,38}, Jian'an Luan¹, John P Kemp⁹, Kati Kristiansson^{39,40}, Claes Ladenvall²⁴, Mattias Lorentzon⁴¹, May E Montasser⁴², Omer T Njajou⁴³, Paul F O'Reilly^{44,45}, Sandosh Padmanabhan⁴⁶, Beate St. Pourcain⁹, Tuomo Rankinen⁴⁷, Perttu Salo^{39,48}, Toshiko Tanaka⁴⁹, Nicholas J Timpson⁹, Veronique Vitart³², Lindsay Waite⁵⁰, William Wheeler⁵¹, Weihua Zhang⁴⁴, Harmen H M Draisma⁵², Mary F Feitosa⁵³, Kathleen F Kerr⁵⁴, Penelope A Lind⁵⁵, Evelin Mihailov^{4,6}, N Charlotte Onland-Moret^{56,57}, Ci Song⁵⁸, Michael N Weedon⁵⁹, Weijia Xie⁵⁹, Loic Yengo⁶⁰, Devin Absher⁵⁰, Christine M Albert^{61,62}, Alvaro Alonso⁶³, Dan E Arking^{64,65}, Paul I W de Bakker^{12,57,66,67}, Beverley Balkau^{68,69}, Cristina Barlassina⁷⁰, Paola Benaglio³⁷, Joshua C Bis^{71,72}, Nabila Bouatia-Naji^{60,73,74}, Søren Brage¹, Stephen J Chanock^{75,76}, Peter S Chines⁷⁷, Mina Chung^{78,79}, Dawood Darbar⁸⁰, Christian Dina^{81,82}, Marcus Dörr^{83,84}, Paul Elliott^{44,45}, Stephan B Felix^{83,84}, Krista Fischer⁵, Christian Fuchsberger³⁵, Eco J C de Geus⁵², Philippe Goyette^{26,27}, Vilmundur Gudnason^{85,86}, Tamara B Harris⁸⁷, Anna-liisa Hartikainen⁸⁸, Aki S Havulinna⁴⁸, Susan R Heckbert^{71,89}, Andrew A Hicks⁹⁰, Albert Hofman^{3,91}, Suzanne Holewijn⁹², Femke Hoogstra-Berends^{7,93}, Jouke-Jan Hottenga⁵², Majken K Jensen²⁸, Åsa Johansson^{34,94}, Juhani Juntila^{95,96}, Stefan Kääh^{97,98}, Bart Kanon⁹⁹, Shamika Ketkar⁵³, Kay-Tee Khaw¹⁰⁰, Joshua W Knowles¹⁰¹, Angrad S Kooner¹⁰², Jan A Kors¹⁰³, Meena Kumari¹⁰⁴, Lili Milani⁵, Päivi Laiho¹⁰⁵, Edward G Lakatta¹⁰⁶, Claudia Langenberg¹, Maarten Leusink^{57,107}, Yongmei Liu¹⁰⁸, Robert N Luben¹⁰⁰, Kathryn L Lunetta^{109,110}, Stacey N Lynch⁸, Marcello R P Markus¹¹¹, Pedro Marques-Vidal¹¹², Irene Mateo Leach³¹, Wendy L McArdle¹¹³, Steven A McCarroll^{12,14}, Sarah E Medland^{20,55}, Kathryn A Miller⁴⁴, Grant W Montgomery¹¹⁴, Alanna C Morrison¹¹⁵, Martina Müller-Nurasyid^{97,116,117}, Pau Navarro³², Mari Nelis^{4,5,118}, Jeffrey R O'Connell⁴², Christopher J O'Donnell^{110,119,120}, Ken K Ong^{1,121}, Anne B Newman¹²², Annette Peters^{98,123}, Ozren Polasek¹²⁴, Anneli Pouta^{88,125}, Peter P Pramstaller^{90,126,127}, Bruce M Psaty^{71,72,89,128,129}, Dabeeru C Rao¹³⁰, Susan M Ring¹¹³, Elizabeth J Rossin^{20,131,132,133}, Diana Rudan¹³⁴, Serena Sanna¹³⁵, Robert A Scott¹, Jaban S Sehmi^{102,136}, Stephen Sharp¹, Jordan T Shin⁸, Andrew B Singleton¹³⁷, Albert V Smith⁸⁵, Nicole Soranzo^{19,138}, Tim D Spector¹³⁸, Chip Stewart^{20,139}, Heather M Stringham^{35,36}, Kirill V Tarasov¹⁰⁶, André G Uitterlinden^{3,91,140}, Liesbeth Vandenput⁴¹, Shih-Jen Hwang¹¹⁰, John B Whitfield¹⁴¹, Cisca Wijmenga¹⁵, Sarah H Wild¹⁴², Gonneke Willemsen⁵², James F Wilson¹⁴², Jacqueline C M Witteman^{3,91}, Andrew Wong¹²¹, Quenna Wong⁵⁴, Yalda Jamshidi^{138,143}, Paavo Zitting¹⁴⁴, Jolanda M A Boer¹⁴⁵, Dorret I Boomsma⁵², Ingrid B Borecki⁵³, Cornelia M Van

Duijn^{3,18}, Ulf Ekelund^{1,146}, Nita G Forouhi¹, Philippe Froguel^{60,73,147}, Aroon Hingorani¹⁰⁴, Erik Ingelsson^{2,58,148}, Mika Kivimaki¹⁰⁴, Richard A Kronma⁵⁴, Diana Kuh¹²¹, Lars Lind¹⁴⁹, Nicholas G Martin¹⁴¹, Ben A Oostra¹⁵⁰, Nancy L Pedersen⁵⁸, Thomas Quertermous¹⁰¹, Jerome I Rotter¹⁵¹, Yvonne T van der Schouw⁵⁷, W M Monique Verschuren¹⁴⁵, Mark Walker¹⁵², Demetrius Albanes⁷⁵, David O Arnar^{153,154}, Themistocles L Assimes¹⁰¹, Stefania Bandinelli¹⁵⁵, Michael Boehnke^{35,36}, Rudolf A de Boer³¹, Claude Bouchard⁴⁷, W L Mark Caulfield^{16,17}, John C Chambers^{44,102}, Gary Curhan^{156,157}, Daniele Cusi⁷⁰, Johan Eriksson^{48,158,159,160}, Luigi Ferrucci⁴⁹, Wiek H van Gilst³¹, Nicola Glorioso¹⁶¹, Jacqueline de Graaf⁹², Leif Groop²⁴, Ulf Gyllensten³⁴, Wen-Chi Hsueh⁴³, Frank B Hu²⁸, Heikki V Huikuri⁹⁶, David J Hunter¹⁶², Carlos Iribarren¹⁶³, Bo Isomaa^{159,164}, Marjo-Riitta Jarvelin^{44,45,125,165,166}, Antti Jula¹⁶⁷, Mika Kähönen¹⁶⁸, Lambertus A Kiemeny^{169,170}, Melanie M van der Klauw¹⁷¹, Jaspal S Kooner^{102,136}, Peter Kraft¹⁶², Licia Iacoviello¹⁷², Terho Lehtimäki¹⁶⁸, Marja-Liisa L Lokki¹⁷³, Braxton D Mitchell⁴², Gerjan Navis¹⁷⁴, Markku S Nieminen¹⁷⁵, Claes Ohlsson⁴¹, Neil R Poulter¹⁷⁶, Lu Qi²⁸, Olli T Raitakari^{177,178}, Eric B Rimm²⁸, John D Rioux^{26,27}, Federica Rizzi¹⁷⁹, Igor Rudan¹⁴², Veikko Salomaa⁴⁸, Peter S Sever¹⁷⁶, Denis C Shields^{180,181,182}, Alan R Shuldiner^{42,183}, Juha Sinisalo¹⁷⁵, Alice V Stanton¹⁸⁴, Ronald P Stolk¹⁰, David P Strachan²⁹, Jean-Claude Tardif^{26,27}, Unnur Thorsteinsdottir^{13,154}, Jaako Tuomilehto^{185,186,187,188}, Dirk J van Veldhuisen³¹, Jarmo Virtamo⁴⁸, Jorma Viikari¹⁸⁹, Peter Vollenweider¹⁹⁰, Gérard Waeber¹⁹⁰, Elisabeth Widen¹⁹¹, Yoon Shin Cho¹⁹², Jesper V Olsen²², Peter M Visscher¹⁹³, Cristen Willer¹⁹⁴, Lude Franke^{15,195}, Global BPgen Consortium¹⁹⁶, CARDIoGRAM Consortium¹⁹⁶, Jeanette Erdmann^{90,197,198}, John R Thompson¹⁹⁹, PR GWAS Consortium¹⁹⁶, Arne Pfeufer^{200,201}, QRS GWAS Consortium¹⁹⁶, Nona Sotoodehnia^{71,202}, QT-IGC Consortium¹⁹⁶, Christopher Newton-Cheh^{12,110,203}, CHARGE-AF Consortium¹⁹⁶, Patrick T Ellinor^{62,204,205}, Bruno H Ch Stricker^{3,91,103,140,206}, Andres Metspalu^{4,5,6}, Markus Perola^{5,39,48}, Jacques S Beckmann^{37,207}, George Davey Smith⁹, Kari Stefansson^{13,154}, Nicholas J Wareham¹, Patricia B Munroe^{16,17}, Ody C M Sibon⁹⁹, David J Milan^{8,208}, Harold Snieder¹⁰, Nilesh J Samani^{209,210}, and Ruth J F Loos^{1,211,212,213}

Affiliations

¹Medical Research Council (MRC) Epidemiology Unit, Institute of Metabolic Science, Addenbrooke's Hospital, Cambridge, UK ²Department of Medical Sciences, Molecular Epidemiology and Science for Life Laboratory, Uppsala University, Uppsala, Sweden ³Department of Epidemiology, Erasmus University Medical Center, Rotterdam, The Netherlands ⁴Estonian Biocenter, Tartu, Estonia ⁵Estonian Genome Center, University of Tartu, Tartu, Estonia ⁶Institute of Molecular and Cell Biology, University of Tartu, Tartu, Estonia ⁷Department of Clinical Pharmacology, University Medical Center Groningen, University of Groningen, Groningen, The Netherlands ⁸Cardiovascular Research Center, Massachusetts General Hospital, Boston, Massachusetts, USA ⁹MRC Centre for Causal Analyses in Translational Epidemiology (CAiTE), School of Social and Community Medicine, University of Bristol, Bristol, UK ¹⁰Department of Epidemiology, University Medical

Center Groningen, University of Groningen, Groningen, The Netherlands
¹¹Department of Molecular Biology, Massachusetts General Hospital, Boston, Massachusetts, USA ¹²Program in Medical and Population Genetics, Broad Institute of Harvard and MIT, Cambridge, Massachusetts, USA ¹³deCODE Genetics, Reykjavik, Iceland ¹⁴Department of Genetics, Harvard Medical School, Boston, Massachusetts, USA ¹⁵Department of Genetics, University Medical Center Groningen, University of Groningen, Groningen, The Netherlands ¹⁶Clinical Pharmacology, National Institute for Health Research (NIHR) Cardiovascular Biomedical Research Unit, William Harvey Research Institute, Barts and The London School of Medicine and Dentistry, Queen Mary University of London, London, UK ¹⁷Genome Centre, Barts and The London School of Medicine and Dentistry, Queen Mary University of London, London, UK ¹⁸Centre for Medical Systems Biology, Leiden, The Netherlands ¹⁹Wellcome Trust Sanger Institute, Hinxton, Cambridge, UK ²⁰Broad Institute of MIT and Harvard, Cambridge, Massachusetts, USA ²¹The Danish National Research Foundation Centre for Cardiac Arrhythmia, Copenhagen, Denmark ²²Novo Nordisk Foundation Center for Protein Research, Faculty of Health Sciences, University of Copenhagen, Copenhagen, Denmark ²³Center for Genome Science, National Institute of Health, Osong Health Technology Administration Complex, Daejeon, The Republic of Korea ²⁴Department of Clinical Sciences, Diabetes and Endocrinology, Lund University and Lund University Diabetes Centre, Malmö, Sweden ²⁵Community Prevention Unit, Institute of Social and Preventive Medicine, Lausanne University Hospital, Lausanne, Switzerland ²⁶Université de Montréal, Montreal, Quebec, Canada ²⁷Montreal Heart Institute, Montreal, Quebec, Canada ²⁸Nutrition, Harvard School of Public Health, Boston, Massachusetts, USA ²⁹Division of Population Health Sciences and Education, St. George's, University of London, London, UK ³⁰Pediatric Epidemiology Center, University of South Florida, Tampa, Florida, USA ³¹Department of Cardiology, University Medical Center Groningen, University of Groningen, Groningen, The Netherlands ³²MRC Human Genetics Unit, Institute of Genetics and Molecular Medicine, University of Edinburgh, Edinburgh, UK ³³Department of Internal Medicine, VU Medical Center, Amsterdam, The Netherlands ³⁴Department of Immunology, Genetics and Pathology, Uppsala University, Uppsala, Sweden ³⁵Department of Biostatistics, University of Michigan, Ann Arbor, Michigan, USA ³⁶Center for Statistical Genetics, University of Michigan, Ann Arbor, Michigan, USA ³⁷Department of Medical Genetics, University of Lausanne, Lausanne, Switzerland ³⁸Swiss Institute of Bioinformatics, Lausanne, Switzerland ³⁹The Institute of Molecular Medicine, University of Helsinki, Helsinki, Finland ⁴⁰Department of Molecular Medicine, National Public Health Institute, Helsinki, Finland ⁴¹Department of Internal Medicine, Institute of Medicine, Sahlgrenska Academy, University of Gothenburg, Gothenburg, Sweden ⁴²Department of Medicine, Division of Endocrinology, Diabetes and Nutrition, University of Maryland, School of Medicine, Baltimore, Maryland, USA ⁴³Department of Medicine, Institute for Human Genetics, University of California, San Francisco, San Francisco, California, USA ⁴⁴Department of Epidemiology and

Biostatistics, Imperial College London, London, UK ⁴⁵MRC–Health Protection Agency (MRC-HPA) Centre for Environment and Health, Imperial College London, London, UK ⁴⁶Institute of Cardiovascular and Medical Sciences, College of Medical, Veterinary and Life Sciences, University of Glasgow, Glasgow, UK ⁴⁷Human Genomics Laboratory, Pennington Biomedical Research Center, Baton Rouge, Louisiana, USA ⁴⁸Department of Chronic Disease Prevention, National Institute for Health and Welfare, Helsinki, Finland ⁴⁹Clinical Research Branch, National Institute on Aging, Baltimore, Maryland, USA ⁵⁰HudsonAlpha Institute for Biotechnology, Huntsville, Alabama, USA ⁵¹Information Management Services, Inc., Rockville, Maryland, USA ⁵²Department of Biological Psychology, VU University Amsterdam and Institute for Health and Care Research (EMGO+), VU Medical Center, Amsterdam, The Netherlands ⁵³Division of Statistical Genomics, Washington University School of Medicine, St. Louis, Missouri, USA ⁵⁴Department of Biostatistics, University of Washington, Seattle, Washington, USA ⁵⁵Quantitative Genetics Laboratory, Queensland Institute of Medical Research, Brisbane, Queensland, Australia ⁵⁶Complex Genetics Section, Department of Medical Genetics–de Bakker Group, University Medical Center Utrecht, Utrecht, The Netherlands ⁵⁷Julius Center for Health Sciences and Primary Care, University Medical Center Utrecht, Utrecht, The Netherlands ⁵⁸Department of Medical Epidemiology and Biostatistics, Karolinska Institutet, Stockholm, Sweden ⁵⁹Peninsula College of Medicine and Dentistry, University of Exeter, Exeter, UK ⁶⁰Centre National de la Recherche Scientifique (CNRS) Unité Mixte de Recherche (UMR) 8199, Institut Pasteur de Lille, Lille, France ⁶¹Division of Preventive Medicine, Cardiovascular Division, Brigham and Women’s Hospital, Boston, Massachusetts, USA ⁶²Harvard Medical School, Boston, Massachusetts, USA ⁶³Division of Epidemiology and Community Health, School of Public Health, University of Minnesota, Minneapolis, Minnesota, USA ⁶⁴Department of Medicine, Division of Cardiology, Johns Hopkins University School of Medicine, Baltimore, Maryland, USA ⁶⁵McKusick-Nathans Institute of Genetic Medicine, Johns Hopkins University School of Medicine, Baltimore, Maryland, USA ⁶⁶Department of Medicine, Division of Genetics, Brigham and Women’s Hospital, Boston, Massachusetts, USA ⁶⁷Department of Medical Genetics, Division of Biomedical Genetics, University Medical Center Utrecht, Utrecht, The Netherlands ⁶⁸Institut National de la Santé et de la Recherche Médicale (INSERM), CESP Centre for Research in Epidemiology and Population Health, U1018, Epidemiology of Diabetes, Obesity and Chronic Kidney Disease over The Lifecourse, Villejuif, France ⁶⁹UMRS 1018, Université Paris–Sud 11, Villejuif, France ⁷⁰Department of Health Sciences, Milan University and Filarete Foundation, Milan, Italy ⁷¹Cardiovascular Health Research Unit, University of Washington, Seattle, Washington, USA ⁷²Department of Medicine, University of Washington, Seattle, Washington, USA ⁷³Lille Nord de France University, Lille, France ⁷⁴INSERM U970, Paris Cardiovascular Research Centre, Hôpital Européen Georges Pampidou, Paris, France ⁷⁵Division of Cancer Epidemiology and Genetics, US National Cancer Institute, Bethesda, Maryland, USA ⁷⁶Laboratory of Translational Genomics, Division of Cancer Epidemiology and

Genetics, US National Cancer Institute, Bethesda, Maryland, USA ⁷⁷National Human Genome Research Institute, US National Institutes of Health, Bethesda, Maryland, USA ⁷⁸Department of Cardiovascular Medicine, Heart and Vascular Institute, Cleveland Clinic, Cleveland, Ohio, USA ⁷⁹Department of Molecular Cardiology, Lerner Research Institute, Cleveland Clinic, Cleveland, Ohio, USA ⁸⁰Department of Medicine, Vanderbilt University School of Medicine, Nashville, Tennessee, USA ⁸¹INSERM U1087, CNRS U6291 and University of Nantes, l'Institut du Thorax, Nantes, France ⁸²Centre Hospitalier Universitaire (CHU) Nantes, l'Institut du Thorax, Nantes, France ⁸³Department of Internal Medicine B, University Medicine Greifswald, Greifswald, Germany ⁸⁴DZHK (German Centre for Cardiovascular Research), partner site Greifswald, Greifswald, Germany ⁸⁵Icelandic Heart Association Research Institute, Kopavogur, Iceland ⁸⁶University of Iceland, Reykjavik, Iceland ⁸⁷Laboratory for Epidemiology, Demography, and Biometry, National Institute on Aging, Bethesda, Maryland, USA ⁸⁸Department of Clinical Sciences, Obstetrics and Gynecology, University of Oulu, Oulu, Finland ⁸⁹Department of Epidemiology, University of Washington, Seattle, Washington, USA ⁹⁰Center for Biomedicine, European Academy Bozen/Bolzano (EURAC), Bolzano, Italy (affiliated institute of the University of Lübeck) ⁹¹Netherlands Genomics Initiative-sponsored Netherlands Consortium for Healthy Ageing, Rotterdam, The Netherlands ⁹²Department of General Internal Medicine, Division of Vascular Medicine, Radboud University Nijmegen Medical Centre, Nijmegen, The Netherlands ⁹³Nyken, Groningen, The Netherlands ⁹⁴Uppsala Clinical Research Center, Uppsala University Hospital, Uppsala, Sweden ⁹⁵Division of Cardiology, Miller School of Medicine, University of Miami, Miami, Florida, USA ⁹⁶Institute of Clinical Medicine, Department of Internal Medicine, University of Oulu, Oulu, Finland ⁹⁷Department of Medicine I, University Hospital Grosshadern, Ludwig-Maximilians- Universität, Munich, Germany ⁹⁸Munich Heart Alliance, Munich, Germany ⁹⁹Department of Cell Biology, University Medical Center Groningen, University of Groningen, Groningen, The Netherlands ¹⁰⁰Department of Public Health and Primary Care, Institute of Public Health, University of Cambridge, Cambridge, UK ¹⁰¹Department of Medicine, Stanford University School of Medicine, Stanford, California, USA ¹⁰²Cardiology Department, Ealing Hospital National Health Service (NHS) Trust, London, UK ¹⁰³Department of Medical Informatics, Erasmus University Medical Center, Rotterdam, The Netherlands ¹⁰⁴Genetic Epidemiology Group, Department of Epidemiology and Public Health, University College London, London, UK ¹⁰⁵National Public Health Institute, Biomedicum Helsinki, Helsinki, Finland ¹⁰⁶Laboratory of Cardiovascular Science, National Institute on Aging, Baltimore, Maryland, USA ¹⁰⁷Division Pharmacoeconomics & Clinical Pharmacology, Utrecht University, Utrecht, The Netherlands ¹⁰⁸Department of Epidemiology & Prevention, Division of Public Health Sciences, Wake Forest University, Winston-Salem, North Carolina, USA ¹⁰⁹Department of Biostatistics, School of Public Health, Boston University, Boston, Massachusetts, USA ¹¹⁰National Heart, Lung, and Blood Institute's Framingham Heart Study, Framingham, Massachusetts, USA ¹¹¹Institute for Community Medicine, University

Medicine Greifswald, Greifswald, Germany ¹¹²Institut Universitaire de Médecine Sociale et Préventive (IUMSP), Centre Hospitalier Universitaire Vaudois (CHUV) and University of Lausanne, Lausanne, Switzerland ¹¹³School of Social and Community Medicine, University of Bristol, Bristol, UK ¹¹⁴Molecular Epidemiology Laboratory, Queensland Institute of Medical Research, Brisbane, Queensland, Australia ¹¹⁵Human Genetics Center, University of Texas Health Science Center, Houston, Texas, USA ¹¹⁶Chair of Genetic Epidemiology, Institute of Medical Informatics, Biometry and Epidemiology, Ludwig-Maximilians-Universität, Munich, Germany ¹¹⁷Institute of Genetic Epidemiology, Helmholtz Zentrum München–German Research Center for Environmental Health, Neuherberg, Germany ¹¹⁸Department of Medical Genetics and Development, University of Geneva Medical School, Geneva, Switzerland ¹¹⁹Cardiology Division, Massachusetts General Hospital, Boston, Massachusetts, USA ¹²⁰National Heart, Lung, and Blood Institute, Bethesda, Maryland, USA ¹²¹MRC Unit for Lifelong Health and Ageing, London, UK ¹²²Department of Epidemiology, University of Pittsburgh, Pittsburgh, Pennsylvania, USA ¹²³Institute of Epidemiology II, Helmholtz Zentrum München–German Research Center for Environmental Health, Neuherberg, Germany ¹²⁴Department of Public Health, Faculty of Medicine, University of Split, Split, Croatia ¹²⁵National Institute for Health and Welfare, Oulu, Finland ¹²⁶Department of Neurology, General Central Hospital, Bolzano, Italy ¹²⁷Department of Neurology, University of Lübeck, Lübeck, Germany ¹²⁸Group Health Research Institute, Seattle, Washington, USA ¹²⁹Department of Health Services, University of Washington, Seattle, Washington, USA ¹³⁰Division of Biostatistics, Washington University School of Medicine, St. Louis, Missouri, USA ¹³¹Analytic and Translational Genetics Unit, Massachusetts General Hospital, Boston, Massachusetts, USA ¹³²Harvard Biological and Biomedical Sciences Program, Harvard University, Boston, Massachusetts, USA ¹³³Health Science and Technology M.D. Program, Harvard University and Massachusetts Institute of Technology, Boston, Massachusetts, USA ¹³⁴Department of Pathophysiology, Faculty of Medicine, University of Split, Split, Croatia ¹³⁵Istituto di Ricerca Genetica e Biomedica, Consiglio Nazionale delle Ricerche (CNR), Monserrato, Italy ¹³⁶National Heart and Lung Institute, Imperial College London, London, UK ¹³⁷Laboratory of Neurogenetics, National Institute on Aging, Bethesda, Maryland, USA ¹³⁸Department of Twin Research and Genetic Epidemiology Unit, St. Thomas' Campus, King's College London, St. Thomas' Hospital, London, UK ¹³⁹Biology Department, Boston College, Chestnut Hill, Massachusetts, USA ¹⁴⁰Department of Internal Medicine, Erasmus Medical Center, Rotterdam, The Netherlands ¹⁴¹genetic Epidemiology Laboratory, Queensland Institute of Medical Research, Brisbane, Queensland, Australia ¹⁴²Centre for Population Health Sciences, University of Edinburgh, Edinburgh, UK ¹⁴³Division of Biomedical Sciences, St. George's University of London, London, UK ¹⁴⁴Department of Physiatrics, Lapland Central Hospital, Rovaniemi, Finland ¹⁴⁵Centre for Nutrition, Prevention and Health Services, National Institute for Public Health and the Environment (RIVM), Bilthoven, The Netherlands ¹⁴⁶Department of Sports Medicine, Norwegian School of Sport Sciences, Oslo, Norway ¹⁴⁷Genomic Medicine,

Hammersmith Hospital, Imperial College London, London, UK ¹⁴⁸Wellcome Trust Centre for Human Genetics, University of Oxford, Oxford, UK ¹⁴⁹Department of Medical Sciences, Uppsala University, Akademiska Sjukhuset, Uppsala, Sweden ¹⁵⁰Department of Clinical Genetics, Erasmus University Medical Center, Rotterdam, The Netherlands ¹⁵¹Medical Genetics Institute, Cedars-Sinai Medical Center, Los Angeles, California, USA ¹⁵²Institute of Cellular Medicine, Newcastle University, Newcastle, UK ¹⁵³Department of Medicine, Landspítali University Hospital, Reykjavik, Iceland ¹⁵⁴Faculty of Medicine, University of Iceland, Reykjavik, Iceland ¹⁵⁵Geriatric Unit, Azienda Sanitaria Firenze (ASF), Florence, Italy ¹⁵⁶Channing Laboratory and Renal Division, Brigham and Women's Hospital, Boston, Massachusetts, USA ¹⁵⁷Department of Medicine, Harvard Medical School, Boston, Massachusetts, USA ¹⁵⁸Department of General Practice and Primary Health Care, University of Helsinki, Helsinki, Finland ¹⁵⁹Folkhälsan Research Centre, Helsinki, Finland ¹⁶⁰Unit of General Practice, Helsinki University Central Hospital, Helsinki, Finland ¹⁶¹Hypertension and Related Diseases Centre—Azienda Ospedaliero Universitaria (AOU), University of Sassari, Sassari, Italy ¹⁶²Program in Molecular and Genetic Epidemiology, Harvard School of Public Health, Boston, Massachusetts, USA ¹⁶³Division of Research, Kaiser Permanente of Northern California, Oakland, California, USA ¹⁶⁴Department of Social Services and Health Care, Jakobstad, Finland ¹⁶⁵Biocenter Oulu, University of Oulu, Oulu, Finland ¹⁶⁶Institute of Health Sciences, University of Oulu, Oulu, Finland ¹⁶⁷Department of Chronic Disease Prevention, National Institute for Health and Welfare, Turku, Finland ¹⁶⁸Department of Clinical Chemistry, University of Tampere and Tampere University Hospital, Tampere, Finland ¹⁶⁹Department for Health Evidence, Radboud University Medical Centre, Nijmegen, The Netherlands ¹⁷⁰Department of Urology, Radboud University Nijmegen Medical Centre, Nijmegen, The Netherlands ¹⁷¹Department of Endocrinology, University Medical Center Groningen, University of Groningen, Groningen, The Netherlands ¹⁷²Laboratory of Genetic and Environmental Epidemiology, Fondazione di Ricerca e Cura Giovanni Paolo II, Catholic University, Campobasso, Italy ¹⁷³transplantation Laboratory, Haartman Institute, University of Helsinki, Helsinki, Finland ¹⁷⁴Department of Internal Medicine, University Medical Center Groningen, University of Groningen, Groningen, The Netherlands ¹⁷⁵Department of Medicine, Division of Cardiology, Helsinki University Central Hospital, Helsinki, Finland ¹⁷⁶International Centre for Circulatory Health (ICCH), Imperial College London, London, UK ¹⁷⁷Research Centre of Applied and Preventive Cardiovascular Medicine, University of Turku, Turku, Finland ¹⁷⁸Department of Clinical Physiology and Nuclear Medicine, University of Turku and Turku University Hospital, Turku, Finland ¹⁷⁹Kos Genetic, Milan, Italy ¹⁸⁰Complex and Adaptive Systems Laboratory, University College Dublin, Belfield, Dublin, Ireland ¹⁸¹Conway Institute of Biomolecular and Biomedical Research, University College Dublin, Belfield, Dublin, Ireland ¹⁸²School of Medicine and Medical Sciences, University College Dublin, Belfield, Dublin, Ireland ¹⁸³Geriatric Research and Education Clinical Center, Veterans Administration Medical Center, Baltimore, Maryland, USA ¹⁸⁴Molecular and Cellular Therapeutics, Royal College of Surgeons

in Ireland, Dublin, Ireland ¹⁸⁵Diabetes Unit, National Institute for Health and Welfare, Helsinki, Finland ¹⁸⁶Centre for Vascular Prevention, Danube-University Krems, Krems, Austria ¹⁸⁷Red Temática de Investigación Cooperativa en Enfermedades Cardiovasculares (Red RECAVA), Grupo RD06/0014/0015, Hospital Universitario La Paz, Madrid, Spain ¹⁸⁸King Abdulaziz University, Jeddah, Saudi Arabia ¹⁸⁹Department of Medicine, University of Turku and Turku University Hospital, Turku, Finland ¹⁹⁰Department of Internal Medicine, University Hospital of Lausanne, Lausanne, Switzerland ¹⁹¹Institute for Molecular Medicine Finland (FIMM), University of Helsinki, Helsinki, Finland ¹⁹²Department of Biomedical Science, Hallym University, Chuncheon, Korea ¹⁹³Queensland Brain Institute, University of Queensland, Brisbane, Queensland, Australia ¹⁹⁴Department of Internal Medicine, University of Michigan, Ann Arbor, Michigan, USA ¹⁹⁵Blizard Institute of Cell and Molecular Science, Barts and The London School of Medicine and Dentistry, Queen Mary University of London, London, UK ¹⁹⁶A full list of members appears in the Supplementary Note ¹⁹⁷Institute for Integrative and Experimental Genomics, University of Lübeck, Lübeck, Germany ¹⁹⁸DZHK, partner site Hamburg/Kiel/Lübeck, Lübeck, Germany ¹⁹⁹Department of Health Sciences, University of Leicester, Leicester, UK ²⁰⁰Institute of Human Genetics, Helmholtz Zentrum München—German Research Center for Environmental Health, Munich, Germany ²⁰¹Institute of Human Genetics, Klinikum Rechts der Isar der Technischen Universität München, Munich, Germany ²⁰²Department of Medicine, Division of Cardiology, University of Washington School of Medicine, Seattle, Washington, USA ²⁰³Center for Human Genetic Research, Cardiovascular Research Center, Massachusetts General Hospital, Boston, Massachusetts, USA ²⁰⁴Cardiac Arrhythmia Service, Massachusetts General Hospital, Boston, Massachusetts, USA ²⁰⁵Cardiovascular Research Center, Massachusetts General Hospital, Charlestown, Massachusetts, USA ²⁰⁶Inspectorate of Health Care, The Hague, The Netherlands ²⁰⁷Service of Medical Genetics, CHUV University Hospital, Lausanne, Switzerland ²⁰⁸Center for Human Genetic Research, Massachusetts General Hospital, Boston, Massachusetts, USA ²⁰⁹Department of Cardiovascular Sciences, University of Leicester, Glenfield Hospital, Leicester, UK ²¹⁰NIHR Leicester Biomedical Research Unit in Cardiovascular Disease, Glenfield Hospital, Leicester, UK ²¹¹The Genetics of Obesity and Related Metabolic Traits Program, The Icahn School of Medicine at Mount Sinai, New York, New York, USA ²¹²The Charles Bronfman Institute of Personalized Medicine, The Icahn School of Medicine at Mount Sinai, New York, New York, USA ²¹³The Mindich Child Health and Development Institute, The Icahn School of Medicine at Mount Sinai, New York, New York, USA

Acknowledgments

A full list of acknowledgments appears in the Supplementary Note. Funding sources had no involvement in the collection, analysis and interpretation of the data.

AUTHOR CONTRIBUTIONS

Steering committee (oversaw the project): M. den Hoed (lead) and R.J.F.L. (chair).
Writing group (drafted the manuscript): M.E., M. den Hoed (chair), R.J.F.L. and N.J.S.
Editing group (edited the manuscript): B.J.J.M.B., P.T.E., T.E., D.M.E., E.J.C.d.G., M. den Hoed (chair), E.I., D.J.M., R.J.F.L., A.L., D.J.M., I.M.N., A.V. Segrè, O.C.M.S., H.S., J.R.T. and N.J.T. **Meta-analysis working group (performed stage 1 and stage 2 meta-analyses):** T.E. and M. den Hoed (chair). **Data preparation working group (prepared data from contributing cohorts for meta-analyses):** M.E., T.E., M. den Hoed (lead) and R.J.F.L. (chair). **Conditional analyses:** M. den Hoed, R.J.F.L., P.M.V. (chair) and J.Y. (lead). **Genetic predisposition score analyses:** D.M.E., M. den Hoed (chair) and I.M.N.
Association analyses with related traits: C.M.A., P.I.W.d.B., CARDIOGRAM Consortium, CHARGE-AF Consortium, Y.S.C., M.C., D.D., P.T.E., J. Erdmann, Global BPgen Consortium, M.J.G., M. den Hoed (chair), H.H., A.I., T.J., S. Kääh, Y.J.K., K.L.L., P.B.M., C.N.-C., A. Pfeufer, PR GWAS Consortium, QRS GWAS Consortium, QT-IGC Consortium, N.J.S., S. Sharp, N. Sotoodehnia and J.R.T. **Copy number variant analyses:** R.E.H. (lead), M. den Hoed (chair), S.A.M. and C. Stewart. **Gene eQTL analyses:** L. Franke (chair), M. den Hoed and H.-J.W. (lead). **Proteomics experiments and genetic enrichment analyses:** M. den Hoed, R.J.F.L., A.L. (lead), E.J.R. and J.V.O. (chair). **SNIPPER analyses for selection of positional candidate genes:** M. den Hoed (chair), R.J.F.L. and C. Willer (lead). **Pathway analyses:** M. den Hoed (chair), R.J.F.L. and A.V. Segrè (lead). **D. melanogaster experiments:** B.J.J.M.B. (lead), M. den Hoed, F.H.-B., B.K., R.J.F.L., O.C.M.S. (chair) and H.S. **D. rerio experiments:** M. den Hoed, R.J.F.L., S.N.L., D.J.M. (chair), D.S.P. (lead) and J.T.S.

Project design, management and coordination of contributing cohorts

Stage 1–GWAS: (ADVANCE) T.L.A., C.I. and T.Q.; (ALSPAC) G.D.S.; (ASCOT cases) N.R.P., P.S.S., D.C.S. and A.V. Stanton; (ATBC) D. Albanes and J. Virtamo; (B58C) W.L.M.C. and D.P.S.; (BLSA) S. Bandinelli and L. Ferrucci; (BRIGHT) M.C., T.J., P.B.M. and N.J.S.; (CoLaus) J.S.B., P.V. and G. Waeber; (COROGENE) M.-L.L.L., M.S.N., M.P. and J.S.; (deCODE) D.O.A., K.S. and U.T.; (DGI) L.G. and B.I.; (EGCUT) A.M.; (EPIC-Norfolk) N.J.W.; (Fenland) U.E., N.G.F., R.J.F.L. and N.J.W.; (Fingesture) H.V.H., J.D.R. and J.-C.T.; (Finrisk07) M.P. and V.S.; (FUSION) M. Boehnke and J.T.; (GOOD) C.O.; (HAPI) B.D.M. and A.R.S.; (HBSC) J. Eriksson, M.P. and E.W.; (Health 2000) A.J. and M.P.; (Health ABC) W.-C.H.; (HERITAGE) C. Bouchard, T.R. and D.C.R.; (HPFS) G.C., F.B.H., D.J.H., P.K., L.Q. and E.B.R.; (Hypergenes) D.C., N.G., L.I. and F.R.; (InCHIANTI) S. Bandinelli and L. Ferrucci; (Korcula) I.R.; (LifeLines) R.A.d.B., M.M.v.d.K., H.S. and R.P.S.; (Lolipop) J.C.C. and J.S.K.; (NBS) J.d.G. and L.A.K.; (NFBC1966) M.-R.J.; (NHS) G.C., F.B.H., D.J.H., P.K., L.Q. and E.B.R.; (NSPHS) U.G.; (PREVEND) W.H.v.G., G.N. and D.J.v.V.; (SPLIT) I.R.; and (YFS) M. Kähönen, T.L., M.P., O.T.R. and J. Viikari. **Stage 2–in silico replication studies:** (AGES, RRgen) V.G. and T.B.H.; (ACTS) N.G.M.; (ALSPAC) G.D.S.; (ARIC, RRgen) A.A.; (CHS, RRgen) B.M.P.; (DESIR) N.B.-N.; (EGCUT) A.M.; (Ely) N.J.W.; (EPIC-NL) J.M.A.B., Y.T.v.d.S. and W.M.M.V.; (EPIC-Norfolk) N.J.W.; (ERF) C.M.v.D. and B.A.O.; (FamHS) I.B.B.;

(Fenland) U.E., N.G.F., R.J.F.L. and N.J.W.; (FHS, RRgen) C.J.O.; (Finrisk07) M.P. and V.S.; (KORA, RRgen) A. Peters and S. Kääb; (LifeLines2) R.A.d.B., M.M.v.d.K., H.S. and R.P.S.; (MESA) R.A.K. and J.I.R.; (MICROS, RRgen) P.P.P.; (NSHD) D.K.; (NTR) D.I.B. and E.J.C.d.G.; (ORCADES, RRgen) J.F.W.; (RISC) M.W.; (PIVUS) E.I. and L.L.; (RS1-3) A. Hofman, B.H.Ch.S. and J.C.M.W.; (SardiNIA, RRgen) E.G.L. and K.V.T.; (SHIP, RRgen) M.D. and S.B.F.; (Stanford IST) T.Q.; (STR) E.I. and N.L.P.; (Twins UK, RRgen) Y.J. and T.D.S.; (ULSAM) E.I.; and (Whitehall II) A. Hingorani and M. Kivimaki.

Genotyping of contributing cohorts

Stage 1–GWAS: (ADVANCE) D. Absher; (ALSPAC) S.M.R. and W.L.M.; (ATBC) S.J.C.; (BLSA) L. Ferrucci and A.B.S.; (BRIGHT) M.C. and P.B.M.; (COROGENE) P.S.; (EGCUT) T.E., L.M. and M.N.; (EPIC-Norfolk) R.J.F.L. and J.H.Z.; (Fenland) J.L.; (Fingesture) P.G. and J.D.R.; (Finrisk07) P.S.; (FUSION) P.S.C.; (GOOD) M. Lorentzon and C.O.; (HBCS) P.S.; (Health2000) P.L. and P.S.; (Health ABC) Y.L.; (HERITAGE) C. Bouchard and T.R.; (HPFS) M.C.C. and M.K.J.; (Hypergenes) C. Barlassina and P.B.; (InCHIANTI) L. Ferrucci and A.B.S.; (Korcula) C.H.; (LifeLines) L. Franke; (Lolipop) J.C.C. and J.S.K.; (NBS) L.A.K.; (NFBC1966) P.E., A.-L.H., M.-R.J. and P.Z.; (NHS) M.C.C. and M.K.J.; (NSPHS) Å.J.; (PREVEND) P.v.d.H.; (SPLIT) C.H. and V.V.; and (YFS) M. Kähönen, T.L., M.P., O.T.R., P.S. and J. Viikari. **Stage 2–*in silico* replication studies:** (ACTS) N.G.M., S.E.M. and G.W.M.; (ALSPAC) S.M.R. and W.L.M.; (ARIC, RRgen) D.E.A.; (DESIR) N.B.-N.; (EGCUT) T.E., L.M. and M.N.; (EPIC-NL) N.C.O.-M. and C. Wijmenga; (ERF) C.M.v.D., A.I. and B.A.O.; (Ely) R.J.F.L. and J.L.; (EPIC-Norfolk) R.J.F.L. and J.H.Z.; (FamHS) I.B.B. and M.F.F.; (Fenland) J.L.; (Finrisk07) P.S.; (LifeLines2) L. Franke; (MESA) J.I.R.; (NSHD) D.K., K.K.O. and A.W.; (NTR) D.I.B. and J.-J.H.; (PIVUS) E.I. and L.L.; (RS1-3) A.G.U.; (Stanford IST) T.L.A. and J.W.K.; (STR) E.I. and N.L.P.; (ULSAM) E.I.; and (Whitehall II) M. Kumari and C. Langenberg.

Phenotyping of contributing cohorts

Stage 1–GWAS: (ADVANCE) C.I.; (ASCOT cases) N.R.P., P.S.S. and A.V. Stanton; (ATBC) D. Albanes and J. Virtamo; (B58C) D.P.S.; (BLSA) S. Bandinelli and L. Ferrucci; (BRIGHT) M.C. and N.J.S.; (CoLaus) P.M.-V.; (COROGENE) M.P.; (deCODE) D.O.A. and H.H.; (DGI) B.I.; (EGCUT) K.F. and A.M.; (EPIC-Norfolk) K.-T.K.; (Fingesture) H.V.H. and J.J.; (Finrisk07) M.P.; (FUSION) H.M.S.; (GOOD) M. Lorentzon, C.O. and L.V.; (HBCS) J. Eriksson, M.P. and E.W.; (Health2000) A.J. and M.P.; (Health ABC) A.B.N.; (HERITAGE) C. Bouchard; (Hypergenes) D.C., N.G., L.I. and F.R.; (InCHIANTI) S. Bandinelli and L. Ferrucci; (Korcula) O.P.; (LifeLines) R.A.d.B., M.M.v.d.K. and R.P.S.; (Lolipop) J.C.C., A.S.K., J.S.K., K.A.M. and J.S.S.; (NBS) S.H.; (NFBC1966) A.-L.H., M.-R.J., A. Pouta and P.Z.; (PREVEND) R.A.d.B., W.H.v.G. and P.v.d.H.; (SPLIT) D.R.; and (YFS) M. Kähönen, T.L., M.P., O.T.R., P.S. and J. Viikari. **Stage 2–*in silico* replication studies:** (AGES, RRgen) V.G.; (ACTS) N.G.M. and J.B.W.; (CHS, RRgen) N. Sotoodehnia; (DESIR) B.B. and P.F.; (EGCUT) K.F. and A.M.; (Ely) S. Brage and U.E.; (EPIC-NL) J.M.A.B., Y.T.v.d.S. and W.M.M.V.; (EPIC-Norfolk) K.-T.K.; (ERF) C.M.v.D., A.I., J.A.K. and B.A.O.; (FamHS) I.B.B. and M.F.F.; (FHA, RRgen) C.N.-C.; (Finrisk07) M.P.; (LifeLines2) R.A.d.B., M.M.v.d.K. and R.P.S.; (MESA) S.R.H. and R.A.K.;

(MICROS, RRgen) A.A.H.; (NSHD) D.K.; (NTR) D.I.B., E.J.C.d.G. and G. Willemsen; (ORCADES, RRgen) S.H.W.; (PIVUS) E.I. and L.L.; (RISC) M.W.; (RS1-3) B.H.Ch.S. and A.G.U.; (SHIP, RRgen) M.D. and M.R.P.M.; (Stanford IST) T.L.A. and J.W.K.; (STR) E.I. and N.L.P.; (ULSAM) E.I.; and (Whitehall II) M. Kumari.

Analyses of contributing cohorts

Stage 1–GWAS: (ADVANCE) T.L.A. and L.W.; (ALSPAC) D.M.E., J.P.K., B.S.P. and N.J.T.; (ASCOT cases) T.J.; (ATBC) W.W.; (B58C) D.H. and D.P.S.; (BLSA) T.T.; (BRIGHT) T.J. and S.P.; (CoLaus) M. Bochud and Z.K.; (COROGENE) P.S.; (deCODE) D.G. and H.H.; (DGI) P.A., C. Ladenvall and R.A.S.; (EGCUT) T.E. and E.M.; (EPIC-Norfolk) M. den Hoed, R.N.L. and J.H.Z.; (Fenland) M. den Hoed and J.L.; (Fingesture) G.B. and P.G.; (Finrisk07) A.S.H., K.K. and P.S.; (FUSION) A.U.J.; (GOOD) M. Lorentzon, C.O. and L.V.; (HAPI) M.E.M. and J.R.O.; (HBCS,) P.S.; (Health2000) P.S.; (Health ABC) W.-C.H. and O.T.N.; (HERITAGE) C. Bouchard, T.R. and D.C.R.; (HPFS) M.C.C. (InCHIANTI) T.T.; (Korcula) C.H.; (LifeLines) I.M.N. and H.S.; (Lolipop) J.C.C., J.S.K., J.S.S. and W.Z.; (NBS) M. den Heijer; (NFBC1966) P.F.O.; (NHS) M.C.C.; (Hypergenes) D.C.; (NSPHS) W.I.; (PREVEND) P.v.d.H. and I.M.L.; (SPLIT) C.H. and V.V.; and (YFS) P.S. **Stage 2– *in silico* replication studies:** (AGES, RRgen) A.V. Smith; (ACTS) P.A.L.; (ALSPAC) D.M.E., J.P.K., B.S.P. and N.J.T.; (ARIC, RRgen) A.C.M.; (CHS, RRgen) J.C.B. and N. Sotoodehnia; (DESIR) C.D., N.B.-N. and L.Y.; (EGCUT) T.E. and E.M.; (Ely) M. den Hoed and J.L.; (EPIC-NL) M. Leusink and N.C.O.-M.; (EPIC-Norfolk) M. den Hoed, R.N.L. and J.H.Z.; (ERF) A.I.; (FamHS) M.F.F. and S. Ketkar; (Fenland) M. den Hoed and J.L.; (FHS, RRgen) C.N.-C. and S.-J.H.; (Finrisk07) A.S.H. and K.K., P.S.; (KORA, RRgen) M.M.-N.; (LifeLines2) I.M.N. and H.S.; (MESA) K.F.K. and Q.W.; (MICROS, RRgen) C.F.; (NSHD) M. den Hoed, J.L. and A.W.; (NTR) H.H.M.D. and J.-J.H.; (ORCADES, RRgen) P.N.; (PIVUS) E.I. and C. Song; (RISC) M.N.W. and W.X.; (RRgen) P.I.W.d.B.; (RS1-3) P.I.W.d.B. and M.E.; (SardinIA, RRgen) S. Sanna; (Stanford IST) W.X.; (STR) E.I. and C. Song; (Twins UK, RRgen) N. Soranzo; (ULSAM) E.I. and C. Song; and (Whitehall II) M. den Hoed and J.L.

The corresponding author (R.J.F.L.) had full access to all the data in the study and had final responsibility for the decision to submit for publication.

References

1. Dyer AR, et al. Heart rate as a prognostic factor for coronary heart disease and mortality: findings in three Chicago epidemiologic studies. *Am J Epidemiol.* 1980; 112:736–749. [PubMed: 7457467]
2. Gillum RF, Makuc DM, Feldman JJ. Pulse rate, coronary heart disease, and death: the NHANES I Epidemiologic Follow-up Study. *Am Heart J.* 1991; 121:172–177. [PubMed: 1985358]
3. Nauman J, Janszky I, Vatten LJ, Wisloff U. Temporal changes in resting heart rate and deaths from ischemic heart disease. *J Am Med Assoc.* 2011; 306:2579–2587.
4. Schwartz PJ, Billman GE, Stone HL. Autonomic mechanisms in ventricular fibrillation induced by myocardial ischemia during exercise in dogs with healed myocardial infarction. An experimental preparation for sudden cardiac death. *Circulation.* 1984; 69:790–800. [PubMed: 6697463]
5. Beere PA, Glagov S, Zarins CK. Retarding effect of lowered heart rate on coronary atherosclerosis. *Science.* 1984; 226:180–182. [PubMed: 6484569]

6. Beere PA, Glagov S, Zarins CK. Experimental atherosclerosis at the carotid bifurcation of the cynomolgus monkey. Localization, compensatory enlargement, and the sparing effect of lowered heart rate. *Arterioscler Thromb*. 1992; 12:1245–1253. [PubMed: 1420083]
7. Böhm M, et al. Heart rate as a risk factor in chronic heart failure (SHIFT): the association between heart rate and outcomes in a randomised placebo-controlled trial. *Lancet*. 2010; 376:886–894. [PubMed: 20801495]
8. Dalageorgou C, et al. Heritability of QT interval: how much is explained by genes for resting heart rate? *J Cardiovasc Electrophysiol*. 2008; 19:386–391. [PubMed: 18031506]
9. De Geus EJ, Kupper N, Boomsma DI, Snieder H. Bivariate genetic modeling of cardiovascular stress reactivity: does stress uncover genetic variance? *Psychosom Med*. 2007; 69:356–364. [PubMed: 17510291]
10. Russell MW, Law I, Sholinsky P, Fabsitz RR. Heritability of ECG measurements in adult male twins. *J Electrocardiol*. 1998; 30(suppl):64–68. [PubMed: 9535482]
11. Cho YS, et al. A large-scale genome-wide association study of Asian populations uncovers genetic factors influencing eight quantitative traits. *Nat Genet*. 2009; 41:527–534. [PubMed: 19396169]
12. Eijgelsheim M, et al. Genome-wide association analysis identifies multiple loci related to resting heart rate. *Hum Mol Genet*. 2010; 19:3885–3894. [PubMed: 20639392]
13. Holm H, et al. Several common variants modulate heart rate, PR interval and QRS duration. *Nat Genet*. 2010; 42:117–122. [PubMed: 20062063]
14. Yang J, et al. Conditional and joint multiple-SNP analysis of GWAS summary statistics identifies additional variants influencing complex traits. *Nat Genet*. 2012; 44:369–375. [PubMed: 22426310]
15. Inoue T, et al. Higher heart rate predicts the risk of developing hypertension in a normotensive screened cohort. *Circ J*. 2007; 71:1755–1760. [PubMed: 17965497]
16. Pfeufer A, et al. Genome-wide association study of PR interval. *Nat Genet*. 2010; 42:153–159. [PubMed: 20062060]
17. Sotoodehnia N, et al. Common variants in 22 loci are associated with QRS duration and cardiac ventricular conduction. *Nat Genet*. 2010; 42:1068–1076. [PubMed: 21076409]
18. Holm H, et al. A rare variant in *MYH6* is associated with high risk of sick sinus syndrome. *Nat Genet*. 2011; 43:316–320. [PubMed: 21378987]
19. Garrison RJ, Kannel WB, Stokes J III, Castelli WP. Incidence and precursors of hypertension in young adults: the Framingham Offspring Study. *Prev Med*. 1987; 16:235–251. [PubMed: 3588564]
20. Levy RL, et al. Transient tachycardia; prognostic significance alone and in association with transient hypertension. *Med Press Egypt*. 1946; 38:207–212. [PubMed: 20278752]
21. Newton-Cheh C, et al. Genome-wide association study identifies eight loci associated with blood pressure. *Nat Genet*. 2009; 41:666–676. [PubMed: 19430483]
22. Schunkert H, et al. Large-scale association analysis identifies 13 new susceptibility loci for coronary artery disease. *Nat Genet*. 2011; 43:333–338. [PubMed: 21378990]
23. Kusumoto FM, Goldschlager N. Cardiac pacing. *N Engl J Med*. 1996; 334:89–97. [PubMed: 8531965]
24. Segrè AV, et al. Common inherited variation in mitochondrial genes is not enriched for associations with type 2 diabetes or related glycemic traits. *PLoS Genet*. 2010; 6:pii: e1001058.
25. Ellinor PT, et al. Meta-analysis identifies six new susceptibility loci for atrial fibrillation. *Nat Genet*. 2012; 44:670–675. [PubMed: 22544366]
26. Li Y, Willer C, Sanna S, Abecasis G. Genotype imputation. *Annu Rev Genomics Hum Genet*. 2009; 10:387–406. [PubMed: 19715440]
27. Marchini J, Howie B, Myers S, McVean G, Donnelly P. A new multipoint method for genome-wide association studies by imputation of genotypes. *Nat Genet*. 2007; 39:906–913. [PubMed: 17572673]
28. Guan Y, Stephens M. Practical issues in imputation-based association mapping. *PLoS Genet*. 2008; 4:e1000279. [PubMed: 19057666]
29. Ellinghaus D, Schreiber S, Franke A, Nothnagel M. Current software for genotype imputation. *Hum Genomics*. 2009; 3:371–380. [PubMed: 19706367]

30. Pei YF, Zhang L, Li J, Deng HW. Analyses and comparison of imputation-based association methods. *PLoS ONE*. 2010; 5:e10827. [PubMed: 20520814]
31. Abecasis GR, Wigginton JE. Handling marker-marker linkage disequilibrium: pedigree analysis with clustered markers. *Am J Hum Genet*. 2005; 77:754–767. [PubMed: 16252236]
32. Aulchenko YS, Struchalin MV, van Duijn CM. ProbABEL package for genomewide association analysis of imputed data. *BMC Bioinformatics*. 2010; 11:134. [PubMed: 20233392]
33. Aulchenko YS, Ripke S, Isaacs A, van Duijn CM. GenABEL: an R library for genome-wide association analysis. *Bioinformatics*. 2007; 23:1294–1296. [PubMed: 17384015]
34. Purcell S, et al. PLINK: a tool set for whole-genome association and population-based linkage analyses. *Am J Hum Genet*. 2007; 81:559–575. [PubMed: 17701901]
35. Willer CJ, Li Y, Abecasis GR. METAL: fast and efficient meta-analysis of genomewide association scans. *Bioinformatics*. 2010; 26:2190–2191. [PubMed: 20616382]
36. Voight BF, et al. The metabochip, a custom genotyping array for genetic studies of metabolic, cardiovascular, and anthropometric traits. *PLoS Genet*. 2012; 8:e1002793. [PubMed: 22876189]
37. Newton-Cheh C, et al. Common variants at ten loci influence QT interval duration in the QTGEN Study. *Nat Genet*. 2009; 41:399–406. [PubMed: 19305408]
38. Thomas PD, et al. PANTHER: a library of protein families and subfamilies indexed by function. *Genome Res*. 2003; 13:2129–2141. [PubMed: 12952881]
39. Lundby A, et al. Quantitative maps of protein phosphorylation sites across 14 different rat organs and tissues. *Nat Commun*. 2012; 3:876. [PubMed: 22673903]
40. Dubois PC, et al. Multiple common variants for celiac disease influencing immune gene expression. *Nat Genet*. 2010; 42:295–302. [PubMed: 20190752]
41. Johnson AD, et al. SNAP: a web-based tool for identification and annotation of proxy SNPs using HapMap. *Bioinformatics*. 2008; 24:2938–2939. [PubMed: 18974171]
42. Handsaker RE, Korn JM, Nemesh J, McCarroll SA. Discovery and genotyping of genome structural polymorphism by sequencing on a population scale. *Nat Genet*. 2011; 43:269–276. [PubMed: 21317889]
43. Clemens JC, et al. Use of double-stranded RNA interference in *Drosophila* cell lines to dissect signal transduction pathways. *Proc Natl Acad Sci USA*. 2000; 97:6499–6503. [PubMed: 10823906]
44. Dietzl G, et al. A genome-wide transgenic RNAi library for conditional gene inactivation in *Drosophila*. *Nature*. 2007; 448:151–156. [PubMed: 17625558]
45. Zhang D, et al. Effects of different small HSPB members on contractile dysfunction and structural changes in a *Drosophila melanogaster* model for atrial fibrillation. *J Mol Cell Cardiol*. 2011; 51:381–389. [PubMed: 21745477]
46. Burns CG, et al. High-throughput assay for small molecules that modulate zebrafish embryonic heart rate. *Nat Chem Biol*. 2005; 1:263–264. [PubMed: 16408054]
47. Shin JT, Pomerantsev EV, Mably JD, MacRae CA. High-resolution cardiovascular function confirms functional orthology of myocardial contractility pathways in zebrafish. *Physiol Genomics*. 2010; 42:300–309. [PubMed: 20388839]




Figure 1.

Manhattan plot of SNPs after meta-analysis of stage 1. The plot shows the significance of associations between all SNPs and heart rate in stage 1. The 7 loci that were previously identified are highlighted in light blue; the 14 newly associated loci are highlighted in dark blue. Loci that reached $P < 3 \times 10^{-5}$ after stage 1 but did not reach $P < 5 \times 10^{-8}$ after meta-analysis of stages 1 and 2 combined are highlighted in red.




Figure 2.

Effect size as a function of effect allele frequency. The plot shows the effect sizes of the 21 heart rate-associated SNPs after joint meta-analysis of stage 1 and stage 2 results as a function of their effect allele frequencies. Light-blue circles represent the 7 previously identified heart rate loci; dark blue circles represent the 14 newly identified heart rate loci.




Figure 3. Combined effect of heart rate-increasing alleles on heart rate. **(a)** Combined effect of the 19 available heart rate loci in adults of European descent (LifeLines2, $n = 5,053$). **(b)** Combined effect of the 21 heart rate loci in 12-year-old children of European descent (ALSPAC, $n = 4,000$). In each plot, the number of heart rate-increasing alleles was summed across the heart rate-associated SNPs. The number of heart rate-increasing alleles is shown (x axis), grouped at the extremes, and mean heart rate \pm s.e.m. is plotted (right y axis). The lines represent the regression of the mean heart rate values across the GPS distribution. The histogram shows the number of individuals in each GPS window (left y axis).




Figure 4.

Effects on heart rate of reduced or ablated expression of orthologs of positional candidate genes from GWAS in *D. melanogaster* and *D. rerio*. Bars show the heart rate (\pm s.e.m.) of (i) *D. melanogaster* pupae with orthologs of positional candidate genes located within 500 kb of GWAS associations downregulated using RNAi compared with control pupae (blue bars) and (ii) *D. rerio* (zebrafish) embryos with expression of orthologs of positional candidate genes reduced by injecting morpholino oligonucleotides versus PBS (red bars). Darker coloring indicates that heart rate is significantly different in targeted animals compared with controls after Bonferroni correction for 23 tests in *D. melanogaster* ($P < 2 \times 10^{-3}$) and 12 tests in *D. rerio* ($P < 4 \times 10^{-3}$); lighter coloring indicates that differences do not reach significance. Results are ordered by availability (*D. melanogaster* and *D. rerio*, *D. melanogaster* only, *D. rerio* only) and by effect size.

Table 1

Stage 1 and stage 2 results for the loci that showed association with heart rate at genome-wide significance ($P < 5 \times 10^{-8}$)

Locus	Chr.	Nearest gene	Annotation	Heart rate SNP	Position (bp) ^a	Alleles		EAF	Per-allele change in heart rate ^b			Stage 1		Stage 2		Stages 1 + 2		
						Effect	Other		β	SE	P	P	N	P	N	P		
Previously identified heart rate loci																		
1	14	<i>MYH6</i>	N,P,S	rs365990	22931651	G	A	0.353	0.564	0.047	5.66 × 10 ⁻²⁶	2.19 × 10 ⁻²¹	148,972	5.39 × 10 ⁻⁴⁵				
2	6	<i>GJA1</i>	S	rs1015451	122173184	C	T	0.102	0.713	0.072	7.06 × 10 ⁻²⁰	2.14 × 10 ⁻¹⁷	155,303	1.14 × 10 ⁻³²				
3	7	<i>ACHE</i>	N,Q,S,B	rs13245899	100335067	G	A	0.195	0.447	0.055	1.54 × 10 ⁻¹⁵	3.27 × 10 ⁻¹⁴	176,643	7.67 × 10 ⁻²⁷				
4	1	<i>CD46</i>	S,B	rs11118555	206007476	A	T	0.124	0.612	0.069	4.35 × 10 ⁻¹⁴	7.49 × 10 ⁻¹⁵	166,654	3.88 × 10 ⁻²⁶				
5	11	<i>FADS1</i>	Q,S,B	rs174549	61327958	A	G	0.310	0.358	0.047	1.13 × 10 ⁻¹²	1.57 × 10 ⁻¹²	172,847	1.38 × 10 ⁻²²				
6	6	<i>SLC35F1</i>	T,Q,P,S	rs1153730	118774215	T	C	0.509	0.381	0.044	7.01 × 10 ⁻¹⁵	8.79 × 10 ⁻⁹	156,783	7.55 × 10 ⁻²¹				
7	12	<i>LINC00477 (C12orf67)</i>		rs17287293	24662145	A	G	0.850	0.444	0.062	6.90 × 10 ⁻¹²	3.98 × 10 ⁻¹¹	151,085	3.07 × 10 ⁻²⁰				
Newly identified heart rate loci																		
8	20	<i>KIAA1755</i>	N,S	rs6127471	36277452	C	T	0.540	0.429	0.045	5.96 × 10 ⁻²³	2.98 × 10 ⁻¹⁰	162,593	5.22 × 10 ⁻²⁹				
9	2	<i>CCDC141</i>	N,P,S	rs17362588	179429291	A	G	0.114	0.736	0.077	3.25 × 10 ⁻¹⁸	4.22 × 10 ⁻¹¹	136,061	3.57 × 10 ⁻²⁶				
10	12	<i>SYT10</i>		rs7980799	33468257	A	C	0.401	0.377	0.046	7.91 × 10 ⁻¹⁴	7.87 × 10 ⁻¹³	166,043	6.22 × 10 ⁻²⁴				
11	15	<i>HCN4</i>	S	rs4489968	71452559	T	G	0.843	0.513	0.060	8.89 × 10 ⁻¹⁵	3.34 × 10 ⁻⁸	160,858	3.82 × 10 ⁻²⁰				
12	3	<i>GNB4</i>	S	rs7612445	180655673	G	T	0.816	0.358	0.060	2.41 × 10 ⁻⁷	7.78 × 10 ⁻¹⁰	140,395	1.86 × 10 ⁻¹⁴				
13	14	<i>FLRT2</i>	S	rs17796783	84879664	T	C	0.716	0.334	0.049	1.65 × 10 ⁻⁹	5.76 × 10 ⁻⁶	145,835	2.69 × 10 ⁻¹³				
14	7	<i>CHRM2</i>	S	rs2350782	136293174	C	T	0.116	0.505	0.078	1.57 × 10 ⁻⁷	2.63 × 10 ⁻⁷	131,781	1.26 × 10 ⁻¹²				
15	5	<i>NKX2-5</i>	Q,S	rs6882776	172596769	G	A	0.680	0.301	0.051	9.67 × 10 ⁻⁶	7.43 × 10 ⁻⁹	158,807	2.29 × 10 ⁻¹²				
16	7	<i>GNG11</i>	C,Q,S,B	rs180242	93387532	T	A	0.333	0.316	0.053	7.52 × 10 ⁻⁷	5.59 × 10 ⁻⁷	148,111	6.78 × 10 ⁻¹²				
17	2	<i>B3GNT7</i>	N,Q,S,P,B	rs13030174	231979528	A	C	0.733	0.300	0.051	1.13 × 10 ⁻⁶	3.66 × 10 ⁻⁴	144,810	1.04 × 10 ⁻¹⁰				
18	3	<i>FNDC3B</i>	S	rs9647379	173267862	C	G	0.400	0.206	0.047	2.59 × 10 ⁻⁷	3.07 × 10 ⁻⁴	137,314	1.17 × 10 ⁻⁹				
19	12	<i>RFX4</i>	S	rs2067615	105673552	A	T	0.490	0.278	0.044	1.05 × 10 ⁻⁷	8.49 × 10 ⁻⁴	151,197	1.58 × 10 ⁻⁹				
20	12	<i>CPNE8</i>		rs826838	37392998	C	T	0.443	0.234	0.045	3.32 × 10 ⁻⁷	6.20 × 10 ⁻⁴	166,632	3.73 × 10 ⁻⁹				
21	2	<i>TFPI</i>	N,S	rs4140885	188041309	A	G	0.317	0.217	0.049	1.85 × 10 ⁻⁶	1.40 × 10 ⁻³	170,395	4.72 × 10 ⁻⁸				

Chr., chromosome. EAF, effect allele frequency based on meta-analysis of stages 1 and 2 combined. Annotation shows whether the heart rate-associated SNP is (i) in strong LD ($r^2 > 0.8$) with a copy number variant (C), nonsynonymous variant (N) or variant in a transcription factor binding site (T); (ii) associated with an eQTL (Q); (iii) expressed at the protein level in mouse heart and phosphorylated

upon stimulation of the β_1 AR (P); (iv) located in or near a gene that was identified as being of potential relevance for heart rate using the automated literature search program SNIPPER (S); or (v) located in or near a biological candidate gene for heart rate (B).

^a Positions are according to HapMap Build 36, and allele coding is based on the positive strand.

^b Effect sizes in bpm per effect allele obtained from stage 1 and stage 2 cohorts with heart rate as only outcome ($N > 96,790$; excludes data from the RRgen Consortium and ERF)

Table 2

Association of the heart rate loci with cardiac conduction and rhythm disorders

Locus	Nearest gene	Alleles		PR duration			QRS duration			QT duration			Atrial fibrillation			SSS				
		Heart rate SNP	Effect	Other	β (ms)	SE (ms)	<i>P</i>	<i>N</i>	β (ms)	SE (ms)	<i>P</i>	<i>N</i>	β (ms)	SE (ms)	<i>P</i>	OR (95% CI)	<i>P</i>	OR (95% CI)	<i>P</i>	
Previously identified heart rate loci																				
1	<i>MYH6</i>	rs365990	G	A	22,711	-0.53	0.23	0.02	36,680	-0.00	0.07	0.98	67,414	0.05	0.10	0.65	1.027	0.12	0.931	0.18
2	<i>GJA1</i>	rs1015451	C	T	26,528	0.66	0.34	0.05	40,090	-0.38	0.11	6.51×10^{-4}	72,806	-0.50	0.16	1.67×10^{-3}	0.903	1.58×10^{-4}	0.961	0.63
3	<i>ACHE</i>	rs3245899	G	A	25,317	0.35	0.27	0.19	37,902	-0.07	0.09	0.41	72,980	0.11	0.12	0.39	0.985	0.46	0.951	0.46
4	<i>CD46</i>	rs11118555	A	T	27,701	-0.03	0.32	0.93	39,530	-0.00	0.11	0.97	73,601	-0.25	0.15	0.09	0.964	0.15	0.895	0.16
5	<i>FADS1</i>	rs174549	A	G	28,482	-0.53	0.22	0.02	40,085	-0.27	0.07	1.85×10^{-4}	74,604	-0.57	0.10	4.33×10^{-6}	1.003	0.85	0.971	0.57
6	<i>SLC35F1</i>	rs1153730	T	C	27,585	0.56	0.20	6.20×10^{-3}	39,200	-0.59	0.07	1.26×10^{-16}	74,932	-1.65	0.10	2.23×10^{-67}	1.061	5.19×10^{-4}	0.973	0.59
7	<i>LINC00477 (C12orf67)</i>	rs17287293	A	G	27,912	2.06	0.29	4.91×10^{-13}	39,548	-0.10	0.09	0.31	74,180	-0.01	0.14	0.93	1.132	2.38×10^{-7}	0.811	2.83×10^{-3}
Newly identified heart rate loci																				
8	<i>KIAA1755</i>	rs6127471	C	T	27,459	0.47	0.21	0.02	39,775	-0.06	0.07	0.40	72,570	-0.28	0.10	4.17×10^{-2}	0.978	0.18	0.973	0.61
9	<i>CCDC141</i>	rs17362588	A	G	24,111	-0.83	0.33	0.01	35,425	0.55	0.11	4.75×10^{-7}	64,780	0.27	0.16	0.09	1.016	0.55	0.873	0.09
10	<i>SYT10</i>	rs7980799	A	C	24,965	0.52	0.22	0.02	36,338	0.03	0.07	0.64	69,431	-0.18	0.10	0.08	0.990	0.57	1.027	0.60
11	<i>HCN4</i>	rs4489968	T	G	25,991	0.67	0.29	0.02	38,579	0.09	0.09	0.33	72,387	0.05	0.13	0.69	0.856	1.85×10^{-12}	0.924	0.26
12	<i>GNB4</i>	rs7612445	G	T	23,153	-0.37	0.28	0.18	38,398	-0.14	0.09	0.11	67,376	-0.11	0.13	0.40	0.959	0.04	0.912	0.14
13	<i>FLRT2</i>	rs17796783	T	C	27,993	-0.12	0.22	0.61	38,467	-0.05	0.07	0.54	73,380	-0.24	0.11	0.02	0.959	0.02	0.898	0.05
14	<i>CHRM2</i>	rs2350782	C	T	23,414	-0.26	0.37	0.48	34,417	-0.07	0.12	0.57	58,549	0.25	0.19	0.17	0.941	0.05	0.911	0.32
15	<i>NXN2-5</i>	rs6882776	G	A	18,484	1.29	0.27	1.46×10^{-6}	30,877	-0.27	0.09	1.87×10^{-2}	60,768	-0.49	0.12	2.27×10^{-5}	1.110	4.41×10^{-6}	1.056	0.33
16	<i>GNG11</i>	rs180242	T	A	21,101	0.33	0.24	0.17	34,141	0.00	0.08	0.96	63,609	0.02	0.11	0.84	0.995	0.80	1.025	0.63

Locus	Nearest gene	Heart rate SNP	Alleles		PR duration			QRS duration			QT duration			Atrial fibrillation			SSS			
			Effect	Other	N	β (ms)	SE (ms)	P	N	β (ms)	SE (ms)	P	N	β (ms)	SE (ms)	P	OR (95% CI)	P	OR (95% CI)	P
17	<i>B3GN77</i>	rs13030174	A	C	27,169	0.18	0.24	0.45	38,446	-0.23	0.08	4.14×10^{-3}	72,111	-0.14	0.11	0.19	0.997 (0.96-1.03)	0.88	1.025 (0.91-1.15)	0.67
18	<i>FND33B</i>	rs9647379	C	G	21,503	0.65	0.23	4.23×10^{-3}	35,642	-0.02	0.07	0.74	63,885	0.03	0.10	0.74	1.002 (0.97-1.04)	0.89	0.969 (0.87-1.08)	0.55
19	<i>RFX4</i>	rs2067615	A	T	27,782	-0.17	0.20	0.40	39,494	-0.05	0.07	0.42	74,116	-0.02	0.10	0.83	1.029 (1.00-1.06)	0.08	1.054 (0.96-1.16)	0.29
20	<i>CPNE8</i>	rs826838	C	T	28,059	0.62	0.21	2.78×10^{-3}	39,313	0.02	0.07	0.80	74,402	-0.04	0.10	0.71	0.999 (0.97-1.03)	0.93	0.926 (0.84-1.02)	0.12
21	<i>TFPI</i>	rs4140885	A	G	26,384	-0.15	0.23	0.50	37,447	0.05	0.08	0.55	68,919	-0.18	0.11	0.10	1.011 (0.98-1.05)	0.55	1.126 (1.02-1.25)	0.02
Multi-SNP predisposition score						2.55	0.67	1.33×10^{-4}		-0.95	0.22	1.83×10^{-5}		-2.66	0.31	1.12×10^{-17}	0.921 (0.83-1.02)	0.11	0.589 (0.44-0.78)	2.34×10^{-4}

Association analyses were performed in data from the PR GWAS Consortium (PR duration), the QRS GWAS Consortium (QRS duration), the QT-IGC Consortium (QT duration), deCODE Genetics and the CHARGE-AF Consortium (atrial fibrillation), and deCODE Genetics (SSS). Analyses were performed in data from 9,183 cases and 91,625 controls for atrial fibrillation and in data from 903 cases and 40,722 controls for SSS. CI, confidence interval; effect allele, the allele that is associated with higher heart rate. *P* values for association with PR duration are not adjusted for population stratification, as are the ones for the remaining traits. The multi-SNP predisposition score shows the combined effect of the 21 heart rate-associated SNPs per genetically predicted 5-bpm increase in heart rate, based on single-SNP summary statistics and weighting by effect sizes for heart rate. *P* values smaller than 2.0×10^{-3} are considered statistically significant.

den Hoed et al.