

The challenge of prognostic markers in pleural mesothelioma

Andrea Imperatori¹, Massimo Castiglioni¹, Lorenzo Mortara², Elisa Nardecchia¹, Nicola Rotolo¹

¹Center for Thoracic Surgery, Department of Surgical and Morphological Sciences, University of Insubria, Varese, Italy; ²Department of Biotechnology and Life Sciences, University of Insubria, Varese, Italy

ABSTRACT

Malignant pleural mesothelioma (MPM) is a very aggressive tumor, highly resistant to chemo- and radio-therapy. Treatment of MPM patients is often disappointing, regardless of the modality used.

Inter-individual variability of response to multimodal treatment remains a challenge and generally the MPM prognosis continues to be poor. Knowledge of predicting factors of outcome is currently insufficient; therefore, it would be highly desirable to find specific prognostic markers for MPM. Translational research projects are to be implemented.

KEY WORDS

Malignant pleural mesothelioma (MPM); prognostic factors; translational research

J Thorac Dis 2013;5(3):205-206. doi: 10.3978/j.issn.2072-1439.2013.06.12

Malignant pleural mesothelioma (MPM) is a very aggressive tumor, highly resistant to chemo- and radio-therapy. Treatment of MPM patients is often disappointing, regardless of the modality used. Median survival remains less than 1 year, with less than 5% 5-year survivors (1). Currently, therapeutic management of MPM is heavily dependent on patient performance status and is expected to be potentially effective predominantly in the epithelioid subtype. The association of pemetrexed with cisplatin resulted in significantly improved efficacy of chemotherapy (2). The multimodal treatment including surgery (extrapleural pneumonectomy, or pleurectomy/decortication or extended pleurectomy/decortication) combined with chemotherapy that was introduced in the 1990s improved the long-term survival in selected operable patients (3). However, inter-individual variability of response to multimodal treatment remains a challenge and generally the MPM prognosis continues to be poor. Knowledge of predicting factors of outcome is currently insufficient even if many clinico-radiographic and molecular variables have been studied, usually with limited numbers of patients (4); therefore, it would be highly desirable to find specific prognostic markers for MPM.

New biomarkers are needed to improve the three aspects of

the clinical management of MPM: early diagnosis, prognosis, and prediction of response to treatment (5). Some promising biomarkers (osteopontin, mesothelin, fibulin-3) have been described to predict prognosis and likelihood of response to therapy, in order to tailor treatment regimens on the basis of patients' individual features (6-8). However, these biomarkers are yet to be fully validated. For example, despite promising initial results, plasma osteopontin levels did not discriminate between chronic pleural inflammatory disease and MPM (9).

In a recent issue of the *Journal of Thoracic Disease*, Mori *et al.* retrospectively assessed the prognostic value of what they term "N-ERC index" in a small group of inoperable MPM patients (10). ERC, previously identified by the authors in a rat renal carcinoma and also known in humans as human megakaryocyte potentiating factor (MPF) or mesothelin, is a 71-kDa protein that can be found in the serum (11). These authors previously identified serum N-ERC level as a marker for early MPM diagnosis and noted that it increased as a function of disease stage (12). The N-ERC levels of MPM patients at diagnosis show wide inter-individual variance. Using baseline pre-treatment N-ERC level and post-chemotherapy treatment level, Mori *et al.* developed an index they term "N-ERC index". The latter was defined as Log₂ of the post-/pre- N-ERC ratio, which normalizes baseline N-ERC variability and post-chemotherapy changes. From the results of their study, Mori *et al.* concluded that "... *The N-ERC index is considered to be a useful biomarker for predicting not only the chemotherapeutic response, but also the prognosis in patients with advanced MPM.*" (10). While this may represent an important step in the direction of finding a useful MPM biomarker, a limitation of the study of Mori *et al.*, as in similar studies is methodology, reproducibility and small sample size. The importance and validity of a prognostic factor is much

Corresponding to: Andrea Imperatori, MD. Center for Thoracic Surgery, Department of Surgical and Morphological Sciences, University of Insubria, Ospedale di Circolo, Via Guicciardini, 21100 Varese, Italy. Email: andrea.imperatori@uninsubria.it.

Submitted Apr 23, 2013. Accepted for publication Jun 13, 2013.
Available at www.jthoracdis.com

ISSN: 2072-1439

© Pioneer Bioscience Publishing Company. All rights reserved.

greater when it is identified in a prospective randomized trial with univariate and multivariate analysis, rather than in a retrospective series review as was the Mori study. Further, to correctly assess the effects of treatment for MPM, clinical trials should stratify patients according to prognostic group (13). In MPM, detailed staging by imaging is certainly required, but it is not sufficient. Prognostic scoring systems have been proposed as a method for evaluating single patient prognosis and for stratification of risks in MPM clinical trials (14). Because of scarce reproducibility, however, the use of scoring systems so far has been disappointing in clinical practice. Mori *et al.* noted that in their series, the low N-ERC level group, which showed significantly longer overall survival, included 4 stable disease patients and 5 progressive disease patients, possibly due to difficulties in evaluating by imaging tumor reduction (10).

While the N-ERC index seems promising in preliminary studies, it should also be investigated in diverse clinical scenarios as translational research projects (15). It should be tested for early diagnosis in subjects at risk for MPM, such as workers exposed to asbestos, and for differential diagnosis in patients with recurrent undetermined pleural effusion. Moreover, the N-ERC index could be used to stratify MPM patient sub-groups for new therapeutic trials, as recently proposed also for disease-specific genetic mutations (16).

In conclusion, the N-ERC index may be used as a new prognostic factor in the design of MPM clinical trials, and in the implementation of translational research projects.

Acknowledgements

Disclosure: The authors declare no conflict of interest.

References

1. Helland Å, Solberg S, Brustugun OT. Incidence and survival of malignant pleural mesothelioma in Norway: a population-based study of 1686 cases. *J Thorac Oncol* 2012;7:1858-61.
2. Vogelzang NJ, Rusthoven JJ, Symanowski J, et al. Phase III study of

3. Sugarbaker DJ, Heher EC, Lee TH, et al. Extrapleural pneumonectomy, chemotherapy, and radiotherapy in the treatment of diffuse malignant pleural mesothelioma. *J Thorac Cardiovasc Surg* 1991;102:10-4; discussion 14-5.
4. Pass HI. Biomarkers and prognostic factors for mesothelioma. *Ann Cardiothorac Surg* 2012;1:449-56.
5. Greillier L, Baas P, Welch JJ, et al. Biomarkers for malignant pleural mesothelioma: current status. *Mol Diagn Ther* 2008;12:375-90.
6. Pass HI, Lott D, Lonardo F, et al. Asbestos exposure, pleural mesothelioma, and serum osteopontin levels. *N Engl J Med* 2005;353:1564-73.
7. Pass HI, Wali A, Tang N, et al. Soluble mesothelin-related peptide level elevation in mesothelioma serum and pleural effusions. *Ann Thorac Surg* 2008;85:265-72; discussion 272.
8. Pass HI, Levin SM, Harbut MR, et al. Fibulin-3 as a blood and effusion biomarker for pleural mesothelioma. *N Engl J Med* 2012;367:1417-27.
9. Paleari L, Rotolo N, Imperatori A, et al. Osteopontin is not a specific marker in malignant pleural mesothelioma. *Int J Biol Markers* 2009;24:112-7.
10. Mori T, Tajima K, Hiramata M, et al. The N-ERC index is a novel monitoring and prognostic marker for advanced malignant pleural mesothelioma. *J Thorac Dis* 2013;5:145-8.
11. Hino O. Multistep renal carcinogenesis in the Eker (Tsc 2 gene mutant) rat model. *Curr Mol Med* 2004;4:807-11.
12. Shiomi K, Hagiwara Y, Sonoue K, et al. Sensitive and specific new enzyme-linked immunosorbent assay for N-ERC/mesothelin increases its potential as a useful serum tumor marker for mesothelioma. *Clin Cancer Res* 2008;14:1431-7.
13. Steele JP, Rudd RM. Malignant mesothelioma: predictors of prognosis and clinical trials. *Thorax* 2000;55:725-6.
14. Edwards JG, Abrams KR, Leverment JN, et al. Prognostic factors for malignant mesothelioma in 142 patients: validation of CALGB and EORTC prognostic scoring systems. *Thorax* 2000;55:731-5.
15. Cesario A, Russo P, Viaggi S, et al. Malignant pleural mesothelioma: time for translational research. *Lancet Oncol* 2004;5:591; discussion 591-2.
16. McMillan R, Zauderer M, Bott M, et al. Important recent insights into the genetics and biology of malignant pleural mesothelioma. *Ann Cardiothorac Surg* 2012;1:462-5.



Cite this article as: Imperatori A, Castiglioni M, Mortara L, Nardecchia E, Rotolo N. The challenge of prognostic markers in pleural mesothelioma. *J Thorac Dis* 2013;5(3):205-206. doi: 10.3978/j.issn.2072-1439.2013.06.12