

Published in final edited form as:

Folia Histochem Cytobiol. ; 50(2): 312–318.

Expression of mRNAs for PPT, CGRP, NF-200, and MAP-2 in cocultures of dissociated DRG neurons and skeletal muscle cells in administration of NGF or NT-3

Weiwei Zhang¹, Hao Li², Ziyang Xing¹, Hongtu Yuan¹, Mark S. Kindy³, and Zhenzhong Li¹

¹Department of Anatomy, Shandong University School of Medicine, Jinan, China

²Department of Orthopedics, Shandong University Qilu Hospital, Jinan, China

³Department of Neurosciences, Medical University of South Carolina, Charleston, South Carolina, USA

Abstract

Both neurotrophins (NTs) and target skeletal muscle (SKM) cells are essential for the maintenance of the function of neurons and nerve-muscle communication. However, much less is known about the association of target SKM cells with distinct NTs on the expression of mRNAs for preprotachykinin (PPT), calcitonin-gene related peptide (CGRP), neurofilament 200 (NF-200), and microtubule associated protein 2 (MAP-2) in dorsal root ganglion (DRG) sensory neurons. In the present study, a neuromuscular coculture model of dissociated dorsal root ganglion (DRG) neurons and SKM cells was established. The morphology of DRG neurons and SKM cells in coculture was observed with an inverted phase contrast microscope. The effects of nerve growth factor (NGF) or neurotrophin-3 (NT-3) on the expression of mRNAs for PPT, CGRP, NF-200, and MAP-2 was analyzed by real time-PCR assay. The morphology of DRG neuronal cell bodies and SKM cells in neuromuscular coculture at different conditions was similar. The neurons presented evidence of dense neurite outgrowth in the presence of distinct NTs in neuromuscular cocultures. NGF and NT-3 increased mRNA levels of PPT, CGRP, and NF-200, but not MAP-2, in neuromuscular cocultures. These results offer new clues towards a better understanding of the association of target SKM cells with distinct NTs on the expression of mRNAs for PPT, CGRP, NF-200 and MAP-2, and implicate the association of target SKM cells and NTs with DRG sensory neuronal phenotypes.

Keywords

neurotrophins; substance P; calcitonin-gene related peptide; neurofilament 200; microtubule associated protein 2; dorsal root ganglion; skeletal muscle cell

Introduction

Targets of neuronal innervation play a vital role in regulating the survival and differentiation of innervating neurotrophin (NT)-responsive neurons [1]. Sensory nerve cross-anastomosis (sensory protection) provides a modified trophic environment by modulating neurotrophic

© Polish Society for Histochemistry and Cytochemistry

Correspondence addresses: Z. Li, Department of Anatomy, Shandong University School of Medicine, 44 Wenhua Xi Road, Jinan, Shandong Province 250012, China; tel.: 86 158 6379 3602; zli@sdu.edu.cn, M.S. Kindy, Department of Neurosciences, Medical University of South Carolina, Charleston 29425, South Carolina, USA; tel.: 1 843 792 0808, fax: 1 843 876 5099; kindyms@musc.edu.

factor synthesis in muscles [2]. Target tissues contribute to the phenotype and function of sensory neurons [3–5]. The neuropeptide-immunoreactive (IR) and neurofilament-IR neurons are two major phenotypical classes in DRG. Neuropeptide-IR neurons are considered to be unmyelinated or thinly myelinated nociceptive afferent neurons which are thought to innervate skin and viscera. Neurofilament (NF)-IR neurons typically have myelinated axons, many of which innervate muscle spindles [6].

Substance P (SP) is a nociceptive peptide [7] and a member of the tachykinin family of peptide neurotransmitters that are derived from the preprotachykinin (PPT) gene by alternative splicing [8]. Calcitonin gene-related peptide (CGRP) is a potent neuropeptide vasodilator [9] and controls sensory transmission [10]. Both SP and CGRP are expressed in primary sensory neurons [10, 11].

Neurofilaments (NFs) are neuron-specific intermediate filaments. Neurofilament heavy chain (NF-H) plays an important role in healthy neurons [12]. The appearance of NF-H represents a critical event in the stabilization of axons that accompanies their maturation [13]. Microtubule associated protein 2 (MAP-2) has been tentatively implicated in neuronal outgrowth and polarity of neuronal cells [14]. Within the neuronal cell, MAP-2 proteins are known to interact with microtubules, NFs and actin, and contribute to the maintenance of neuronal cytoarchitecture [15, 16].

Neurotrophins (NTs) are a family of growth factors that regulate the peripheral and central nervous system [17]. Nerve growth factor (NGF) and neurotrophin-3 (NT-3) are two members of the neurotrophin (NT) family, and all play important roles in the development of the peripheral sensory nervous system. Additionally, these growth factors are proposed to modulate the properties of the sensory system in the adult under pathological conditions [18, 19]. NGF utilizes two receptors, the NGF-specific tyrosine kinase receptor, TrkA, and also the non-specific neurotrophin receptor, p75(NTR) (p75). The specific contribution of each receptor to SP signaling remains to be determined [7]. Some neuronal phenotypes of neuropeptide are retained and can remain sensitive to NGF regulation in aged DRG neuronal cultures [20]. It has been demonstrated that NGF upregulates CGRP [21, 22]. NT-3, an important component of the Schwann cell autocrine survival loop, plays an important role in survival and differentiation of sensory neurons sprouting of neurites, synaptic reorganization, and axonal growth [19, 23].

Whether these factors regulate the expression of mRNAs for PPT, CGRP, NF-200, and MAP-2 in DRG neurons in the presence of target skeletal muscle (SKM) cells *in vitro* remains to be seen. In the present study, neuromuscular cocultures of dissociated DRG neurons and SKM cells were established. Using this unique culture system, the levels of mRNAs for PPT, CGRP, NF-200, and MAP-2 were determined after the administration of different NTs.

Material and methods

Neuromuscular coculture preparations and NT treatment

Animals used in this experiment were obtained from the Experimental Animal Center at Shandong University in China. All experiments involving animals were approved by the Ethical Committee of Shandong University. DRG cell cultures were prepared using Wistar rat pups at embryonic day 15 (E15). Under aseptic conditions, dorsal root ganglia (DRGs) were removed bilaterally using a sharp pair of forceps from each embryo and placed in Dulbecco's Modified Eagle Medium with F-12 supplement (DMEM//F-12) media (Gibco) in a half petri dish. After that, DRG explants were digested with 0.25% trypsin (Sigma) in D-Hanks solution at 37°C for 10 min and centrifuged for 5 min at 1×10^3 rpm. The

supernatants were removed and the pellets were resuspended in DMEM/F-12 media and triturated using a sterile modified Pasteur's glass pipette. DRG cells were then filtered using a Millipore filtering apparatus with a 74 μm mesh, followed by counting. Dissociated DRG cells were plated at a density of 2×10^5 cells/ml in six-well clusters (Costar, Corning, NY, USA) which would contain a single layer of SKM cells prepared as follows.

SKM cell culture preparations utilized newborn Wistar rats. SKM cell cultures were prepared three days before DRG preparation. Under aseptic conditions and using the newborn rat, SKM was removed from the hind limb of each animal, minced with fine dissecting scissors into fragments approximately 0.5 mm in diameter, digested with 0.25% trypsin (Sigma) in D-Hanks solution at 37°C for 20 min, centrifuged, and triturated in growth media supplemented with 5% fetal bovine serum (Gibco). Isolated SKM cells were plated at a density of 2×10^5 cells/ml in six-well clusters (Costar, Corning, NY, USA) which would contain 24 mm diameter coverslips precoated with poly-L-lysine (0.1 mg/ml) and then incubated at 37°C in a 5% CO₂ incubator.

The neuromuscular cocultures were prepared as follows. The newly prepared DRG cell suspension was plated at three-day old SKM culture and coculture with SKM cell for an additional six days, with a medium change every two days.

All the cocultures were randomly divided into three groups. (1) NGF group: the neuromuscular cocultures were treated with NGF (10 ng/ml) during the six days of coculture. (2) NT-3 group: the neuromuscular cocultures were treated with NT-3 (10 ng/ml) during the six days of coculture. (3) control group: the neuromuscular cocultures were continuously exposed to growth media as control. The composition of the culture medium was DMEM/F-12 (1:1) supplemented with 10% fetal bovine serum, 2% B-27 supplement (Gibco), L-glutamine (0.1 mg/ml, Sigma), penicillin (100 U/ml), and streptomycin (100 μg /ml).

Living cell observation by inverted phase contrast microscopy

All cultures were observed using inverted phase contrast microscopy at different culture ages. The observers were blinded to the treatment protocol for the evaluation. The pictures were taken for monitoring morphological characteristics of neuromuscular cocultures of DRG and SKM cells with different NTs.

Real time-PCR analysis of mRNAs for PPT, CGRP, NF-200, and MAP-2

The mRNA levels of PPT, CGRP, NF-200, or MAP-2 in neuromuscular cocultures at six days of culture age with different NTs were analyzed by real time-PCR. The expression of glyceraldehyde-3-phosphate dehydrogenase (GAPDH) mRNA was also determined as an internal control. The GAPDH gene is highly conserved and expressed in a wide variety of tissue types and so is useful as a reference 'house-keeping' gene. It is a stable reference gene for application in cell culture research using the rat. It allows determination of relative expression levels of other genes of interest. Total DRG cell RNA of each well of the clusters was isolated by TRIzol (TakaRa). cDNA was synthesized using a cDNA synthesis kit (Fermentas) according to the manufacturer's instructions.

The synthetic oligonucleotide primer sequences for PPT, CGRP, NF-200, MAP-2 and GAPDH were as follows: PPT 5'- CGA CAG TGA CCA AAT CAA GG -3' (coding sense) and 5'- CAA AGA ACT GCT GAG GCT TG -3' (coding antisense). CGRP 5'- CCT TTC CTG GTT GTC AGC ATC TT -3' (coding sense) and 5'- CAG TAG GCG AGC TTC TTC TTC AC -3' (coding antisense). NF-200 5'- AAA GTG AAC ACG GAT GCT ATG C -3' (coding sense) and 5'- GTG CTT TTC AGT GCC TCC AAC -3' (coding antisense). MAP-2 5'- GGC ACT CCT CCA AGC TAC TCT -3' (coding sense) and 5'- CTT GAC GTT CTT

CAG GTC TGG -3' (coding antisense). GAPDH 5'- GGC ACA GTC AAG GCT GAG AAT G -3' (coding sense) and 5'- ATG GTG GTG AAG ACG CCA GTA -3' (coding antisense).

Real time-PCR was performed using SYBR green dye (Fermantas) according to the manufacturer's instructions. PCR was performed at 50°C for two minutes, 94°C for 15 minutes, followed by 40 cycles at 94°C for 15 seconds, 58°C for 30 seconds, and 72°C for 30 seconds.

A comparative cycle of threshold fluorescence (Ct) method was used and the relative transcript amount of the target gene was normalized to that of GAPDH using the $2^{-\Delta\Delta C_t}$ method based on a previous study [24]. The final results of real time-PCR were expressed as the ratio of mRNA of control.

Statistical analysis

Data is expressed as mean \pm SD. All the data was processed for verifying normality test for variables. If the normality test failed, the data was analyzed with SPSS software by non-parametric methods (Mann–Whitney test). If the normality test passed, statistical analysis was evaluated with SPSS software by one-way ANOVA followed by the Student–Newman–Keuls test (homogeneity of variance) or Dunnett's T3 test (heterogeneity of variance) for significance to compare the differences among various groups. Significance was determined as $p < 0.05$.

Results

Morphology of DRG neurons and SKM cells in coculture under inverted phase contrast microscope

The DRG neuronal cell bodies send axons passing over, or terminating on, the surfaces of SKM cells. The fusing SKM cells were fusiform, rod, bifurcate or 'starfish' in shape. The contraction of the fused SKM cells could be seen under an inverted phase contrast microscope because of the characteristic of spontaneous contraction. Axons from several neurons would cross each other, forming a fine network in areas of high cell density. The crossing axons adhered to each other; hence, displacement of a terminal axon on a contracting muscle cell would also oscillate the proximally crossing axonal network. The morphology of DRG neuronal cell bodies and SKM cells in coculture at different conditions was similar. The neurons exhibited evidence of dense neurite outgrowth in the presence of distinct NTs (Figure 1).

PPT mRNA expression at different culture conditions

To determine the mRNA levels of PPT, the neuromuscular cocultures at six days of culture age with different NT treatment were analyzed by real-time PCR. PPT mRNA levels increased in NGF (2.60 fold, $p < 0.01$) or NT-3 (2.02 fold, $p < 0.01$) treated cocultures compared to those in the control group (Figure 2).

CGRP mRNA expression at different culture conditions

To determine the mRNA levels of CGRP, the neuromuscular cocultures at six days of culture age with different NT treatment were analyzed by real-time PCR. CGRP mRNA levels increased in NGF (2.33 fold, $p < 0.05$) or NT-3 (1.98 fold, $p < 0.01$) treated cocultures compared to those in the control group (Figure 3).

NF-200 mRNA expression at different culture conditions

To determine the mRNA levels of NF-200, the neuromuscular cocultures at six days of culture age with different NT treatment were analyzed by real-time PCR. NF-200 mRNA levels increased in NGF (1.36 fold, $p < 0.05$) or NT-3 (1.55 fold, $p < 0.001$) treated cocultures compared to those in the control group (Figure 4).

MAP-2 mRNA expression at different culture conditions

To determine the mRNA levels of MAP-2, the neuromuscular cocultures at six days of culture age with different NT treatment were analyzed by real-time PCR. MAP-2 mRNA levels did not change significantly in the presence of NGF or NT-3 compared to those in the control group (Figure 5).

Discussion

During development, neurons extend axons to their targets, then become dependent for their survival on trophic substances secreted by their target cells [25]. The exogenous administration of NTs has protective properties for injured neurons, and stimulates axonal regeneration [26]. The expression of mRNAs for PPT, CGRP, NF-200, and MAP-2 may reflect the responsiveness of DRG sensory neurons to these factors.

In the present study, NGF and NT-3 increased mRNA levels of PPT, CGRP, and NF-200, but not MAP-2, in neuromuscular cocultures.

Many neurotrophic factors have been shown to promote neurite outgrowth by improving the microenvironment that is required for nerve regeneration [27]. It has been suggested that the concentrations of NGF were from 1 to 300 ng/ml for DRG neurons *in vitro* [28]. Serum concentrations of NGF were several pg/ml [29] which is much lower than that used in *in vitro*. The levels of growth factors in fetal bovine serum were much lower than the levels of exogenous growth factors used in this experiment. In the present study, i.e. the exogenous administration of NGF or NT-3, respectively, to the neuromuscular cocultures of DRG neurons and SKM cells, the DRG neurons presented evidence of dense neurite outgrowth in the presence of distinct NT, suggesting that the microenvironment improved by distinct NT is suitable for neurite outgrowth of DRG neurons in the neuromuscular cocultures.

In DRG, NGF signaling is required not only for survival, but also for the differentiation of nociceptors which express SP [30]. NGF is a potent stimulator of SP which is produced in DRG sensory neurons [28, 31]. It has been demonstrated that expression of SP is regulated by NGF and exogenous exposure to high levels of NGF increases its cellular content and release. NGF up-regulation of SP expression requires the involvement of both TrkA and p75, although the specific contribution of each receptor to SP signaling remains to be determined [7, 32]. NGF has a key role in the survival and establishment of the phenotype of responsive primary afferent neurons during development [33]. In the present study, the exposure of exogenous NGF to neuromuscular cocultures of E15 DRG neurons and SKM cells could increase the levels of PPT mRNA encoding for SP, a finding consistent with previous studies. NGF also increased the levels of CGRP mRNA and NF-200 mRNA in neuromuscular cocultures of DRG neurons and SKM cells.

NT-3 signaling is required for the generation and survival of various DRG neuron classes, in particular proprioceptors [30]. NT-3 prevented abnormalities of NF-H in sensory neurons of diabetic rats [34]. The latest research demonstrated that NT-3 may have antinociceptive effects which may be mediated, at least in part, by its cognate receptor tyrosine-kinase C (TrkC) in the medium or large size DRG neurons [35].

In the present study, exposure of exogenous NT-3 to neuromuscular cocultures of E15 DRG neurons and SKM cells could increase the levels of mRNAs for PPT, CGRP, and NF-200. These results suggest that the effects of NT-3 on promoting neuronal survival might be through regulating expression of sensory neuropeptide and protein expressed in DRG neurons.

It has been suggested that at the initiation of radial growth, neurofilaments are likely to consist primarily of neurofilament light and medium, as neurofilament heavy expression is developmentally delayed [36]. In the present study, either NGF or NT-3 has a promotive effect on NF-200 mRNA expression, suggesting that both NGF and NT-3 were involved in regulation of the later stage of DRG development.

MAP-2 is a cytoskeletal protein and plays a regulatory role in neuronal plasticity and in maintaining the morphology of differentiated neurons [37]. MAP-2 has been tentatively implicated in neuronal outgrowth and polarity of neuronal cells [14]. MAP-2 is generally used as a neuronal marker [38]. In the present study, MAP-2 mRNA expression did not depend on the presence of distinct NTs in neuromuscular cocultures of DRG neurons and SKM cells suggested that target SKM cells is sufficiently on maintaining MAP-2 mRNA expression.

In conclusion, the neurons presented evidence of dense neurite outgrowth in the presence of distinct NTs in neuromuscular cocultures of DRG neurons and SKM cells. NGF and NT-3 increased mRNA levels of PPT, CGRP, and NF-200, but not MAP-2, in neuromuscular cocultures. These results offer new clues towards a better understanding of the association of target SKM cells with distinct NTs on the expression of mRNAs for PPT, CGRP, NF-200, and MAP-2 and implicate the association of target SKM cells and NTs with DRG sensory neuronal phenotypes.

Acknowledgments

This work was supported by the National Natural Science Foundation of China (No. 30973051) and the Natural Science Foundation of Shandong Province of China (ZR2010HM027).

References

1. Howe CL, Valletta JS, Rusnak AS, et al. NGF signaling from clathrin-coated vesicles: Evidence that signaling endosomes serve as a platform for the Ras-MAPK pathway. *Neuron*. 2001; 32:801–814. [PubMed: 11738027]
2. Zhao C, Veltri K, Li S, et al. NGF, BDNF, NT-3, and GDNF mRNA expression in rat skeletal muscle following denervation and sensory protection. *J Neurotrauma*. 2004; 21:1468–1478. [PubMed: 15672636]
3. Lu J, Zhou XF, Rush RA. Small primary sensory neurons innervating epidermis and viscera display differential phenotype in the adult rat. *Neurosci Res*. 2001; 41:355–363. [PubMed: 11755222]
4. Wang L, Liu Z, Liu H, et al. Neuronal phenotype and tyrosine kinase receptor expression in cocultures of dorsal root ganglion and skeletal muscle cells. *Anat Rec*. 2009; 292:107–112.
5. Wang L, Li H, Liu Z, et al. Co-administration of monosialogan-glioside and skeletal muscle cells on dorsal root ganglion neuronal phenotypes in vitro. *Cell Mol Neurobiol*. 2010; 30:43–49. [PubMed: 19582569]
6. Hall AK, Ai X, Hickman GE, et al. The generation of neuronal heterogeneity in rat sensory ganglion. *J Neurosci*. 1997; 17:2775–2784. [PubMed: 9092599]
7. Skoff AM, Adler JE. Nerve growth factor regulates substance P in adult sensory neurons through both TrkA and p75 receptors. *Exp Neurol*. 2006; 197:430–436. [PubMed: 16300761]

8. Carter MS, Krause JE. Structure, expression and some regulatory mechanisms of the rat preprotachykinin gene encoding substance P, neurokinin A, neuropeptide K and neuropeptide gamma. *J Neurosci.* 1990; 10:2203–2214. [PubMed: 1695945]
9. Zhang ZH, Fang XB, Xi GM, et al. Calcitonin gene-related peptide enhances CREB phosphorylation and attenuates tau protein phosphorylation in rat brain during focal cerebral ischemia/reperfusion. *Biomed Pharmacother.* 2010; 64:430–436. [PubMed: 20537498]
10. Schaeffer V, Meyer L, Patte-Mensah C, et al. Progress in dorsal root ganglion neurosteroidogenic activity: basic evidence and pathophysiological correlation. *Prog Neurobiol.* 2010; 92:33–41. [PubMed: 20438798]
11. Pan XQ, Gonzalez JA, Chang S, et al. Experimental colitis triggers the release of substance P and calcitonin gene-related peptide in the urinary bladder via TRPV1 signaling pathways. *Exp Neurol.* 2010; 225:262–273. [PubMed: 20501335]
12. Gotow T. Neurofilaments in health and disease. *Med Electron Microsc.* 2000; 33:173–199. [PubMed: 11810476]
13. Yabe JT, Wang FS, Chylinski T, et al. Selective accumulation of the high molecular weight neurofilament subunit within the distal region of growing axonal neurites. *Cell Motil Cytoskeleton.* 2001; 50:1–12. [PubMed: 11746668]
14. Sánchez C, Díaz-Nido J, Avila J. Phosphorylation of micro-tubule-associated protein 2 (MAP2) and its relevance for the regulation of the neuronal cytoskeleton function. *Prog Neurobiol.* 2000; 61:133–168. [PubMed: 10704996]
15. Kaech S, Parmar H, Roelandse M, et al. Cytoskeletal micro-differentiation: a mechanism for organizing morphological plasticity in dendrites. *Proc Natl Acad Sci USA.* 2001; 98:7086–7092. [PubMed: 11416192]
16. Ozer RS, Halpain S. Phosphorylation-dependent localization of microtubule-associated protein MAP2c to the actin cytoskeleton. *Mol Biol Cell.* 2000; 11:3573–3587. [PubMed: 11029056]
17. Brahimi F, Malakhov A, Lee HB, et al. A peptidomimetic of NT-3 acts as a TrkC antagonist. *Peptides.* 2009; 30:1833–1839. [PubMed: 19647025]
18. Price TJ, Louria MD, Candelario-Soto D, et al. Treatment of trigeminal ganglion neurons in vitro with NGF, GDNF or BDNF: effects on neuronal survival, neurochemical properties and TRPV1-mediated neuropeptide secretion. *BMC Neurosci.* 2005; 6:4. [PubMed: 15667652]
19. Wang TH, Meng QS, Qi JG, et al. NT-3 Expression in spared DRG and the associated spinal laminae as well as its anterograde transport in sensory neurons following removal of adjacent DRG in cats. *Neurochem Res.* 2008; 33:1–7. [PubMed: 17710544]
20. Jiang ZG, Smith RA. Regulation by nerve growth factor of neuropeptide phenotypes in primary cultured sensory neurons prepared from aged as well as adult mice. *Brain Res Dev Brain Res.* 1995; 90:190–193.
21. Lindsay RM, Harmar AJ. Nerve growth factor regulates expression of neuropeptide genes in adult sensory neurons. *Nature.* 1989; 337:362–364. [PubMed: 2911387]
22. Amann R, Sirinathsinghi DJ, Donnerer J, et al. Stimulation by nerve growth factor of neuropeptide synthesis in the adult rat in vivo: bilateral response to unilateral intraplantar injections. *Neurosci Lett.* 1996; 203:171–174. [PubMed: 8742020]
23. Sahenk Z, Nagaraja HN, McCracken BS, et al. NT-3 promotes nerve regeneration and sensory improvement in CMT1A mouse models and in patients. *Neurology.* 2005; 65:681–689. [PubMed: 16157899]
24. Pfaffl MW. A new mathematical model for relative quantification in real-time RT-PCR. *Nucleic Acids Res.* 2001; 29:e45. [PubMed: 11328886]
25. Wang H, Tessier-Lavigne M. En passant neurotrophic action of an intermediate axonal target in the developing mammalian CNS. *Nature.* 1999; 401:765–769. [PubMed: 10548102]
26. Lykissas MG, Batistatou AK, Charalabopoulos KA, et al. The role of neurotrophins in axonal growth, guidance, and regeneration. *Curr Neurovasc Res.* 2007; 4:143–151. [PubMed: 17504212]
27. Liu JJ, Wang CY, Wang JG, et al. Peripheral nerve regeneration using composite poly(lactic acid-caprolactone)/nerve growth factor conduits prepared by coaxial electrospinning. *J Biomed Mater Res A.* 2011; 96:13–20. [PubMed: 20949481]

28. Winston J, Toma H, Shenoy M, et al. Nerve growth factor regulates VR-1 mRNA levels in cultures of adult dorsal root ganglion neurons. *Pain*. 2001; 89:181–186. [PubMed: 11166474]
29. Sigala S, Martocchia A, Missale C, et al. Increased serum concentration of nerve growth factor in patients with microprolactinoma. *Neuropeptides*. 2004; 38:21–24. [PubMed: 15003712]
30. Ernsberger U. Role of neurotrophin signalling in the differentiation of neurons from dorsal root ganglia and sympathetic ganglia. *Cell Tissue Res*. 2009; 336:349–384. [PubMed: 19387688]
31. Supowit SC, Ethridge RT, Zhao H, et al. Calcitonin gene-related peptide and substance P contribute to reduced blood pressure in sympathectomized rats. *Am J Physiol Heart Circ Physiol*. 2005; 289:H1169–H1175. [PubMed: 15879482]
32. Skoff AM, Resta C, Swamydas M, et al. Nerve growth factor (NGF) and glial cell line-derived neurotrophic factor (GDNF) regulate substance P release in adult spinal sensory neurons. *Neurochem Res*. 2003; 28:847–854. [PubMed: 12718437]
33. Bueno L. Gastrointestinal pharmacology: irritable bowel syndrome. *Curr Opin Pharmacol*. 2005; 5:583–588. [PubMed: 16188501]
34. Sayers NM, Beswick LJ, Middlemas A, et al. Neurotrophin-3 prevents the proximal accumulation of neurofilament proteins in sensory neurons of streptozocin-induced diabetic rats. *Diabetes*. 2003; 52:2372–2380. [PubMed: 12941778]
35. Tender GC, Kaye AD, Li YY, et al. Neurotrophin-3 and tyrosine-kinase C have modulatory effects on neuropathic pain in the rat dorsal root ganglia. *Neurosurgery*. 2011; 68:1048–1055. [PubMed: 21221027]
36. Shen H, Barry DM, Garcia ML. Distal to proximal development of peripheral nerves requires the expression of neurofilament heavy. *Neuroscience*. 2010; 170:16–21. [PubMed: 20633607]
37. Di Stefano G, Casoli T, Fattoretti P, et al. Distribution of map2 in hippocampus and cerebellum of young and old rats by quantitative immunohistochemistry. *J Histochem Cytochem*. 2001; 49:1065–1066. [PubMed: 11457938]
38. Liu Y, Saad RS, Shen SS, et al. Diagnostic value of microtubule-associated protein-2 (MAP-2) for neuroendocrine neoplasms. *Adv Anat Pathol*. 2003; 10:101–106. [PubMed: 12605092]




Figure 1. Inverted phase contrast microscopic photomicrographs of the living cells in neuromuscular coculture at six days of culture age. DRG neuronal cell body (thin arrows) sends axons across or terminating on SKM cells (thick arrows). Panel A: neuromuscular coculture with NGF treatment; Panel B: neuromuscular coculture with NT-3 treatment; Panel C: neuromuscular coculture without NT treatment. Scale bar = 50 μ m




Figure 2. PPT mRNA expression at different culture conditions. Normalized levels of PPT mRNA were higher in NGF treated cocultures than control. NT-3 treatment increased PPT mRNA levels. Bar graphs with error bars represent mean \pm SD (n = 12). *p < 0.01




Figure 3. CGRP mRNA expression at different culture conditions. Normalized levels of CGRP mRNA were higher in NGF treated cocultures than control. NT-3 treatment increased CGRP mRNA levels. Bar graphs with error bars represent mean \pm SD (n = 12). *p < 0.05, **p < 0.01




Figure 4. NF-200 mRNA expression at different culture conditions. Normalized levels of NF-200 mRNA were higher in NGF treated cocultures than control. NT-3 treatment increased NF-200 mRNA levels. Bar graphs with error bars represent mean \pm SD (n = 12). *p < 0.05, **p < 0.001




Figure 5. MAP-2 mRNA expression at different culture conditions. Normalized levels of MAP-2 mRNA were not changed significantly in NGF or NT-3 treated cocultures than control. Bar graphs with error bars represent mean \pm SD (n = 6)