

Published in final edited form as:

Cancer Res. 2010 August 15; 70(16): 6649–6658. doi:10.1158/0008-5472.CAN-10-0909.

Metastasis associated protein 1 and its short form variant stimulates *Wnt1* transcription via promoting its derepression from *Six3* corepressor

Rakesh Kumar^{1,2,3}, Seetharaman Balasenthil^{2,*,#}, Bramanandam Manavathi^{2,*,%}, Suresh K Rayala^{2,*,+}, and Suresh B Pakala¹

¹Department of Biochemistry and Molecular Biology, The George Washington University Medical Center, Washington, District of Columbia 20037, USA

²Departments of Molecular and Cellular Oncology, University of Texas M. D. Anderson Cancer Center, Houston, Texas 77030, USA

Abstract

Although Wnt1 downstream signaling components have been well studied and activated in human cancer, the pathways that regulate *Wnt1* itself have not been explored in-depth. Here we provide gain-of-function, loss-of function, and molecular evidence supporting functional interactions between metastasis associated protein 1 short-form (MTA1s), metastasis associated protein 1 (MTA1) and Wnt1 signaling components during mammary gland development and tumorigenesis. Using multiple model systems involving overexpression or knockdown of MTA1s or MTA1, we discovered that MTA1s and MTA1 hyperactivate the Wnt1 pathway due to increased expression of *Wnt1* transcription. MTA1s and MTA1 physically interact with *Six3* chromatin, protein product of which is a direct HDAC-dependent repressor of *Wnt1* transcription. Deletion of the MTA1s and MTA1 allele in murine embryonic fibroblasts resulted in upregulation of *Six3* and downregulation of Wnt signaling. In addition, mammary glands from the MTA1s/MTA1^{-/-} mice exhibited increased recruitment of *Six3* corepressor complex to the *Wnt1* promoter and inhibition of Wnt1 pathway in mammary glands. These findings identify MTA1s and MTA1 as important upstream modifiers of the *Wnt1* transcription, and consequently its functions, by directly inhibiting the transcription of *Six3* allowing de-repression of *Wnt1* transcription.

Introduction

The Wingless (*Wnt* genes encode a family of secreted glycoproteins with roles in normal and pathologic processes, including cancer. For example, Wnt1 represents one of the earliest pathways, linked with the development of hyperplasia and cancer (1). Once upregulated, secreted Wnt1 acts as an autocrine and/or paracrine factor and initiates a cascade of cytoplasmic signaling events leading to phosphorylation of glycogen synthase kinase 3 β (GSK-3 β) and inhibition of its ability to phosphorylate β -catenin. Stabilized β -catenin translocates to the nucleus, leading to stimulation of Wnt-target genes (2, 3) .

Although Wnt1 is known to be widely upregulated in human cancer including, breast cancer, much of the work on Wnt signaling research mainly focused on the action of Wnt1 from the

³Corresponding author. bcmrxk@gwumc.edu.

*Contributed equally to this work

#Present address: Department of Genetics, M.D. Anderson Cancer Center, Houston, Texas 77030, USA.

%Present address: Department of Biochemistry, University of Hyderabad, Hyderabad- 500 046, India.

+Present address: Department of Biotechnology, IIT Madras, Chennai – 600 036, India.

plasma membrane to the nucleus. However, the regulation of *Wnt1* transcription continues to be poorly understood, particularly, in the context of mammary gland. One of the best characterized direct coregulators of *Wnt1* transcription is the Six3 homeodomain protein in the retinal or neuronal cells (4). Six3 interacts with Groucho family of corepressors which associate with HDACs (5). Interestingly, *Six3* transcription has been shown to be tightly regulated by metastatic tumor antigen 1 (MTA1) containing Mi-2/nucleosome remodeling and deacetylase (NuRD) complex (6) in retinal cells. The NuRD complexes are abundant deacetylase complexes in mammalian cells and have been implicated in chromatin remodeling in normal as well as in cancerous cells (7). The NuRD complex couples histone deacetylation and ATP-dependent chromatin remodeling in the same complex and is involved in chromatin compaction and transcriptional repression. The MTA1 was initially cloned from highly metastatic mammary adenocarcinomas (8) and its expression correlated with the aggressiveness of several human cancers (7). MTA1 acts a potent repressor of estrogen receptor-alpha (9) and of BRCA1 (10). In contrast of MTA1, its naturally occurring variant MTA1s primarily localizes in the cytoplasm (11) and participates in the stimulation of canonical Wnt1-signaling in breast cancer cells (Companion manuscript # 1). MTA3, another member of the MTA family, was reported to physically interact with the *Wnt4* chromatin in a histone deacetylase-dependent manner resulting in the suppression of the Wnt4-dependent morphogenesis (12).

Most of our current understanding of the Wnt1 functions in mammary epithelial cells and in other systems is derived from the membrane-initiated signaling pathways feeding into target gene expression. However, in spite of widely reported increased Wnt1 expression in cancer, the nature of coregulators that regulate the transcription of *Wnt1* gene itself in mammary epithelial cells remains unknown. Herein we provide gain-of-function, loss-of function, and molecular evidence supporting regulatory roles for MTA1 and its variant isoform MTA1s in the transcriptional stimulation of the *Wnt1* gene via Six3 pathway in mammary epithelial and cancer cells.

Materials and methods

Cell line authorization Statement

All the cell lines used in this study are from Dr. Rakesh Kumar's laboratory and have been tested, authenticated, previously used in the peer-reviewed papers from the laboratory (6, 9, 10, 11, 12).

Cell culture

HC11, MCF-7, SKBR3 and MDA-MB-435 cells, MEFs from mice WT, heterozygous, or homozygous for MTA1s cells were cultured in as described in Supplementary Methods section.

siRNA transfection and luciferase reporter assays

The Δ Six3-luc was constructed by deleting the regions containing the three clustered Six3 recognition sequences (5). siRNA transfections and luciferase reporter assays were carried out as described elsewhere (11)

Immunofluorescence and confocal microscopy studies

Cellular localization of proteins was determined by using indirect immunofluorescence as described previously (11). Confocal scanning analysis was performed as described in supplementary methods section.

Immunohistochemistry and Mammary gland whole mounts and histology

Detailed experimental procedures were described in the Supplementary Methods section.

Chromatin immunoprecipitation assay (ChIP) and ChIP -qPCR assay

ChIP analysis and ChIP-qPCR was carried out as described by Kumar et al (11) and detailed procedure is described in supplementary methods. Primers used for ChIP analysis for Six3 promoter walk, and for ChIP analysis on Wnt1 promoter and enhancer regions are presented in Supplementary Table 2. Primers and Taqman probes used for ChIP-qPCR assay are presented in Supplementary Table 3.

Statistical analysis and reproducibility

Statistical analysis of the data was performed using Graphpad Prism software (GraphPad Software Inc., San Diego, CA). Statistical analysis of reporter assays was performed using Student's *t*-test and the results were presented as mean \pm standard error.

Results and Discussion

MTA1 and MTA1s are upstream positive modifier of the Wnt1 transcription

Although MTA1s predominantly localizes in the cytoplasm (11), an albeit but easily detectable amount of T7-MTA1s was also found in the nucleus of HC11 murine mammary epithelial cells expressing MTA1s cells (Fig. 1A). This observation combined with the fact that MTA1s over expression in HC11 cells was accompanied by increased secretion of Wnt1 (Companion manuscript # 1) suggested the possibility of increased expression of *Wnt1* by MTA1s. Indeed, we found that MTA1s over expression in the HC11 cells was accompanied by increased expression of *Wnt1* mRNA and protein (Fig. 1B). As MTA1s is a differentially spliced naturally-occurring variant of MTA1 (11), we next wished to evaluate whether Wnt1 pathway and targets are also regulated by MTA1, a predominantly nuclear protein. We found that stable overexpression of T7-MTA1 in the HC11 murine mammary epithelial cells (HC11/MTA1) also resulted in a significant increased level of Wnt1 (Fig. 1C, D) as well as β -catenin-associated increased phosphorylated- GSK-3 β signaling (Fig. 1D) and Top-TK luc activity as readouts of β -catenin transactivation (Fig. 1E). Conversely, silencing of MTA1 by MTA1-specific siRNA in the MCF-7 cells resulted in downregulation of the Wnt-signaling components (Fig. 1F), suggesting that MTA1-mediated increased *Wnt1* expression is also accompanied by increased stimulation of the cytoplasmic Wnt-signaling. Together, these findings suggested that both MTA1s and MTA1 may be upstream modifiers of the *Wnt1* transcription in mammary epithelial and breast cancer cells.

MTA1 and MTA1s target Six3 expression in the mammary epithelial and breast cancer cells

Since Wnt1 expression itself is upregulated in breast cancer (3), we next explored the possibility whether MTA1 and MTA1s stimulate the Wnt1 expression by facilitating the stimulation of *Wnt1* transcription via repressing the expression of its negative regulator, We turned our attention to a direct known repressor of *Wnt1* transcription, namely Six3, in the neuronal and retinal systems (4, 6).

We found that upregulation of MTA1 or MTA1s in the HC11 cells led to a substantial reduction in the level of Six3 mRNA and protein (Fig. 2A), suggesting corepressive activity of MTA1 and MTA1s for Six3. Indeed, we found that stable upregulation of MTA1 or MTA1s in mammalian cells led to a substantial reduction in the *Six3* promoter activity (Fig. 2B). We next explored the roles of MTA1 or MTA1s complexes in the repression of *Six3*. Treatment of HC11/MTA1 and HC11/MTA1s cells with a general HDAC inhibitor TSA resulted in an increased Six3 promoter activity (Fig. 2C), suggesting that MTA1 and MTA1s

repress *Six3* expression in an HDAC-dependent manner in mammary epithelial cells. Since previous studies (11) have not delineated the basis of albeit levels of MTA1s in the nucleus (Fig. 1A), we decided to understand the biochemical basis of nuclear localization of MTA1s and found a role of endoplasmic reticulum (ER) membrane retention signal motif RTKP (K428) in this process (Supplementary Fig. S1).

Status of *Six3* in cells depleted of MTA1 and MTA1s

To establish a causative relationship between the levels of MTA1, MTA1s and *Six3* expression, we studied *Six3* expression in murine embryonic fibroblasts (MEFs) from WT, heterozygous, and homozygous mice for MTA1 and MTA1s (6). Results indicate that loss of MTA1 and MTA1s expression in MEF resulted in a distinct upregulation of *Six3* protein levels (Fig. 3A) and decreased TOP-Flash luciferase activity (Fig. 3B). In addition, selective silencing of MTA1s in MDA-MB-435 (Fig. 3C), and of MTA1 silencing in MCF-7 cells (Fig. 3D), also resulted in a distinct upregulation of *Six3*. To implicate a role for *Six3* expression in the observed increased Wnt1 pathway by MTA1 or MTA1s, we selectively silenced endogenous *Six3* by using *Six3* specific siRNA in HC11 cells, which expresses high levels of *Six3*, and studied the ability of transfected MTA1s or MTA1 to upregulate β -catenin protein expression as well as TOPFLASH activity. Results showed that selective silencing of endogenous *Six3* by *Six3* specific siRNA potentiated the ability of both MTA1 and MTA1s to upregulate β -catenin protein expression as well as TOP-FLASH activity (Fig. 3E). Because Wnt1 pathway could be activated at multiple levels, these results suggested that MTA1s and MTA1 contribute to observed enhancement of the Wnt1 pathway by repressing the expression of a direct *Wnt1* transcriptional repressor *Six3*, in addition to the stabilization of β -catenin. Furthermore, these data also showed for the first time a significant biological role for *Six3* outside of the retinal and neuronal systems, and identified MTA1 and MTA1s as regulators of Wnt1 expression in mammary epithelial cells.

MTA1s and MTA1 as upstream activators of the Wnt1- β -catenin pathway

To conclusively establish a temporal relationship between MTA1s and Wnt1 signaling components such as P-GSK-3 β , P-ERK and β -catenin, we found that the loss of MTA1s and MTA1 expression in MEFs from MTA1/MTA1s homozygous mice resulted in a clearly reduced levels of phosphorylated GSK-3 β , ERK as well as β -catenin and *Wnt1* expression (Fig. 4A). Interestingly, reconstitution of Myc-MTA1s or T7-MTA1 but not control pcDNA in MTA1s/MTA1-null MEFs was accompanied by increased expression of phosphorylated GSK-3 β , β -catenin, *Wnt1* and downregulation of *Six3* (Fig. 4B), implicating MTA1s and MTA1 as upstream determinants of the β -catenin. Next we generated MTA1s/MTA1^{-/-}MEFs stably expressing V5-tagged MTA1s and MTA1 protein. Stable expression of V5-tagged MTA1 and MTA1s resulted in upregulation of β -catenin, phosphorylated GSK-3 β , *Wnt1* and downregulation of *Six3* (Fig. 4C). These findings identified MTA1s and MTA1 as upstream activators of the Wnt1- β -catenin pathway.

Mechanism of *Six3* regulation by MTA1 and MTA1s

Next, we wished to gain a deeper insight about the mechanism by which MTA1 and MTA1s downregulates *Six3* expression. The negative regulatory region of the mouse *Six3* has been mapped up to 5 kb upstream of the start site (4). To examine the physical interaction of MTA1 and MTA1s with the *Six3* chromatin, we carried out a promoter walk for the 5.5 kb region upstream of the *Six3* gene using chromatin immunoprecipitation (ChIP) analysis, with primers specifically designed against every 500-bp region of the *Six3* promoter. Initially, ChIP lysates from the HC11/T7-MTA1 and HC11/T7-MTA1s cells were immunoprecipitated with monoclonal anti-T7 antibody to pull down T7-MTA1s/MTA1 or with IgG as a control and eluted DNA was tested by PCR analysis. This analysis identified one *Six3* promoter MTA1sinteracting region, encompassing roughly a 1.5 kb region (+1 bp

to -1434 bp i.e. region 1, 2 and 3), and two *Six3* promoter MTA1-interacting regions a 1.0 kb region (+1 bp to -954 bp i.e. regions 1 and 2), and a 0.5 kb region (-3834 bp to -4314 bp i.e. region 9 (Fig. 5A, B). There was no other specific binding to other regions. Region 6 of the *Six3* chromatin, where no association of was seen in ChIP assays were used as negative controls for MTA1s and MTA1 respectively. In addition to this, ChIP - qPCR analysis also showed similar results when carried out under identical conditions (Supplementary Fig. S2A).

Since HDACs could interact with MTA1s or MTA1 and because MTA1s or MTA1 represses *Six3* promoter activity in an HDAC-dependent manner (Fig. 2C), we next tested whether HDAC2 is also recruited to the MTA1s/MTA1-interacting *Six3* chromatin. ChIP analysis with an HDAC2-specific antibody showed that HDAC2 was present on the same region 1-3 (+1 bp to -1434 bp i.e. region 1, 2 and 3), of the *Six3* chromatin as MTA1s and also present on the same region 1, 2 (+1 bp to -954 bp i.e. regions 1 and 2) and region 9 (-3834 bp to -4314 bp) of the *Six3* chromatin as MTA1 (Fig. 5C), raising the possibility of the recruitment of the MTA1s/HDAC or MTA1/HDAC complex to the above mentioned regions of the *Six3* chromatin. Similar results were obtained when we carried out ChIP analysis followed by a qPCR analysis under identical conditions (Supplementary Fig. S2B). Further ChIP analysis for Mi-2, the largest subunit of the NuRD complex showed the recruitment of Mi-2 to region 1-3 (+1 bp to -1434 bp i.e. region 1, 2 and 3), of the *Six3* chromatin as MTA1s and to region 1, 2 (+1 bp to -954 bp i.e. regions 1 and 2) and region 9 (-3834 bp to -4314 bp) of the *Six3* chromatin as MTA1 (Fig. 5D). ChIP analysis followed by a qPCR analysis carried out under the same conditions also yielded similar results (Supplementary Fig. S2C). These results suggests that possibility that MTA1s/MTA1 in association with Mi-2 and HDAC as a complex can be recruited to the *Six3* chromatin.

MTA1 or MTA1s corepressor complexes inhibit *Six3*-transcription

To understand the basis of MTA1- and MTA1s-regulation of *Six3*-transcription, we next performed a double ChIP assay in the HC11/MTA1s and HC11/MTA1 clones; the initial ChIP analysis was done with anti-T7 antibody to immunoprecipitate the T7-MTA1s or T7-MTA1- bound DNA sequences and the subsequent ChIP analysis was done using the anti-HDAC2 antibody. Results showed the simultaneous co-association of MTA1s and HDAC2 with the region 1-3 (+1 bp to -1434 bp i.e. region 1, 2 and 3), but not to the regions 6 and 9 and coassociation of MTA1 with region 1, 2 (+1 bp to -954 bp i.e. regions 1 and 2) and region 9 (-3834 bp to -4314 bp) but not to the regions 3 and 6 (Fig. 6A). Double ChIP assay followed by a qPCR analysis under the same conditions also yielded similar results (Supplementary Fig. S3A).

The above-mentioned set of ChIP analyses suggested that the MTA1s/NuRD or MTA1/NuRD complexes associate with two distinct regions of *Six3* regulatory chromatin, bringing about histone deacetylation by the virtue of associated HDAC activity leading to chromatin compaction and this would be reflected in the decreased acetylation status of the histones present in these two regions of the *Six3* chromatin. To explore the potential decreased acetylation status of the histones associated with the MTA1s and MTA1-interacting *Six3* regions, we performed a ChIP assay with acetyl-Histone 3 (H3) specific antibodies in HC11/pcDNA, HC11/MTA1s and HC11/MTA1 clones. Results indicated a relative decrease in the association of the acetylation H3 with the 1.5kb region (+1 bp to -1434 bp i.e. region 1, 2 and 3) of the *Six3* chromatin in HC11/MTA1s cells (Fig. 6B). Similarly, the levels of acetylated H3 associated with the 1.0 kb (+1 bp to -954 bp i.e. regions 1 and 2) and 0.5 kb (-3834 bp to -4314 bp) regions of the *Six3* chromatin in HC11/MTA1 cells were considerably decreased in comparison to the same regions in the HC11/pcDNA cells. ChIP followed by qPCR analysis carried out under identical conditions also yielded similar results (Supplementary Fig. S3B).

Overall, the above ChIP analysis provided clear evidence of distinct physical association of the MTA1s/HDAC or MTA1/HDAC complexes with two distinct regions of the *Six3* regulatory chromatin, resulting in greater levels of histone deacetylation and chromatin compaction, ultimately leading to transcriptional repression. Further analysis of the *Six3* promoter regions bound by MTA1s and MTA1 (promoter regions 1- 3, and 9 of *Six3* chromatin), revealed the presence of core ATTA (DNA sequence bound by Six3) motifs in the regions 1, 2 and 9 but not in regions 3 and 6 of the *Six3* chromatin. Previous studies have reported the presence of three clustered ATTA core motifs in the Six3 promoter (5). In addition, Six3 has been shown to bind to its own promoter and regulate its own transcription (5). The presence of core ATTA motifs in regions 1, 2 and 9 of the Six3 promoter bound by MTA1s and MTA1 (Supplementary Fig S4) suggested the possibility that Six3 binds to its own promoter and Six3 interaction with MTA1s/MTA1 family of corepressors along with the components of NuRD complex such as Mi-2 and HDAC2, and thus, could facilitate autorepression of Six3 transcription. Furthermore, the presence of core ATTA motifs in the *Six3* promoter opens up the possibility that MTA1s or MTA1 may mediate *Six3* repression thro ATTA motif. It is interesting to note that the Six3-promoter-luciferase construct used in the present study (Fig. 2B), were constructed by inserting the 1.36 kb mouse *Six3* genomic fragment including the three clustered Six3 recognition sequences (5).

To validate that the repression of *Six3* transcription by MTA1s or MTA1 is indeed mediated through the Six3 recognition sequences and/or other elements in the *Six3* promoter region, we made use of the Six3 reporter gene construct (Δ Six3 pro-luc), which was generated by deleting the region containing the three Six3 recognition sequences. MTA1s or MTA1 showed increased Δ Six3 pro-luc reporter activity compared to the wild type Six3-pro-luciferase, but still reduced as compared to the pcDNA controls, suggest that MTA1s or MTA1 may not only use ATTA motifs to regulate Six3 (Fig. 6C).

Next we determined whether MTA1s or MTA1 and Six3 proteins physically interact in mammary cell lines. Since the MTA1s antibody used in our present study is not suitable for immunoprecipitation assays, we used Six3 antibody in the immunoprecipitation assay and found that the endogenous Six3 can physically interact with transiently transfected T7-tagged MTA1s and T7-MTA1 in the HC11 cells (Fig. 6D, E). In addition, endogenous Six3 was also found to physically interact with endogenous MTA1 and MTA1s in MD-MBA-435 cells (Fig. 6F). These findings provided a mechanistic basis of *Six3* gene repression upon deregulation of MTA1s or MTA1 suggesting that MTA1s- or MTA1-complexes inhibit the expression of Six3 which, in-turn, allows upregulation of the Wnt1 transcription.

Activation of the Wnt pathway in MTA1s-TG mice

We recently generated transgenic mice with expression of MTA1s under the control of the mouse mammary tumor virus (MMTV) promoter long terminal repeat (MTA1s-TG mice) (Companion manuscript # 1). MTA1s-TG mammary glands were accompanied by precocious alveolar development (Companion manuscript # 1) and stimulation of Wnt1-target gene WISP1 in the mammary glands of 12-week-old virgin mice (Supplementary Fig. S5A). In previous studies from this laboratory we also observed increased expression of β -casein and β -catenin- Wnt1 signaling in virgin MTA1-TG mice (13). Similar to MTA1s-TG mice, mammary glands obtained from virgin MTA1-TG also exhibited upregulation of WISP1 (Supplementary Fig. S5B). Furthermore, mammary glands obtained from virgin MTA1s-TG and MTA1-TG mice exhibited significantly reduced expression of Six3, while the levels of Wnt1, β -catenin, P-GSK- 3 β and P-ERK were upregulated (Supplementary Fig. S5C). These findings identified MTA1s and MTA1 as upstream repressor of Six3, leading to increased Wnt1 expression and resulting signaling.

Downregulation of the Wnt1 pathway in MTA1s/MTA1^{-/-} mice

Since both MTA1s and MTA1-TG mice exhibited excessive hyperbranching, we examined the effect of MTA1s/MTA1 knockdown on mammary gland development using recently generated mice homozygous (-/-) for *MTA1/MTA1s* (6). We examined whole-mount preparations from littermates with matching estrous cycles at different developmental stages. We found that mammary glands from MTA1s/MTA1^{-/-} mice exhibited hypobranching as compared to control MTA1s/MTA1^{+/+} mice (Fig. 7A). These findings implicated an important role for MTA1s/MTA1 in mammary gland development. Since both MTA1s-TG and MTA1-TG mice exhibited upregulation of Wnt1 pathway, we examined the status of Wnt1 pathway components in MTA1s/MTA1^{-/-} mice. Immunohistochemical analysis in virgin mammary glands from 12 wk old MTA1s/MTA1^{-/-} mice showed upregulation of Six3 and downregulation of β -catenin (Fig. 7B), further validating a regulatory role for MTA1s and MTA1 in the Wnt1 pathway.

Differential recruitment of Six3 to Wnt1 chromatin in WT and MTA1s/MTA1^{-/-} mice

Results from MTA1s and MTA1 overexpressing cell lines, and MTA1s/MTA1^{-/-} mice implicated an important role for MTA1s and MTA1 in the Wnt1 pathway. To confirm whether *Wnt1* gene chromatin is a direct target of Six3 in the mammary epithelium and to determine whether MTA1s/MTA1 repression of Six3 expression compromises the recruitment of Six3 to the *Wnt1* promoter, we performed ChIP analysis in mammary glands from WT, and MTA1s/MTA1^{-/-} mice. In these assays, we used primers encompassing 720 bp of the *Wnt1* 5'-promoter region and also 280 bp of the *Wnt1* 3'-enhancer element (4) (Fig. 7C). In MTA1s/MTA1^{-/-} mammary glands, we observed an increased recruitment of Six3 to the promoter regions of Wnt1 compared to the corresponding control MTA1s/MTA1^{+/+} mice (Fig. 7D). ChIP followed by qPCR analysis carried out under identical conditions also yielded similar results (Fig. 7D, lower panel). In the *Wnt1* enhancer region, as in the promoter region, strong recruitment of Six3 in MTA1s/MTA1^{-/-} mammary glands compared to its corresponding control MTA1s/MTA1^{+/+} mice (Fig. 7E) was observed. ChIP assay followed by real time quantitative PCR analysis carried out under identical conditions also yielded similar results (Fig. 7E, lower panel). Western analysis of Six3 expression and RT-PCR analysis of *Wnt1* in the same samples showed increased expression of Six3 in MTA1s/MTA1^{-/-} mice compared to its corresponding control mice (Fig. 7F). These findings suggest that Six3 may restrain the activation of *Wnt1* transcription in mammary gland and that MTA1s and MTA1 up regulate Wnt1 expression via down regulating the expression of Six3, a direct repressor of *Wnt1*.

Next we attempted to investigate whether Six3 could be recruited to the *Wnt1* promoter in human breast cancer cell lines in an MTA1s- or MTA1-dependent manner. To examine this possibility we carried out ChIP analysis of Six3 recruitment to the *Wnt1* promoter and enhancer regions in the MDA-MB-435 cell line following treatment with MTA1s specific siRNA or control siRNA. Interestingly, our results showed an enhanced recruitment of Six3 to the *Wnt1* promoter and enhancer regions in MTA1s-siRNA silenced MDA-MB-435 cells compared to control siRNA treated cells (Fig. 7G), suggesting that Six3 may be recruited to *Wnt1* promoter in a MTA1s dependent manner in breast cancer cell lines.

In brief, results presented here provide new insight into the mechanism by which MTA1s and MTA1 regulate WNT signaling and for the first time identify MTA1s and MTA1 as novel upstream regulators of the Wnt1 transcription. These results suggest that MTA1s and MTA1 promotes WNT signaling at multiple levels by inhibiting two known regulators of Wnt pathway namely Six3, as well as GSK3- β (Fig. 7H). As components of the Wnt1 pathway are established contributors to mammary gland tumorigenesis, and considering the fact that MTA1s and MTA1 are upregulated in breast tumors, the newly established

dysfunction of the MTA1s- and MTA1- Wnt1 pathways in mammary epithelium may contribute to the pathogenesis of breast tumors. Aberrant Wnt1 pathway is implicated in a various diseases in addition to the breast cancer, regulation of the Wnt pathway is emerging as an attractive target for molecular intervention.

Supplementary Material

Refer to Web version on PubMed Central for supplementary material.

Acknowledgments

We thank the following past employees and staff in the laboratory: R. Wang for mammary gland mounting, C.J. Barnes for confocal imaging, and R. R Singh for Western blotting. We thank L. Clifton Stephens for histological diagnosis, G. Oliver for providing Six3-pro luc reporter and Δ Six3-pro-luc reporter system. This study was supported by NIH grant CA 98823 to RK.

References

1. Tsukamoto AS, Grosschedl R, Guzman RC, Parslow T, Varmus HE. Expression of the int-1 gene in transgenic mice is associated with mammary gland hyperplasia and adenocarcinomas in male and female mice. *Cell*. 1988; 55(4):619–625. [PubMed: 3180222]
2. Moon RT, Bowerman B, Boutros M, Perrimon N. The promise and perils of wnt signaling through beta-catenin. *Science*. 2002; 296(5573):1644–1646. [PubMed: 12040179]
3. Brown AM. Wnt signaling in breast cancer: Have we come full circle? *Breast Cancer Res*. 2001; 3(6):351–355. [PubMed: 11737884]
4. Lagutin OV, Zhu CC, Kobayashi D, et al. Six3 repression of wnt signaling in the anterior neuroectoderm is essential for vertebrate forebrain development. *Genes Dev*. 2003; 17(3):368–379. [PubMed: 12569128]
5. Zhu CC, Dyer MA, Uchikawa M, Kondoh H, Lagutin OV, Oliver G. Six3-mediated auto repression and eye development requires its interaction with members of the groucho-related family of corepressors. *Development*. 2002; 129(12):2835–2849. [PubMed: 12050133]
6. Manavathi B, Peng S, Rayala SK, et al. Repression of Six3 by a corepressor regulates rhodopsin expression. *Proc Natl Acad Sci U S A*. 2007; 104(32):13128–13133. [PubMed: 17666527]
7. Manavathi B, Kumar R. Metastasis tumor antigens, an emerging family of multifaceted master coregulators. *J Biol Chem*. 2007; 282(3):1529–1533. [PubMed: 17142453]
8. Toh Y, Pencil SD, Nicolson GL. A novel candidate metastasis-associated gene, mta1, differentially expressed in highly metastatic mammary adenocarcinoma cell lines. cDNA cloning, expression, and protein analyses. *J Biol Chem*. 1994; 269(37):22958–22963. [PubMed: 8083195]
9. Mazumdar A, Wang RA, Mishra SK, et al. Transcriptional repression of oestrogen receptor by metastasis-associated protein 1 corepressor. *Nat Cell Biol*. 2001; 3(1):30–37. [PubMed: 11146623]
10. Molli PR, Singh RR, Lee SW, Kumar R. MTA1-mediated transcriptional repression of BRCA1 tumor suppressor gene. *Oncogene*. 2008; 27(14):1971–1980. [PubMed: 17922032]
11. Kumar R, Wang RA, Mazumdar A, et al. A naturally occurring MTA1 variant sequesters oestrogen receptor-alpha in the cytoplasm. *Nature*. 2002; 418(6898):654–657. [PubMed: 12167865]
12. Zhang H, Singh RR, Talukder AH, Kumar R. Metastatic tumor antigen 3 is a direct corepressor of the Wnt4 pathway. *Genes Dev*. 2006; 20(21):2943–2948. [PubMed: 17050676]
13. Bagheri-Yarmand R, Talukder AH, Wang RA, Vadlamudi RK, Kumar R. Metastasis-associated protein 1 deregulation causes inappropriate mammary gland development and tumorigenesis. *Development*. 2004; 131(14):3469–3479. [PubMed: 15226262]




Figure 1. MTA1s overexpression upregulates Wnt1 signaling pathway in mammary epithelial cells

(A) Expression of T7-MTA1s by confocal microscopy, T7-MTA1s (red). Scale bar, 10 μm.

(B) RT-PCR and confocal analysis of Wnt1 in HC11/pcDNA and HC11/MTA1s cells. (C)

RT-PCR and confocal analysis of Wnt1 in HC11/pcDNA and HC11/MTA1 cells. (D)

Western blot analysis of β-catenin, Phospho-GSK3-β, GSK3-β, Wnt1 and T7 expression in

HC11/pcDNA and HC11/MTA1 clones. (E) Transcription status of Top-flash, Fop-flash in

HC11/pcDNA and HC11/MTA1 clones. (F) Western blot analysis of β-catenin, Wnt1,

GSK-3β, Phospho-GSK3-β and MTA1 expression in MCF-7 cell line following transfection

with MTA1 or control siRNA for 48h.




Figure 2. MTA1s and MTA1 targets Six3 expression in the mammary epithelial and breast cancer cells

(A) RT-PCR and Western blot analysis for Six3 in HC11/pcDNA, HC11/MTA1 and HC11/MTA1s cells. (B) MTA1s and MTA1 downregulates Six3 promoter-luc activity in HC11/MTA1s or HC11/MTA1 cells. (C) Six3 promoter luc activity in HC11/pcDNA and HC11/MTA1 or HC11/MTA1s cells following treatment with HDAC inhibitor, TSA.




Figure 3.

(A) Western blot analysis of Six3, MTA1s and MTA1 and in MEFs from WT (+/+), heterozygous (+/-) and homozygous (-/-) mice for MTA1s/MTA1. (B) Top-Flash luciferase activity in MEFs from WT (+/+) and homozygous (-/-) mice for MTA1s/MTA1. (C) Western blot analysis of Six3 expression in MDA-MB-435 cell line following transfection with MTA1s or control siRNA for 48h. (D) Western blot analysis of Six3 in MCF-7 cell line following transfection with MTA1 or control siRNA for 48h. (E) Western blot analysis and Top-Flash luciferase activity in HC11 cells following transfection with myc-tagged MTA1s or MTA1 and Six3-siRNA.




Figure 4. Status of Wnt signaling components

(A) Western blot analysis of phospho-GSK-3 β , GSK-3 β , phospho-ERK, ERK and β -catenin and RT-PCR analysis of *Wnt1* in MEFs from WT (+/+), heterozygous (+/-) and homozygous (-/-) mice for MTA1s/MTA1. (B) Western blot analysis of phospho-GSK-3 β , GSK-3 β , Six3 and β -catenin and RT-PCR analysis of *Wnt1* in MTA1s/MTA1^{-/-} MEFs transiently transfected with pcDNA, myc-MTA1s and T7-MTA1. (C) Western blot analysis of phospho-GSK-3 β , GSK-3 β , Six3 and β -catenin and RT-PCR analysis of *Wnt1* in MTA1s/MTA1^{-/-} MEFs stably expressing pcDNA, V5-MTA1 and V5-MTA1s.




Figure 5. MTA1s/MTA1/NuRD complex targets *Six3* regulatory chromatin
(A) Schematic diagram of 5.5 kb regulatory chromatin of mouse *Six3* gene. **(B)** ChIP showing differential recruitment of T7-MTA1s to regions 1–3 of *Six3* chromatin, and T7-MTA1 to regions 1, 2 and 9 of *Six3* chromatin in HC11-pcDNA, HC11-MTA1s and HC11-MTA1 cells. **(C, D)** ChIP showing recruitment of HDAC2 and Mi-2 to regions *Six3* chromatin in HC11/pcDNA, HC11/MTA1s and HC11/MTA1 cells.




Figure 6. Transcriptional regulation of *Six3* by MTA1 and MTA1s

(A) A double ChIP assay, first with T7-Ab followed by the HDAC2 Ab for the regions 1–3, 6 and 9 regions of the *Six3* chromatin in HC11/pcDNA, HC11/MTA1s and HC11/MTA1 cells. (B) Status of the acetylated histone H3 in the MTA1s/MTA1-NuRD complex bound *Six3* chromatin in the HC11/pcDNA, HC11/MTA1s and HC11/MTA1 by ChIP analysis. (C) *Six3* promoter-luc and Δ*Six3*-promoter-luc activity in HC11/pcDNA and HC11/MTA1s cells. Schematic representation of the 1.36 kb *Six3* promoter-luciferase construct and the three *Six3* recognition (ATTA) sequences are presented as boxes. Schematic representation of the 1.36 kb Δ*Six3* promoter-luciferase construct without the three *Six3* recognition (ATTA) sequences. (D) Endogenous *Six3* interacts with transiently transfected T7-MTA1s in the HC11 cells. (E) Transiently transfected T7-MTA1 interacts with endogenous *Six3* in the HC11 cells. (F) Endogenous interaction of MTA1, MTA1s with *Six3* in MDA-MB-435 cells.




Figure 7. Mammary gland development and status of Wnt pathway in MTA1s/MTA1^{-/-} mice
(A) Carmine Red-stained whole mounts of inguinal mammary glands obtained from MTA1s/MTA1^{+/+} mice and MTA1s/MTA1^{-/-} mice at 12 weeks of age. The images in the bottom panels are the images of the top panels at increased magnification. **(B)** Immunohistochemical analysis of Six3 and β -catenin expression in mammary glands of 12-week old virgin MTA1s/MTA1^{+/+} and MTA1s/MTA1^{-/-} mice. **(C)** Schematic representation of the Six3 binding sites in *Wnt1* promoter. **(D)** ChIP analysis showing differential recruitment of Six3 to the *Wnt1* proximal promoter region in MTA1s/MTA1^{-/-} mice and its corresponding MTA1s/MTA1^{+/+} mice. Bars below represents the relative

quantitation of Six3 recruitment to the *Wnt1* promoter using ChIP- q PCR assay. **(E)** ChIP analysis showing differential recruitment of Six3 to the *Wnt1* enhancer region in MTA1s/MTA1^{-/-} mice and its corresponding MTA1s/MTA1^{+/+} mice. Bars below represents the quantitation of Six3 recruitment to the *Wnt1* enhancer using ChIP-qPCR assay. **(F)** Western blot analysis of Six3 expression and RT-PCR analysis of *Wnt1* expression in MTA1s/MTA1^{-/-} mice and its corresponding MTA1s/MTA1^{+/+} mice. **(G)** ChIP analysis showing enhanced recruitment of Six3 to both *Wnt1* promoter (top panel) and *Wnt1* enhancer region in MTA1s-specific siRNA transfected MDA-MB-435 cells as compared to control siRNA transfected cells for 48h. **(H)** Schematic representation of Wnt1 expression and signaling by MTA1s or MTA1. Dotted lines going to transformation represents other possible mechanisms not investigated here.