

RESEARCH ARTICLE

Open Access

Soluble TWEAK independently predicts atherosclerosis in renal transplant patients

Kultigin Turkmen^{1*}, Halil Zeki Tonbul², Fatih Mehmet Erdur², Aysun Toker⁴, Zeynep Biyik², Huseyin Ozbiner³, Abduzhappar Gaipov², Elvin Enes Gul⁵, Mehmet Kayrak⁵, Yalcin Solak², Orhan Ozbek³, Suleyman Turk² and Adrian Covic⁶

Abstract

Background: Cardiovascular risk is increased in the early stages of chronic kidney disease (CKD) and also found to be ongoing in renal transplant (Rtx) patients. As a sign of atherosclerosis, increased carotid intima-media thickness (CIMT) has been widely accepted as a strong predictor of cardiovascular disease (CVD) and mortality in CKD patients. A novel markers, soluble tumor necrosis factor-like weak inducer of apoptosis (sTWEAK) and neutrophil-to-lymphocyte ratio (NLR) were introduced as potential markers in inflammatory disorders including CKD. The role of Rtx in terms of atherogenesis is still unclear. We aimed to investigate the relationship between sTWEAK, NLR and CIMT in Rtx patients without overt CVD and to compare these results with those obtained from healthy subjects.

Methods: Cross-sectional analysis in which CIMT measurements, NLR and serum TWEAK levels were assessed in 70 Rtx patients (29 females; mean age, 40.6 ± 12.4 years) and 25 healthy subjects (13 females, mean age; 37.4±8.8 years).

Results: sTWEAK levels were significantly decreased ($p=0.01$) and hs-CRP, NLR and CIMT levels of Rtx patients were significantly increased compared to healthy subjects ($p<0.0001$, $p=0.001$, $p<0.0001$, respectively). sTWEAK was also found to be decreased when eGFR was decreased ($p=0.04$ between all groups). CIMT was positively correlated with sTWEAK and NLR in Rtx patients ($r=0.81$, $p<0.0001$ and $r=0.33$, $p=0.006$, respectively). sTWEAK was also positively correlated with NLR ($r=0.37$, $p=0.002$). In the multivariate analysis only sTWEAK was found to be an independent variable of increased CIMT.

Conclusion: sTWEAK might have a role in the pathogenesis of ongoing atherosclerosis in Rtx patients.

Keywords: sTWEAK, Neutrophil-to-lymphocyte ratio, Carotid intima-media thickness, Renal transplantation

Background

Despite improvements in diagnostic tools and therapeutic approaches, premature death related to cardiovascular disease (CVD) remains the most common cause of death in patients with chronic kidney disease (CKD) [1]. Cardiovascular risk is increased even in the early stages of CKD and this heightened risk is also found after renal transplantation (Rtx) [2,3]. Besides traditional risk factors including hypertension, diabetes, dyslipidemia, advanced age and left ventricular hypertrophy (LVH), novel risk factors such as endothelial dysfunction (ED), vascular calcifications, oxidative stress, and inflammation

are highly prevalent and seem to play a more important role in renal patients compared to healthy subjects [4-7]. Several studies suggested that persistent systemic inflammation could be the main factor responsible for the increased risk in ESRD patients, regardless of the renal replacement therapy [8]. To prove this hypothesis, researchers analyzed a large panel of biomarkers to fully characterize the relation between inflammation and CVD, including C-reactive protein, interleukin (IL)-1 β , IL-6, tumor necrosis factor- α (TNF- α) [8,9], as well as several interesting new biomarkers.

Neutrophil-to-lymphocyte ratio (NLR) is a potential marker for inflammation in cardiac and non-cardiac disorders [10-12] that was also shown to be a predictor of long term mortality in patients who underwent percutaneous coronary intervention [13]. We recently demonstrated that

* Correspondence: mdk2010@yahoo.com

¹Department of Nephrology, Erzincan University Mengucek Gazi Training and Research Hospital, Erzincan, Turkey

Full list of author information is available at the end of the article

NLR could predict inflammation in ESRD patients [14]. Recently, a novel marker - the soluble tumor necrosis factor-like weak inducer of apoptosis (sTWEAK, TNFSF12) was introduced as a tumor necrosis factor (TNF) related cytokine in various inflammatory and non-inflammatory disorders [15]. To date, a transmembrane protein, fibroblast growth factor-inducible 14 (Fn14) and a scavenger receptor, CD163 were discovered as receptors of sTWEAK [16,17]. Binding of sTWEAK to Fn14 mediates multiple effects, including cellular growth, proliferation, migration, differentiation, apoptosis, angiogenesis, fibrogenesis and inflammation [18]. A soluble form of CD163 was also discovered and introduced as a risk factor for long term mortality in patients with peripheral artery disease [19].

Since the association between TWEAK and subclinical atherosclerosis was demonstrated in general and CKD populations [20,21], we aimed to investigate the relationship between sTWEAK, NLR and carotid IMT (CIMT) in Rtx patients without overt CVD and to compare these results with those obtained from healthy subjects.

Methods

The study protocol was approved by the Medical Ethics Committee of Selcuk University (Meram School of Medicine, Konya, Turkey). Written informed consent was obtained from all subjects included in the study.

This was a cross-sectional study involving 70 Rtx patients (29 females, 41 males; mean age, 40.6 ± 12.4 years) followed for at least 6 months in the transplantation unit of the Necmettin Erbakan University and 25 healthy subjects (13 females; mean age, 37.4 ± 8.8 years) between January and November 2011.

Patients aged 18–70 years willing to participate in the assessment of CIMT by carotid duplex ultrasonography were screened. A review of medical records (including information on age; sex; weight; medications; primary disease of ESRD) was undertaken. Exclusion criteria were: (i) angina pectoris and/or documented coronary artery disease, (ii) congestive heart failure; (iii) active infection; (iv) autoimmune disease; (v) severe secondary hyperparathyroidism (patients with iPTH > 500 pg/mL); (vi) nephrotic-range proteinuria and (vii) patients with HIV, HBV and HCV history. Eighty-six Rtx patients were screened and 16 patients were excluded from the study. Of the 16 excluded patients - 7 had documented coronary artery disease, 4 congestive heart failure (New York Heart Association (NYHA) class III–IV); 2 active infection; 2 secondary hyperparathyroidism; and 1 patient had autoimmune disease. None of the patients included in the study had nephrotic-range proteinuria and arrhythmia based on electrocardiography (ECG).

Twenty-five age and sex-matched normotensive healthy individuals (13 females, mean age; 37.4 ± 8.8 years) referred from outpatient clinics of the Internal Medicine

Department of Necmettin Erbakan University were enrolled as control subjects. They were subjected to the same inclusion and exclusion criteria as the transplanted patients.

Blood pressure was measured in the upright sitting position after ≥ 5 min of rest using an Erka sphygmomanometer (PMS Instruments Limited, Berkshire, UK) with an appropriate cuff size. Two readings were recorded for each individual. The mean value of the two readings was defined as the blood pressure. Patients with SBP and DBP ≥ 140 mmHg and ≥ 90 mmHg respectively were assumed to be hypertensive. Forty-two patients were taking anti-hypertensive drugs (all patients were on verapamil). Eleven patients were taking an oral calcium-vitamin D combination, and 9 patients were on active vitamin D. Ten patients (14.3%) were on intensive insulin and 16 (22.9%) were on oral anti-diabetic medication. All 70 Rtx patients, were on prednisolone and mycophenolate mofetil (MMF), 60 (85.7%) were on tacrolimus, 4 (5.7%) were on cyclosporine-A (Cyc-A), 2 were on sirolimus (2.9%) and 4 (5.7%) were on everolimus therapy. The donor source was living-related in 32 (45.7%), living-unrelated in 4 (5.7%) and deceased-donor in 34 (48.6%) patients. None of the patients had transplant nephrectomy.

Biochemical analyses

Venous blood samples were drawn after an overnight fast and stored at -80°C for biochemical analyses. Serum creatinine, urea, aspartate aminotransferase (AST), alanine aminotransferase (ALT), calcium, albumin, uric acid, total cholesterol (TC), high density lipoprotein cholesterol (HDL-C), triglyceride (TG) and phosphorus (P) were determined using a Synchron LX20 system (Beckman Coulter, USA) with original Beckman reagents. HDL-C levels were determined by a direct enzymatic method, without precipitation. LDL-C levels were calculated using the Friedewald formula [22]. Serum hs-CRP levels were measured with a high sensitive immunoturbidimetric assay (Diasis Diagnostic System) using an automated clinical chemistry analyzer. Normal range reference interval of hs-CRP for adults was accepted as <10 mg/L.

STWEAK measurements

Serum TWEAK levels were measured, in both patients and healthy controls, by a commercially available, kit based on enzyme-linked immunosorbent assay (eBioscience, Human TWEAK Instant Elisa, Cat no: BMS2006INST). The results were expressed as pg/mL. The calculated overall intra-assay coefficient of variation (CV) was 7.9%.

Definition of NLR

Complete blood counts with automated differential counts, which included total white blood cells, neutrophils and lymphocytes, obtained at the time of admission.

NLR was calculated as the ratio between neutrophils and lymphocytes, obtained from the same automated blood sample.

Glomerular filtration rate (GFR) assesment

GFR was calculated according to the simplified version of the Modification of Diet in Renal Disease (MDRD) Study prediction equation formula, $GFR = 186 * Creatinine^{-1.154} * Age^{-0.203} * 1.212$ (if African-American) * 0.742 (if female), as defined by Levey [23].

Carotid intima-media thickness measurements

The carotid intima media thickness recordings were performed by a single investigator who was blinded for patient characteristics. The carotid arteries were evaluated with the Vivid 7 echocardiography device (General Electrics, Horten, Norway) by using a 10-MHz linear probe. The acquired images were recorded for playback analysis and were later measured off-line. The common carotid artery, the carotid bulb, and internal and external carotid arteries were visualized on both sides. The IMT of the carotid arteries were measured in the distal common carotid artery at a level 15 to 20 mm proximal to the carotid bulb. The two bright echogenic lines in the arterial wall were identified as the intima and the media. Three measurements were made for each side of the body; separate means were calculated and recorded as the right and left IMT. The intraobserver coefficient of variation for carotid IMT was 2.0%.

Abnormal CIMT was defined as $CIMT \geq 0.82$ mm to characterize atherosclerosis [24].

Statistical analyses

Statistical analyses were carried out using the Statistical Package for Social Sciences for Windows version 15.0 (SPSS, Chicago, IL, USA). Data are expressed as the mean \pm SD, with a significance level of $P < 0.05$. The normal distribution of all variables was tested using the Kolmogorov-Smirnov Test. Dichotomous variables were compared using the chi-square test. Statistical differences between parametric data of two groups were analyzed using the Student's *t*-test. The Mann-Whitney U test was used to determine differences between non-parametric data. The non-parametric Spearman coefficient of correlation was used to assess correlations between variables without normal distribution. The Rtx patients were divided into tertiles of eGFR. Differences among tertiles were analyzed by one-way ANOVA.

Multivariate linear regression analyses were undertaken to determine independent associations among CIMT and other variables. Age, serum phosphorus, LDL-cholesterol, hs-CRP, NLR, and sTWEAK were entered into the regression model as independent variables, and CIMT was entered as a dependent variable. The backward elimination

method was preferred in the stepwise regression analysis and $p > 0.1$ used as a criterion for elimination in this model. At the end of the fifth step, only sTWEAK was remained statistically significant in the model. $p < 0.05$ was considered significant for all tests.

Results

Patients' baseline characteristics

The baseline characteristics of the 70 Rtx patients and 25 healthy subjects are depicted in Table 1. The etiology of renal disease in Rtx patients was chronic glomerulonephritis (n=6, 8.6%), diabetic nephropathy (n=6, 8.6%), hypertensive nephropathy (n=13, 18.6%), polycystic kidney disease (n=2, 3.3%), tubulointerstitial nephritis (n=4, 5.7%), amyloidosis (1, 1.4%) and unknown etiology (n=38, 53.8%). Of 70 Rtx patients, 42 (60%) had post-transplant hypertension, 47 (67.1%) had post-transplant dyslipidemia, and 20 (28.6%) had new-onset diabetes after transplantation (NODAT). There were no differences in age, gender, biochemical parameters including serum LDL and HDL cholesterol, AST, ALT and phosphorus between Rtx patients and healthy subjects. Spot urine protein to creatinine ratio (PCR) values of healthy controls and Rtx patients were 0.01 ± 0.1 and 0.3 ± 0.2 , respectively ($p < 0.0001$). Serum glucose, urea, creatinine, uric acid, triglyceride and calcium were significantly higher in Rtx patients compared to controls (Table 1). Estimated GFR measurements were also found to be significantly lower in Rtx patients than healthy subjects (61.7 ± 22.2 vs 119.2 ± 32.9 , respectively, $p < 0.0001$).

STWEAK, NLR, hs-CRP and carotid intima media thickness measurements

Results of inflammation and atherosclerosis markers including sTWEAK, hs-CRP, NLR and CIMT are shown in Table 2 with univariate correlates of CIMT and sTWEAK in Rtx patients. sTWEAK levels were significantly decreased ($p = 0.01$) and hs-CRP levels were significantly increased ($p < 0.0001$) in Rtx patients compared to healthy subjects. Neutrophil count and NLR were also significantly higher in Rtx patients compared to healthy controls ($p < 0.0001$, $p = 0.001$, respectively). The mean CIMT of Rtx patients and healthy subjects were 0.72 ± 0.28 mm and 0.41 ± 0.07 mm, respectively ($p < 0.0001$).

In order to analyze the values of sTWEAK, NLR, hs-CRP, and CIMT according to renal function of Rtx patients, we defined a high eGFR group (group 1) with levels above the 66th percentile (upper tertile) and a low eGFR group (group 3) with levels below the 33th percentile (lower tertile) (Table 3). NLR and CIMT were highest and sTWEAK was lowest in eGFR group 3 (Table 3). Furthermore, sTWEAK are significantly decreasing with decreasing levels / tertile of eGFR ($p = 0.04$ between all groups). There was also significant

Table 1 The baseline characteristics, clinic and laboratory features of the healthy subjects and renal transplant patients

Parameters	Healthy subjects (n=25) Median (IQR)	Transplant patients (n=70) Median (IQR)	P value
Age (years)	34 (29–43)	38 (31–47)	0.17
Female	13	29	0.48
BMI (kg/m ²)	27 (24–29)	24.8 (21.4–28.4)	0.02
SBP (mmHg)	115 (110–120)	122 (110–137)	<0.0001
DBP (mmHg)	75 (60–80)	80 (70–90)	0.001
Glucose (mg/dL)	92 (86–105)	92 (86–104)	0.04
Urea(mg/dL)	25 (19–34)	32 (24–39)	0.002
Creatinine (mg/dL)	0.7 (0.6–0.8)	1.1 (0.8–1.4)	<0.0001
Albumin (g/dL)	4.4 (4.3–4.5)	4.3 (4.1–4.5)	0.02
Uric acid (mg/dL)	3.7 (3.5–4.6)	5.5 (3.9–7.0)	<0.0001
AST(IU/L)	21 (17–22)	21 (17–26)	0.25
ALT(IU/L)	18 (15–20)	19 (15–25)	0.18
LDL Cholesterol (mg/dL)	125 (105–137)	118 (92–134)	0.38
HDL Cholesterol (mg/dL)	46 (38–50)	46 (38–53)	0.69
Triglyceride (mg/dL)	64 (88–147)	144 (101–181)	0.001
Calcium (mg/dL)	8.5 (8.5–9.4)	9.5 (9.1–9.9)	0.0001
Phosphorus (mg/dL)	3.0 (2.7–3.3)	3.0 (2.7–3.6)	0.68
eGFR (ml/min)	107 (91–144)	71 (53–90)	<0.0001
Spot urine protein/creatinine ratio (g/mg)	0.08 (0.01–0.2)	0.2 (0.05–0.34)	<0.0001
Hemoglobin (mg/dL)	13.5 (13.3–14.3)	14 (12.6–14.7)	0.70
Neutrophil (mm ³)	3100 (2520–3515)	4260 (3287–5937)	<0.0001
Lymphocyte (mm ³)	2190 (1690–2583)	2100 (1582–2710)	0.29

BMI body mass index, SBP systolic blood pressure, DBP diastolic blood pressure, ALT alanine aminotransferase, AST aspartate aminotransferase, LDL low-density lipoprotein, HDL high-density lipoprotein, eGFR estimated glomerular filtration rate.

difference in terms of sTWEAK between healthy subjects and eGFR group 2 and group 3 patients ($p=0.02$ and $p=0.03$, respectively).

Predictors of CIMT in Rtx patients

To investigate the relationship between atherosclerosis and inflammation, we performed a univariate correlation analysis. CIMT was positively correlated with sTWEAK

and NLR in Rtx patients ($r=0.81$, $p<0.0001$ and $r=0.33$, $p=0.006$, respectively). Serum sTWEAK was also positively correlated with NLR ($r=0.37$, $p=0.002$).

We also checked the relation between serum uric acid and inflammatory markers. There were no relationship between serum uric acid and inflammatory markers including hs-CRP and NLR ($r=0.01$, $p=0.89$ and $r=0.05$, $p=0.66$, respectively).

Table 2 sTWEAK, hs-CRP, NLR and CIMT values of the healthy subjects and renal transplant patients

Parameters	Healthy subjects Median Median (IQR) (n=25)	Renal transplant patients (n=70) Median (IQR)	P	ρ (P) ^a	ρ (P) ^b
hs-CRP (mg/L)*	3,3 (2.25–5.8)	8.9 (4.97–14)	<0.0001	0,15 ($p=0.25$)	0.09 ($p=0.46$)
sTWEAK (pg/mL)*	457 (320.5–537.8)	388.9 (324.5–426.9)	0.01	-	0.81 ($p<0.0001$)
NLR	1.16 (0.97–1.5)	2.18 (1.4–2.8)	0.001	0.37 ($p=0.002$)	0.33 ($p=0.006$)
CIMT (mm)	0.4 (0.4–0.5)	0.5 (0.4–0.9)	<0.0001	0.81 ($p<0.0001$)	-

sTWEAK soluble TNF-like weak inducer of apoptosis, NLR neutrophil-to-lymphocyte ratio, CIMT carotid intima media thickness, hs-CRP high sensitive C-reactive protein, IQR interquartile range, * represents the variables in median (IQR).

^a Spearman rank univariate analysis for sTWEAK in renal transplant patients.

^b Spearman rank univariate analysis for CIMT in renal transplant patients.

Table 3 sTWEAK, NLR, CIMT results according to eGFR groups of renal transplant patients and healthy subjects

Parameters	Healthy subjects (n=25) Median (IQR)	eGFR group 1 (n=23) Median (IQR)	eGFR group 2 (n=24) Median (IQR)	eGFR group 3 (n=23) Median (IQR)	p*
MDRD (mL/min)	107 (52.7)	83 (13.56)	63 (8.7)	42 (22)	<0.0001
hs-CRP (mg/L)	3.3 (2.25-5.8)	8.7 (5.6-12.6)	9.45 (4.92-14)	9.0 (3.3-18)	<0.0001
NLR	1.16 (0.54)	2.25 (1)	2.16 (1.5)	2.58 (2.5)	0.005
CIMT (mm)	0.4 (0.1)	0.7 (0.4)	0.6 (0.4)	0.7 (0.5)	<0.001
sTWEAK (pg/mL)	457 (320.5-537.8)	399.5 (329-439)	382.9 (308.6-421.5)	361.2 (323.4-426.5)	0.04

sTWEAK soluble TNF-like weak inducer of apoptosis, NLR neutrophil-to-lymphocyte ratio, CIMT carotid intima-media thickness, MDRD modification of diet in renal disease, hs-CRP high sensitive C-reactive protein, IQR interquartile range, eGFR estimated glomerular filtration rate, *p value between all groups.

In the multivariate analysis, only sTWEAK was found to be an independent variable of increased CIMT. Regression results are shown in Table 4.

Discussion

The major findings of the present study were as follows; i) sTWEAK levels were significantly decreased and CIMT, NLR and hs-CRP were significantly increased in Rtx patients compared to healthy subjects, ii) sTWEAK was positively correlated with CIMT and NLR but not with hs-CRP in Rtx patients, iii) sTWEAK was found to be an independent predictor of increased CIMT in Rtx patients.

In recent years, sTWEAK was discovered as a multi-functional cytokine that was related to the TNF super family. Description of TWEAK in kidney was reported for the first time by Justo et al. [25] in a mouse model of folate-induced acute kidney injury. In the following years,

Yilmaz et al. [26] demonstrated that a decline in eGFR was accompanied by a gradual reduction in sTWEAK in CKD patients. The same group also showed that endothelial dysfunction and decreased sTWEAK were independently associated with cardiovascular outcomes in predialytic CKD patients [21]. In addition, treatment of type 1 hypertensive diabetic CKD with amlodipin and/or valsartan improves endothelial dysfunction and normalizes sTWEAK [27]. During the preparation of the present manuscript, Gungor et al. [28] described a possible role for sTWEAK in the pathogenesis of atherogenesis in the Rtx population.

In our cohort, Rtx patients had lower sTWEAK levels compared to healthy subjects. However, median values of sTWEAK levels of our Rtx patient were higher compared to CKD patients in the cohorts of Yilmaz et al. [21] and Carrero et al. [29] (388.97 pg/mL vs. 245.5 pg/mL and 208 pg/mL, respectively). We also investigated the relation between sTWEAK and kidney function in Rtx patients. After the classification of Rtx patients according to their eGFR, we found that sTWEAK levels were significantly decreased when eGFR values were decreased. These findings are in accord with those obtained by Yilmaz et al. [26]. Therefore, we hypothesized that the decrease of sTWEAK among eGFR groups in Rtx patients might be associated with ongoing inflammation in this population.

Inflammation seen after Rtx has a complex etiology, including a role for innate and acquired immunity. Macrophages and neutrophils play an important role in the pathogenesis of inflammation and acute and chronic allograft rejection [30]. However, ongoing inflammation in Rtx patients is much less intense compared to CKD and ESRD patients. Neutrophil-to-lymphocyte ratio was introduced as a potential marker to determine inflammation in cardiac and non-cardiac disorders [10-12]. In a previous study, we demonstrated that NLR was positively correlated with TNF- α in ESRD patients [14]. In the present study, we found that both hs-CRP and NLR were significantly increased in Rtx patients suggesting that these patients had infra-clinic, smoldering inflammation. We also showed that NLR was positively correlated with sTWEAK in this population. This finding might represent the link between inflammation and

Table 4 The stepwise regression analysis to determine the independent predictors of carotid-intima media thickness in renal transplant patients

Parameters	Standardized Beta [‡]	P value	95% CI
Step 1			
Age (years)	0.17	0.18	-0.003-0.016
Phosphorus (mg/dL)	-0.03	0.78	-0.179-0.135
LDL (mg/dL)	-0.11	0.37	-0.006-0.002
hs-CRP (mg/L)	0.13	0.30	-0.007-0.023
NLR	0.03	0.84	-0.089-0.109
TWEAK (pg/mL)	0.42	0.001	0.001-0.004
adjusted r ² =0.18			
*p=0.008			
Step 5			
TWEAK (pg/mL)	0.42	<0.001	0.001-0.004
adjusted r ² =0.20			
*p<0.0001			

*It was define to model statistic for each step. CI Confidence interval, LDL low density lipoprotein, hs-CRP high sensitive C-reactive protein, NLR neutrophil-to-lymphocyte ratio, TWEAK tumor necrosis factor-like weak inducer of apoptosis.

[‡] Standardized Beta means standardized regression coefficients of the variables.

sTWEAK in Rtx patients. For better understanding, we illustrated this potential pathophysiological relationship in Figure 1.

Recently, Hornum et al. investigated the effect of Rtx on arterial function and they showed a significant improvement in endothelial function, mean arterial pressure and plasma vWF levels but not in plasma CRP and albumin levels [31]. Seyahi et al. also demonstrated that renal transplantation did not reverse or halt the progression of coronary artery calcification [32]. Our results are in accord with these studies - Rtx patients had higher CIMT compared to healthy subjects. Moreover, sTWEAK and NLR were found to be significantly and positively correlated with CIMT. Since sTWEAK is also related to eGFR, we hypothesize that decreased allograft function is associated with inflammation and atherosclerosis in a vicious circle. Therefore, sTWEAK is most probably a pathogenic cytokine rather than only a biomarker in Rtx patients.

The role of TWEAK in the renal tubulointerstitial inflammation was highlighted by Ortiz et al. [33] as an activator of nuclear factor kappa β (NF κ B), which has effects on transcription of inflammatory cytokines including MCP-1, RANTES, and IL-6 [33]. Tacrolimus, widely used as an immunosuppressive agent in Rtx patients, has been shown to down-regulate NF κ B pathway and induce apoptosis of both activated T cells and macrophages, in IL-10 knockout mice [34]. Recently, Du et al. [35] showed the importance of suppression of NF κ B by cyclosporin A and tacrolimus in renal tubular cells. In our study, out of 70 Rtx patients, 60 were receiving


tacrolimus as a main immunosuppressive therapy. Despite the absence of any correlation between tacrolimus usage and sTWEAK it is plausible to suggest that decreased sTWEAK levels might be a compensatory response to the inhibitory effects of tacrolimus on the NF κ B pathway (Figure 1).

Our study had two main limitations. First, this was a cross-sectional analysis of Rtx patients focusing on CIMT and sTWEAK. Second, the sample size was relatively small. This was not a prospective controlled study and therefore, we cannot draw cause-and-effect relationships from our findings.

None of our renal transplant patient died after they enrolled in this study. Hence, we could not analyze the relation between inflammation markers and mortality. However, a prospective study looking at the significance of these markers in mortality should be done.

Conclusions

sTWEAK might have a role in the pathogenesis of ongoing atherosclerosis in Rtx patients. Rtx may not completely reverse non-traditional risk factors including inflammation and atherosclerosis. When considered together, ongoing inflammation and interactions between drugs and macrophages might explain why CKD and ESRD patients had lower sTWEAK levels than Rtx patients. Renal transplant patients should be assessed to their inflammatory and uremic status individually. There are many missing pieces in the puzzle and nothing is yet known about the clinical implications of sTWEAK in Rtx



patients. Further randomized controlled trials and experimental studies investigating the exact roles of sTWEAK in the pathogenesis of atherosclerosis in renal transplantation patients are needed.

Competing interests

The authors declare that they have no conflict of interest.

Authors' contributions

KT: Participated in research design and performance of the research, data analysis, participated in the writing of the paper. HZT: Participated in research design. FME: Participated in the performance of the research. AT: Participated in laboratory analysis. ZB: Participated in the performance of the research. HO: Participated in radiological measurements. AG: Participated in data analysis. EEG: Participated in the data analysis. MK: Participated in the research design. YS: Participated in data analysis. OO: Participated in data analysis of radiological measurements. ST: Participated in the research design. AC: Participated in the editing of the manuscript. All authors read and approved the final manuscript.

Acknowledgements

Dr. A. Gaipov received grant support from the ERA-EDTA fellowship program.

Author details

¹Department of Nephrology, Erzincan University Mengucek Gazi Training and Research Hospital, Erzincan, Turkey. ²Department of Nephrology, Necmettin Erbakan University Meram School of Medicine, Konya, Turkey. ³Department of Radiology, Necmettin Erbakan University Meram School of Medicine, Konya, Turkey. ⁴Department of Biochemistry, Necmettin Erbakan University Meram School of Medicine, Konya, Turkey. ⁵Department of Cardiology, Necmettin Erbakan University Meram School of Medicine, Konya, Turkey. ⁶Nephrology Clinic, Dialysis and Renal Transplant Center, 'C.I. PARHON' University Hospital, 'Gr. T. Popa' University of Medicine and Pharmacy, Iasi, Romania.

Received: 8 December 2012 Accepted: 8 July 2013

Published: 12 July 2013

References

- Collins AJ: Cardiovascular mortality in end-stage renal disease. *Am J Med Sci* 2003, **325**(4):163-7.
- Morris ST, McMurray JJ, Rodger RS, Farmer R, Jardine AG: Endothelial dysfunction in renal transplant recipients maintained on cyclosporine. *Kidney Int* 2000, **57**(3):1100-6.
- Vanholder R, Massy Z, Argiles A, Spasovski G, Verbeke F, Lameire N: Chronic kidney disease as cause of cardiovascular morbidity and mortality. *Nephrol Dial Transplant* 2005, **20**(6):1048-56.
- Parfrey PS, Foley RN, Harnett JD, Kent GM, Murray DC, Barre PE: Outcome and risk factors for left ventricular disorders in chronic uraemia. *Nephrol Dial Transplant* 1996, **11**(7):1277-85.
- Cheung AK, Sarnak MJ, Yan G, Dwyer JT, Heyka RJ, Rocco MV, et al: Atherosclerotic cardiovascular disease risks in chronic hemodialysis patients. *Kidney Int* 2000, **58**(1):353-62.
- Muntner P, He J, Astor BC, Folsom AR, Coresh J: Traditional and nontraditional risk factors predict coronary heart disease in chronic kidney disease: results from the atherosclerosis risk in communities study. *J Am Soc Nephrol* 2005, **16**(2):529-38.
- Himmelfarb J, Stenvinkel P, Ikizler TA, Hakim RM: The elephant in uremia: oxidant stress as a unifying concept of cardiovascular disease in uremia. *Kidney Int* 2002, **62**(5):1524-38.
- Stenvinkel P, Carrero JJ, Axelsson J, Lindholm B, Heimbürger O, Massy Z: Emerging biomarkers for evaluating cardiovascular risk in the chronic kidney disease patient: how do new pieces fit into the uremic puzzle? *Clin J Am Soc Nephrol* 2008, **3**(2):505-21.
- Turkmen K, Gorgulu N, Uysal M, Ozkok A, Sakaci T, Unsal A, et al: Fetuin-A, inflammation, and coronary artery calcification in hemodialysis patients. *Indian J Nephrol* 2011, **21**(2):90-4.
- Tamhane UU, Aneja S, Montgomery D, Rogers EK, Eagle KA, Gurm HS: Association between admission neutrophil to lymphocyte ratio and outcomes in patients with acute coronary syndrome. *Am J Cardiol* 2008, **102**(6):653-7.
- Nunez J, Nunez E, Bodi V, Sanchis J, Minana G, Mainar L, et al: Usefulness of the neutrophil to lymphocyte ratio in predicting long-term mortality in ST segment elevation myocardial infarction. *Am J Cardiol* 2008, **101**(6):747-52.
- Walsh SR, Cook EJ, Goulder F, Justin TA, Keeling NJ: Neutrophil-lymphocyte ratio as a prognostic factor in colorectal cancer. *J Surg Oncol* 2005, **91**(3):181-4.
- Duffy BK, Gurm HS, Rajagopal V, Gupta R, Ellis SG, Bhatt DL: Usefulness of an elevated neutrophil to lymphocyte ratio in predicting long-term mortality after percutaneous coronary intervention. *Am J Cardiol* 2006, **97**(7):993-6.
- Turkmen K, Guney I, Yerlikaya FH, Tonbul HZ: The Relationship Between Neutrophil-to-Lymphocyte Ratio and Inflammation in End-Stage Renal Disease Patients. *Ren Fail* 2012, **34**(2):155-9.
- Burkly LC, Michaelson JS, Zheng TS: TWEAK/Fn14 pathway: an immunological switch for shaping tissue responses. *Immunol Rev* 2011, **244**(1):99-114.
- Wiley SR, Winkles JA: TWEAK, a member of the TNF superfamily, is a multifunctional cytokine that binds the TweakR/Fn14 receptor. *Cytokine Growth Factor Rev* 2003, **14**(3-4):241-9.
- Polek TC, Talpaz M, Darnay BG, Spivak-Kroizman T: TWEAK mediates signal transduction and differentiation of RAW264.7 cells in the absence of Fn14/TweakR. Evidence for a second TWEAK receptor. *J Biol Chem* 2003, **278**(34):32317-23.
- Ortiz A, Sanz AB, Munoz Garcia B, Moreno JA, Sanchez Nino MD, Martin-Ventura JL, et al: Considering TWEAK as a target for therapy in renal and vascular injury. *Cytokine Growth Factor Rev* 2009, **20**(3):251-8.
- Urbonaviciene G, Martin-Ventura JL, Lindholt JS, Urbonavicius S, Moreno JA, Egido J, et al: Impact of soluble TWEAK and CD163/TWEAK ratio on long-term cardiovascular mortality in patients with peripheral arterial disease. *Atherosclerosis* 2011, **219**(2):892-9.
- Blanco-Colio LM, Martin-Ventura JL, Munoz-Garcia B, Orbe J, Paramo JA, Michel JB, et al: Identification of soluble tumor necrosis factor-like weak inducer of apoptosis (sTWEAK) as a possible biomarker of subclinical atherosclerosis. *Arterioscler Thromb Vasc Biol* 2007, **27**(4):916-22.
- Yilmaz MI, Sonmez A, Ortiz A, Saglam M, Kilib S, Eyleten T, et al: Soluble TWEAK and PTX3 in nondialysis CKD patients: impact on endothelial dysfunction and cardiovascular outcomes. *Clin J Am Soc Nephrol* 2011, **6**(4):785-92.
- Friedewald WT, Levy RI, Fredrickson DS: Estimation of the concentration of low-density lipoprotein cholesterol in plasma, without use of the preparative ultracentrifuge. *Clin Chem* 1972, **18**(6):499-502.
- Levey AS, Berg RL, Gassman JJ, Hall PM, Walker WG: Creatinine filtration, secretion and excretion during progressive renal disease. Modification of Diet in Renal Disease (MDRD) Study Group. *Kidney Int Suppl* 1989, **27**:S73-80.
- Benedetto FA, Mallamaci F, Tripepi G, Zoccali C: Prognostic value of ultrasonographic measurement of carotid intima media thickness in dialysis patients. *J Am Soc Nephrol* 2001, **12**(11):2458-64.
- Justo P, Sanz AB, Sanchez-Nino MD, Winkles JA, Lorz C, Egido J, et al: Cytokine cooperation in renal tubular cell injury: the role of TWEAK. *Kidney Int* 2006, **70**(10):1750-8.
- Yilmaz MI, Carrero JJ, Ortiz A, Martin-Ventura JL, Sonmez A, Saglam M, et al: Soluble TWEAK plasma levels as a novel biomarker of endothelial function in patients with chronic kidney disease. *Clin J Am Soc Nephrol* 2009, **4**(11):1716-23.
- Yilmaz MI, Qureshi AR, Carrero JJ, Saglam M, Suliman ME, Caglar K, et al: Predictors of carotid artery intima-media thickness in chronic kidney disease and kidney transplant patients without overt cardiovascular disease. *Am J Nephrol* 2010, **31**(3):214-21.
- Gungor O, Kismali E, Sisman AR, Kircelli F, Carrero JJ, Tatar E, et al: The relationships between serum sTWEAK, FGF-23 levels, and carotid atherosclerosis in renal transplant patients. *Ren Fail* 2013, **35**(1):77-81.
- Carrero JJ, Ortiz A, Qureshi AR, Martin-Ventura JL, Barany P, Heimbürger O, et al: Additive effects of soluble TWEAK and inflammation on mortality in hemodialysis patients. *Clin J Am Soc Nephrol* 2009, **4**(1):110-8.
- Wood KJ, Goto R: Mechanisms of rejection: current perspectives. *Transplantation* 2012, **93**(1):1-10.
- Hornum M, Clausen P, Idorn T, Hansen JM, Mathiesen ER, Feldt-Rasmussen B: Kidney transplantation improves arterial function measured by pulse

- wave analysis and endothelium-independent dilatation in uraemic patients despite deterioration of glucose metabolism. *Nephrol Dial Transplant* 2011, **26**(7):2370–7.
32. Seyahi N, Cebi D, Altiparmak MR, Akman C, Ataman R, Pekmezci S, *et al.* Nephrol Dial Transplant: Progression of coronary artery calcification in renal transplant recipients; 2011.
 33. Sanz AB, Justo P, Sanchez-Nino MD, Blanco-Colio LM, Winkles JA, Kretzler M, *et al.* The cytokine TWEAK modulates renal tubulointerstitial inflammation. *J Am Soc Nephrol* 2008, **19**(4):695–703.
 34. Yoshino T, Nakase H, Honzawa Y, Matsumura K, Yamamoto S, Takeda Y, *et al.* Immunosuppressive effects of tacrolimus on macrophages ameliorate experimental colitis. *Inflamm Bowel Dis* 2010, **16**(12):2022–33.
 35. Du S, Hiramatsu N, Hayakawa K, Kasai A, Okamura M, Huang T, *et al.* Suppression of NF-kappaB by cyclosporin a and tacrolimus (FK506) via induction of the C/EBP family: implication for unfolded protein response. *J Immunol* 2009, **182**(11):7201–11.

doi:10.1186/1471-2369-14-144

Cite this article as: Turkmen *et al.*: Soluble TWEAK independently predicts atherosclerosis in renal transplant patients. *BMC Nephrology* 2013 **14**:144.

**Submit your next manuscript to BioMed Central
and take full advantage of:**

- Convenient online submission
- Thorough peer review
- No space constraints or color figure charges
- Immediate publication on acceptance
- Inclusion in PubMed, CAS, Scopus and Google Scholar
- Research which is freely available for redistribution

Submit your manuscript at
www.biomedcentral.com/submit

 **BioMed** Central